


F3 The reproducibility floor is a property of the ARM (and of the statistic, and of the view - T9) same configuration, same global step, only the training seed differs



arm	seed 42	seed 43	seed 44	seed 45	seed 46	spread
programme_only	110.765	110.879	111.078	94.952	112.078	1.18x
programme_free	7.545	45.646	12.194	18.881	22.518	6.05x

n = 3 (seeds 42/43/44 only), as printed in paper/P2_FIGURES.md F3: programme_only 1.003x, programme_free 6.05x.

SUPERSEDED. Seeds 45 and 46 landed afterwards in `~e0_run/d1_seeds4546/` and are included above; the stable arm's floor is 1.18x, not 1.003x.

F3. Statistic R3 - the in-run tripwire statistic, NOT R1 - and the block is a TRAINING batch at global step 200, not a held-out set. The states behind these numbers were never saved, so they are [NOT RECOMPUTABLE] under R1, and no number in this figure may share an axis with any R1 value in this paper. The key is ``train_rank_tripwire_observed`` with the ``train_`` prefix, because querying it without the prefix returns [] and an empty list reads as a confident negative. The two rows are not connected and share only the axis: they are two arms, not a series. The reading is that rank's reproducibility is worst exactly where the arm is interesting - which is the configuration a practitioner is looking at when they reach for rank. F3 and F4(a) are not in conflict: F4(a) holds the model fixed and varies the patient sample; F3 holds the step fixed and varies the seed.