

# Simple Analysis of Sparse, Sign-Consistent JL

Meena Jagadeesan

Harvard University

RANDOM 2019

## Linear dimensionality reduction: $\ell_2$ -to- $\ell_2$

**Informal goal:** Project vectors in  $\mathbb{R}^n$  to  $\mathbb{R}^m$  (for  $m \ll n$ ) with a linear map while “preserving geometry” (i.e.  $\|f(x) - f(y)\|_2 \approx \|x - y\|_2$ ).

# Linear dimensionality reduction: $\ell_2$ -to- $\ell_2$

**Informal goal:** Project vectors in  $\mathbb{R}^n$  to  $\mathbb{R}^m$  (for  $m \ll n$ ) with a linear map while “preserving geometry” (i.e.  $\|f(x) - f(y)\|_2 \approx \|x - y\|_2$ ).

Many applications:

- ▶ Feature hashing (Weinberger et al. '09, Dahlgaard et al. '17, Freksen et al. '18, etc.)
- ▶ Numerical linear algebra (Clarkson and Woodruff '12, Nelson and Nguyen '14, etc.)
- ▶ Approximate nearest neighbors (Ailon and Chazelle '09, etc.)
- ▶ k-means/k-medians (Makarychev, Makarychev, Razenshteyn '18)

# Linear dimensionality reduction: $\ell_2$ -to- $\ell_2$

**Informal goal:** Project vectors in  $\mathbb{R}^n$  to  $\mathbb{R}^m$  (for  $m \ll n$ ) with a linear map while “preserving geometry” (i.e.  $\|f(x) - f(y)\|_2 \approx \|x - y\|_2$ ).

Many applications:

- ▶ Feature hashing (Weinberger et al. '09, Dahlgaard et al. '17, Freksen et al. '18, etc.)
- ▶ Numerical linear algebra (Clarkson and Woodruff '12, Nelson and Nguyen '14, etc.)
- ▶ Approximate nearest neighbors (Ailon and Chazelle '09, etc.)
- ▶ k-means/k-medians (Makarychev, Makarychev, Razenshteyn '18)
- ▶ Compression in the brain (Allen-Zhu, Gelashvili, Micali, Shavit '15)

# Mathematical framework

# Mathematical framework

Use a probability distribution  $\mathcal{M}$  over linear maps  $\mathbb{R}^n \rightarrow \mathbb{R}^m$  ( $m \ll n$ ).

**Geometry-preserving property:** for each  $x \in \mathbb{R}^n$

$$\mathbb{P}_{M \in \mathcal{M}}[(1 - \epsilon) \|x\|_2 \leq \|Mx\|_2 \leq (1 + \epsilon) \|x\|_2] > 1 - \delta.$$

# Mathematical framework

Use a probability distribution  $\mathcal{M}$  over linear maps  $\mathbb{R}^n \rightarrow \mathbb{R}^m$  ( $m \ll n$ ).

**Geometry-preserving property:** for each  $x \in \mathbb{R}^n$

$$\mathbb{P}_{M \in \mathcal{M}}[(1 - \epsilon) \|x\|_2 \leq \|Mx\|_2 \leq (1 + \epsilon) \|x\|_2] > 1 - \delta.$$

Fundamental result of linear dimensionality reduction:

Lemma (Distributional Johnson-Lindenstrauss Lemma)

*Can obtain  $m = \Theta(\epsilon^{-2} \log(1/\delta))$  using  $\mathcal{M}$  with i.i.d gaussian entries.*

This dimension is actually optimal for any distribution over linear maps (Kane et al. '11, Jayram and Woodruff '11).


Optimality for  $N$ -point version (Larsen and Nelson '17)

# Application to information compression in the brain



# Application to information compression in the brain

Convergent pathways compress information w/o losing the ability to perform computations.




A model (Ganguli, Sompolinsky '12)

- ▶ Source information:  $x \in \mathbb{R}^n$
- ▶ Target information:  $y \in \mathbb{R}^m$
- ▶ Synaptic connections: a random matrix  $M \in \mathbb{R}^{m \times n}$


# Biological Constraints on $\text{Supp}(\mathcal{M})$

(As per Ganguli and Sompolinsky's model)



# Biological Constraints on $\text{Supp}(\mathcal{M})$

(As per Ganguli and Sompolinsky's model)



**Sparsity:** every column has  $\leq s$  nonzero entries.

- Neurons connected to few post-synaptic neurons

**Sign-consistency:** in each column, nonzero entries are *all positive* or *all negative*

- Neurons are excitatory or inhibitory

# JL for sparse matrices

Sparsity is also more generally useful for reducing projection time.

## Informal Construction (Sparse JL)

*Uniformly choose  $s$  nonzero entries per column; i.i.d signs for nonzero entries*

Can set  $m = \Theta(\epsilon^{-2} \log(1/\delta))$  and  $s = \Theta(\epsilon^{-1} \log(1/\delta))$  (Kane and Nelson, J. ACM '12)

Can set  $m = \min(2\epsilon^{-2}/\delta, \Theta(\epsilon^{-2} \log(1/\delta)B))$  and  $s = \Theta(\epsilon^{-1} \log(1/\delta)/\log B)$  (Cohen, SODA '16)

# Sparse, Sign-Consistent JL (Allen-Zhu et al.)

# Sparse, Sign-Consistent JL (Allen-Zhu et al.)

Uniformly choose  $s$  nonzero entries per column; i.i.d signs for each column:

# Sparse, Sign-Consistent JL (Allen-Zhu et al.)

Uniformly choose  $s$  nonzero entries per column; i.i.d signs for each column:

## Informal Construction (Sparse, Sign-Consistent JL)

$\mathcal{M}$  is defined so the  $(r, i)$ th entry is  $\sigma_i \eta_{r,i} / \sqrt{s}$  where:

- ▶  $\sigma_i$  are i.i.d. Rademachers (random signs)
- ▶  $\eta_{r,i}$  are  $\{0, 1\}$  rvs s.t.  $\sum_{r=1}^m \eta_{r,i} = s$  and w/ mild assumptions

# Sparse, Sign-Consistent JL (Allen-Zhu et al.)

Uniformly choose  $s$  nonzero entries per column; i.i.d signs for each column:

## Informal Construction (Sparse, Sign-Consistent JL)

$\mathcal{M}$  is defined so the  $(r, i)$ th entry is  $\sigma_i \eta_{r,i} / \sqrt{s}$  where:

- ▶  $\sigma_i$  are i.i.d. Rademachers (random signs)
- ▶  $\eta_{r,i}$  are  $\{0, 1\}$  rvs s.t.  $\sum_{r=1}^m \eta_{r,i} = s$  and w/ mild assumptions

Can set  $m = \Theta(\epsilon^{-2} \log^2(1/\delta))$ ,  $s = \Theta(\epsilon^{-1} \log(1/\delta))$  (Allen-Zhu, Gelashvili, Micali, and Shavit, PNAS '15)



# This work

Simplify and generalize the analysis of sparse, sign-consistent JL.

## Theorem (Informal)

*For any  $\epsilon \leq B \leq 1/\delta$ , can set  $m = \Theta(\epsilon^{-2} \log^2(1/\delta) B / \log^2(B))$  and  $s = \Theta(\epsilon^{-1} \log(1/\delta) / \log B)$  for sparse, sign-consistent JL.*

# This work

Simplify and generalize the analysis of sparse, sign-consistent JL.

## Theorem (Informal)

*For any  $\epsilon \leq B \leq 1/\delta$ , can set  $m = \Theta(\epsilon^{-2} \log^2(1/\delta) B / \log^2(B))$  and  $s = \Theta(\epsilon^{-1} \log(1/\delta) / \log B)$  for sparse, sign-consistent JL.*

(Bears resemblance to dim/sparsity tradeoffs for sparse JL by Cohen '16.)

# This work

Simplify and generalize the analysis of sparse, sign-consistent JL.

## Theorem (Informal)

*For any  $e \leq B \leq 1/\delta$ , can set  $m = \Theta(\epsilon^{-2} \log^2(1/\delta) B / \log^2(B))$  and  $s = \Theta(\epsilon^{-1} \log(1/\delta) / \log B)$  for sparse, sign-consistent JL.*

(Bears resemblance to dim/sparsity tradeoffs for sparse JL by Cohen '16.)

**Proof method:** Different techniques for analyzing moments of JL error terms which turn to be “needed” for this setting

# This work

Simplify and generalize the analysis of sparse, sign-consistent JL.

## Theorem (Informal)

*For any  $\epsilon \leq B \leq 1/\delta$ , can set  $m = \Theta(\epsilon^{-2} \log^2(1/\delta) B / \log^2(B))$  and  $s = \Theta(\epsilon^{-1} \log(1/\delta) / \log B)$  for sparse, sign-consistent JL.*

(Bears resemblance to dim/sparsity tradeoffs for sparse JL by Cohen '16.)

**Proof method:** Different techniques for analyzing moments of JL error terms which turn to be “needed” for this setting

- ▶ Intuition critical in my follow-up work on analyzing sparse JL for feature hashing (NeurIPS '19, to appear).

# This work

Simplify and generalize the analysis of sparse, sign-consistent JL.

## Theorem (Informal)

*For any  $e \leq B \leq 1/\delta$ , can set  $m = \Theta(\epsilon^{-2} \log^2(1/\delta) B / \log^2(B))$  and  $s = \Theta(\epsilon^{-1} \log(1/\delta) / \log B)$  for sparse, sign-consistent JL.*

(Bears resemblance to dim/sparsity tradeoffs for sparse JL by Cohen '16.)

**Proof method:** Different techniques for analyzing moments of JL error terms which turn to be “needed” for this setting

- ▶ Intuition critical in my follow-up work on analyzing sparse JL for feature hashing (NeurIPS '19, to appear).

Remainder of the talk will focus on the proof method.

# High-level approach

Need to show  $\mathbb{P}_{M \in \mathcal{M}}[(1 - \epsilon) \|x\|_2 \leq \|Mx\|_2 \leq (1 + \epsilon) \|x\|_2] > 1 - \delta.$

# High-level approach

Need to show  $\mathbb{P}_{M \in \mathcal{M}}[(1 - \epsilon) \|x\|_2 \leq \|Mx\|_2 \leq (1 + \epsilon) \|x\|_2] > 1 - \delta$ .

Analyze high moment of  $Z := \|Mx\|_2^2 - 1$  for all  $x \in B_{\ell_2}$  (like in previous work).

# High-level approach

Need to show  $\mathbb{P}_{M \in \mathcal{M}}[(1 - \epsilon) \|x\|_2 \leq \|Mx\|_2 \leq (1 + \epsilon) \|x\|_2] > 1 - \delta$ .

Analyze high moment of  $Z := \|Mx\|_2^2 - 1$  for all  $x \in B_{\ell_2}$  (like in previous work).

For sparse, sign-consistent JL:

$$Z := \|Mx\|_2^2 - 1 = \frac{1}{s} \sum_{i \neq j} \sum_{r=1}^m \sigma_i \sigma_j \eta_{r,i} \eta_{r,j} x_i x_j.$$



# Analyzing the moments of $Z$

$$Z := \|M_X\|_2^2 - 1 = \frac{1}{s} \sum_{i \neq j} \sum_{r=1}^m \sigma_i \sigma_j \eta_{r,i} \eta_{r,j} x_i x_j.$$

# Analyzing the moments of $Z$

$$Z := \|M_X\|_2^2 - 1 = \frac{1}{s} \sum_{i \neq j} \sum_{r=1}^m \sigma_i \sigma_j \eta_{r,i} \eta_{r,j} x_i x_j.$$

Limitations of existing approaches for bounding moments of  $Z$ :

# Analyzing the moments of $Z$

$$Z := \|M_X\|_2^2 - 1 = \frac{1}{s} \sum_{i \neq j} \sum_{r=1}^m \sigma_i \sigma_j \eta_{r,i} \eta_{r,j} x_i x_j.$$

Limitations of existing approaches for bounding moments of  $Z$ :

1. Combinatorics (Allen-Zhu et al. '15, Kane and Nelson '12, Freksen et al. '18, etc.)

# Analyzing the moments of $Z$

$$Z := \|M_X\|_2^2 - 1 = \frac{1}{s} \sum_{i \neq j} \sum_{r=1}^m \sigma_i \sigma_j \eta_{r,i} \eta_{r,j} x_i x_j.$$

Limitations of existing approaches for bounding moments of  $Z$ :

1. Combinatorics (Allen-Zhu et al. '15, Kane and Nelson '12, Freksen et al. '18, etc.)
  - Intricate; unclear how to get dim-sparsity tradeoffs w/ Allen-Zhu et al.

# Analyzing the moments of $Z$

$$Z := \|M_X\|_2^2 - 1 = \frac{1}{s} \sum_{i \neq j} \sum_{r=1}^m \sigma_i \sigma_j \eta_{r,i} \eta_{r,j} x_i x_j.$$

Limitations of existing approaches for bounding moments of  $Z$ :

1. Combinatorics (Allen-Zhu et al. '15, Kane and Nelson '12, Freksen et al. '18, etc.)
  - ▶ Intricate; unclear how to get dim-sparsity tradeoffs w/ Allen-Zhu et al.
2. (Sub)-gaussian quadratic form bounds (Cohen et al. '18, for sparse JL)

# Analyzing the moments of $Z$

$$Z := \|M_X\|_2^2 - 1 = \frac{1}{s} \sum_{i \neq j} \sum_{r=1}^m \sigma_i \sigma_j \eta_{r,i} \eta_{r,j} x_i x_j.$$

Limitations of existing approaches for bounding moments of  $Z$ :

1. Combinatorics (Allen-Zhu et al. '15, Kane and Nelson '12, Freksen et al. '18, etc.)
  - ▶ Intricate; unclear how to get dim-sparsity tradeoffs w/ Allen-Zhu et al.
2. (Sub)-gaussian quadratic form bounds (Cohen et al. '18, for sparse JL)
  - ▶ Using the gaussian bound is too weak for this setting!

# Analyzing the moments of $Z$

$$Z := \|Mx\|_2^2 - 1 = \frac{1}{s} \sum_{i \neq j} \sum_{r=1}^m \sigma_i \sigma_j \eta_{r,i} \eta_{r,j} x_i x_j.$$

Limitations of existing approaches for bounding moments of  $Z$ :

1. Combinatorics (Allen-Zhu et al. '15, Kane and Nelson '12, Freksen et al. '18, etc.)
  - ▶ Intricate; unclear how to get dim-sparsity tradeoffs w/ Allen-Zhu et al.
2. (Sub)-gaussian quadratic form bounds (Cohen et al. '18, for sparse JL)
  - ▶ Using the gaussian bound is too weak for this setting!

My key ingredient: more precise quadratic form bounds

# Expressing $Z$ as a Rademacher quadratic form

$$Z = \frac{1}{s} \sum_{i \neq j} \sigma_i \sigma_j x_i x_j \left( \sum_{r=1}^m \eta_{r,i} \eta_{r,j} \right)$$



# Expressing $Z$ as a Rademacher quadratic form

$$Z = \frac{1}{s} \sum_{i \neq j} \sigma_i \sigma_j x_i x_j \left( \sum_{r=1}^m \eta_{r,i} \eta_{r,j} \right) = \sigma^T A_\eta \sigma.$$

# Expressing $Z$ as a Rademacher quadratic form

$$Z = \frac{1}{s} \sum_{i \neq j} \sigma_i \sigma_j x_i x_j \left( \sum_{r=1}^m \eta_{r,i} \eta_{r,j} \right) = \sigma^T A_\eta \sigma.$$

$$\implies \mathbb{E}[Z^p] = \mathbb{E}_\eta \left[ \mathbb{E}_\sigma \left[ (\sigma^T A_\eta \sigma)^p \right] \right].$$

$\mathbb{E}_\sigma [(\sigma^T A_\eta \sigma)^p]$  is a moment of a Rademacher quadratic form.

# Expressing $Z$ as a Rademacher quadratic form

$$Z = \frac{1}{s} \sum_{i \neq j} \sigma_i \sigma_j x_i x_j \left( \sum_{r=1}^m \eta_{r,i} \eta_{r,j} \right) = \sigma^T A_\eta \sigma.$$

$$\implies \mathbb{E}[Z^p] = \mathbb{E}_\eta \left[ \mathbb{E}_\sigma [(\sigma^T A_\eta \sigma)^p] \right].$$

$\mathbb{E}_\sigma [(\sigma^T A_\eta \sigma)^p]$  is a moment of a Rademacher quadratic form.

CJN '18 (for sparse JL) uses  $\mathbb{E}_\sigma [(\sigma^T A_\eta \sigma)^p] \lesssim \mathbb{E}_g [(g^T A_\eta g)^p]$ .

# Expressing $Z$ as a Rademacher quadratic form

$$Z = \frac{1}{s} \sum_{i \neq j} \sigma_i \sigma_j x_i x_j \left( \sum_{r=1}^m \eta_{r,i} \eta_{r,j} \right) = \sigma^T A_\eta \sigma.$$

$$\implies \mathbb{E}[Z^p] = \mathbb{E}_\eta \left[ \mathbb{E}_\sigma [(\sigma^T A_\eta \sigma)^p] \right].$$

$\mathbb{E}_\sigma [(\sigma^T A_\eta \sigma)^p]$  is a moment of a Rademacher quadratic form.

CJN '18 (for sparse JL) uses  $\mathbb{E}_\sigma [(\sigma^T A_\eta \sigma)^p] \lesssim \mathbb{E}_g [(g^T A_\eta g)^p]$ .

But, we show using  $\mathbb{E}_g [(g^T A_\eta g)^p]$  is too weak for this setting!

- ▶  $x = [1/\sqrt{2}, 1/\sqrt{2}, 0, \dots, 0]$  already yields sub-optimal dimension  $m$

# Expressing $Z$ as a Rademacher quadratic form

$$Z = \frac{1}{s} \sum_{i \neq j} \sigma_i \sigma_j x_i x_j \left( \sum_{r=1}^m \eta_{r,i} \eta_{r,j} \right) = \sigma^T A_\eta \sigma.$$

$$\implies \mathbb{E}[Z^p] = \mathbb{E}_\eta \left[ \mathbb{E}_\sigma [(\sigma^T A_\eta \sigma)^p] \right].$$

$\mathbb{E}_\sigma [(\sigma^T A_\eta \sigma)^p]$  is a moment of a Rademacher quadratic form.

CJN '18 (for sparse JL) uses  $\mathbb{E}_\sigma [(\sigma^T A_\eta \sigma)^p] \lesssim \mathbb{E}_g [(g^T A_\eta g)^p]$ .

But, we show using  $\mathbb{E}_g [(g^T A_\eta g)^p]$  is too weak for this setting!

- ▶  $x = [1/\sqrt{2}, 1/\sqrt{2}, 0, \dots, 0]$  already yields sub-optimal dimension  $m$

Thus, we need a Rademacher-specific bound for  $\mathbb{E}_\sigma [(\sigma^T A_\eta \sigma)^p]$ .

# A digression on Rademachers vs. gaussians for linear forms

$\|Y\|_p := (\mathbb{E}[Y^p])^{1/p}$ ;  $\sigma_i$  i.i.d. Rademachers;  $g_i$  i.i.d gaussians;  
 $|a_1| \geq \dots \geq |a_n|$  scalar coefficients

# A digression on Rademachers vs. gaussians for linear forms

$\|Y\|_p := (\mathbb{E}[Y^p])^{1/p}$ ;  $\sigma_i$  i.i.d. Rademachers;  $g_i$  i.i.d gaussians;  
 $|a_1| \geq \dots \geq |a_n|$  scalar coefficients

**Khintchine bound:**  $\|\sum_{i=1}^n a_i \sigma_i\|_p \lesssim \|\sum_{i=1}^n a_i g_i\|_p \sim \sqrt{p} \sqrt{\sum_{i=1}^n a_i^2}$

# A digression on Rademachers vs. gaussians for linear forms

$\|Y\|_p := (\mathbb{E}[Y^p])^{1/p}$ ;  $\sigma_i$  i.i.d. Rademachers;  $g_i$  i.i.d. gaussians;  
 $|a_1| \geq \dots \geq |a_n|$  scalar coefficients

**Khintchine bound:**  $\|\sum_{i=1}^n a_i \sigma_i\|_p \lesssim \|\sum_{i=1}^n a_i g_i\|_p \sim \sqrt{p} \sqrt{\sum_{i=1}^n a_i^2}$

$\lim_{p \rightarrow \infty} \|\sum_{i=1}^n a_i \sigma_i\|_p \leq \sum_{i=1}^n |a_i| < \infty$ , but  $\lim_{p \rightarrow \infty} \sqrt{p} \sqrt{\sum_{i=1}^n a_i^2} = \infty$ .



# A digression on Rademachers vs. gaussians for linear forms

$\|Y\|_p := (\mathbb{E}[Y^p])^{1/p}$ ;  $\sigma_i$  i.i.d. Rademachers;  $g_i$  i.i.d. gaussians;  
 $|a_1| \geq \dots \geq |a_n|$  scalar coefficients

**Khinchine bound:**  $\|\sum_{i=1}^n a_i \sigma_i\|_p \lesssim \|\sum_{i=1}^n a_i g_i\|_p \sim \sqrt{p} \sqrt{\sum_{i=1}^n a_i^2}$

$\lim_{p \rightarrow \infty} \|\sum_{i=1}^n a_i \sigma_i\|_p \leq \sum_{i=1}^n |a_i| < \infty$ , but  $\lim_{p \rightarrow \infty} \sqrt{p} \sqrt{\sum_{i=1}^n a_i^2} = \infty$ .

Khinchine is *not tight* for Rademachers for large  $p$ .

# A digression on Rademachers vs. gaussians for linear forms

$\|Y\|_p := (\mathbb{E}[Y^p])^{1/p}$ ;  $\sigma_i$  i.i.d. Rademachers;  $g_i$  i.i.d. gaussians;  
 $|a_1| \geq \dots \geq |a_n|$  scalar coefficients

**Khinchine bound:**  $\|\sum_{i=1}^n a_i \sigma_i\|_p \lesssim \|\sum_{i=1}^n a_i g_i\|_p \sim \sqrt{p} \sqrt{\sum_{i=1}^n a_i^2}$

$\lim_{p \rightarrow \infty} \|\sum_{i=1}^n a_i \sigma_i\|_p \leq \sum_{i=1}^n |a_i| < \infty$ , but  $\lim_{p \rightarrow \infty} \sqrt{p} \sqrt{\sum_{i=1}^n a_i^2} = \infty$ .

Khinchine is *not tight* for Rademachers for large  $p$ .

Turns out  $\|\sum_{i=1}^n a_i \sigma_i\|_p \sim \left| \sum_{i=1}^p a_i \right| + \sqrt{p} \sqrt{\sum_{i>p} a_i^2}$  (Hitzchenko '93).

# Generalizing to quadratic forms of Rademachers

# Generalizing to quadratic forms of Rademachers

## Lemma

*If  $(A_{i,j})$  is a symmetric  $n \times n$  matrix with zero diagonal and  $p$  even, then*

$$\left\| \sum_{i=1}^n \sum_{j=1}^n A_{i,j} \sigma_i \sigma_j \right\|_p \lesssim \left( \sum_{i=1}^{\min(p,n)} \sum_{j=1}^{\min(p,n)} |A_{i,j}| \right) + \sqrt{p} \sqrt{\sum_{i=1}^n \left\| \sum_{j>p} A_{i,j} \sigma_j \right\|_p^2}.$$

# Generalizing to quadratic forms of Rademachers

## Lemma

*If  $(A_{i,j})$  is a symmetric  $n \times n$  matrix with zero diagonal and  $p$  even, then*

$$\left\| \sum_{i=1}^n \sum_{j=1}^n A_{i,j} \sigma_i \sigma_j \right\|_p \lesssim \left( \sum_{i=1}^{\min(p,n)} \sum_{j=1}^{\min(p,n)} |A_{i,j}| \right) + \sqrt{p} \sqrt{\sum_{i=1}^n \left\| \sum_{j>p} A_{i,j} \sigma_j \right\|_p^2}.$$

- ▶ An interpolation of  $\ell_1$  and  $\ell_2$  bounds.

# Generalizing to quadratic forms of Rademachers

## Lemma

*If  $(A_{i,j})$  is a symmetric  $n \times n$  matrix with zero diagonal and  $p$  even, then*

$$\left\| \sum_{i=1}^n \sum_{j=1}^n A_{i,j} \sigma_i \sigma_j \right\|_p \lesssim \left( \sum_{i=1}^{\min(p,n)} \sum_{j=1}^{\min(p,n)} |A_{i,j}| \right) + \sqrt{p} \sqrt{\sum_{i=1}^n \left\| \sum_{j>p} A_{i,j} \sigma_j \right\|_p^2}.$$

- ▶ An interpolation of  $\ell_1$  and  $\ell_2$  bounds.
- ▶ When  $p \rightarrow \infty$ , bounded by  $\sum_{i=1}^{\min(p,n)} \sum_{j=1}^{\min(p,n)} |A_{i,j}| < \infty$ .

# Generalizing to quadratic forms of Rademachers

## Lemma

If  $(A_{i,j})$  is a symmetric  $n \times n$  matrix with zero diagonal and  $p$  even, then

$$\left\| \sum_{i=1}^n \sum_{j=1}^n A_{i,j} \sigma_i \sigma_j \right\|_p \lesssim \left( \sum_{i=1}^{\min(p,n)} \sum_{j=1}^{\min(p,n)} |A_{i,j}| \right) + \sqrt{p} \sqrt{\sum_{i=1}^n \left\| \sum_{j>p} A_{i,j} \sigma_j \right\|_p^2}.$$

- ▶ An interpolation of  $\ell_1$  and  $\ell_2$  bounds.
- ▶ When  $p \rightarrow \infty$ , bounded by  $\sum_{i=1}^{\min(p,n)} \sum_{j=1}^{\min(p,n)} |A_{i,j}| < \infty$ .

(Tight bound on  $\|\sigma^T A \sigma\|_p$  (Latała '99) messy when  $A$  a *random matrix*.)

# Follow-up work that uses intuition from these methods



# Follow-up work that uses intuition from these methods

## “Understanding Sparse JL for Feature Hashing” (NeurIPS 2019)

I study sparse JL on feature vectors.

- ▶ Model: limit to vectors  $x$  with “small”  $\ell_\infty$ -to- $\ell_2$  norm ratio
- ▶  $s = 1$  understood (Weinberger et al '09, Dahlgaard et al. '17, Freksen et al. '18, etc.)

My main result: Generalization to  $s > 1$ .

- ▶ Tight tradeoff between  $\ell_\infty$ -to- $\ell_2$  ratio,  $s$ ,  $m$ ,  $\epsilon$ , and  $\delta$  for sparse JL
- ▶  $\implies$  Even (small)  $s > 1$  can be much better than  $s = 1$ .

# Follow-up work that uses intuition from these methods

## “Understanding Sparse JL for Feature Hashing” (NeurIPS 2019)

I study sparse JL on feature vectors.

- ▶ Model: limit to vectors  $x$  with “small”  $\ell_\infty$ -to- $\ell_2$  norm ratio
- ▶  $s = 1$  understood (Weinberger et al '09, Dahlgaard et al. '17, Freksen et al. '18, etc.)

My main result: Generalization to  $s > 1$ .

- ▶ Tight tradeoff between  $\ell_\infty$ -to- $\ell_2$  ratio,  $s$ ,  $m$ ,  $\epsilon$ , and  $\delta$  for sparse JL
- ▶  $\implies$  Even (small)  $s > 1$  can be much better than  $s = 1$ .

Similarly unclear how to adapt combinatorics; gaussian bounds too weak.

Tractable Rademacher-specific bounds are the key technical tool.

# Conclusion

- ▶ Simplified and generalized the analysis of sparse, sign-consistent JL (Allen-Zhu, Gelashvili, Micali, Shavit '15).
- ▶ Specifically obtained dimensionality-sparsity tradeoffs  
 $m = \Theta(\epsilon^{-2} \log^2(1/\delta) B / \log^2(B))$  and  $s = \Theta(\epsilon^{-1} \log(1/\delta) / \log B)$ .
- ▶ Introduced a simple moment bound for Rademacher quadratic forms which enables a simpler analysis of sparse, sign-consistent JL, and could be of broader use.