

Filtered products and boundary-preserving compression in complex cobordism

OpenAI

September 25, 2026

Abstract

We prove that, in sufficiently large smash powers, an MU -null restriction to a finite pointed subcomplex becomes stably null on the entire union of products with a fixed positive proportion of restricted factors. This coherent vanishing theorem yields boundary-preserving compression of cube-valued maps on every compact metrizable input space. When the torus slot bundle embeds continuously into a trivial bundle of rank less than twice the source rank, the compression places a positive proportion of slots on their boundaries in arbitrarily large powers and fixes every original boundary slot exactly. An example at equality shows that the strict rank inequality cannot be removed.

1 Introduction

We study when a map into a product of cubes can be moved onto many coordinate boundaries while retaining every boundary value already present. The input spaces may have arbitrarily large, or infinite, covering dimension. The obstruction we use is a relative point class in complex cobordism, tested against a family of vectors supported in only a few cube factors. Our main results convert the vanishing of this class into compression on arbitrarily large Cartesian powers.

The proof has two coherence requirements. A cobordism-null map on one subcomplex must give a single stable nullhomotopy on a union of products, including all their intersections. Subsequently, stable avoidance data on small coordinate boxes must agree on adjacent simplices. We prove both compatibility statements explicitly. A padding construction then removes the dimension restriction needed to turn stable avoidance into an actual map.

1.1 Coherent products

Write $\mathbb{S}$ for the sphere spectrum and MU for the complex cobordism spectrum. A based map is *stably null* if its suspension spectrum is nullhomotopic. The unit $\mathbb{S} \rightarrow MU$ sends a stable cohomotopy class to its complex-cobordism class. Our first result concerns the difference between vanishing on a subcomplex and vanishing on an entire family of products.

Theorem 1.1 (Filtered products). *Let $G \subset E$ be a finite pointed CW pair, let $d \geq 0$ be an integer, and let $v : E \rightarrow S^d$ be a based map. Suppose that the composite*

$$\Sigma^\infty G \longrightarrow \Sigma^\infty E \xrightarrow{\Sigma^\infty v} \Sigma^d \mathbb{S} \longrightarrow \Sigma^d MU$$

is nullhomotopic. For $\ell \geq 1$, let $U_\ell \subset E^{\wedge \ell}$ be the union of the subcomplexes in which at least $\lceil \ell/2 \rceil$ factors are required to belong to G . Then, for every sufficiently large ℓ , the restriction

$$v^{\wedge \ell}|_{U_\ell} : U_\ell \longrightarrow S^{d\ell}$$

is stably nullhomotopic.

The conclusion concerns the union itself. It therefore supplies a nullhomotopy that can be pulled back along any map into that union. The fraction $1/2$ may be replaced by any fixed positive fraction; see [Remark 2.6](#). The proof also gives a more general compatible lift for an arbitrary commutative ring spectrum, before invoking a vanishing theorem for MU . Its target is the total fiber of a cube of unit maps. To produce one null cone over the punctured cube, we use simplex parameters that collapse the entire constrained union at once; the number of restricted factors determines how many simplex degrees collapse.

1.2 Cube slots and the torus criterion

For positive integers t, n , put

$$C(t, n) = ([-1, 1]^t)^n, \quad T(t, n) = ([-2, 2]^t)^n.$$

A point has n ordered *slots*, each a vector in $\mathbb{R}^t$. A boundary slot belongs to the whole boundary $\partial[-1, 1]^t$. Preserving it means preserving its entire vector, including scalar coordinates that lie in $(-1, 1)$. All distances between cube points use the maximum norm. For a real number s , define the sparse parameter arrangement

$$Z_s(t, n) = \{z \in T(t, n) : \#\{j : z_j \neq 0\} \leq s\}.$$

Thus it permits at most $\lfloor s \rfloor$ nonzero whole slots; $Z_s = \emptyset$ for $s < 0$, and $Z_s = T(t, n)$ for $s \geq n$. For $0 \leq s < n$ its dimension is $t\lfloor s \rfloor$. For $f : Y \rightarrow C(t, n)$, the map $f^{\times u} : Y^u \rightarrow C(t, un)$ concatenates the values of f , in copy order and then slot order.

The following torus data will turn a bundle embedding into a vanishing point class. Fix $c = 1/32$. For each positive integer p , choose a smooth embedding

$$L \cong (\mathbb{T}^2)^p \subset (-c, c)^{2p+1}$$

with a framed real normal line. Choose line bundles λ_i , each pulled back from the i th two-torus factor and embedded in $\mathbb{C}^2$, so that

$$H = \bigoplus_{i=1}^p \lambda_i \subset L \times \mathbb{C}^{2p}, \quad \left\langle \prod_{i=1}^p c_1(\lambda_i), [L] \right\rangle = 1.$$

[Lemma 5.1](#) constructs these data. Here p is a rank parameter, with no primality assumption.

Notation 1.2 (Torus locus and slot bundle). On L^n put

$$H_n = \bigoplus_{j=1}^n \text{pr}_j^* H \subset L^n \times \mathbb{C}^{2pn}.$$

For $f : Y \rightarrow C(2p+1, n)$ and $Z \subset T(2p+1, n)$, let

$$\mathcal{T}(f, Z) = \{(y, z) \in Y \times Z : f(y) - z \in L^n\}, \quad d_f(y, z) = f(y) - z.$$

The *slot bundle* on this locus is $E_f = d_f^* H_n$, of complex rank pn , with its fixed embedding in the trivial bundle of rank $2pn$.

Theorem 1.3 (Torus compression). *Let $p, n \geq 1$ be integers, put $t = 2p + 1$, and choose L, H as above. Let $0 < a < 1/16$, let Y be any compact metrizable space, and let $f : Y \rightarrow C(t, n)$ be continuous. Suppose there is one continuous complex bundle monomorphism*

$$E_f \longrightarrow \mathcal{T}(f, Z_{an}(t, n)) \times \mathbb{C}^K$$

over the entire locus $\mathcal{T}(f, Z_{an}(t, n))$, for a nonnegative integer $K < 2pn$. Then for every integer $U \geq 1$ there are an integer $u \geq U$ and a continuous map

$$G : Y^u \longrightarrow C(t, un)$$

such that at least $(a/8)un$ slots of $G(\mathbf{y})$ lie in $\partial[-1, 1]^t$ for every $\mathbf{y} = (y_0, \dots, y_{u-1})$. Moreover, for $0 \leq b < u$ and $1 \leq j \leq n$,

$$f_j(y_b) \in \partial[-1, 1]^t \implies G_{b,j}(\mathbf{y}) = f_j(y_b).$$

The bundle hypothesis includes continuous dependence on the sparse parameter as its support changes. Since the boundary count is an integer, its stated lower bound means at least $\lceil (a/8)un \rceil$ slots. The theorem also includes an empty torus locus; on a nonempty locus the embedding itself forces $K \geq pn$. The power u can be arbitrarily large, as stated, and the theorem requires no dimension bound on Y . The strict rank inequality is essential for this general criterion: already for $n = 1$ and $K = 2p$, the identity map of a cube admits the canonical bundle embedding but no boundary-preserving compression with a positive boundary count; see [Remark 5.4](#). This example does not address the optimal boundary fraction under the strict inequality.

1.3 Background and the proof mechanism

The nilpotence theorem of Devinatz–Hopkins–Smith [4] provides the stable-homotopy background for detecting smash-nilpotence by complex cobordism. Here we need the quantitative connective form developed by Mathew [10, Section 3.4] and recorded for BP in Burklund’s appendix [3, Appendix B]. Its cutoff is sublinear in the degree and uniform over all connective spectra at a fixed prime. This uniform statement, together with explicit upper cell bounds, converts a linearly growing Adams filtration into vanishing. The relation between unit-fiber powers, Adams towers and the cobar construction is standard [11, Section 2.1, Propositions 2.11 and 2.14]. The compatible null cone on the whole constrained product union is constructed in [Section 2](#).

The geometric step belongs to the collapse-and-duality framework of Thom and Atiyah [15, 1]. We prove the precise local duality needed for finite unions of coordinate planes, including its variance under changes of boxes and arrangements. A single relative nullhomotopy then gives a compatible stable lift over all simplices. Related stable obstruction criteria for sections in a connectivity range appear in [8, Theorems 4.2–4.3] and [7, Corollary 3.5]. Here relative support and nested boxes keep the lift compatible and control its displacement. Freudenthal stabilization, in the form recalled in [5, Corollary 4.24], turns it into a small actual displacement when the parameter dimension is in the indicated range. The singular arrangement and diagram compatibility assertions used here are proved in [Section 3](#).

The bundle criterion is related to the complementary-bundle Chern obstructions used by Hirshberg–Phillips in their lower bounds for the radius of comparison; see the proof of Theorem 3.3 in their author preprint [6]. We work with a supported point class in MU . The characteristic-class framework is classical; see Quillen [13] and the precise orientation and Thom formulas in [9, Lectures 5–6]. If $r = pn$, the complementary bundle has rank less than r , which forces the product of the r square-zero Chern classes to vanish. A framed real

Thom class and relative excision then give vanishing of the supported point class. These steps retain the whole sparse parameter locus.

The resulting theorem can be used in cover constructions. Fix a homeomorphism from the cube to a simplex carrying boundary to boundary. A boundary output then has at least one zero simplex coordinate. Preserving an original boundary vector retains the entire corresponding simplex point, and hence every coordinate that was already zero; this is the agreement used in the companion's reconstruction lemma [12, Lemma 4.3]. The companion [12, Theorem 1.1] uses [Theorem 1.3](#), together with dynamical comparison and cover reconstruction, to prove $\text{rc}(C(X) \rtimes_h \mathbb{Z}) = \frac{1}{2} \text{mdim}(X, h)$ for every minimal homeomorphism h of an infinite compact metrizable space X , with both sides allowed to be infinite. Here rc denotes radius of comparison and mdim denotes mean dimension.

Here is the sequence of constructions in the proof.

- (i) A point test for $f : P \rightarrow C(t, n)$ collapses the difference $f(x) - z$ near the origin, with z in a prescribed parameter arrangement. Product point tests are smash products. [Theorem 1.1](#) therefore promotes MU -vanishing of a sparse test to stable vanishing for a large batch of copies.
- (ii) Compatible local duality converts stable test vanishing into a map avoiding the sparse arrangement. An elementary radial normalization creates boundary slots and restores all original boundary vectors exactly.
- (iii) At each new batch, we keep only the old output as a parameter. Its dimension is bounded by its ambient cube. A fixed number of initial unchanged batches then keeps every step within the stable avoidance range. The resulting boundary fraction is at least $a/8$ for arbitrarily large powers.
- (iv) The torus-bundle embedding forces the MU point test to vanish. For an arbitrary compact metrizable Y , finitely many auxiliary coordinates give a polyhedral model, while including every coordinate of f preserves f exactly. Compression on that model pulls back with the same boundary agreement.

[Section 2](#) proves the filtered-product theorem. [Section 3](#) develops the point tests, local duality and stable avoidance. [Section 4](#) proves the general MU -null boundary-compression theorem by padding. [Section 5](#) establishes the bundle criterion and the exact compact-to-polyhedral reduction, completing [Theorem 1.3](#).

1.4 Conventions

Finite polyhedra and their pairs may be subdivided as needed. A quotient K/L means K_+/L_+ , where $+$ adjoins a disjoint basepoint; in particular, $K/\emptyset = K_+$. Suspension spectra, smash products, cofibers and limits of spectra are derived. The notation $F(K, E)$ denotes the function spectrum. We retain suspension shifts whenever connectivity or cell dimensions are used. A spectrum is b -connective when its homotopy groups in degrees below b vanish.

2 Compatible smash products and quantitative nilpotence

The topological argument will produce a complex-cobordism class which vanishes on a finite pointed subspace. We need the corresponding stable cohomotopy class to vanish on a union of products. The compatibility on the intersections of these products is essential. The following result provides it with a fixed positive proportion of restricted factors.

Our goal is [Theorem 1.1](#). We first build a lift through a power of the unit fiber on the entire union. We then use quantitative nilpotence at finitely many primes and a separate connectivity estimate at all larger primes. The latter step makes the eventual product threshold independent of the prime.

All constructions with spectra in this section are derived, and a commutative ring spectrum means an $\mathbb{E}_\infty$ ring spectrum. A spectrum is b -connective if its homotopy groups below degree b vanish. We write $[K, Y]$ for the group of homotopy classes of maps of spectra and $\text{Map}(K, Y)$ for their mapping space. For a ring spectrum R under the sphere, put

$$I_R = \text{fib}(\mathbb{S} \longrightarrow R), \quad \varphi_{R,n} : I_R^{\wedge n} \longrightarrow \mathbb{S}.$$

The second map is the smash product of the n fiber inclusions, with $I_R^{\wedge 0} = \mathbb{S}$. In a prime-local category, the sphere in this notation is $\mathbb{S}_{(p)}$.

We use the commutative Thom-spectrum model of MU and its canonical complex orientation as recalled in [9, Lecture 6, Remarks 5 and 7 and Theorem 8].

A map $K \rightarrow Y$ has R -Adams filtration at least n if it lifts through

$$\varphi_{R,n} \wedge \text{id}_Y : I_R^{\wedge n} \wedge Y \longrightarrow Y.$$

2.1 A compatible lift on the union

A map to the sphere which becomes null after applying the unit $\mathbb{S} \rightarrow R$ lifts through I_R . On a product with n restricted factors, this suggests a lift through $I_R^{\wedge n}$. The difficulty is making these lifts agree where different choices of restricted factors meet. We construct one lift on the whole union.

Proposition 2.1 (Compatible Adams lift). *Let R be a commutative ring spectrum, let $G \subset E$ be a finite pointed CW pair, let $d \in \mathbb{Z}$, and let $v : \Sigma^\infty E \rightarrow \Sigma^d \mathbb{S}$ be a stable map whose restriction to $\Sigma^\infty G$ becomes null in $\Sigma^d R$. For integers $1 \leq n \leq \ell$, let $U_{\ell,n} \subset E^{\wedge \ell}$ be the union requiring at least n factors in G . Then the restriction of $v^{\wedge \ell}$ admits a lift*

$$\Sigma^\infty U_{\ell,n} \longrightarrow \Sigma^{d\ell} I_R^{\wedge n}$$

through $\Sigma^{d\ell} \varphi_{R,n}$.

We describe the finite diagram in which the lift will be constructed. Let $\mathcal{P}_n^\times$ be the poset of nonempty subsets of $\{1, \dots, n\}$. Give a subset S the value $R^{\wedge |S|}$, with maps inserting units along subset inclusions. Adjoining the sphere at the empty subset gives the n -fold cube of units. The fiber of its unit-cone map

$$\mathbb{S} \longrightarrow \text{holim}_{S \in \mathcal{P}_n^\times} R^{\wedge |S|}$$

is $I_R^{\wedge n}$, as we verify at the end of the proof. It therefore suffices to null the composite of $v^{\wedge \ell}|_{U_{\ell,n}}$ with the $d\ell$ -fold suspension of this map.

We will produce this nullhomotopy by introducing simplex parameters. The given R -nullhomotopy on G will supply compatible nullhomotopies at the vertices of every simplex. We distribute one simplex parameter among the ℓ product factors so that in degree j at most j of the resulting parameters are away from vertices. When $j < n$, at least one of the n restricted factors then receives a vertex and is collapsed. The following construction makes this distribution natural in the simplex. The relation between unit cubes and the cosimplicial cobar construction is recalled in [11, Section 2.1, Propositions 2.11 and 2.14].

Let Δ denote the simplex category, with objects $[j] = \{0, \dots, j\}$ and monotone maps. The topological simplices Δ^j form a covariant functor: a monotone map sends a distribution on its source vertices to the pushforward distribution on its target vertices. Write $V^j \subset \Delta^j$ for the set of vertices.

Lemma 2.2 (Slicing a simplex). *For every $\ell \geq 1$ there is a natural continuous map*

$$\rho = (\rho_1, \dots, \rho_\ell) : \Delta^j \longrightarrow (\Delta^j)^\ell$$

in the variable $[j] \in \Delta$, such that at most j of the points $\rho_r(x)$ are nonvertices, for each $x \in \Delta^j$.

Proof. For $x = (x_0, \dots, x_j)$ put $s_a = \sum_{b < a} x_b$, so that $s_0 = 0$ and $s_{j+1} = 1$. For $1 \leq r \leq \ell$ define

$$\rho_r(x)_a = \ell \max \left\{ 0, \min \left(s_{a+1}, \frac{r}{\ell} \right) - \max \left(s_a, \frac{r-1}{\ell} \right) \right\}, \quad 0 \leq a \leq j. \quad (2.1)$$

This is ℓ times the length of the intersection of $[s_a, s_{a+1}]$ with the r th interval of length $1/\ell$. The coordinates are continuous, nonnegative, and sum to one.

Every nonempty fiber of a monotone map $\alpha : [j] \rightarrow [k]$ is a consecutive block. Passing from x to $\alpha_* x$ merges the adjacent mass intervals in that block. The intersection length of the merged interval with a fixed slice equals the sum of its constituent intersection lengths. Empty fibers contribute zero. It follows that $\alpha_* \rho_r(x) = \rho_r(\alpha_* x)$ for every r . This verifies naturality for all maps of Δ , including its surjections.

A nonvertex value $\rho_r(x)$ requires at least one of the internal breaks $s_1, \dots, s_j$ to lie strictly between $(r-1)/\ell$ and r/ℓ . Distinct such slices have disjoint interiors and therefore require distinct breaks. There are at most j of them. Breaks lying at slice endpoints do not create nonvertex values. $\square$




Figure 1: Simplex slicing for $j = 2$ and $\ell = 4$. Each slice records the proportions of the three mass intervals that it meets. A nonvertex value requires an internal mass break in the interior of that slice, so there can be at most j such values.

Figure 1 shows the counting mechanism in Lemma 2.2. We now combine this distribution with one chosen nullhomotopy to construct the lift.

Proof of Proposition 2.1. Work in the stable functor category $\mathcal{D} = \text{Fun}(\Delta, \text{Sp})$. Its equivalences, cofibers, and smash products are computed objectwise. Consider the cosimplicial commutative ring spectrum

$$B^j = R^{\wedge(j+1)}.$$

For a monotone map $\alpha : [j] \rightarrow [k]$, its structure map multiplies the factors in each fiber of α and inserts a unit for each empty fiber. This describes its cofaces and codegeneracies, together with their coherences. In particular, the maps $(B^j)^{\wedge \ell} \rightarrow B^j$ given by multiplication are natural in $[j]$.

The nullhomotopy on the vertices. The functor V is the covariant representable $\text{Hom}_\Delta([0], -)$. Thus $\Sigma^\infty(G \wedge V_+^\bullet)$ is the left Kan extension of $\Sigma^\infty G$ from the object $[0]$. The resulting adjunction is an equivalence of mapping spaces

$$\text{Map}_{\mathcal{D}}(\Sigma^\infty(G \wedge V_+^\bullet), \Sigma^d B^\bullet) \simeq \text{Map}_{\text{Sp}}(\Sigma^\infty G, \Sigma^d R). \quad (2.2)$$

Indeed, the left Kan extension in degree j is the wedge of copies of $\Sigma^\infty G$ indexed by the $j+1$ maps $[0] \rightarrow [j]$; the adjunction says that a map from this diagram is determined coherently by its value on the copy in degree zero.

There is a natural map

$$a : \Sigma^\infty(E \wedge \Delta_+^\bullet) \longrightarrow \Sigma^d B^\bullet$$

which forgets the simplex coordinate and applies v followed by the unit. Its restriction to $G \wedge V_+^\bullet$ corresponds under Equation (2.2) to the assumed null map into $\Sigma^d R$. Choose a nullhomotopy there. The mapping-space adjunction supplies a single natural nullhomotopy on the vertex diagram. The cofiber universal property therefore gives a map

$$\psi : Q^\bullet \longrightarrow \Sigma^d B^\bullet, \quad Q^j = \Sigma^\infty \left(\frac{E \wedge \Delta_+^j}{G \wedge V_+^j} \right),$$

and a homotopy $\psi q \simeq a$ in $\mathcal{D}$, where q is the quotient map. The displayed ordinary quotients compute the derived cofibers: the inclusion of the indicated pointed CW subspace is a cofibration in each degree. This also permits us to use the actual collapse maps in the next step.

Collapse on the whole union. Put $U = U_{\ell,n}$. The maps of Lemma 2.2 give a natural map of pointed spaces

$$U \wedge \Delta_+^j \longrightarrow (E \wedge \Delta_+^j)^{\wedge \ell}, \quad (x_1 \wedge \cdots \wedge x_\ell, x) \longmapsto \bigwedge_{r=1}^{\ell} (x_r, \rho_r(x)).$$

Follow it by the smash of the quotient maps to obtain, after suspension, a map into $(Q^j)^{\wedge \ell}$. If $j < n$, at least n of the factors x_r belong to G , while at most j of the simplex parameters $\rho_r(x)$ are nonvertices. Some factor is consequently in $G \wedge V_+^j$ and is collapsed. The resulting map

$$\Sigma^\infty(U \wedge \Delta_+^j) \longrightarrow (Q^j)^{\wedge \ell} \tag{2.3}$$

is the zero map for every $j < n$. It is the zero natural transformation on these degrees, since it is already a literal collapse on the diagrams of spaces.

Compose Equation (2.3) with $\psi^{\wedge \ell}$ and multiplication to obtain a natural map into $\Sigma^{d\ell} B^j$. Smashing the chosen factorization homotopy $\psi q \simeq a$ and then multiplying identifies this map with the map which forgets the simplex coordinate and applies $v^{\wedge \ell}|_U$ followed by the unit. This identification is natural: each map a forgets its own simplex parameter, so it is independent of the particular values of $\rho_r(x)$.

The natural projection

$$\Sigma^\infty(U \wedge \Delta_+^\bullet) \longrightarrow \text{const}(\Sigma^\infty U)$$

is an objectwise equivalence, since every Δ^j is contractible. Invert this equivalence in the functor category and restrict to the full subcategory $\Delta_{\leq n-1}$. We have proved that the cone

$$\text{const}(\Sigma^\infty U) \longrightarrow \text{const}(\Sigma^{d\ell} \mathbb{S}) \longrightarrow \Sigma^{d\ell} B^\bullet$$

is null on that subcategory. Inverting an objectwise equivalence here does not require a natural choice of a simplex vertex or a natural contraction of the simplices.

Restriction to the unit cube. The rule

$$S \longmapsto [|S| - 1]$$

defines a functor from the punctured cube $\mathcal{P}_n^\times$ to $\Delta_{\leq n-1}$: an inclusion of subsets induces the ordered injection on their enumerations. Restricting $B^\bullet$ gives the diagram $S \mapsto R^{\wedge|S|}$, whose maps insert units. The null cone just constructed therefore yields a null composite

$$\Sigma^\infty U \xrightarrow{v^{\wedge \ell}} \Sigma^{d\ell} \mathbb{S} \longrightarrow \Sigma^{d\ell} \operatorname{holim}_{S \in \mathcal{P}_n^\times} R^{\wedge|S|}. \quad (2.4)$$

Only restriction of cones is used; no assertion about cofinality or an identification of partial totalizations is needed.

Adjoin the value $\mathbb{S}$ at the empty subset. The resulting n -cube is the external smash product of n copies of $\mathbb{S} \rightarrow R$. Its total fiber is $I_R^{\wedge n}$. To verify this, compute fibers successively in its n directions. Taking the fiber in the last direction replaces every map $C \wedge \mathbb{S} \rightarrow C \wedge R$ by $C \wedge I_R$, because smash with C preserves finite fiber sequences in spectra. Induction on the remaining directions gives $I_R^{\wedge n}$. Successive fibers compute the fiber of the initial vertex over the limit of the punctured cube. For two directions the relevant identity is

$$\operatorname{fib}(A \longrightarrow B \times_D C) \simeq \operatorname{fib}(\operatorname{fib}(A \rightarrow B) \longrightarrow \operatorname{fib}(C \rightarrow D)),$$

obtained by expressing each fiber as a pullback over zero. Repeated application gives the assertion for any number of directions, because finite limits commute. Thus there is a fiber sequence

$$I_R^{\wedge n} \xrightarrow{\varphi_{R,n}} \mathbb{S} \longrightarrow \operatorname{holim}_{S \in \mathcal{P}_n^\times} R^{\wedge|S|}.$$

The chosen nullhomotopy in Equation (2.4) gives the required lift through its suspended fiber. $\square$

2.2 The quantitative nilpotence input

The preceding proposition gives Adams filtration at least the number of restricted factors. To turn this lift into vanishing, we use a quantitative form of nilpotence for MU . Its cutoff is sublinear in the degree and uniform over connective spectra at a fixed prime. We state it as a null action on a truncated sphere, the form needed below for finite sources which may have negative-degree cells. We write $\tau_{<k} X$ for the Postnikov truncation of X retaining homotopy groups in degrees below k .

Theorem 2.3 (Quantitative nilpotence). *Fix a prime p , and let R be either $MU_{(p)}$ or the Brown–Peterson spectrum BP at that prime. For $k \geq 1$, let $f_R(k)$ be the least integer $n \geq 0$ for which*

$$I_R^{\wedge n} \wedge \tau_{<k} \mathbb{S}_{(p)} \xrightarrow{\varphi_{R,n} \wedge \operatorname{id}} \tau_{<k} \mathbb{S}_{(p)} \quad (2.5)$$

is nullhomotopic. These integers exist and satisfy

$$\lim_{k \rightarrow \infty} \frac{f_R(k)}{k} = 0.$$

Equivalently, $f_R(k)$ is the least n such that, for every connective p -local spectrum X and every $i < k$, a class in $\pi_i X$ of R -Adams filtration at least n is zero.

For $MU_{(p)}$, this is [10, Definitions 2.28 and 3.26, Proposition 3.28, and Theorem 3.30]. For BP , the same formulation and its sublinear bound are recorded in [3, Definition B.3, the discussion following Remark B.4, and Proposition B.15]. Both rings are connective, have degree-zero homotopy $\mathbb{Z}_{(p)}$, and have finitely generated homotopy in every degree, as required in those statements.

The truncated action gives the formulation for all connective spectra directly. If Equation (2.5) is null, smash it with a connective p -local spectrum X . The map

$$X \longrightarrow X \wedge \tau_{<k}\mathbb{S}_{(p)}$$

is an isomorphism on homotopy in degrees below k : its fiber is $X \wedge \tau_{\geq k}\mathbb{S}_{(p)}$, which is k -connective. A class with the asserted lift maps to zero on the right and therefore was zero in $\pi_i X$.

For our application, the key consequence is a null map into the truncated sphere itself. Precomposing Equation (2.5) with the truncation unit $\eta_k : \mathbb{S}_{(p)} \rightarrow \tau_{<k}\mathbb{S}_{(p)}$ in its second factor gives

$$\eta_k \circ \varphi_{R,n} = 0 \quad (n \geq f_R(k)). \quad (2.6)$$

The assertion for $n > f_R(k)$ follows by applying the fiber inclusions to the additional smash factors.

The next observation explains how truncation detects maps from the finite complexes occurring here. It uses an upper bound on cell degrees, without requiring the source to be connective.

Lemma 2.4 (Detection by truncation). *Suppose that a finite spectrum K has a finite cell presentation with all cell degrees at most D , either in spectra or in p -local spectra. If Y is $(D+1)$ -connective, then $[K, Y] = 0$. In particular, for every integer $k > D$, the map*

$$[K, \mathbb{S}_{(p)}] \longrightarrow [K, \tau_{<k}\mathbb{S}_{(p)}]$$

is injective.

Proof. For a single cell in degree $s \leq D$, the mapping group is $[S^s, Y] = \pi_s Y = 0$. Attaching finitely many such cells proves the first assertion by induction: the cofiber of an attaching step is a finite wedge of these spheres, and the exact sequence obtained by mapping into Y makes the mapping group of the enlarged complex zero when those of the preceding complex and the cofiber are zero. For the second assertion, apply the first to $Y = \tau_{\geq k}\mathbb{S}_{(p)}$ and use the exact sequence obtained from the fiber sequence

$$\tau_{\geq k}\mathbb{S}_{(p)} \longrightarrow \mathbb{S}_{(p)} \longrightarrow \tau_{<k}\mathbb{S}_{(p)}.$$

□

2.3 From local vanishing to integral vanishing

Proposition 2.1 supplies a single lift of filtration proportional to the number of factors. We now show that the restricted product class vanishes once the product is large. At large primes a uniform connectivity estimate suffices, leaving only finitely many prime-dependent nilpotence thresholds.

Lemma 2.5 (The unit fiber at large primes). *For every odd prime p , the spectrum BP is a retract of $MU_{(p)}$ by unit-preserving maps of spectra, and*

$$I_{BP} \in \mathrm{Sp}_{\geq 2p-3}.$$

Consequently a lift through $(I_{MU})_{(p)}^{\wedge n}$ induces a lift through a spectrum which is $n(2p-3)$ -connective.

Proof. The unit-preserving retract follows, for example, from the $\mathbb{E}_4$ splitting of $MU_{(p)}$ in [2, Theorem 1.1, the following paragraph, and Section 5]. We use the underlying retract of spectra for ordinary homology and the forward map under $\mathbb{S}_{(p)}$ to transfer unit fibers.

Let C be the cofiber of $\mathbb{S}_{(p)} \rightarrow BP$. The mod- p homology calculation in [2, Section 5] gives

$$H_0(BP; \mathbb{F}_p) = \mathbb{F}_p, \quad H_i(BP; \mathbb{F}_p) = 0 \quad (0 < i < 2p - 2).$$

The unit induces the degree-zero isomorphism, and BP is connective. Its unit cofiber therefore has zero mod- p homology in every degree below $2p - 2$.

The splitting above exhibits BP as a retract of $MU_{(p)}$. The ordinary integral homology of MU is $\mathbb{Z}[b_1, b_2, \dots]$, with $|b_i| = 2i$ [9, Lecture 7, Proposition 2 and Corollary 3]; in particular, it is finitely generated in every degree. Taking the retract and then the homology sequence of the unit cofiber shows that $H_i(C; \mathbb{Z}_{(p)})$ is finitely generated in every degree. For each $i < 2p - 2$, the universal coefficient sequence contains an injection

$$H_i(C; \mathbb{Z}_{(p)})/pH_i(C; \mathbb{Z}_{(p)}) \hookrightarrow H_i(C; \mathbb{F}_p) = 0.$$

Nakayama's lemma therefore gives $H_i(C; \mathbb{Z}_{(p)}) = 0$ for every $i < 2p - 2$. The spectrum C is connective. If it had a nonzero homotopy group below degree $2p - 2$, let $b \geq 0$ be the least such degree. The stable Hurewicz isomorphism in the first nonzero degree would give $\pi_b(C) \cong H_b(C; \mathbb{Z}_{(p)}) = 0$, a contradiction. Thus C is $(2p - 2)$ -connective, and $I_{BP} \simeq \Sigma^{-1}C$ is $(2p - 3)$ -connective.

The map under the sphere induces a map $(I_{MU})_{(p)} \rightarrow I_{BP}$ commuting with the fiber inclusions. Smashing n times proves the last assertion, since smash products add connectivity bounds. $\square$

Proof of Theorem 1.1. Put $e = \dim E$, $n_\ell = \lceil \ell/2 \rceil$, and

$$K_\ell = \Sigma^{-d\ell} \Sigma^\infty U_\ell.$$

The finite subcomplex $U_\ell \subset E^{\wedge \ell}$ has cell dimension at most $e\ell$, so K_ℓ has upper cell degree

$$D_\ell = (e - d)\ell. \tag{2.7}$$

The class under consideration is now a map $\beta_\ell : K_\ell \rightarrow \mathbb{S}$. By Proposition 2.1, it lifts through $I_{MU}^{\wedge n_\ell}$.

If $e < d$, Lemma 2.4 applied directly to the connective sphere shows that $\beta_\ell = 0$ for every ℓ . Assume henceforth that $e \geq d$ and put

$$k_\ell = (e - d)\ell + 1.$$

For a fixed prime p , localization of the lift gives a lift through $I_{MU_{(p)}}^{\wedge n_\ell}$. If $e > d$, Theorem 2.3 gives

$$f_{MU_{(p)}}(k_\ell) = o(\ell),$$

so that $n_\ell \geq f_{MU_{(p)}}(k_\ell)$ for all sufficiently large ℓ . If $e = d$, the same inequality eventually holds because $k_\ell = 1$ is fixed and $n_\ell \rightarrow \infty$. By Equation (2.6), the composite of $(\beta_\ell)_{(p)}$ with $\tau_{< k_\ell} \mathbb{S}_{(p)}$ is zero. By Lemma 2.4 and Equation (2.7), $(\beta_\ell)_{(p)} = 0$. This proves eventual vanishing for each fixed prime.

To obtain a single threshold, choose an integer $P \geq 2$ such that

$$\frac{2p - 3}{2} > \max(e - d, 0) \quad \text{for every prime } p > P. \tag{2.8}$$

For such a prime, [Lemma 2.5](#) turns the lift of $(\beta_\ell)_{(p)}$ into a lift through $I_{BP}^{\wedge n_\ell}$. Its target has connectivity at least

$$n_\ell(2p-3) \geq \frac{\ell}{2}(2p-3) > (e-d)\ell = D_\ell.$$

The first assertion of [Lemma 2.4](#) makes this lift zero, for every ℓ . Only the finitely many primes at most P remain. Take the maximum of their eventual thresholds from the preceding paragraph. Above that maximum, β_ℓ vanishes after localization at every prime.

Because K_ℓ is finite, localization of its stable mapping group agrees with maps from its localization. One way to see this is the duality identity

$$F(K_\ell, \mathbb{S}) \simeq DK_\ell, \quad (DK_\ell)_{(p)} \simeq F((K_\ell)_{(p)}, \mathbb{S}_{(p)}),$$

where DK_ℓ denotes the finite Spanier–Whitehead dual. An element of an abelian group which vanishes in every localization at a prime is zero. Indeed, an element of infinite order remains nonzero in every such localization, and a nonzero element of finite order remains nonzero at any prime dividing its order. Applying this to $[K_\ell, \mathbb{S}]$ gives $\beta_\ell = 0$ integrally, as required. $\square$

Remark 2.6. The same proof works with $\lceil \lambda \ell \rceil$ restricted factors for any fixed $0 < \lambda \leq 1$. In [Proposition 2.1](#), take $n = \lceil \lambda \ell \rceil$; in [Equation \(2.8\)](#), replace $1/2$ by λ . The finite pair affects the eventual value of ℓ , but imposes no lower bound on the positive proportion λ .

3 Point tests and controlled avoidance

The next step turns a stable vanishing statement into a small change of a map. The parameter in the test records the points to be avoided. Keeping this parameter throughout the argument supplies the compatibility needed to make the changes on adjacent simplices agree.

3.1 Point tests and their products

We use the cubes $C(t, n)$ and $T(t, n)$ and the sparse arrangements $Z_s(t, n)$ from [Section 1](#), and write $M = tn$. Coordinate subspaces below may also be specified by setting individual scalar coordinates to zero, so the terminology includes whole-slot conditions. A point test records equality with a parameter z as a relative stable cohomotopy class. Its relative support is essential for the later control on displacement.

Definition 3.1 (Point test). Recall $c = 1/32$. Let P be a finite polyhedron, let $f : P \rightarrow C(t, n)$ be PL, and let $Z \subseteq T(t, n)$ be a subpolyhedron. Define

$$\begin{aligned} A_c(f, Z) &= \{(x, z) \in P \times Z : \|f(x) - z\|_\infty \geq c\}, \\ Q_c(f, Z) &= (P \times Z) / A_c(f, Z). \end{aligned}$$

For $0 < \rho < c$, take the oriented collapse $\kappa_\rho : \mathbb{R}^M \rightarrow S^M$ of the complement of $(-\rho, \rho)^M$. The *point test* is the based map

$$\varphi_\rho(f, Z) : Q_c(f, Z) \longrightarrow S^M, \quad [x, z] \longmapsto \kappa_\rho(f(x) - z). \quad (3.1)$$

Its stable class is denoted by

$$\omega_{\mathbb{S}}(f, Z) \in [\Sigma^\infty Q_c(f, Z), \Sigma^M \mathbb{S}].$$

Composing with the unit $\mathbb{S} \rightarrow MU$ gives $\omega_{MU}(f, Z) \in \widetilde{MU}^M(Q_c(f, Z))$.

One concrete choice of κ_ρ is the smash product of M oriented one-dimensional interval collapses, in the order of the scalar coordinates. Varying ρ gives a homotopy through maps constant outside $(-c, c)^M$. More generally, any positively oriented collapse supported in that cube represents its relative generator and gives the same class. We may therefore shrink the support whenever necessary. The difference map is PL, so $A_c(f, Z)$ is a subpolyhedron after a finite subdivision. The quotient in [Definition 3.1](#) is consequently a finite pointed polyhedron.

Lemma 3.2 (Restrictions and products). *Point tests are natural under pullback in the input and under restriction of the parameter set. In particular, if $Z \subseteq Z'$ are subpolyhedra of the same parameter cube, the induced map $Q_c(f, Z) \rightarrow Q_c(f, Z')$ identifies the former with a pointed subpolyhedron after a common subdivision.*

Suppose $f_i : P_i \rightarrow [-1, 1]^{M_i}$ and $Z_i \subseteq [-2, 2]^{M_i}$ are finitely many such data. With coordinates concatenated in order, there is a canonical based homeomorphism

$$Q_c\left(\prod_i f_i, \prod_i Z_i\right) \cong \bigwedge_i Q_c(f_i, Z_i). \quad (3.2)$$

Under the corresponding identification $S^{\sum_i M_i} \cong \bigwedge_i S^{M_i}$, the point test is the smash product of the point tests. The analogous equality for MU uses the multiplication of MU .

Proof. All restriction maps are induced by maps of the defining pairs. For the subpolyhedron assertion, triangulate $P \times Z'$ so that $P \times Z$ and $A_c(f, Z')$ are subpolyhedra. Collapsing the latter identifies no two points of the former except those already collapsed in $Q_c(f, Z)$. Quotients of this finite polyhedral pair give the assertion, including the stated basepoint convention.

For products the relative subset is exactly the union of the subsets on which at least one factor belongs to $A_c(f_i, Z_i)$. Its quotient is the smash product of the individual quotients. Choosing all collapses as products of the same one-dimensional collapse makes the identity of maps literal. Composing with units and multiplying in MU gives the last assertion. $\square$

3.2 Coordinate complements in boxes

We record explicitly both the local duality and its compatibility. The equality-collapse pairing is a local version of the classical collapse and stable-duality constructions [\[15, 1\]](#). We prove it here for coordinate arrangements, starting with one plane and passing to unions by excision. The proof will also identify the variance needed to assemble maps over the simplices of P . Write $T = [-2, 2]^M$. A *coordinate arrangement* in T is a finite union of sets

$$L_J = \{z \in T : z_i = 0 \text{ for all } i \in J\}, \quad J \subseteq \{1, \dots, M\}.$$

For a PL map $f : P \rightarrow C(t, n)$ from a finite polyhedron and a coordinate arrangement Z , our goal is to turn $\omega_{\mathbb{S}}(f, Z) = 0$ into a displacement of size less than $1/2$ avoiding Z , provided $\dim Z \leq M - q$ and $\dim P \leq 2q - 5$ for an integer $q \geq 1$. The local complements will be $(q - 2)$ -connected. Equality-collapse duality will identify the stable obstruction, and this connectivity will let us realize a compatible stable lift by actual maps in [Lemma 3.5](#).

Consider finitely many pairs of full-dimensional closed axis-parallel boxes

$$B_\alpha \subset \text{int } B_\alpha^+ \subset (-2, 2)^M. \quad (3.3)$$

Assume that no coordinate endpoint is zero, and that corresponding coordinate intervals of B_α and B_α^+ either both contain zero or both miss zero. In particular they meet precisely

the same coordinate subspaces. Whenever the boxes are indexed by a partial order, we also require both families to be increasing in that order.

Choose $\epsilon > 0$ smaller than the absolute value of every endpoint of every box. If Z is a nonempty arrangement other than T , write $Z = \bigcup_{J \in \mathcal{J}} L_J$ with every J nonempty, and put

$$d_Z(y) = \min_{J \in \mathcal{J}} \max_{i \in J} |y_i|, \quad U_\epsilon(Z) = \{y \in T : d_Z(y) < \epsilon\}, \quad V_\alpha(Z) = B_\alpha \setminus U_\epsilon(Z).$$

Here d_Z is exactly the maximum-norm distance to Z : setting the specified coordinates to zero stays inside T . For the empty and full arrangements put $U_\epsilon(\emptyset) = \emptyset$ and $U_\epsilon(T) = T$.

Lemma 3.3 (Thickened complements). *For sufficiently small ϵ as above, the inclusion*

$$V_\alpha(Z) \longrightarrow B_\alpha \setminus Z$$

is a strong deformation retract equivalence, simultaneously for all the boxes. The homotopy fixes $V_\alpha(Z)$ pointwise, preserves each box, and preserves the set of zero coordinates. If $\dim Z \leq M - q$ for an integer $q \geq 1$, then a nonempty such complement is $(q - 2)$ -connected.

Proof. The empty and full arrangements are immediate. In the remaining case, for $y \notin Z$ set

$$\lambda(y) = \max\{1, \epsilon/d_Z(y)\}, \quad \lambda_u(y) = 1 + u(\lambda(y) - 1), \quad 0 \leq u \leq 1.$$

Define a coordinatewise homotopy by

$$H_u(y)_i = \operatorname{sgn}(y_i) \min\{\lambda_u(y)|y_i|, \max\{|y_i|, \epsilon\}\}, \quad (3.4)$$

with value zero when $y_i = 0$. It is continuous on $T \setminus Z$: λ_u is locally bounded there, which also gives continuity at zero coordinates. Coordinates of magnitude at least ϵ are fixed. Each smaller coordinate is expanded, without changing sign, by at most enough to reach magnitude ϵ .

Every coordinate interval of a box either contains $[-\epsilon, \epsilon]$ or lies outside that interval. Thus Equation (3.4) preserves even asymmetric boxes. Coordinate magnitudes never decrease, and zero coordinates remain zero, so the homotopy stays in the complement and preserves $V_\alpha(Z)$ at every time. In fact it fixes that subspace pointwise, since $d_Z(y) \geq \epsilon$ implies $\lambda(y) = 1$. If $d_Z(y) < \epsilon$, each set J contains an index with $|y_i| \geq d_Z(y)$, and that coordinate has magnitude at least ϵ at time one. Hence $d_Z(H_1(y)) \geq \epsilon$. This proves the retraction. The formula is the same for every box. In particular it applies to any compact family in the unthickened complement, with no need to move arbitrarily small nonzero coordinates discontinuously to a fixed radius.

For connectivity, first move a compact family from the closed box into its interior without changing its zero coordinates: shrink intervals containing zero by a fixed factor in $(0, 1)$, and move the other intervals toward an interior point of the same sign. A map from S^i into the complement, $i \leq q - 2$, can then be approximated by a PL map in the interior and extended over a triangulated D^{i+1} into the convex box. A sufficiently small generic perturbation of its vertices misses all the excluded subspaces, since

$$(i + 1) + (M - q) < M.$$

The perturbed boundary map is homotopic to the original within the complement. This proves the claimed connectivity, including path connectedness in the applicable range. Existence of a point in the complement when $q \geq 1$ follows from the same dimension observation. The retraction transfers connectivity to $V_\alpha(Z)$. $\square$

The local problem is to replace a map into B_α by one with values in $V_\alpha(Z)$. We first seek a lift after stabilization. The cofiber of the complement inclusion records its obstruction; the corresponding point test retains the parameters of Z inside B_α^+ . Define finite spectra and finite pointed test spaces by

$$\mathcal{C}_\alpha(Z) = \text{cofib}(\Sigma^\infty V_\alpha(Z)_+ \longrightarrow \Sigma^\infty (B_\alpha)_+), \quad (3.5)$$

$$T_\alpha(Z) = Z/(Z \setminus \text{int } B_\alpha^+). \quad (3.6)$$

The letter T_α in Equation (3.6) denotes a pointed quotient, rather than the parameter cube T .

Lemma 3.4 (Compatible local duality). *For the finite box families in Equation (3.3), one sufficiently small collapse support gives equivalences*

$$\mathcal{C}_\alpha(Z) \simeq F(\Sigma^\infty T_\alpha(Z), \Sigma^M \mathbb{S}). \quad (3.7)$$

They are induced by collapse around equality and are natural both under box inclusions and under inclusions of coordinate arrangements. In the box variable the cofibers are covariant and the test spaces are contravariant, by quotient collapse. In the arrangement variable, the cofibers are contravariant and the test spaces are covariant.

Proof. Choose one $\rho > 0$ with

$$\rho < \epsilon, \quad \rho < \min \text{dist}_\infty(B_\alpha, T \setminus \text{int } B_\alpha^+). \quad (3.8)$$

All minima are positive, by finiteness and strict containment. The formula $(y, z) \mapsto \kappa_\rho(y - z)$ descends to a pairing

$$\mathcal{C}_\alpha(Z) \wedge \Sigma^\infty T_\alpha(Z) \longrightarrow \Sigma^M \mathbb{S}. \quad (3.9)$$

Indeed, if $y \in V_\alpha(Z)$ and $z \in Z$, their distance is at least ϵ . If $z \notin \text{int } B_\alpha^+$ and $y \in B_\alpha$, the second inequality in Equation (3.8) applies. These are exactly the two collapsed subspaces. Since $V_\alpha(Z)$ is a subpolyhedron, its ordinary quotient by inclusion in B_α represents the cofiber in Equation (3.5).

First take $Z = L_J$, with $r = |J|$ and $b = M - r$. If it misses B_α , some required coordinate is bounded away from zero by more than ϵ , so $V_\alpha(Z) = B_\alpha$. It also misses B_α^+ , and both sides of Equation (3.7) are zero. Otherwise split the box into its normal and tangential factors, $B_\alpha = B_N \times B_T$. Then

$$B_\alpha/V_\alpha(L_J) \cong (B_N/(B_N \setminus (-\epsilon, \epsilon)^r)) \wedge (B_T)_+ \simeq S^r, \quad T_\alpha(L_J) \cong S^b.$$

The conventions include $r = 0$ and $b = 0$. Fixing an interior point $w \in B_T$ identifies the pairing with the collapse of $(u, z_T) \mapsto (u, w - z_T)$, up to a coordinate permutation. It has degree ± 1 : the origin has one preimage, and its local linear map is $\text{diag}(I_r, -I_b)$. Its adjoint is therefore an equivalence $S^r \simeq F(S^b, \Sigma^M \mathbb{S})$.

Here is the excision argument for unions. For coordinate planes,

$$U_\epsilon(L_J) \cap U_\epsilon(L_K) = U_\epsilon(L_{J \cup K}).$$

Distributing finite unions gives, for arrangements Z, Z' , the identities

$$\begin{aligned} V_\alpha(Z \cup Z') &= V_\alpha(Z) \cap V_\alpha(Z'), \\ V_\alpha(Z \cap Z') &= V_\alpha(Z) \cup V_\alpha(Z'). \end{aligned}$$

A rectangular subdivision followed by triangulation makes all these closed complements subpolyhedra. Their union square is a homotopy pushout. Applying suspension spectra and cofibers into the constant spectrum $\Sigma^\infty(B_\alpha)_+$ gives the homotopy pullback description

$$\mathcal{C}_\alpha(Z \cup Z') \simeq \mathcal{C}_\alpha(Z) \underset{\mathcal{C}_\alpha(Z \cap Z')}{\overset{h}{\times}} \mathcal{C}_\alpha(Z'). \quad (3.10)$$

We used here that a pushout of spectra is also a pullback. On the test-space side there is a pushout of pointed polyhedra along cofibrations

$$T_\alpha(Z \cup Z') \cong T_\alpha(Z) \cup_{T_\alpha(Z \cap Z')} T_\alpha(Z').$$

Its function spectra give the same pullback as in Equation (3.10), and the pairings commute with these maps. Induction on the number of planes proves Equation (3.7): the intersection of one plane with a union of fewer planes is again a union of no more than that many coordinate planes. The empty arrangement contributes zero throughout.

Finally, for an arrangement inclusion $Z \subseteq Z'$, the maps are

$$\mathcal{C}_\alpha(Z') \longrightarrow \mathcal{C}_\alpha(Z), \quad T_\alpha(Z) \longrightarrow T_\alpha(Z').$$

For a box inclusion indexed by $\alpha \leq \beta$, the maps are

$$\mathcal{C}_\alpha(Z) \longrightarrow \mathcal{C}_\beta(Z), \quad T_\beta(Z) \longrightarrow T_\alpha(Z),$$

the second being quotient collapse. On representatives, each required compatibility says that the pairing is $\kappa_\rho(y - z)$. If the smaller test quotient additionally collapses z , the pairing with $y \in B_\alpha$ was already the basepoint by Equation (3.8). Thus the adjoints are natural equivalences of diagrams, not just objectwise identifications. $\square$

3.3 From a stable lift to a map of complements

The local duality is now available as an equivalence of diagrams. We use a single global nullhomotopy to lift all simplex maps simultaneously, and then use connectivity to realize the lift by actual maps into the complements.

Lemma 3.5 (Stable avoidance). *Let P be a finite polyhedron and $f : P \rightarrow C(t, n)$ a PL map. Let $Z \subseteq T(t, n)$ be a coordinate arrangement, and let $q \geq 1$ be an integer. Suppose that*

$$\omega_\mathbb{S}(f, Z) = 0, \quad \dim Z \leq M - q, \quad \dim P \leq 2q - 5.$$

There is a continuous map

$$g : P \longrightarrow (-2, 2)^M \setminus Z \quad \text{such that} \quad \|g(x) - f(x)\|_\infty < \frac{1}{2} \quad (x \in P).$$

Proof. The empty domain or empty arrangement needs no argument. Otherwise the dimension hypotheses imply $q \geq 3$.

Choice of boxes. Subdivide P so finely that $f(\sigma)$ has maximum-norm diameter less than c for each simplex. Thicken its coordinate bounding box by a radius between $2c$ and $3c$, increasing strictly with the dimension of σ . A face box is then contained in the interior of its coface box. Choose the finitely many radii generically so that no endpoint is zero. The resulting boxes B_σ have the properties

$$\text{dist}_\infty(f(\sigma), T \setminus \text{int } B_\sigma) > c, \quad \sup_{x \in \sigma, y \in B_\sigma} \|y - f(x)\|_\infty < \frac{1}{2}. \quad (3.11)$$

They all lie in $(-2, 2)^M$; indeed the latter supremum is less than $4c$. Enlarge them by one sufficiently small uniform amount to obtain B_σ^+ , preserving the coordinate incidences and all inclusions. Choose a common ϵ and then a common collapse radius $\rho < c$ as in [Lemmas 3.3](#) and [3.4](#). Write $V_\sigma = V_\sigma(Z)$ and $T_\sigma = T_\sigma(Z)$.

The stable lift as a map of diagrams. Index covariant diagrams by the nonempty simplices of P , with maps from faces to cofaces. Set

$$\mathcal{D}_\sigma = \Sigma^\infty \sigma_+, \quad \mathcal{B}_\sigma = \Sigma^\infty (B_\sigma)_+, \quad \mathcal{V}_\sigma = \Sigma^\infty (V_\sigma)_+.$$

The maps $f|_\sigma$ define a natural map $\mathcal{D} \rightarrow \mathcal{B}$. Its obstruction to lifting through $\mathcal{V} \rightarrow \mathcal{B}$ is the composite to the cofiber diagram $\mathcal{C}$.

For clarity, the source cell diagram has latching maps

$$\operatorname{colim}_{\tau \subsetneq \sigma} \tau_+ = \partial \sigma_+ \longrightarrow \sigma_+. \quad (3.12)$$

They are cofibrations; at a vertex the latching object is the basepoint. Consequently this is a cofibrant diagram, and compatible mapping out of its suspension spectrum computes the derived mapping spectrum. This fact can also be seen directly: add the simplices in increasing dimension. At each step a compatible map is an extension over σ of the already specified map on $\partial \sigma$. Restriction along [Equation \(3.12\)](#) is a mapping-space fibration after fibrant replacement. Induction therefore makes compatible mapping invariant under objectwise equivalences and preserves the fiber sequences of mapping spectra used here.

The duality in [Lemma 3.4](#) identifies the obstruction with the adjoint of the equality test on $\sigma_+ \wedge T_\sigma$. There is a well-defined based map

$$\sigma_+ \wedge T_\sigma \longrightarrow Q_c(f, Z), \quad [x, z] \longmapsto [x, z]. \quad (3.13)$$

Indeed, if z lies outside $\operatorname{int} B_\sigma^+$, then [Equation \(3.11\)](#) puts (x, z) in the relative set of the global test. For a face $\sigma \subset \tau$, compare the two maps on $\sigma_+ \wedge T_\tau$: including σ in τ or collapsing T_τ to T_σ gives the same based map to $Q_c(f, Z)$. Every additional collapse already lands in the global relative set by [Equation \(3.11\)](#). After adjunction these equalities give the required natural transformation. Pulling back one stable nullhomotopy of $\varphi_\rho(f, Z)$ thus gives a compatible nullhomotopy of the obstruction diagram. This uses the assumed relative point-test vanishing, rather than separately chosen local nullhomotopies.

The category of spectrum diagrams is stable. Mapping out of $\mathcal{D}$ into the objectwise cofiber sequence $\mathcal{V} \rightarrow \mathcal{B} \rightarrow \mathcal{C}$ now gives a stable lift $\mathcal{D} \rightarrow \mathcal{V}$, together with a compatible homotopy of its composite to the map given by f .

Augmentation and desuspension. For an unbased space V , the augmentation $\Sigma^\infty V_+ \rightarrow \mathbb{S}$ sends every point to the nonbasepoint of S^0 . A map of spaces into V therefore has augmentation equal to the unit, whereas an arbitrary stable map need not. Each B_σ is contractible, so its augmentation is an equivalence. After adjunction, augmenting the compatible homotopy in the box diagram gives a homotopy from the stable lift's augmentation to the constant unit map. We retain the lift together with this homotopy by setting

$$E_\sigma = \operatorname{hofib}_1 (\Omega^\infty \Sigma^\infty (V_\sigma)_+ \longrightarrow \Omega^\infty \mathbb{S}), \quad (3.14)$$

where 1 is the unit point. Naturality of the augmentations makes these data a single derived map from the cell diagram σ into the homotopy-fiber diagram E . We keep the lift and its augmentation homotopy together when choosing functorial models.

There is a natural comparison $j_\sigma : V_\sigma \rightarrow E_\sigma$. Its augmentation is literally the unit, so the path in the homotopy fiber can be taken constant. To compute its connectivity, choose a point $v_\sigma \in V_\sigma$. The resulting splitting of the augmentation identifies E_σ with $\Omega^\infty \Sigma^\infty (V_\sigma, v_\sigma)$

and identifies j_σ with stabilization. The choices are used only for this objectwise connectivity calculation; the comparison itself was defined naturally. By [Lemma 3.3](#), V_σ is $(q - 2)$ -connected. Freudenthal therefore makes j_σ at least $(2q - 3)$ -connected [[5](#), Corollary 4.24]. In particular, the weaker bound $2q - 4$ is available with room for the stated dimension hypothesis.

Here is an explicit way to realize the lift without making compatible basepoint choices. Replace j by its natural mapping-path factorization. Objectwise this is

$$W_\sigma = \{(v, \gamma) : v \in V_\sigma, \gamma : [0, 1] \rightarrow E_\sigma, \gamma(0) = j_\sigma(v)\}, \quad p_\sigma(v, \gamma) = \gamma(1).$$

The maps p_σ are fibrations of connectivity at least $2q - 3$. Projection $W_\sigma \rightarrow V_\sigma$ is a natural homotopy equivalence, with section given by constant paths. The cofibrant cell diagram allows the preceding compatible derived map to E to be represented by an actual map of diagrams, using objectwise fibrant models.

Inductively lift this map to W . At a simplex σ , the chosen lifts on proper faces give a map $\partial\sigma \rightarrow W_\sigma$ over the prescribed map $\sigma \rightarrow E_\sigma$. Its extension is possible because the fibers of p_σ are $(2q - 4)$ -connected and $\dim \sigma \leq 2q - 5$. This is the ordinary lifting obstruction for a disk and its boundary. Projection back to V gives an actual compatible family $\sigma \rightarrow V_\sigma$. No dimension of a separate indexing space is added: the extension domains are exactly the simplices in [Equation \(3.12\)](#).

The family glues to a continuous map $g : P \rightarrow (-2, 2)^M \setminus Z$. Its image on σ lies in B_σ , so [Equation \(3.11\)](#) gives the claimed displacement bound. $\square$

3.4 Restoring cube boundaries

The next elementary step is stated for compact inputs, since no polyhedral hypothesis is needed. Its preservation clause is what allows successive applications of [Lemma 3.5](#).

Lemma 3.6 (Boundary restoration). *Let K be compact, let $0 \leq s < n$ be an integer, and suppose that $f : K \rightarrow C(t, n)$ and $g : K \rightarrow (-2, 2)^M \setminus Z_s(t, n)$ are continuous and satisfy $\|g(x) - f(x)\|_\infty < 1/2$ for every x . There is a continuous map $G : K \rightarrow C(t, n)$ such that at least $s + 1$ of its slots belong to $\partial[-1, 1]^t$ at every point, and*

$$f_j(x) \in \partial[-1, 1]^t \implies G_j(x) = f_j(x) \quad (1 \leq j \leq n).$$

Proof. If K is empty the assertion is immediate. Otherwise, choose a continuous $\chi : [0, 1] \rightarrow [0, 1]$ which is zero on $[0, 3/4]$ and equals one at 1. Define slotwise

$$g'_j(x) = (1 - \chi(\|f_j(x)\|_\infty))g_j(x) + \chi(\|f_j(x)\|_\infty)f_j(x).$$

Whenever the cutoff is nonzero, one scalar coordinate of $f_j(x)$ has magnitude greater than $3/4$. The corresponding coordinate of $g_j(x)$ has the same sign and magnitude greater than $1/4$. The convex interpolation consequently cannot turn that slot into zero. All other slots are unchanged. Hence g' still has at least $s + 1$ nonzero slots everywhere, and it already agrees with f at each original boundary slot.

The $(s + 1)$ -st largest of the finitely many continuous functions $\|g'_j(x)\|_\infty$ is continuous and positive on K . By compactness choose $0 < \delta < 1$ smaller than its minimum. Set

$$G_j(x) = \frac{g'_j(x)}{\max\{\delta, \|g'_j(x)\|_\infty\}}.$$

This is continuous even at zero, belongs to the unit cube, and sends every slot of radius at least δ to the boundary. It fixes every slot already of radius one. Both conclusions follow. $\square$

4 Boundary compression by padding

We now turn a vanishing point test into a uniform loss of interior cube slots. The parameter polyhedron can have arbitrarily large dimension. The device that makes this possible is to retain the previous *output*, whose dimension is bounded by its ambient cube, when a new batch of parameters is introduced. The resulting theorem applies to every finite polyhedron, without an upper bound on its dimension. In [Section 5](#), the torus-bundle criterion and exact factorization will extend this construction to arbitrary compact metrizable parameter spaces.

We use the cubes $C(t, n)$ and $T(t, n)$, the sparse sets $Z_s(t, n)$, and the point tests of [Definition 3.1](#). Thus $Z_s(t, n)$ allows at most $\lfloor s \rfloor$ nonzero slots, and the fixed relative threshold is $c = 1/32$. For convenience put $Z_{-1}(t, n) = \emptyset$. A boundary slot means a slot in $\partial[-1, 1]^t$; in particular, it is a nonzero slot. The notation $f^{\times u}$ means the product of u copies of f , with the slots ordered first by copy and then by their original index.

4.1 A stable test for a large batch

Lemma 4.1 (Batch vanishing). *Let P be a finite polyhedron, let $f : P \rightarrow C(t, n)$ be PL, and let $0 < a < 1/16$. Suppose*

$$\omega_{MU}(f, Z_{an}(t, n)) = 0.$$

For every sufficiently large integer ℓ , with $b = n\ell$ and $H = \lfloor ab/4 \rfloor$, one has

$$\omega_S(f^{\times \ell}, Z_H(t, b)) = 0.$$

Proof. Take the finite pointed pair

$$E = Q_c(f, T(t, n)), \quad G = Q_c(f, Z_{an}(t, n)) \subset E.$$

A common polyhedral subdivision of the domain, sparse subsets, and relative subsets makes this a pointed CW pair. The origin collapse gives a based map $v : E \rightarrow S^{tn}$ whose restriction to G is zero after applying the unit into MU . By [Theorem 1.1](#), for all sufficiently large ℓ the map $v^{\wedge \ell}$ is stably null on the union of smash products having at least $\lceil \ell/2 \rceil$ factors in G .

Split a parameter in $T(t, b)$ into ℓ consecutive groups of n slots. Every group outside $Z_{an}(t, n)$ uses strictly more than an nonzero slots. Consequently a parameter with at most $H \leq an\ell/4$ nonzero slots has fewer than $\ell/4$ such groups. In particular, at least $\lceil \ell/2 \rceil$ groups belong to $Z_{an}(t, n)$. By [Lemma 3.2](#), the quotient for the batch test maps into the preceding union and the batch test is the restriction of $v^{\wedge \ell}$. Pulling back its single stable nullhomotopy proves the assertion. $\square$

4.2 Padding by previous outputs

Batch vanishing alone does not put the growing product domain within the stable avoidance range. The next argument replaces that domain at each stage by the previous output and one new batch. Only the new batch contributes the original parameter dimension.

Theorem 4.2 (Boundary compression). *Let P be a finite polyhedron, let $f : P \rightarrow C(t, n)$ be PL, and fix $0 < a < 1/16$. If*

$$\omega_{MU}(f, Z_{an}(t, n)) = 0,$$

then for arbitrarily large positive integers u there is a continuous map

$$G : P^u \longrightarrow C(t, un)$$

with the following properties. At every point of P^u , at least $(a/8)un$ output slots are on their cube boundaries. Moreover, if any slot of $f^{\times u}$ is on its boundary, the corresponding slot of G equals that original slot exactly.

Proof. The assertion is immediate for an empty P , so assume P is nonempty and write $d_P = \dim P$. Fix ℓ as in Lemma 4.1, large enough that $ab \geq 16$, where

$$b = n\ell, \quad H = \lfloor ab/4 \rfloor.$$

In particular $H \geq 1$. Fix a positive integer B of padding batches so large that

$$tb(B+1) \geq \max\{20, 2\ell d_P\}. \quad (4.1)$$

The initial B batches will have output $f^{\times \ell B}$ and no required boundary count.

For $N \geq 1$ and $0 \leq s \leq N$, let

$$Q_s(t, N) = \{y \in C(t, N) : \text{at least } s \text{ slots of } y \text{ lie in } \partial[-1, 1]^t\}.$$

This is a finite polyhedron: it is a finite union of products in which specified factors are cube boundaries. Its dimension is at most tN .

Suppose $j-1$ ordinary batches have already been added. Put

$$N_{j-1} = b(B+j-1), \quad s = (j-1)H.$$

The previous output lies in $Q_s(t, N_{j-1})$. To add the next batch we solve a new problem on the finite polyhedron

$$W_j = Q_s(t, N_{j-1}) \times P^\ell, \quad F_j(y, x) = (y, f^{\times \ell}(x)). \quad (4.2)$$

After a product subdivision F_j is PL. We first show that its sphere point test vanishes on

$$Z_{jH-1}(t, b(B+j)).$$

Write the sparse parameter as (z_o, z_n) , for the old and new slots. On this parameter set the two closed conditions

$$\begin{aligned} A_j : \quad & z_o \in Z_{s-1}(t, N_{j-1}), \\ B_j : \quad & z_n \in Z_{H-1}(t, b) \end{aligned}$$

cover the whole set: failure of both would give at least $s+H=jH$ nonzero slots. If $s=0$, the first set is empty. On A_j , one of the at least s boundary slots of y has zero old parameter, and therefore

$$\|y - z_o\|_\infty \geq 1. \quad (4.3)$$

Thus the old relative quotient is at its basepoint there.

Here is the precise gluing of the relative maps. Set

$$E_o = Q_c(\text{id}_{Q_s(t, N_{j-1})}, T(t, N_{j-1})), \quad G_n = Q_c(f^{\times \ell}, Z_{H-1}(t, b)).$$

On the part satisfying B_j send $((y, x), (z_o, z_n))$ to $[y, z_o] \wedge [x, z_n]$ in $E_o \wedge G_n$. On the part satisfying A_j send it to the basepoint. These maps agree on their intersection by Equation (4.3), and the closed pasting lemma gives a continuous map on the full domain. It is constant on the relative subset $\|F_j - (z_o, z_n)\|_\infty \geq c$: on B_j at least one smash factor is then the basepoint. It therefore descends to a based map

$$Q_c(F_j, Z_{jH-1}(t, b(B+j))) \longrightarrow E_o \wedge G_n. \quad (4.4)$$

Composing with the smash of the two point tests gives exactly the point test of F_j , by Lemma 3.2; on A_j both descriptions are constant by Equation (4.3). The new factor is stably null, since its parameter set is contained in the one in Lemma 4.1. Its smash with the old factor is therefore stably null. This proves vanishing on the entire sparse union by one relative factorization.

We check the avoidance range before carrying out the induction. The current ambient dimension and the codimension of the excluded sparse set are

$$D_j = tb(B+j), \quad q_j = D_j - t(jH-1). \quad (4.5)$$

Indeed $0 \leq jH-1 < b(B+j)$, so that sparse set has dimension $t(jH-1)$. Since $H \leq ab/4$ and $a < 1/16$,

$$t(jH-1) < \frac{a}{4}D_j < \frac{1}{8}D_j, \quad q_j > \frac{7}{8}D_j.$$

On the other hand, Equations (4.1) and (4.2) give

$$\dim W_j \leq tb(B+j-1) + \ell d_P < \frac{3}{2}D_j.$$

For all $j \geq 1$ we have $D_j \geq 20$, and hence

$$2q_j - 5 > \frac{7}{4}D_j - 5 \geq \frac{3}{2}D_j > \dim W_j. \quad (4.6)$$

Thus Lemma 3.5 applies to the null point test just constructed. It gives an output avoiding Z_{jH-1} within distance $1/2$ of F_j . By Lemma 3.6, this output can be changed to a continuous map

$$g_j : W_j \longrightarrow Q_{jH}(t, b(B+j))$$

that fixes every boundary slot of F_j .

Starting with $G_0 = f^{\times \ell B}$, define recursively

$$G_j(\mathbf{x}, x) = g_j(G_{j-1}(\mathbf{x}), x), \quad \mathbf{x} \in P^{\ell(B+j-1)}, \quad x \in P^\ell.$$

This is well-defined because G_{j-1} takes values in the first factor of W_j . It has at least jH boundary slots. Boundary slots in the new batch are fixed when that batch enters; boundary slots already present in the old output are fixed at every later step. The padding output initially equals the original input. Induction therefore proves the required agreement with every original boundary slot.

Finally the floor in H and the padding loss have explicit bounds:

$$\frac{H}{b} \geq \frac{a}{4} - \frac{1}{b} \geq \frac{3a}{16}, \quad \frac{jH}{b(B+j)} \geq \frac{2}{3} \frac{3a}{16} = \frac{a}{8} \quad (j \geq 2B).$$

Taking $u = \ell(B+j)$ for these arbitrarily large j proves the theorem. The dimension estimate was applied only to W_j ; no bound on the dimension of the full original parameter product is needed. $\square$

5 Torus bundles and compact parameter spaces

The remaining task is to verify the MU -vanishing hypothesis from a bundle embedding and then remove the polyhedral assumption on the input. We first construct the torus data used in [Notation 1.2](#). The Chern-class calculation takes place on a tube, so that its vanishing controls the relative point class rather than only its image in absolute cohomology.

5.1 A framed torus detecting the point class

Fix $p \geq 1$ and set $t = 2p + 1$. We describe once and for all the geometric bundle data used in the criterion.

Lemma 5.1 (Torus and line bundles). *There is a smooth embedding*

$$L \cong (\mathbb{T}^2)^p \subset (-c, c)^{2p+1}$$

with a framed real normal line bundle. There are complex line bundles $\lambda_1, \dots, \lambda_p$ on L , each pulled back from its indicated two-torus factor, such that

$$H = \bigoplus_{i=1}^p \lambda_i \subset L \times \mathbb{C}^{2p}, \quad \left\langle \prod_{i=1}^p c_1(\lambda_i), [L] \right\rangle = 1$$

for a choice of orientation on L .

Proof. Begin with a circle embedded in $\mathbb{R}^2$ with its outward normal framing. If $\mathbb{T}^k$ has been embedded in $\mathbb{R}^{k+1}$ with a framed normal line, its inclusion into $\mathbb{R}^{k+2}$ has a framed normal two-plane bundle. The boundary of a sufficiently small tubular disk bundle is consequently a smooth embedded hypersurface $\mathbb{T}^k \times S^1$ in $\mathbb{R}^{k+2}$. Its outward normal is a framing. Iteration produces $\mathbb{T}^{2p}$ in $\mathbb{R}^{2p+1}$, and translation and rescaling put it in the specified cube.

Choose a smooth degree-one map $\mathbb{T}^2 \rightarrow S^2 = \mathbb{CP}^1$; one may collapse the complement of an oriented disk and then take a smooth approximation. The orthogonal complement of the tautological line over $\mathbb{CP}^1$ is a line subbundle of $\mathbb{C}^2$ with first Chern number one. Pull it back to $\mathbb{T}^2$, and then to each indicated factor of L . Taking their direct sum gives the stated embedding. The evaluation follows from the product orientation and the Chern number on each factor. $\square$

Retain the slot bundle H_n and the locus $\mathcal{T}(f, Z)$ of [Notation 1.2](#). The next lemma is valid for every real a ; the range $0 < a < 1/16$ is needed later for compression. Its complementary-bundle argument is related to the ordinary Chern-class obstruction in [6, proof of Theorem 3.3 in the author preprint]. We supply the additional MU square-zero calculation and supported Thom-class argument here.

Lemma 5.2 (The cobordism criterion). *Fix positive integers p, n and a real number a . Let P be a finite polyhedron, let $f : P \rightarrow C(2p + 1, n)$ be PL , and let $Z = Z_{an}(2p + 1, n)$. If the slot bundle on $\mathcal{T}(f, Z)$ embeds into a trivial complex bundle of rank $0 \leq K < 2pn$, then*

$$\omega_{MU}(f, Z) = 0.$$

Proof. Put $M = (2p + 1)n$, $r = pn$, $D = P \times Z$, and $d = d_f$. The rank bound will force the pulled-back top MU Chern class of H_n to vanish. We need this vanishing on the preimage of a neighborhood of L^n , where multiplication by the normal Thom class will recover the relative point test.

Fix the normal framing of L^n , of real rank n , obtained by taking products in [Lemma 5.1](#). A tubular neighborhood retraction extends H_n and its orthogonal projection $\Pi : L^n \rightarrow M_{2r}(\mathbb{C})$ to a neighborhood of L^n .

We first extend the embedding to the pullback of a whole tube. On the compact closed subset $\mathcal{T}(f, Z) \subset D$, compose the assumed embedding with the source projection $\Pi \circ d$. This represents it by a continuous K -by- $2r$ matrix. Extend the real and imaginary parts of its entries to D by the Tietze extension theorem [[14](#), Theorem 1.10.5]. On the original test locus its restriction to the source fibers has a uniform positive lower bound on unit vectors, by compactness. This lower bound persists on the pullback of a sufficiently thin tube: otherwise a sequence of unit source vectors over points tending to the central locus would converge to a vector contradicting injectivity there. If the central locus is empty, compactness instead gives a tube with empty pullback. We may thus choose an open tube U with compact closure in $(-c, c)^M$ and retraction $\rho : U \rightarrow L^n$ such that on $d^{-1}(U)$ the pulled-back bundle $d^*\rho^*H_n$ embeds into $\mathbb{C}^K$.

We use the complex orientation of MU and its Chern and Thom classes, with the direct-sum Whitney formula [[9](#), Lecture 5, Proposition 6 and the following discussion; Lecture 6, Remark 7]. List the r line summands of H_n and write

$$x_i = c_1^{MU}(\lambda_i) \in MU^2(L^n), \quad 1 \leq i \leq r, \quad \alpha = \prod_{i=1}^r x_i.$$

Here the indices include both the slot and the two-torus factor. Each x_i is pulled back from a two-dimensional torus. Since MU is connective, $MU^4(\mathbb{T}^2) = 0$: filtering that torus by its cells reduces this assertion to $MU^q(\text{pt}) = 0$ for $q > 0$. Consequently $x_i^2 = 0$.

Pull these classes to $d^{-1}(U)$. If this space is nonempty, the image of the embedded rank- r bundle has an orthogonal complement F in $d^{-1}(U) \times \mathbb{C}^K$. Thus F has rank $K - r < r$. With a formal variable v recording Chern index, the Whitney formula and the square-zero identities give

$$c_v^{MU}(F) = \prod_{i=1}^r (1 + d^*\rho^*x_i v)^{-1} = \prod_{i=1}^r (1 - d^*\rho^*x_i v). \quad (5.1)$$

The coefficient of v^r on the left is zero by rank. The coefficient on the right is $(-1)^r d^*\rho^*\alpha$. Hence

$$d^*\rho^*\alpha = 0 \quad \text{in } MU^{2r}(d^{-1}(U)). \quad (5.2)$$

The same assertion is automatic if the tube pullback is empty. This uses only the direct-sum Chern formula; no formula for tensor products of lines is involved.

For the support pairs below, relative MU means reduced cohomology of homotopy cofibers. The collared ambient tube pairs and the final polyhedral pair also admit the literal quotient descriptions used here. Choose a closed smaller tubular disk bundle V with $L^n \subset \text{int } V$ and $V \subset U$. The normal framing gives a Thom class

$$\tau \in MU^n(U, U \setminus \text{int } V).$$

Indeed the relative quotient is the Thom space of a trivial real rank- n bundle, whose Thom class is the n -fold suspension of the unit. This construction works for odd as well as even n [[9](#), Lecture 5, Example 1]. The relative product

$$\eta = (\rho^*\alpha)\tau \in MU^M(U, U \setminus \text{int } V)$$

extends by excision to a class supported in V in $MU^M(\mathbb{R}^M, \mathbb{R}^M \setminus \text{int } V)$. Its image in

$$MU^M(\mathbb{R}^M, \mathbb{R}^M \setminus (-c, c)^M) \cong \widetilde{MU}^M(S^M) \cong \pi_0 MU = \mathbb{Z} \quad (5.3)$$

is a generator, up to the sign of the normal orientation. To check the integer, apply the ordinary orientation map $MU \rightarrow H\mathbb{Z}$. It induces an isomorphism on the group in Equation (5.3). The ordinary image of η has evaluation

$$\left\langle \prod_{i=1}^r c_1(\lambda_i), [L^n] \right\rangle = 1$$

by Lemma 5.1 and the product orientation. Thus the image of η is exactly the MU origin class up to sign.

We give the pullback with its relative sets to retain this supported information. On $d^{-1}(U)$ the pullback of η vanishes, because its absolute factor is zero by Equation (5.2). Excision for the smaller closed tube therefore gives zero in

$$MU^M(D, D \setminus d^{-1}(\text{int } V)).$$

This excision is valid after pullback: the closure of $D \setminus d^{-1}(U)$ lies in the interior of $D \setminus d^{-1}(\text{int } V)$, since $V \subset U$ and V is compact. In particular the excision isomorphisms commute with d and with the relative products used above. As $V \subset (-c, c)^M$, the relative sets satisfy

$$A_c(f, Z) \subset D \setminus d^{-1}(\text{int } V).$$

The induced map on relative cohomology sends this zero class to

$$MU^M(D, \{(x, z) : \|f(x) - z\|_\infty \geq c\}). \quad (5.4)$$

The image is, up to sign, the pullback of the origin class in Equation (5.3), so it is the MU point test and is zero. Finally the pair in Equation (5.4) is a finite polyhedral pair: f is PL, Z is a finite union of coordinate cubes, and the relative condition is a finite union of linear inequalities after subdivision. Its relative cohomology is therefore the reduced cohomology of the exact quotient in Definition 3.1. This proves the required vanishing. $\square$

5.2 Compact parameter spaces and exact factorization

Lemma 5.3 (Finite parameter reduction). *Fix positive integers p, n and a real number a . Let Y be compact metrizable, let $f : Y \rightarrow C(2p+1, n)$ be continuous, and put $Z = Z_{an}(2p+1, n)$. Suppose the slot bundle on $\mathcal{T}(f, Z)$ embeds into a trivial complex bundle of nonnegative integer rank K . There are a finite polyhedron P , a continuous map $j : Y \rightarrow P$, and a PL map $f_P : P \rightarrow C(2p+1, n)$ such that*

$$f = f_P \circ j,$$

and the slot bundle on $\mathcal{T}(f_P, Z)$ also embeds into a trivial bundle of rank K .

Proof. If Y is empty, take P empty and the unique maps. Otherwise put $M = (2p+1)n$ and $r = pn$. Represent the embedding on the compact test locus by a K -by- $2r$ matrix A , by first projecting onto the fixed source bundle. Extend its entries to $Y \times Z$ by the Tietze theorem [14, Theorem 1.10.5]. Choose $\delta > 0$ such that on the central test locus

$$\|A(y, z)v\| \geq 4\delta\|v\| \quad \text{for } v \in (H_n)_{f(y)-z}. \quad (5.5)$$

Compactness of its unit sphere bundle supplies this number. If the locus is empty the condition is vacuous and any $\delta > 0$ may be used.

The real algebra on $Y \times Z$ generated by continuous real functions on Y and the Euclidean coordinates of z contains the constants and separates points. By the real Stone–Weierstrass

theorem [14, Theorem 1.10.18], the real and imaginary entries of A can be uniformly approximated by elements of this algebra. Only finitely many functions on Y occur in these approximants. Rescale them into $[-1, 1]$ and denote them by $\varphi_1, \dots, \varphi_q$. Include every coordinate of f explicitly, by setting

$$j_0(y) = (f(y), \varphi_1(y), \dots, \varphi_q(y)) \in W = [-1, 1]^{M+q}.$$

The approximating matrix has the form $B(j_0(y), z)$, where B is a matrix of polynomials on $W \times Z$. Choose the approximation so that its operator-norm error is less than δ everywhere on $Y \times Z$. By Equation (5.5),

$$\|B(j_0(y), z)v\| \geq 3\delta\|v\| \quad \text{on the original slot bundle.} \quad (5.6)$$

Let $C = j_0(Y)$ and let $f_0 : W \rightarrow [-1, 1]^M$ be projection onto the first M coordinates. The bundle over a point (u, z) with $f_0(u) - z \in L^n$ depends only on $f_0(u) - z$. Consequently Equation (5.6) gives the same bound at every such point of $C \times Z$, regardless of the choice of a preimage of u in Y .

For completeness, consider the following compact bad set:

$$\mathcal{B} = \{(u, z, v) : u \in W, z \in Z, \|v\| = 1, f_0(u) - z \in L^n, \\ \Pi(f_0(u) - z)v = v, \|B(u, z)v\| \leq 2\delta\}.$$

Here Π is the continuous orthogonal projection onto H_n . The projection of $\mathcal{B}$ onto W is compact and disjoint from C , by Equation (5.6). Its complement is therefore a relative open neighborhood of C in W . Subdivide W into sufficiently small closed cubes and take the union P of all cubes meeting C . Finiteness and the positive distance from C to the bad projection ensure that P is contained in this neighborhood. If the bad projection is empty, any such subdivision suffices. This construction is relative to W ; it does not extend the first M coordinates beyond the unit cube.

Regard j_0 as a map $j : Y \rightarrow P$ and put $f_P = f_0|_P$. The map f_P is coordinate-linear and hence PL. On $\mathcal{T}(f_P, Z)$, the restriction of $B(u, z)$ to the slot bundle is bounded below by 2δ on unit vectors, and thus is the required continuous bundle embedding. Finally $f_P \circ j = f$ by the choice of the first M coordinates. $\square$

5.3 Completion of the compact-input theorem

The factorization just proved retains the cube map exactly, while moving the auxiliary embedding data to a finite polyhedron. We can therefore apply boundary compression there and pull it back without losing its pointwise preservation clause.

Proof of Theorem 1.3. Apply Lemma 5.3 to obtain $j : Y \rightarrow P$ and f_P with the same bundle embedding and $f = f_P \circ j$. By Lemma 5.2 the MU point test of f_P vanishes. Theorem 4.2 gives the stated compression maps on P^u for arbitrarily large u . Composing with $j^{\times u}$ gives the maps on Y^u . The factorization of f is exact, so the boundary agreement also pulls back exactly. $\square$

Remark 5.4 (The strict rank threshold). The bound $K < 2pn$ in Theorem 1.3 cannot in general be replaced by $K \leq 2pn$. Take $n = 1$, let $Y = C(2p+1, 1)$, and let f be the identity. For every $0 < a < 1/16$ the sparse set is $Z_a(2p+1, 1) = \{0\}$, so the torus locus is L and the canonical inclusion $H \subset L \times \mathbb{C}^{2p}$ supplies an embedding at the endpoint $K = 2p$.

Suppose that, for some $u \geq 1$, a continuous map $G : B \rightarrow B$, where $B = C(2p+1, u)$, has at least one boundary slot everywhere and fixes every original boundary slot. Its image

lies in the total boundary ∂B . For $x \in \partial B$, at least one boundary slot of x remains fixed throughout the straight-line homotopy $(1-s)x + sG(x)$, $0 \leq s \leq 1$. This homotopy therefore stays in ∂B , showing that $G|_{\partial B}$ has degree one. But $G : B \rightarrow \partial B$ extends this restriction over the cube, forcing its degree to be zero. This contradiction rules out such a map for every u . In particular, the positive bound $(a/8)u$ in [Theorem 1.3](#), which requires at least one boundary slot, is impossible at this endpoint.

References

- [1] M. F. Atiyah. Thom complexes. *Proceedings of the London Mathematical Society*, 11:291–310, 1961. <https://doi.org/10.1112/plms/s3-11.1.291>.
- [2] Maria Basterra and Michael A. Mandell. The multiplication on BP. *Journal of Topology*, 6(2):285–310, 2013. <https://doi.org/10.1112/jtopol/jts032>.
- [3] Robert Burklund, Jeremy Hahn, and Andrew Senger. On the boundaries of highly connected, almost closed manifolds. *Acta Mathematica*, 231(2):205–344, 2023. Appendix B, *Vanishing curves in Adams spectral sequences*, by Robert Burklund. <https://doi.org/10.4310/ACTA.2023.v231.n2.a1>.
- [4] Ethan S. Devinatz, Michael J. Hopkins, and Jeffrey H. Smith. Nilpotence and stable homotopy theory I. *Annals of Mathematics*, 128(2):207–241, 1988.
- [5] Allen Hatcher. *Algebraic Topology*. Cambridge University Press, Cambridge, 2002. <https://pi.math.cornell.edu/~hatcher/AT/AT.pdf>.
- [6] Ilan Hirshberg and N. Christopher Phillips. Radius of comparison and mean cohomological independence dimension. *Advances in Mathematics*, 406:108563, 2022. <https://doi.org/10.1016/j.aim.2022.108563>. Author preprint: <https://arxiv.org/abs/2009.13045>.
- [7] John R. Klein and E. Bruce Williams. Homotopical intersection theory I. *Geometry & Topology*, 11(2):939–977, 2007. <https://doi.org/10.2140/gt.2007.11.939>.
- [8] Lawrence L. Larmore. Twisted cohomology theories and the single obstruction to lifting. *Pacific Journal of Mathematics*, 41(3):755–769, 1972. <https://msp.org/pjm/1972/41-3/pjm-v41-n3-p14-s.pdf>.
- [9] Jacob Lurie. Chromatic homotopy theory. Course notes, Harvard University, 2010. Lectures 5–7. <https://www.math.ias.edu/~lurie/252x.html>.
- [10] Akhil Mathew. Examples of descent up to nilpotence. In Jon F. Carlson, Srikanth B. Iyengar, and Julia Pevtsova, editors, *Geometric and Topological Aspects of the Representation Theory of Finite Groups*, volume 242 of *Springer Proceedings in Mathematics & Statistics*, pages 269–311. Springer, Cham, 2018. https://doi.org/10.1007/978-3-319-94033-5_11. Author preprint: <https://arxiv.org/abs/1701.01528>.
- [11] Akhil Mathew, Niko Naumann, and Justin Noel. Nilpotence and descent in equivariant stable homotopy theory. *Advances in Mathematics*, 305:994–1084, 2017.
- [12] OpenAI. Radius of comparison equals half the mean dimension. OpenAI Math Release preprint OAI:Radius-of-comparison-equals-half-the-mean-dimension-September-25-2026, 2026.
- [13] Daniel Quillen. Elementary proofs of some results of cobordism theory using Steenrod operations. *Advances in Mathematics*, 7(1):29–56, 1971. [https://doi.org/10.1016/0001-8708\(71\)90041-7](https://doi.org/10.1016/0001-8708(71)90041-7).
- [14] Terence Tao. *An Epsilon of Room, I: Real Analysis: Pages from Year Three of a Mathematical Blog*, volume 117 of *Graduate Studies in Mathematics*. American Mathematical Society, 2010. Author’s preliminary version: <https://terrytao.wordpress.com/wp-content/uploads/2012/12/gsm-117-tao3-epsilon1.pdf>.
- [15] René Thom. Quelques propriétés globales des variétés différentiables. *Commentarii Mathematici Helvetici*, 28:17–86, 1954. <https://doi.org/10.1007/BF02566923>.