arXiv:2607.23423v1 [math.GR] 26 Jul 2026

AN INFINITE FAMILY OF COUNTEREXAMPLES TO THE POLYCIRCULANT CONJECTURE

SAUL D. FREEDMAN AND MELISSA LEE

ABSTRACT. We disprove the Polycirculant Conjecture, which states that every transitive 2-closed permutation group is non-elusive, i.e. contains a derangement of prime order. In fact, we prove a stronger result, answering a long-standing question of Marušič and Jordan: there exists a vertex-transitive graph admitting no semiregular automorphism. To do so, we employ recently developed methods of Chen et al. for constructing elusive groups via non-split extensions, allowing us to construct an elusive group $7^{6}{\cdot}\mathrm{PSU}_3(3)$ of degree $16{,}464$. We show that this group is the full automorphism group of seven of its orbital graphs and hence is 2-closed. Our example extends to infinitely many counterexamples of the Polycirculant Conjecture, and infinitely many vertex-transitive graphs admitting no semiregular automorphism.

1. INTRODUCTION

A *derangement* is a finite permutation that has no fixed points. A classical theorem of Jordan asserts that every finite transitive group contains a derangement. Using the classification of finite simple groups, Fein, Kantor and Schacher [8] showed that Jordan's theorem may be strengthened to require the existence of a derangement of prime power order. However, perhaps surprisingly, it is false that every finite transitive permutation group must contain a derangement of prime order. For instance, the Mathieu group M_{11} acting on 12 points has derangements only of orders 4 and 8. Due to the seeming rarity of finite transitive permutation groups containing no derangements of prime order, such a group is termed *elusive*.

Elusive groups have attracted significant interest (see for example [4, 5, 11]), in part due to their connection to the well-known Polycirculant Conjecture, posed by Klin [6, Problem BCC15.12] in 1997. Recall that the 2-*closure* of a permutation group $G \leqslant \mathrm{Sym}(\Omega)$ is the largest subgroup of $\mathrm{Sym}(\Omega)$ with the same orbits on $\Omega \times \Omega$ as G, and that G is 2-*closed* if it is its own 2-closure.

Conjecture 1.1 (The Polycirculant Conjecture)**.** *No elusive group is* 2*-closed.*

This conjecture extends the earlier Semiregularity Problem, formulated by Marušič [17] and Jordan [16] in the 1980s. A non-trivial permutation is called *semiregular* if all of its cycles are the same length (so in particular, it has no trivial cycles).

Date: July 28, 2026.

Key words and phrases. polycirculant conjecture, elusive group, derangement, vertex-transitive graph, semiregular automorphism.

The authors would like to thank the Isaac Newton Institute for Mathematical Sciences, Cambridge, for support and hospitality during the programme *Algebraic groups, geometry, invariants and related topics*, where work on this paper was undertaken. This work was supported by EPSRC grant EP/Z000580/1.

Problem 1.2 (The Semiregularity Problem)**.** Does there exist a vertex-transitive directed graph admitting no semiregular automorphism?

This problem includes the case where the arc set of the directed graph is symmetric, so that the graph can be considered undirected. Observe that each derangement of prime order is semiregular, and each semiregular permutation has a power equal to a derangement of prime order. Hence the Semiregularity Problem asks if there is a vertex-transitive directed graph whose full automorphism group is elusive. Note that, by definition, the automorphism group of any finite directed graph is 2-closed. Hence a positive answer to the Semiregularity Problem would imply that the Polycirculant Conjecture is false. On the other hand, there exist 2-closed groups that are not the full automorphism groups of any directed graph, such as the subgroup $C_2 \times C_2$ of S_4. Therefore, a counterexample to the Polycirculant Conjecture might not provide an answer to the Semiregularity Problem.

The Polycirculant Conjecture and Semiregularity Problem have received considerable attention; see [1] for a survey of results. Perhaps most notably, Giudici [10] determined all elusive permutation groups with at least one transitive minimal normal subgroup (hence all elusive quasiprimitive groups) and showed that each satisfies the Polycirculant Conjecture. A few years later, Giudici and Xu [12] proved that the conjecture is satisfied for every group that is *biquasiprimitive*, i.e. transitive with each minimal normal subgroup having at most two orbits. Hence in any counterexample to the conjecture, each minimal normal subgroup must be intransitive, and at least one must have at least three orbits.

Recently, Chen et al. [7] described a novel method of constructing elusive groups via non-split extensions, and used the method to construct the first known examples of elusive groups with odd and twice-odd degree. Their examples again satisfy the Polycirculant Conjecture.

In this paper, we further the methods of Chen et al. to construct the first known groups that are counterexamples to the Polycirculant Conjecture, and the first known graphs that are positive answers to the Semiregularity Problem. In fact, we construct infinitely many such groups and graphs.

To state our main results, we require the following notation. We will write $N{\cdot}G$ to denote a non-split extension of a group N by a group G, with $N : G$ denoting a split extension of N by G (not to be confused with the index $|G : H|$ in G of a subgroup H, nor the set $[G : H]$ of right cosets of H in G). Additionally, for a prime p and a positive integer a, we write p^a to denote the elementary abelian group of order p^a. Recall also that an *orbital* O of a permutation group G on a set Ω is an orbit of G on $\Omega \times \Omega$, and that the corresponding *orbital graph* Γ_O is the directed graph with vertex set Ω and arc set O. Clearly, each element $g \in G$ is an automorphism of Γ_O, with $(\alpha, \beta)^g = (\alpha^g, \beta^g)$ for each $(\alpha, \beta) \in O$, and if G is transitive, then Γ_O is vertex-transitive. If O is *self-paired*, i.e. if $(\beta, \alpha) \in O$ for all $\alpha, \beta \in \Omega$ with $(\alpha, \beta) \in O$, then we can consider the self-paired orbital graph Γ_O as an undirected graph.

Theorem 1.3. *There is an elusive permutation group X of degree $16{,}464 = 2^4.3.7^3$ isomorphic to $7^6{\cdot}\mathrm{PSU}_3(3)$, with point stabiliser of shape $7^4 : (C_3 \times S_3)$, such that X has (at least) seven pairwise non-isomorphic, connected, self-paired orbital graphs Γ_O of valency 63, with $\mathrm{Aut}(\Gamma_O) = X$ for each Γ_O. Therefore, each of these vertex-transitive graphs of order $16{,}464$ admits no semiregular automorphism.*

This theorem, together with the elementary result Proposition 2.4 on direct products and wreath products of 2-closed groups and elusive groups, immediately yields the following corollary.

Corollary 1.4. *The elusive permutation group X is 2-closed. Moreover, for each integer $k \geqslant 1$, any group constructed from k copies of X via direct products and/or wreath products is isomorphic to a 2-closed elusive group of degree $16{,}464^k$. These groups are therefore all counterexamples to the Polycirculant Conjecture.*

As discussed in Section 3, this corollary also yields a positive answer to [7, Problem 5.2] on counterexamples to the Polycirculant Conjecture constructed via non-split extensions with quasiprimitive quotients.

Remark 1.5. The group X in fact has a second faithful transitive elusive action of degree $16{,}464$ (the point stabiliser in the action from Theorem 1.3 has exactly 16 normal subgroups, while the point stabiliser in the second action, which also has shape $7^4 : (C_3 \times S_3)$, has exactly nine normal subgroups). However, in this second action, the hypotheses of Theorem 5.3 in [5] apply to the socle 7^6 of X. Thus this theorem implies that the 2-closure of this socle, which lies in the 2-closure of X, contains a derangement of prime order. Therefore, this action of X satisfies the Polycirculant Conjecture. Furthermore, writing $\mathrm{P\Gamma U}_3(3)$ to denote the automorphism group $\mathrm{PSU}_3(3) : 2$ of $\mathrm{PSU}_3(3)$, there is a unique elusive permutation group U of degree $16{,}464$ isomorphic to $7^6{\cdot}\mathrm{P\Gamma U}_3(3)$ (with $\mathrm{P\Gamma U}_3(3)$ acting irreducibly on 7^6). This group contains a copy of X in its second elusive action, and so U again satisfies the Polycirculant Conjecture.

We also use a graph product construction to extend the graphs from Theorem 1.3 to an infinite family of graphs that give a positive answer to the Semiregularity Problem.

Corollary 1.6. *There exists an infinite family of graphs of orders $16{,}464^k$ for each $k \geqslant 1$ that are vertex-transitive and do not admit semiregular automorphisms.*

As mentioned above, any counterexample to the Polycirculant Conjecture must have each minimal normal subgroup intransitive, with at least one having at least three orbits. Our method of construction of the counterexample X to the conjecture involves the fact that the elementary abelian normal subgroup $N = 7^6$ of X is an irreducible $\mathrm{PSU}_3(3)$-module. Since $\mathrm{PSU}_3(3)$ is simple and X is non-split, this implies that N is the unique minimal normal subgroup of X. We note that N has 336 orbits, and that X has order $711{,}541{,}152 = 2^5.3^3.7^7$ (and degree $16{,}464 = 2^4.3.7^3$, as mentioned above). Additionally, the graphs Γ_O from Theorem 1.3 each have valency 63. Recall also that there exist 2-closed groups that are not the full automorphism groups of any directed graphs. The following open problems therefore naturally arise. Given a permutation group G, we write $\xi(G)$ to denote the maximum number of orbits of M among all minimal normal subgroups M of G.

Problem 1.7. Among all counterexamples G to the Polycirculant Conjecture, determine:

(1) the smallest degree of G;
(2) the smallest order of G;
(3) the least number of orbits in a minimal normal subgroup of G; and
(4) the smallest value of $\xi(G)$.

Similarly, among all vertex-transitive directed graphs Γ admitting no semiregular automorphism, determine:

(5) the smallest order of Γ; and
(6) the smallest valency (or out-valency) of Γ.

Note that the aforementioned survey paper [1] includes results on necessary conditions on the valency and other properties of such a graph Γ.

Problem 1.8. Does there exist a counterexample G to the Polycirculant Conjecture, such that G is not the full automorphism group of a vertex-transitive directed graph?

If no such G exists, then Parts (1) and (5) of Problem 1.7 have the same answer.

The remainder of the paper is organised as follows. Section 2 outlines some preliminary results, and we discuss constructions of elusive groups from non-split extensions in Section 3. The proofs of Theorem 1.3 and Corollary 1.6 are then detailed in Section 4. We note that the former proof is essentially entirely computational. As such, Section 4 also includes a proof of Corollary 1.4, combining theoretical and computational techniques, that does not rely on Theorem 1.3. The techniques in this proof may be useful when considering further potential counterexamples to the Polycirculant Conjecture that are not amenable to the computational methods used to prove Theorem 1.3, in particular those that may not be the full automorphism group of any vertex-transitive directed graph.

2. Preliminaries

We first introduce some notation. For a permutation group $G \leqslant \mathrm{Sym}(\Omega)$, and Σ a G-invariant partition of G on Ω, we write G^Σ to denote the subgroup of $\mathrm{Sym}(\Sigma)$ induced by G on Σ. For $\Delta \in \Sigma$, we write G_Δ for the setwise stabiliser of Δ in G, and G^Δ_Δ for the subgroup of $\mathrm{Sym}(\Delta)$ induced by G_Δ on Δ. The 2-closure of G is denoted by $\kappa(G)$, where the action is clear from context.

Our proof of Corollary 1.4 requires the following structural results.

Lemma 2.1 ([2, Lemma 3.1])**.** *Let $G \leqslant \mathrm{Sym}(\Omega)$ be a transitive group, and suppose that Σ is a non-trivial G-invariant partition of Ω. Then the following statements hold.*

(1) *The partition Σ is also $\kappa(G)$-invariant.*
(2) *The induced group $\kappa(G)^\Sigma$ is a subgroup of $\kappa(G^\Sigma)$.*
(3) *For $\Delta \in \Sigma$, the group $\kappa(G)^\Delta_\Delta$ is a subgroup of $\kappa(G^\Delta_\Delta)$.*

Theorem 2.2 (Imprimitive Wreath Embedding Theorem)**.** *Let $G \leqslant \mathrm{Sym}(\Omega)$, let Σ be a G-invariant partition of Ω with finitely many blocks, and let $\Delta \in \Sigma$. Further assume that the group G^Σ induced by G on Σ is transitive. Then G is permutationally isomorphic to a subgroup of $G^\Delta_\Delta \wr G^\Sigma$ acting on $\Delta \times \Sigma$.*

See [18, Theorem 5.5] for a proof, which details a permutational isomorphism that can be used to embed the group G in the specified wreath product. A more direct description of the permutational embedding of G is given in [14, Section II.4].

Our next result is the case $k = 2$ of the theorem [19, Theorem 5.6] on *k-closures* of permutation groups for each $k \geqslant 1$.

Proposition 2.3. *Let $G \leqslant \mathrm{Sym}(\Omega)$. A permutation $\sigma \in \mathrm{Sym}(\Omega)$ lies in $\kappa(G)$ if and only if for all $\alpha, \beta \in \Omega$, there exists $g \in G$ with $\alpha^\sigma = \alpha^g$ and $\beta^\sigma = \beta^g$.*

The following proposition shows that any 2-closed elusive group gives rise to an infinite family of 2-closed elusive groups.

Proposition 2.4. *Let $G_1 \leqslant \mathrm{Sym}(\Omega_1)$ and $G_2 \leqslant \mathrm{Sym}(\Omega_2)$ be transitive permutation groups.*

(1) *If G_1 and G_2 are elusive, then $G_1 \times G_2$ is elusive in its (product) action on $\Omega_1 \times \Omega_2$, and $G_1 \wr G_2$ is elusive in its (imprimitive) action on $\Omega_1 \times \Omega_2$.*
(2) *If G_1 and G_2 are 2-closed, then $G_1 \times G_2$ and $G_1 \wr G_2$ are 2-closed in their actions on $\Omega_1 \times \Omega_2$.*

Proof. Part (1) is [5, Theorem 4.1(c)], while Part (2) is [5, Theorem 5.1]. □

We also need some theory concerning graph products to provide the machinery for our proof of Corollary 1.6. Let $\Gamma_1 = (V_1, E_1)$ and $\Gamma_2 = (V_2, E_2)$ be two simple graphs, i.e. without loops or multiple edges. We recall two well-known graph products. The *Cartesian product* $\Gamma_1 \,\square\, \Gamma_2$ and *strong product* $\Gamma_1 \boxtimes \Gamma_2$ each have vertex set

$$V(\Gamma_1 \,\square\, \Gamma_2) = V(\Gamma_1 \boxtimes \Gamma_2) = \{(v, w) \mid v \in V(\Gamma_1), w \in V(\Gamma_2)\}.$$

Two distinct vertices (v, w) and (v', w') are adjacent in the Cartesian product if and only if either $v = v'$ and w is adjacent to w' in Γ_2, or $w = w'$ and v is adjacent to v' in Γ_1. They are adjacent in the strong product if and only if v and v' are either equal or adjacent in Γ_1 and w and w' are either equal or adjacent in Γ_2. Hence $\Gamma_1 \,\square\, \Gamma_2$ is a subgraph of $\Gamma_1 \boxtimes \Gamma_2$.

A graph is called *prime* with respect to one of these graph products if it is non-trivial and cannot be written as the product of two non-trivial graphs. If a graph is non-trivial and not prime, we say that it is *decomposable* with respect to that product. We further say that two graphs are *relatively prime* with respect to a graph product if there is no non-trivial graph that (up to isomorphism) is a factor of both of them. We now prove some properties of graphs constructed from Cartesian and strong products.

Proposition 2.5. *Let $\Gamma_1 = (V_1, E_1)$ and $\Gamma_2 = (V_2, E_2)$ be simple graphs on at least two vertices. Then the following statements hold.*

(1) *The products $\Gamma_1 \,\square\, \Gamma_2$ and $\Gamma_1 \boxtimes \Gamma_2$ are connected if and only if Γ_1 and Γ_2 are, and vertex-transitive if and only if Γ_1 and Γ_2 are.*
(2) *If, for each $i \in \{1, 2\}$, no two vertices in Γ_i have the same closed neighbourhoods, then no two vertices in $\Gamma_1 \,\square\, \Gamma_2$ have the same closed neighbourhoods. Separately, if Γ_1 and Γ_2 are connected and relatively prime with respect to the Cartesian product, then $\Gamma_1 \,\square\, \Gamma_2$ admits semiregular automorphisms only if at least one of Γ_1 and Γ_2 do.*
(3) *If, for each $i \in \{1, 2\}$, no two vertices in Γ_i have the same closed neighbourhoods, then no two vertices in $\Gamma_1 \boxtimes \Gamma_2$ have the same closed neighbourhoods. If, in addition to this property, Γ_1 and Γ_2 are connected and relatively prime with respect to the strong product, then $\Gamma_1 \boxtimes \Gamma_2$ admits semiregular automorphisms only if at least one of Γ_1 and Γ_2 do.*

Proof. Part (1) follows immediately from Corollaries 5.3 and 5.6 and Theorems 6.17 and 7.19 in [13]. Next, we consider Part (2). The statement about closed neighbourhoods in $\Gamma_1 \,\square\, \Gamma_2$ is immediate from the definition of the Cartesian product. Suppose now that Γ_1 and Γ_2 are connected and relatively prime with respect to the Cartesian product. Then [13, Corollary 6.12] yields $\mathrm{Aut}(\Gamma_1 \,\square\, \Gamma_2) = \mathrm{Aut}(\Gamma_1) \times \mathrm{Aut}(\Gamma_2)$. Hence if $\Gamma_1 \,\square\, \Gamma_2$

admits a semiregular automorphism, then by considering this automorphism's projection onto each of Γ_1 and Γ_2, we see that at least one of these two factors also admits such an automorphism. The proof of Part (3) is similar, instead using [13, Lemma 7.2, Corollary 7.17]. □

We now provide the key result that will enable us to construct our desired family of graphs.

Proposition 2.6 ([13, Exercise 7.6])**.** *Every connected graph that is decomposable with respect to the Cartesian product is prime with respect to the strong product and vice versa.*

3. Constructing elusive groups via non-split extensions

In this section, we briefly formalise and generalise methods used by Chen et al. [7] to construct elusive groups via non-split extensions. We then discuss how these constructions apply to the groups discussed in Theorem 1.3 and Remark 1.5.

Given a set π of primes, we will say that a transitive permutation group G is *π-elusive* if none of its derangements have orders in π, and *π'-elusive* if each derangement g of G of prime order has $|g| \in \pi$ (this is a slight generalisation of notation used by Chen et al.). Additionally, for an extension $N.G$ (with N specified), let $\overline{\ } : X \to G$ be the natural projection, so that $\overline{Y} \cong NY/N$ for all $Y \leqslant X$.

Recall that a non-split extension $N{\cdot}G$, corresponding to a specific action of a group G on an elementary abelian group N (so that N is a specific G-module), exists if and only if the corresponding second cohomology group $\mathrm{H}^2(G, N)$ has positive dimension. By the Schur–Zassenhaus Theorem, this can only be the case if the prime dividing $|N|$ also divides $|G|$.

Proposition 3.1. *Let G be a finite group, H a core-free subgroup of G, and N an elementary abelian group on which G acts irreducibly, such that $\mathrm{H}^2(G, N)$ has positive dimension. Additionally, let p be the prime dividing $|N|$, let X be a non-split extension $N{\cdot}G$, and let L be the preimage of H under the projection $\overline{\ }$. Suppose furthermore that the following conditions all hold.*

(a) *In its action on the set $[G : H]$ of right cosets of H, the group G is π-elusive for some set π of primes.*
(b) *The order of G is not divisible by p^2.*
(c) *The second cohomology group $\mathrm{H}^2(H, N)$ (corresponding to the restriction to H of the G-module N) has dimension 0.*

Then there is a one-to-one correspondence between:

(1) *subgroups Y of X (up to L-conjugacy) with $\overline{Y} = H$, such that X is faithful and $(\pi \cup \{p\})$-elusive in its action on $[X : Y]$; and*
(2) *proper H-submodules M of N satisfying $\bigcup_{g \in G} M^g = N$,*

given by $Y \cap N = M$.

Proof. Since $N \leqslant L$ and $L/N \cong H$, the group L is an extension of N by H. Additionally, $\mathrm{H}^2(H, N)$ has dimension 0, and so $L = N : H$ (with H identified with the appropriate subgroup of L). As each subgroup Y of L with $\overline{Y} = H$ satisfies $Y \cap N \trianglelefteqslant Y$, it follows

that (up to L-conjugacy) these subgroups are precisely $M:H$ for the H-submodules M of N. (Note that if $M=N$, then $Y=L$ contains N and so is not core-free in X.)

Now, fix a proper H-submodule M of N, and let $Y:=M:H$. As N lies in the kernel of the action of X on N, the action of G on N induces the action of X on N. Since N is an irreducible G-module and $\overline{Y}=H$ is core-free in G, we deduce that Y is core-free in X. From now on, we will consider X as a permutation group acting faithfully and transitively on $[X:Y]$.

As p is the only prime dividing N, [7, Lemma 2.1] shows that X is $(\pi\backslash\{p\})$-elusive. Hence it remains to consider whether X has a derangement of order p. Observe that this is the case if and only if an element of order p in X lies in no X-conjugate of Y. Since G is not divisible by p^2, a theorem of Gaschütz (see [15, Main Theorem I.17.4]) implies that $X\backslash N$ has no elements of order p, by the reasoning given in the proof of [7, Lemma 2.5]. Thus X contains no derangement of order p if and only if $\bigcup_{x\in X}(Y\cap N)^x=N$. Since the action of X on $Y\cap N=M$ is induced by the action of G, this is the case if and only if $\bigcup_{g\in G}M^g=N$, as required. □

We note that Proposition 3.1 does not depend on the specific non-split extension of the G-module N by G, and that the condition $\bigcup_{g\in G}M^g=N$ also appears in [5, Theorem 3.1] on the construction of elusive groups of affine type. Additionally, the Schur–Zassenhaus Theorem implies that Condition (c) holds whenever p does not divide $|H|$. Furthermore, $|\{M^g \mid g\in G\}|\leqslant |G:H|$ for each H-submodule M of N. Hence if M and N have dimension m and n, respectively, over $\mathbb{F}_p$ and $|G:H|\,(p^m-1)<p^n-1$, then $\bigcup_{g\in G}M^g\subsetneq N$.

Suppose now that G is π'-elusive for some set π of primes, and that there exist a fixed subgroup H of G and irreducible G-modules N_p for each prime $p\in\pi$ that satisfy the conditions of Proposition 3.1. If each N_p contains a proper H-submodule M_p with $\bigcup_{g\in G}M_p^g=N_p$, then the proposition implies that each non-split extension $(\prod_{p\in\pi}N_p)\cdot G$ (corresponding to the relevant actions of G on each N_p) is elusive (when acting on the set of cosets of an appropriate subgroup that projects onto H). By [5, Theorem 5.3] (see also the discussion in [7, Section 5]), a counterexample to the Polycirculant Conjecture constructed via this method must have $\dim(N_p)-\dim(M_p)\geqslant 2$ for all $p\in\pi$.

Our constructions of the elusive groups described in Theorem 1.3 and Remark 1.5 appeal to Proposition 3.1. Indeed, $G_1:=\mathrm{PSU}_3(3)$ has a unique (up to conjugacy) subgroup H_1 isomorphic to $C_3\times S_3$, and a unique irreducible 6-dimensional G-module N_1 over $\mathbb{F}_7$. Similarly, $G_2:=\mathrm{P\Gamma U}_3(3)$ has a unique (up to conjugacy) subgroup H_2 isomorphic to $S_3\times S_3$, and a unique irreducible 6-dimensional G-module N_2 over $\mathbb{F}_7$ satisfying $\dim(\mathrm{H}^2(G,N))>0$. It is easy to show computationally that, for each $i\in\{1,2\}$, the group G_i is $\{7\}'$-elusive on $[G_i:H_i]$, that G_i, H_i and N_i satisfy the hypotheses and conditions of Proposition 3.1, and that there is a unique non-split extension $N_i{\cdot}G_i$. The elusive groups mentioned in Theorem 1.3 and Remark 1.5 correspond to the H_i-submodules M_i of N_i satisfying $\bigcup_{g\in G_i}M_i^g=N_i$ and $\dim(N_i)-\dim(M_i)=2$. Since the 2-closures of the groups in Remark 1.5 are not elusive, the necessary condition given at the end of the previous paragraph is not sufficient for the constructed groups to be counterexamples to the Polycirculant Conjecture.

We conclude this section by noting that Problem 5.2 in [7] asks if a counterexample to the Polycirculant Conjecture can be constructed as such a non-split extension $N{\cdot}G$,

with G quasiprimitive and $\{p\}'$-elusive on $[G : H]$ for some prime p. Since the $\{7\}'$-elusive permutation group G_1 is simple, it is quasiprimitive. Therefore, the 2-closed elusive group $X = N_1{\cdot}G_1$ from Theorem 1.3 and Corollary 1.4 provides a positive answer to this problem.

4. Proofs of Theorem 1.3 and Corollaries 1.4 and 1.6

We now embark on the proofs of Theorem 1.3 and Corollaries 1.4 and 1.6. While the proof of the theorem is primarily computational, our proofs of the corollaries combine theoretical ideas and computations. As such, we will begin by theoretically describing the setup of all three proofs. Our computations are performed in Magma [3] via the accompanying code [9], which requires approximately 35 minutes of CPU time with a 3.85 GHz processor and 15 GB of RAM (or just 13 minutes and 0.8 GB if only the computations necessary to prove Theorem 1.3 and Corollary 1.6 are performed). The main results shown by our code will be emphasised via numbered Computational Propositions.

4.1. **Proof of Theorem 1.3 and Corollary 1.4.** Let $G = \mathrm{PSU}_3(3)$ and $N = 7^6$. There is a unique non-split extension $X := N{\cdot}G$ with G acting irreducibly on N, and (up to conjugacy) X has exactly three subgroups of order $43{,}218 = 2.3^2.7^4$ (each isomorphic to groups of shape $7^4 : (C_3 \times S_3)$) whose images under the natural projection from X to G are equal to a subgroup $H \cong C_3 \times S_3$ (cf. the discussion at the end of Section 3). Of these three subgroups of X, one has exactly nine normal subgroups, while the other two have 16 (cf. Remark 1.5). Let Y be either of the latter two subgroups, and consider the action of X on the set $\Omega := [X : Y]$ of right cosets of Y in X.

Since G is simple and acts irreducibly on N, we observe that N is the unique proper non-trivial normal subgroup of the non-split extension X. Thus Y is core-free in X, and so the action of X is faithful and transitive of degree $16{,}464 = 2^4.3.7^3$.

Computational Proposition 4.1. X is elusive.

We also observe computationally that there is an X-invariant partition Σ of Ω consisting of 56 blocks of size 294 (as well as three other non-trivial X-invariant partitions). Let $\Delta \in \Sigma$, let X^Σ be the permutation group induced by X on Σ, and let X_Δ^Δ be the permutation group induced by X_Δ on Δ. As X acts transitively on Ω, Lemma 2.1 shows that Σ is a $\kappa(X)$-invariant partition, that $\kappa(X)^\Sigma \leqslant \kappa(X^\Sigma)$, and that $\kappa(X)_\Delta^\Delta \leqslant \kappa(X_\Delta^\Delta)$.

Now, by Theorem 2.2, $\kappa(X)$ is a subgroup (up to permutational isomorphism) of the group $\kappa(X)_\Delta^\Delta \wr \kappa(X)^\Sigma$, in its natural imprimitive action on $\Xi := \Delta \times \Sigma$. It follows from the previous paragraph that $X \leqslant \kappa(X) \leqslant \kappa(X_\Delta^\Delta) \wr \kappa(X^\Sigma)$. In fact, we check computationally that $\kappa(X_\Delta^\Delta) = X_\Delta^\Delta$, and so $X \leqslant \kappa(X) \leqslant X_\Delta^\Delta \wr \kappa(X^\Sigma)$. We will write $A := X_\Delta^\Delta$, $B := \kappa(X^\Sigma)$ and $W := A \wr B$.

Note that B is a subgroup of the symmetric group of degree $|\Sigma| = 56$, and so $W = (A_1 \times A_2 \times \cdots \times A_{56}) : B$, where $A_i \cong A$ for each i. Let $C := A_1 \times A_2 \times \cdots \times A_{56}$. We will identify Σ with the set $\{1, \ldots, 56\}$. For $(a_1, \ldots, a_{56}) \in C$, $b \in B$, $\delta \in \Delta$ and $i \in \Sigma$, the image $(\delta, i)^{(a_1,\ldots,a_{56})}$ is equal to (δ^{a_i}, i), while $(\delta, i)^b = (\delta, i^b)$. Writing $\Delta_i := \Delta \times \{i\}$, it follows that $\Delta_i^{(a_1,\ldots,a_{56})b} = \Delta_{i^b}$. It is therefore clear that each A_i acts transitively on Δ_i, and trivially on Δ_j for each $i \neq j$. Furthermore, the partition $\{\Delta_1, \ldots, \Delta_{56}\}$ of Σ is W-invariant, and hence X-invariant.

Let $S := C \cap X$, let $D := N_W(S)$, and let $\alpha \in \Xi$. Since $C \trianglelefteq W$, it is clear that $S \trianglelefteq X$, and so $X \leqslant D$. (In fact, S is the unique proper non-trivial normal subgroup N of X). Straightforward computations show that X_α has exactly 14 orbits on Ξ (each of size 63) that extend to larger orbits of D_α (each of size 441). Of the 14 corresponding X-orbitals of size $63 \cdot 16{,}464$, exactly eight are self-paired, and of these eight, only one is D-conjugate to a non-self-paired X-orbital. Let $\mathcal{U}$ be the set of the remaining seven self-paired X-orbitals, which all have valency 63.

Computational Proposition 4.2. The group X is the full automorphism group of the orbital graph of each orbital in $\mathcal{U}$, and these seven graphs are connected and pairwise non-isomorphic.

Thus we have proved Theorem 1.3, and it immediately follows that X is 2-closed. However, as promised at the end of Section 1, we will continue to prove this fact without relying on the above automorphism group calculation. Note also that we have not been able to compute the automorphism groups of the orbital graphs of the remaining seven X-orbitals that are not D-orbitals. (The orbital graph of any X-orbital that is also a D-orbital clearly has automorphism group containing D.)

Computational Proposition 4.3. For each transitive overgroup M of S in D with $|X|$ dividing but not equal to $|M|$, the number of orbits on Ξ of the point stabiliser M_α is less than the number of orbits of X_α.

Observe that $D \cap \kappa(X)$ contains X, and therefore contains S; that the transitivity of X implies the transitivity of $D \cap \kappa(X)$; and that $(D \cap \kappa(X))_\alpha$ has the same orbits on Ξ as X_α. We therefore deduce from Computational Proposition 4.3 that $D \cap \kappa(X) = X$.

Next, fix $c \in C \cap \kappa(X)$, and write $c = (c_1, \ldots, c_{56})$, with $c_i \in A_i$ for each i. By Proposition 2.3, for each $\alpha, \beta \in \Xi$, there exists $x \in X$ such that $(\alpha, \beta)^c = (\alpha, \beta)^x$. Consider the case where $\alpha \in \Delta_i$ and $\beta \in \Delta_j$ for $i \neq j$. Since C stabilises each of Δ_i and Δ_j setwise and $\{\Delta_1, \ldots, \Delta_{56}\}$ is an X-invariant partition, x lies in the intersection T_{ij} of the setwise stabilisers X_{Δ_i} and X_{Δ_j}. Therefore (cf. the proof of [2, Lemma 3.5(2)]), for all $(\alpha, \beta) \in \Delta_i \times \Delta_j$, the pair $(\alpha, \beta)^c = (\alpha^{c_i}, \beta^{c_j})$ lies in the orbit $(\alpha, \beta)^{T_{ij}}$. Equivalently, letting $A_{ij} := \langle A_i, A_j \rangle$ and letting $\mathcal{O}_{ij}$ be the set of orbits of T_{ij} on $\Delta_i \times \Delta_j$, the tuple $(1, \ldots, 1, c_i, 1, \ldots, 1, c_j, 1, \ldots, 1)$ lies in the intersection $\bigcap_{O \in \mathcal{O}_{ij}} (A_{ij})_O$ of setwise stabilisers. Therefore, an element $c \in C$ lies in $\kappa(X)$ only if this condition holds for all distinct i and j. (It does not take much more work to show that this is also a sufficient condition, but we will not need this fact.)

For each $i \in \{1, \ldots, 56\}$, let F_i be the subgroup of A_i consisting of all elements c_i such that, for all $j \neq i$, there exists $c_j \in A_j$ such that $(1, \ldots, 1, c_i, 1, \ldots, 1, c_j, 1, \ldots, 1)$ lies in $\bigcap_{O \in \mathcal{O}_{ij}} (A_{ij})_O$. Then $C \cap \kappa(X) \leqslant \langle F_1, \ldots, F_{56} \rangle$.

Computational Proposition 4.4. There exists $k \in \{1, \ldots, 56\}$ such that $F_k \leqslant D$.

Let $x \in X$. Then $(T_{ij})^x = T_{i^x j^x}$ and $(A_i)^x = A_{i^x}$ for each i and j, and so $(F_i)^x = F_{i^x}$. Since X is transitive on Ξ, and since D contains $\langle F_k, X \rangle$ by Computational Proposition 4.4, it follows that D contains $\langle F_1, \ldots, F_{56} \rangle$, which contains $C \cap \kappa(X)$. Thus $C \cap \kappa(X) \leqslant D \cap \kappa(X) = X$, and so the normal subgroup $C \cap \kappa(X)$ of $\kappa(X)$ is equal to $C \cap X = S$. Therefore, $\kappa(X) \leqslant N_W(S) = D$. Again using the fact that $D \cap \kappa(X) = X$, we conclude that $\kappa(X) = X$, i.e. X is 2-closed. Combining this fact with Computational Proposition 4.1, we see that X is a counterexample to the Polycirculant Conjecture.

Finally, Proposition 2.4 shows that, for each integer $k \geqslant 1$, every group constructed from k copies of X via direct products and/or wreath products has a 2-closed elusive action of degree $16{,}464^k$ on the set Ω^k. Each of these groups is a counterexample to the Polycirculant Conjecture. This completes the proof of Corollary 1.4.

4.2. **Proof of Corollary 1.6.** Let Γ_1 and Γ_2 be any two distinct graphs from Computational Proposition 4.2. We define a family $\mathcal{G} := \{\mathcal{G}_i \mid i \in \mathbb{N}\}$ of graphs by $\mathcal{G}_1 := \Gamma_1$ and, for each $i \in \mathbb{N}$,

$$\mathcal{G}_{2i} := \mathcal{G}_{2i-1} \square \Gamma_2, \qquad \text{and} \qquad \mathcal{G}_{2i+1} := \mathcal{G}_{2i} \boxtimes \Gamma_1.$$

We claim that each member of $\mathcal{G}$ is a connected, vertex-transitive graph admitting no semiregular automorphism. Our computations reveal that no two vertices in Γ_1 or Γ_2 have the same closed neigbourhoods. Moreover, [13, Theorem 6.13] shows that if Γ_1 or Γ_2 is not prime with respect to the Cartesian product, then its automorphism group is a direct product of wreath products. By [13, Lemma 7.2, Theorem 7.18] and the above closed neighbourhood property, the same is true for the strong product. However, Computational Proposition 4.2 shows that X is the full automorphism group of each of Γ_1 and Γ_2. The unique proper non-trivial normal subgroup of X is $N \cong 7^6$, which has no complement in X. Hence X cannot be written as a direct product, nor as a single wreath product. Therefore, Γ_1 and Γ_2 must be prime with respect to the Cartesian and strong products.

We now proceed via induction. Suppose that, for some $i \in \mathbb{N}$, the graph $\mathcal{G}_i$ is connected and vertex-transitive, that no two of its vertices have equal closed neighbourhoods, and that it does not admit any semiregular automorphism. We immediately deduce from Proposition 2.5 that $\mathcal{G}_{i+1}$ is connected and vertex-transitive, and that no two of its vertices have equal closed neighbourhoods. By applying the previous paragraph when $i = 1$, or Proposition 2.6 and the definition of $\mathcal{G}_i$ when $i > 1$, we see that $\mathcal{G}_i$ is prime with respect to the product used to construct $\mathcal{G}_{i+1}$, and also relatively prime to Γ_2 and (if $i > 1$) Γ_1. Since Γ_1 and Γ_2 admit no semiregular automorphisms, it follows from Proposition 2.5(2)–(3) and our inductive hypothesis that $\mathcal{G}_{i+1}$ also admits no semiregular automorphism. Observing that $|V(\mathcal{G}_i)| = 16{,}464^i$ for each $i \geqslant 1$ completes the proof of Corollary 1.6.

Remark 4.5. One can replace Γ_1 and Γ_2 at each step of the construction of $\mathcal{G}$ with any of the seven orbital graphs from Computational Proposition 4.2, provided that the two factors of $\mathcal{G}_2$ are chosen to be distinct. It is easy to see that each graph in the resulting infinite family of graphs will be vertex-transitive and will not admit semiregular automorphisms. As a result, this construction produces countably many families of positive answers to the Semiregularity Problem. There are also likely to be other constructions, perhaps also based on graph products, that extend Computational Proposition 4.2 to further infinite families of vertex-transitive graphs that do not admit semiregular automorphisms.

REFERENCES

[1] M. Arezoomand, A. Abdollahi, and P. Spiga. On problems concerning fixed-point-free permutations and on the polycirculant conjecture—a survey. *Trans. Comb.*, 8(1):15–40, 2019.

[2] M. Arezoomand, M. A. Iranmanesh, C. E. Praeger, and G. Tracey. Totally 2-closed finite groups with trivial Fitting subgroup. *Bull. Math. Sci.*, 14(1):Paper No. 2350004, 76, 2024.

[3] W. Bosma, J. Cannon, and C. Playoust. The Magma algebra system. I. The user language. *J. Symbolic Comput.*, 24(3-4):235–265, 1997. Computational algebra and number theory (London, 1993).

[4] T. C. Burness and M. Giudici. Locally elusive classical groups. *Israel J. Math.*, 225(1):343–402, 2018.

[5] P. J. Cameron, M. Giudici, G. A. Jones, W. M. Kantor, M. H. Klin, D. Maruˇsič, and L. A. Nowitz. Transitive permutation groups without semiregular subgroups. *J. London Math. Soc. (2)*, 66(2):325–333, 2002.

[6] P. J. Cameron (ed.). Research problems from the 15th British Combinatorial Conference. *Discrete Mathematics*, 167/168:605–615, 1997.
[7] J. Chen, M. Lee, Đ. Mitrović, E. A. O'Brien, and B. Xia. Elusive groups from non-split extensions. *J. Pure Appl. Algebra*, 230(4):Paper No. 108228, 18, 2026.
[8] B. Fein, W. M. Kantor, and M. Schacher. Relative Brauer groups. II. *J. Reine Angew. Math.*, 328:39–57, 1981.
[9] S. D. Freedman and M. Lee. Code accompanying this paper. `https://github.com/SaulDFreedman/PolycirculantCounterexample`.
[10] M. Giudici. Quasiprimitive groups with no fixed point free elements of prime order. *J. London Math. Soc. (2)*, 67(1):73–84, 2003.
[11] M. Giudici and S. Kelly. Characterizing a family of elusive permutation groups. *J. Group Theory*, 12(1):95–105, 2009.
[12] M. Giudici and J. Xu. All vertex-transitive locally-quasiprimitive graphs have a semiregular automorphism. *J. Algebraic Combin.*, 25(2):217–232, 2007.
[13] R. Hammack, W. Imrich, and S. Klavˇ zar. *Handbook of product graphs*. Discrete Mathematics and its Applications (Boca Raton). CRC Press, Boca Raton, FL, second edition, 2011. With a foreword by Peter Winkler.
[14] A. Hulpke. *Notes on Computational Group Theory*. Lecture Notes, 2010.
[15] B. Huppert. *Finite groups. I*, volume 364 of *Grundlehren der mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences]*. Springer, Cham, german edition, [2025] ©2025. With a foreword by Schroeder.
[16] D. Jordan. Eine Symmetrieeigenschaft von Graphen. In *Graphentheorie und ihre Anwendungen (Stadt Wehlen, 1988)*, volume 9 of *Dresdner Reihe Forsch.*, pages 17–20. Päd. Hochsch. Dresden, Dresden, 1988.
[17] D. Marušič. On vertex symmetric digraphs. *Discrete Math.*, 36(1):69–81, 1981.
[18] C. E. Praeger and C. Schneider. *Permutation groups and Cartesian decompositions*, volume 449 of *London Mathematical Society Lecture Note Series*. Cambridge University Press, Cambridge, 2018.
[19] H. Wielandt. Permutation groups through invariant relations and invariant functions [83]. In *Mathematische Werke/Mathematical Works, Volume 1: Group Theory*, pages 237–296. De Gruyter, Berlin, New York, 1994.

(Freedman) DEPARTMENT OF MATHEMATICS, COLORADO STATE UNIVERSITY, FORT COLLINS, CO 80523, USA

(Lee) SCHOOL OF MATHEMATICS, MONASH UNIVERSITY, CLAYTON, AUSTRALIA

Email address: `saul.freedman@colostate.edu, melissa.lee@monash.edu`