

TechNote 13.2

Relative Efficiency of the Latin Square Design

We assume a repeated measures design with a subjects, each tested on a random sequence of a conditions. We also have a Latin square with the same dimensions. Assume that the population error variance for the repeated measures design is $\sigma_{e(RM)}^2$ and that for the Latin square design is

$\sigma_{e(LS)}^2$. Because the Latin square design merely involves a rearrangement of the treatments for

each subject, the expected total sums of squares are the same for the two designs, as are the expected sums of squares for subjects. Therefore,

$$(a-1)^2 \sigma_{e(RM)}^2 + (a-1)(\sigma_{e(RM)}^2 + a\theta_A^2) = (a-1)(a-2)\sigma_{e(LS)}^2 \\ + (a-1)(\sigma_{e(LS)}^2 + a\theta_A^2) + (a-1)(\sigma_{e(LS)}^2 + a\theta_C^2)$$

Canceling θ_A^2 from both sides and combining terms where possible, we have

$$a(a-1)\sigma_{e(RM)}^2 = (a-1)^2 \sigma_{e(LS)}^2 + (a-1)(\sigma_{e(LS)}^2 + a\theta_C^2)$$

Dividing both sides by $a(a-1)$, and converting population variances to mean squares, we have the result in Equation 13.4.