

Impact and influence of the natural *Vibrio*-squid symbiosis in understanding bacterial-animal interactions

1 Mark J. Mandel^{1*}, Anne K. Dunn²

2 ¹ Northwestern University Feinberg School of Medicine, Department of Microbiology-Immunology,
3 320 E Superior St, Chicago, IL 60611, USA

4 ² University of Oklahoma, Department of Microbiology and Plant Biology, 770 Van Vleet Oval,
5 Norman, OK 73019, USA

6 * **Correspondence:**

7 Mark J. Mandel

8 m-mandel@northwestern.edu

9 **Keywords: symbiosis, microbiome, invertebrate model, marine microbiology, epithelial**
10 **colonization, evolution**

11 Abstract

12 Animals are colonized by bacteria, and in many cases partners have co-evolved to perform mutually
13 beneficial functions. An exciting and ongoing legacy of the past decade has been an expansion of
14 technology to enable study of natural associations *in situ/in vivo*. As a result more symbioses are
15 being examined, and additional details are being revealed for well-studied systems with a focus on
16 the interactions between partners in the native context. With this framing, we review recent literature
17 from the *Vibrio fischeri*-*Euprymna scolopes* symbiosis and focus on key studies that have had an
18 impact on understanding bacteria-animal interactions broadly. This is not intended to be a
19 comprehensive review of the system, but rather to focus on the molecular basis to intriguing
20 observations in the field of host-microbe interactions. In this review we discuss the following topics:
21 processes regulating strain and species specificity; bacterial signaling to host morphogenesis;
22 multiple roles for nitric oxide; flagellar motility and chemotaxis; and efforts to understand
23 unannotated and poorly annotated genes. Overall these studies demonstrate how functional
24 approaches *in vivo* in a tractable system have provided valuable insight into general principles of
25 microbe-host interactions.

26 1 Introduction

27 Studies of human, animal, and plant microbiomes have been advanced by novel culture-independent
28 approaches and technological advancements in DNA sequencing. In recent years a prominent role for
29 microbial communities of the gut, skin, and other organs has emerged as modulators of human health
30 (Human Microbiome Project Consortium 2012). These studies followed from influential animal
31 studies in systems that are yielding critical insight into microbiome assembly, stability,
32 communication, and evolution (Ruby 2008, McFall-Ngai et al. 2013). The focus of this review is to
33 examine one model system, The *Vibrio fischeri*-*Euprymna scolopes* symbiosis, and how key findings
34 in that system have enabled an increasingly higher resolution of the processes and principles that
35 underlie microbe-host communication.

When Hawaiian bobtail squid hatch from their eggs, they are exposed to a million bacteria in each milliliter of seawater. Although *V. fischeri* make up less than 1 in 5,000 of these planktonic, environmental bacteria, the “light organ” of the hatchling squid becomes colonized exclusively with *V. fischeri* (Ruby and Lee 1998, Mandel 2010). The microbe-host specificity relies on a series of reciprocal communications between the partners, many of which are detailed in the sections below. Over the course of 48 hours the bacteria establish a mature colonization in epithelium-lined crypts of the squid light organ, and, at high cell density, produce light as a result of quorum-sensing. The bacterial bioluminescence is reflected by host tissue to camouflage the shadow or silhouette that the nocturnal-foraging squid would cast in the moonlight, thus protecting the host in a process termed counter-illumination (Ruby and McFall-Ngai 1992, Jones and Nishiguchi 2004). Initiation of colonization occurs in newly-hatched squid, seeding an individual host’s crypts for its lifetime. The bacteria produce light at night, then at dawn approximately 90-95% of the symbiotic population is expelled into the seawater. The remaining cells grow up during the day, produce light at night, and a diel cycle of growth, light production, and expulsion proceeds for the lifetime of the animal (Wier et al. 2010). Host cellular changes accompany this cycle, e.g. a daily reshaping of the epithelial brush border against which the bacteria reside during the final two hours prior to the daily expulsion (Wier et al. 2010).

As an environmentally-transmitted symbiosis, the *Vibrio*-squid model has a number of valuable characteristics that have served it well as a study system for identifying molecular mechanisms. First, the binary system (two partners) is naturally reduced. Second, both partners can be raised separately and then introduced for experimentation. Third, *V. fischeri* is genetically tractable, and unbiased mutagenesis as well as precise genetic alterations can be introduced with relative ease. Fourth, the bacteria colonize the host light organ directly under the semi-transparent mantle and funnel; this permits imaging of the site of infection and direct analysis of bacterial behaviors and host responses. Fifth, synchronous colonization of hatchlings has permitted developmental staging of the colonization process. For most of the processes described below, many of these benefits were important in the advances described.

2 From pattern to process in the *Vibrio*-squid symbiosis

In each section below, we highlight key discoveries in the *Vibrio*-squid symbiosis with a specific focus on how this model system has revealed molecular processes that underlie mutually beneficial phenotypes.

2.1 Just the two of us

E. scolopes squid are colonized only by *V. fischeri*, and this exclusivity has guided substantial inquiry and discovery in the system. This pattern was first explored by McFall-Ngai and Ruby (McFall-Ngai and Ruby 1991) and extended in subsequent works (Ruby and Lee 1998, Mandel et al. 2009). The ability to image the live animal during colonization enabled the discovery of *V. fischeri* aggregating in close proximity to the ciliated epithelial fields of the light organ (Nyholm et al. 2000). Nyholm discovered that a narrow distance between the green fluorescent protein-expressing bacteria and the squid epithelial tissue was the result of host-produced mucus, which included *N*-acetylneuraminic acid and *N*-acetylgalactosamine. Recent work has demonstrated that *V. fischeri* bind to cilia within this mucus field (Altura et al. 2013). Whereas many bacteria can bind in host mucus, only specific strains and species exhibit a competitive dominance over non-colonizing isolates, and only (some) *V. fischeri* strains proceed to fully initiate colonization (Nyholm et al. 2000, Nyholm and McFall-Ngai 2003, Mandel et al. 2009).

Around this same time, the genetic basis for bacterial aggregation was being discovered and characterized in the laboratory of Karen Visick. A forward genetic screen for colonization factors first identified an orphan histidine kinase, RscS (regulator of symbiotic colonization-sensor), but without a phenotype or target it was difficult to know how this factor connected to the colonization process (Visick and Skoufos 2001). The same screen identified an eighteen gene locus that encoded regulatory proteins, glycosyltransferases, and other factors involved in exopolysaccharide production and export. Mutations in this region, the *syp* locus (symbiosis polysaccharide), conferred dramatic colonization defects in the animal as well as defects in biofilm formation in culture (Yip et al. 2005). A connection between these earlier studies was discovered when it was shown that RscS regulates expression of the *syp* locus (Yip et al. 2006). Overexpression of RscS provided a valuable tool in which bacterial colony formation took on a wrinkled or rugose colony morphology that is typical of biofilm formation (Yip et al. 2006). Phenotypes of *rscS* and *syp* alleles in colony-based biofilm assays map closely to their phenotypes during squid colonization, providing a valuable experimental tool for discovery and characterization of biofilm regulation. Further work has identified multiple layers of regulation, including a negative regulatory pathway that includes SypE and SypA, putative matrix proteins that integrate with the polysaccharide matrix, and a unique phosphorelay pathway (Visick 2009, Morris and Visick 2013, Norsworthy and Visick 2015, Ray et al. 2015).

The genetic approaches described above (and in most studies in this review) were conducted in strain ES114, a squid isolate from Kaneohe Bay, Hawaii, that is used widely as a canonical squid symbiont. In addition to the biofilm regulatory pathway, a number of approaches including forward and reverse genetics studies had identified factors in strain ES114 that were important for squid colonization (Stabb and Visick 2013). However, only some *V. fischeri* strains can colonize squid. Therefore, to examine the genetic basis for this host colonization specificity, Mandel and colleagues conducted a comparative genomic analysis of strains ES114 and MJ11, the latter being a fish symbiont that does not colonize squid robustly (Mandel et al. 2009). The study determined that 91 % of ES114 genes were almost identical between the squid and fish symbiont, but that approximately 400 genes in each strain were unique. Analysis of these factors revealed that the squid biofilm regulator, RscS, was encoded in the squid symbiont but not in the fish symbiont. The known RscS target genes, *sypA*, *sypB*, ..., *sypR* were encoded in both genomes and fairly conserved (>85 % amino acid identity). It was known previously that ES114 mutants that lacked RscS were unable to productively colonize the squid (Visick and Skoufos 2001). Therefore, the study asked whether the absence of the regulator could explain the differential colonization phenotype. Introduction of RscS into strain MJ11 was sufficient to allow it to colonize the squid host. Phylogenetic analyses supported a model in which MJ11 was part of an ancestral group of *V. fischeri* that lacked *rscS*, and that this gene was acquired coincident with colonization of squid in the North Pacific Ocean (i.e., Japan and Hawaii).

The idea that a single gene was sufficient to shift the animal hosts available to a bacterium was extreme but consistent with an emerging literature that individual loci could impact microbe-host specificity. Work in entomopathogenic nematodes showed that symbiotic *Xenorhabdus nematophila* requires the three-gene *niLABC* locus for colonization, and that expression of these factors in a heterologous symbiont is sufficient to enable colonization of *Steinernema carpocapsae*, the worm host that otherwise is specific for *X. nematophila* (Cowles and Goodrich-Blair 2008). Small genetic changes in *Yersinia pestis* have been key to its ability to colonize new niches, including single gene acquisitions and even inactivation of a gene already present (Sun et al. 2008, Sun et al. 2014, Zimbl et al. 2015). In the human gut microbiome there are examples in which single gene changes have been critical; e.g., in *Bacteroides fragilis*, polysaccharide A (PSA) confers a key immunomodulatory benefit that cannot be obtained from the other seven capsular polysaccharides produced (Mazmanian et al. 2008).

Studies on host colonization specificity in general, and biofilm formation in particular, have highlighted many of the strengths of the squid model. Imaging *in situ* was key to the initial discovery of the aggregates, forward genetics identified core exopolysaccharide synthetic and regulatory components, comparative genomics revealed the role of this pathway in the evolution and specificity of the association, and high-throughput genetic approaches are identifying additional levels of regulation. Additionally, this work highlights the value of model systems of beneficial bacteria, including *Vibrio* and *Xenorhabdus* models, to identify mechanistic details that resonate in beneficial and pathogenic colonization models.

2.2 The Codeword is TCT

E. scolopes squid provide a particularly dramatic example of a role for bacteria influencing a specific host developmental process. Development of the host tissue proceeds on different trajectories depending on whether the specific symbiont *V. fischeri* is present. Only once the symbiont has colonized, the ciliated appendages of the host light organ undergo apoptosis, hemocyte infiltration, and tissue regression during the subsequent five days (McFall-Ngai and Ruby 1991, Montgomery and McFall-Ngai 1994). The host morphogenesis is striking, with appendages that begin as outstretched mucus factories to recruit colonizing bacteria being reduced to small stumps (Montgomery and McFall-Ngai 1994). As a result, it seems that initiation of the symbiosis is restricted to the first few days of the animal's life while the appendages are present and secreting mucus.

How does the host know that the bacteria are inside to appropriately time the regression? It turns out that *V. fischeri* sheds envelope components that are received by receptors on the host. In particular, the bacterial peptidoglycan fragment, tracheal cytotoxin (TCT)—previously shown to induce a damaging apoptosis in ciliated epithelia upon release from *Bordetella pertussis*—was identified to perform a similar function in *V. fischeri*, but this time with a resulting beneficial outcome (Koropatnick et al. 2004). To recapitulate the apoptosis phenotype observed when intact *V. fischeri* are presented to the host, in the absence of the bacteria both the Lipid A portion of lipopolysaccharide (LPS) and TCT are required. The cell death from these compounds, in conjunction with hemocyte trafficking that is also induced from TCT, results in the regression phenotype. Previously these compounds had only pathogenic associations, but this work underscored a remarkable conservation to the cell biology of microbial-host interactions, emphasizing the context of the interaction to understand the fitness effects on the partners involved (Koropatnick et al. 2004).

Once the bacteria announce their arrival, how does the host speak back? In addition to regression of the appendages that recruit the bacteria, there are additional mechanisms by which the host receives and likely modulates the bacterial signal. Host nitric oxide production, described in more detail below, is diminished as a result of bacterial signaling (synergistically with LPS) (Altura et al. 2011). The host produces a peptidoglycan recognition protein, EsPGRP2, which is secreted into the crypts containing bacteria and has the ability to degrade TCT (Troll et al. 2010). Additionally, there are data to suggest that host alkaline phosphatase, EsAP, modifies Lipid A after the initial signaling (Rader et al. 2012). In each case the host response is to diminish the potency of the bacterial products, but only after they have exerted their influence on host development.

This work in *V. fischeri* was influenced by studies in invertebrate systems that demonstrated host development in response to symbiont colonization and in vertebrates that showed general responses to consortia (reviewed in Montgomery and McFall-Ngai 1994), and itself has influenced a field in which bacterial products play important roles in animal development. An early mammalian example

by Hooper and Gordon demonstrated that in response to colonization by gut *Bacteroidetes* such as *Bacteroides thetaiotaomicron*, terminal tissue differentiation (e.g., fucosylation) is dependent on the presence of the symbiotic bacteria (Hooper and Gordon 2001). There now exist many examples of bacteria directing specific host development. Recent exciting examples include *Algoriphagus machipongonensis* sulfonolipid signaling for multicellular rosette development in the choanoflagellate *Salpingoeca rosetta*, and *Pseudoalteromonas luteoviolacea* phage tail-like structures that stimulate tubeworm metamorphosis (Alegado et al. 2012, Shikuma et al. 2014).

2.3 NO way in

There is a long history of the study of nitric oxide (NO) in eukaryotes, and this small diffusible molecule has been implicated in many different cellular processes including signaling and innate immunity (Fang 2004). Although the roles for NO in eukaryotic physiology and defense against pathogens was discovered many years ago, the study of this compound in the *Vibrio*-squid system and other symbioses (Damiani et al. 2016) has revealed that NO also influences the establishment and maintenance of mutualistic microbe-host relationships as both a signal and a specificity determinant (Wang and Ruby 2011).

Davidson, et al. (Davidson et al. 2004) first demonstrated that NO is produced in squid host tissue through the activity of nitric oxide synthase (NOS), and this activity was attenuated after successful colonization by *V. fischeri*. Using staining and immunocytochemistry, NOS and NO were found located in the epithelium of the light organ, as well as in vesicles within mucus shed from these cells. It is within this mucus that the bacterial cells aggregate prior to entering the light organ. Normally, *V. fischeri* aggregate in the mucus, colonize the host, and after successful colonization NOS activity and NO production are attenuated. Treatment of the animals with an NO-scavenging compound to diminish NO levels allowed large aggregates of non-symbiotic vibrios to form, but these bacteria did not successfully initiate colonization. (Davidson et al. 2004) The results suggested that NO acts as a specificity determinant, helping to limit aggregation of non-symbiotic vibrios and select for symbiotically competent *V. fischeri* from the mixed microbial population found in seawater.

If NO plays a role in specificity, then how do colonizing *V. fischeri* sense and respond to the host-produced NO to successfully establish the partnership? Using genetic approaches it was demonstrated that a strain lacking the NO-detoxifying enzyme flavohemoglobin (Hmp) displayed a colonization deficiency (Poole and Hughes 2000, Wang et al. 2010b). Expression of *hmp* is regulated by the NO-responsive negative regulator NsrR (Rodionov et al. 2005, Tucker et al. 2010). However, NsrR is not the only important NO-sensing regulator in *V. fischeri*. H-NOX, a heme NO/oxygen-binding protein, also plays a role in symbiotically relevant NO-responsive regulation of genes in *V. fischeri* (Wang et al. 2010a). Although H-NOX-like proteins are widely distributed in bacteria, this was the first report describing bacterial H-NOX function. Interestingly, it appears that one role for H-NOX in *V. fischeri* is to sense NO and correspondingly suppress bacterial hemin uptake during the early stages of host colonization. The authors predicted that early repression of iron uptake would protect the cells from the potentially harmful effects of Fenton chemistry when they are exposed to host-generated oxidants (Davidson et al. 2004, Graf and Ruby 2000, Wang et al. 2010a). Consistent with this model, hemin uptake genes in *V. fischeri* were shown to be induced during the later stages of symbiotic colonization, and deletion of these genes negatively impacted colonization (Septer et al. 2011). Together, these studies support a model whereby host NO stimulates repression of hemin uptake genes; once bacterial colonization leads to an attenuation of host oxidant production, then hemin uptake genes are derepressed to support growth in the iron-limited light organ environment.

Therefore, the ability to sense and detoxify NO is important for symbiotic specificity, and NO acts as a temporal signal to modulate bacterial gene expression and promote successful colonization.

Although these studies have led to a better understanding of the role of a few key proteins and regulators in the response of *V. fischeri* to NO and the initial stages of the symbiosis, there is much yet to be learned about the global effects of NO on *V. fischeri* gene expression and metabolism, how this molecule acts as a specificity determinant, and whether there is a role for NO in the mature symbiosis. For example, the work of Wier et al. has suggested that NO may play a role in the daily symbiotic rhythm in the adult animal (Wier et al. 2010). Their data predicted that nitrate/nitrite respiration is used by the bacterial symbionts throughout the daylight hours. Similarly to *Escherichia coli* (Vine and Cole 2011), it is predicted that NO is produced by *V. fischeri* during respiration of nitrate/nitrite in laboratory culture. Endogenously-produced NO could induce alternative respiratory pathways that likely influence the physiology and metabolism of the bacterium (Dunn et al. 2010). Together these separate lines of evidence suggest that NO may play a role beyond signaling and selection in the initiation of the symbiotic relationship. In the future it will be exciting to combine studies of NO and the bacterial NO response with the more recently developed ability to rear squid to adulthood (Koch et al. 2013; see section below on light production).

The value of further studies of NO in the *Vibrio*-squid system lie not only in providing important information about the role of this molecule in beneficial host-microbe interactions, but also for comparative studies to host-pathogen responses. Our current understanding supports a view that NO is being produced by the host and sensed by the bacteria in similar ways in many of the studied host-microbial interactions, whether the outcome of the relationship is beneficial or detrimental (Fang 2004, Wang and Ruby 2011). The prevalence of NO in host tissues colonized by bacteria suggests that a better understanding of the role of NO in symbiosis may have wide-reaching consequences for microbes at the interface of health and disease.

2.4 Swimming against the flow

In the mucus field that serves as the entry point for bacteria heading into the host, colonizing bacteria enter at one of three pores on either side of the bilaterally symmetrical light organ. Mucus is shed from the pores of the host at the same time that *V. fischeri* aggregates in that mucus. The bacteria proceed to migrate toward the pores, and each aggregate swims into a pore to colonize the ducts and crypts of the host. How do colonizing bacteria travel against this powerful flow? A key role for flagellar motility was identified over twenty years ago (Ruby and Asato 1993). In that work Ruby and Asato confirmed that planktonic *V. fischeri* were motile due to a polar tuft of sheathed flagella. However, by 24 hours-post-inoculation most cells in the light organ crypts were non-flagellated. Upon expulsion of bacteria from the host, the bacteria regrow their flagella in 45-60 min even in nutrient-deplete seawater (Ruby and Asato 1993). Therefore, the bacterial life cycle alternates between a motile planktonic lifestyle and a non-flagellated crypt-colonized state.

Significant details have since been elucidated about the molecular mechanisms that control flagellar development in *V. fischeri*, which in turn has solidified the importance of swimming motility for squid colonization. Random transposon mutagenesis provided evidence that nonmotile mutants could not colonize (Graf et al. 1994), and reverse genetics revealed that mutants defective for flagellar motility or chemotaxis did not establish productive colonization with the squid host (Millikan and Ruby 2003, Millikan and Ruby 2004, DeLoney-Marino and Visick 2012). Together these studies established a model of a hierarchy of flagellar gene expression in *V. fischeri* controlled by the σ^{54} -

dependent regulator FlrA. There is evidence for regulation by quorum sensing and by magnesium, and other sensory inputs are likely (O'Shea et al. 2005, Cao et al. 2012).

Bacterial flagellar motility often occurs in a directed fashion in which rotation of the flagellar bundle results in net movement toward preferred nutrient sources. Given the above information that chemotaxis was required for colonization, it seemed likely that the bacteria were swimming toward a host compound. The first evidence for chitin oligosaccharides as the specific attractant was obtained when addition of exogenous chitobiose, the *N*-acetylglucosamine dimer, blocked colonization, whereas the monomer did not have such an effect (Mandel et al. 2012). Given that *N*-acetylglucosamine is abundant on eukaryotic cell surfaces, yet chitin and its breakdown oligosaccharides are more specialized in their localization, it seemed possible that oligosaccharides may be a specific cue to direct entry into the host crypts. Mutants defective for chemotaxis remained at the outer face of the light organ pore, the same stage at which wild-type *V. fischeri* arrested their symbiotic development in the presence of added chitin oligosaccharides (Mandel et al. 2012). These results strongly suggested that host chitin served as a signal for the bacteria to enter the pore. Direct imaging revealed the presence of insoluble chitin bound to hemocytes within the host (Mandel et al. 2012, Heath-Heckman and McFall-Ngai 2011), which may be released through the action of a host endochitinase (Kremer et al. 2013). Together, this illustrates a specific colonization checkpoint that is regulated by both host and symbiont factors.

Work on bacterial motility at the host interface has provided a valuable toolset to probe mechanisms of symbiosis and reveal novel signaling pathways. Many bacterial strains have dozens of genes that encode chemotactic sensory proteins, the methyl-accepting chemotaxis proteins (MCPs). The set of 43 MCPs in *V. fischeri* is typical in this regard, and despite difficulties in studying a large protein family, functions have now been assigned to three of these proteins. VfcA is the major amino acid chemoreceptor, and VfcB and VfcB2 are fatty acid chemoreceptors (Brennan et al. 2013, Nikolakakis et al. 2016). In addition to providing information directly about colonization, these tools provided insight into the role of LPS during colonization and for the evolution and the generation of torque at the flagellar motor (Post et al. 2012, Beeby et al. 2016). Furthermore, recent work suggests that the rotation of the flagella—which is enclosed in an LPS sheath—stimulates outer membrane vesicle release and triggers the host immune response by promoting LPS release (Brennan et al. 2014, Aschtgen et al. 2016).

Satisfying answers to some of these questions are beginning to be addressed, including a role for cilia in modulating adhesion, as well as chemotaxis toward host-produced and host-cleaved chitin modulating a key developmental checkpoint. Still, important questions remain that suggest novel and interesting biology to be revealed through the symbiosis. Open questions include how bacteria transit through the mucus in a flagellar-independent manner; the molecular basis of chitin oligosaccharide sensing in the symbiont; and the processes that regulate the developmental switch between the aflagellate state in the host versus the swimming state in seawater.

2.5 Light up my life

An important aspect to mutualistic symbioses is the selection of appropriate and cooperative partners. In both the rhizobium-leguminous plant (Kiers et al. 2003) and *Vibrio*-squid symbioses the microbial partners provide costly services to their hosts (nitrogen fixation and light production, respectively). In theory, these relationships could be exploited by symbionts that are less cooperative (i.e. “cheaters”) (Ghoul et al. 2014). However, it is rare to find bacterial symbionts associated with the hosts that do not provide these services. Therefore, the *Vibrio*-squid mutualism provides an excellent model

302 system for studying cooperative partner stability, and studies to date indicate that bacterial light
303 production is required for bacterial cells to persist in the light organ.

304 *V. fischeri* is known to produce light in the squid host, and a key study demonstrated a role for
305 luciferase, the enzyme that produces light, in bacterial symbiotic persistence (Visick et al. 2000).
306 Mutants with defective luminescence structural genes or luminescence regulatory genes colonized
307 juvenile squid to the same levels as wild type in the first 24 hours. However, by 48 hours there was a
308 three- to four-fold reduction in colonization by the dark mutants relative to wild-type controls. In
309 squid co-colonized with both a luminescence mutant and wild type, levels of the mutant strains
310 similarly decreased, indicating that light-producing wild-type cells in the light organ could not
311 complement the colonization defect of the light-deficient cells. These results suggested that the
312 ability of individual bacteria to produce light was important for persistence in the light organ, and
313 that somehow non-luminescent cells are selected against during development of the symbiosis.

314 Interestingly, the light-deficient strains have a specific effect on host development. Although
315 colonization by a luminescence mutant still triggered apoptosis-related developmental changes in the
316 ciliated surface of the light organ, colonization of the tissue by these strains no longer increased cell
317 swelling of the epithelial cells lining the light organ crypt spaces. Therefore, light production
318 appeared to play a specific role in host developmental pathways. Notably, this was the first report of
319 *V. fischeri* genes required for induction of bacterial-triggered differentiation of host tissue (Visick et
320 al. 2000).

321 It was later discovered that the antibiotic markers and method for constructing the early luminescence
322 mutants (Visick et al. 2000) resulted in colonization attenuation and pleiotropic effects. In a later
323 study, newly developed genetic tools were used to construct luminescence mutants that were not
324 negatively affected in growth and colonization (Bose et al. 2008). Using these strains, the early
325 results were confirmed demonstrating that the strain lacking the luminescence structural genes
326 displayed a four-fold reduction in colonization as compared to wild type at 48 hours-post-inoculation.

327 Previous studies suggested that maintenance of the symbiosis over the life of the animal requires a
328 maturation process of several weeks (Montgomery and McFall-Ngai 1998), leaving the question of
329 how production of light influences symbiosis maturation beyond 72 hours. A major breakthrough for
330 the field came with the development of protocols for simplified rearing of newly-hatched juvenile
331 squid through and beyond the maturation process. These methods allowed investigation of how
332 bacterial-produced light affects the development of the symbiosis over four weeks (Koch et al. 2013).
333 In these studies, the levels of the luminescence-deficient mutant associated with the squid light organ
334 continued to diminish over time, to the minimum level of detection after 28 days. Similar results
335 were observed in squid colonized with mixed inocula containing both wild type and the luminescence
336 mutant, where after 15 days the mutant was barely detected. Therefore the persistence defect
337 observed during early colonization becomes more pronounced as the symbiosis matures, with
338 eventual loss (or near loss) of non-luminescent strains in a matter of weeks.

339 Luminescence regulation is one of the hallmarks of the *V. fischeri*-squid symbiosis and has been
340 studied intensively, yet there are still exciting open questions. First, how are the dark mutants
341 removed from the population even in the midst of neighboring bright populations? A clue comes
342 from studies testing the influence of a previous colonization event on recolonization (Koch et al.
343 2013). Juvenile animals were colonized with either wild type or a luminescence mutant. After 1-5
344 days the animals were treated with antibiotics to clear bacteria from the light organ and then exposed
345 again to wild-type *V. fischeri* to test whether light production is a “signal” to the host that influences

symbiotic maturation. Animals treated with antibiotics after one day were readily recolonized, regardless of the strain that initially colonized. However, after five days, wild-type *V. fischeri* induced a refractory state in the animal that prevented recolonization. In contrast, in animals initially colonized by a luminescence mutant, greater than 80% of the animals were recolonized by wild type. These results support the idea that the host is detecting light production by bacterial cells and/or is altering physiological conditions to sanction the non-luminescent strains. In addition, the host apparently is able to “eject” an inappropriate light deficient strain, while allowing future recolonization by a symbiotically appropriate light-producing strain. The exact mechanisms by which the detection, sanctioning, and/or ejection occurs remain to be described.

A second interesting question relates to how bacterial light production is matched to the moonlight in such an exquisite fashion. The squid contains elaborate tissues to physically reflect and modulate bacterial light production (Crookes et al. 2004). This physical response could be triggered through the activity of products of host cryptochrome and eye-specification genes; the expression of these genes appears to be influenced by the light produced by *V. fischeri* (Peyer et al. 2014, Heath-Heckman et al. 2013). The physical reflection and modulation of bacterial luminescence is also coordinated with a molecular signaling response. For example, host epithelial cells swell in response to light-producing strains but not dark mutants (Visick et al. 2000). This swelling could release chemical cues into the light organ environment. Recent evidence indicates that bacterial luminescence in the light organ is controlled not only through quorum sensing, but also through response to environmental signaling (Septer and Stabb 2012). These results suggest there is complex chemical and physical control of light production in the symbiosis. Bacterial luminescence is a particularly intriguing and engaging aspect of the *Vibrio*-squid symbiosis, and it is clear that there are abundant questions remaining to be addressed as to how the interaction with the host and the environment lead to specific phenotypic output in the host.

2.6 Nice to meet you... now what is it you do?

The *Vibrio*-squid symbiosis has provided a useful framework for identifying the function of bacterial genes and studying novel genes *in vivo*. Due to the wealth of genetic tools that have been developed for *V. fischeri* and the ability to access the host interface with direct imaging, it is possible to test the effects of gene loss in the real-world environment of the host. Two examples discussed below are using the *Vibrio*-squid system to broaden understanding of gene function for alternative oxidase (AOX) and for discovering the role of the biofilm inhibitor BinK.

AOX is a terminal respiratory oxidase that is ubiquitous in plants, and is unusual because its activity is not directly linked to generation of the proton motive force (Vanlerberghe and McIntosh 1997). The study of the function of AOX in plants is an active area of research, and AOX function has been linked to both abiotic and biotic stress responses (Vanlerberghe 2013). Only with the explosion of genome and metagenome sequencing was it discovered that certain bacterial genomes also encode this protein (Stenmark and Nordlund 2003), and that *aox*-like genes are abundant in metagenomic sequences from ocean surface waters (McDonald and Vanlerberghe 2005). However, early progress towards understanding the physiological benefit of AOX function in bacteria was limited by the lack of genetic tools for many of the AOX-encoding organisms. A path to revealing a functional role for AOX came with the discovery that the genome of *V. fischeri* strain ES114 encoded AOX (Ruby et al. 2005). A transcriptomic analysis of the *V. fischeri* response to NO revealed that nitric oxide induces expression of *aox* (Wang et al. 2010a). The connection to NO was further clarified through characterization of the role of the NO-responsive negative regulator NsrR in regulation of *aox* expression, and identification of the ability of *V. fischeri* AOX to function as an NO-resistant oxidase

(Dunn et al. 2010). Despite the known connections between *aox* and NO, and between NO and the early stages of host colonization, no discernible phenotypic difference between the *aox* mutant and wild type in early colonization of the squid host has been observed. Although there is the possibility that AOX does not play a role in bacterial physiology during host colonization, an alternative explanation is that the benefit of AOX expression during colonization does not result in a phenotype dramatic enough to be detected in the short time frame of the experiments (1-3 days). Experiments to test this possibility are in progress and would be consistent with studies above described for luminescence mutants in which colonization phenotypes change over the course of symbiosis and effects are magnified over a multi-week time course.

Studying AOX regulation and function in *V. fischeri* as a model organism will provide a framework for understanding how bacteria in ocean surface waters utilize this respiratory pathway in growth and survival. Work is underway to clarify the physiological benefit of AOX function in *V. fischeri* and other *aox*-containing bacteria, with the ultimate goal of better understanding how bacteria cope with changing conditions in the environment. Studying AOX in the context of the symbiosis has provided insight into the expression and function of this interesting protein, and provides a framework for broad studies of how AOX function influences bacterial physiology in the environment.

Study of AOX followed a reverse-genetic approach, starting with identification of an interesting gene through genome sequencing, and through directed experimental approaches leading to a better understanding of gene function. However, in many cases forward genetic approaches have identified genes whose products are relevant for a specific colonization process. An excellent example is *binK*, which encodes a histidine kinase. Above we described a key role for biofilm formation in the colonization process as regulated by RscS and Syp. In a recent global genetic screen for mutants with an advantage in squid colonization, *binK* was identified as a locus that when disrupted resulted in substantially better colonization of the *V. fischeri* strain (Brooks and Mandel 2016). Typical means to predict protein function (e.g., homology, neighboring genes) were not helpful, so phenotypes of cells lacking *binK* were examined in culture and in the host and revealed a substantial increase in symbiotic biofilm formation. BinK (biofilm inhibitor kinase) is therefore a negative regulator of biofilm formation and an additional membrane-bound histidine kinase that is critical for proper regulation of the Syp biofilm.

In the case of both AOX and BinK, the depth of the *V. fischeri*-squid system has provided a means to assign function to novel and poorly-understood proteins. A striking number of genes are poorly understood in bacterial genomes, exemplified by the 149 (32 %) of the minimal 473 genes in the JCVI-syn3.0 genome with functions that remain to be discovered (Hutchison et al. 2016). The ability to study biological function in the context of the host thus provides a useful lens through which to identify and characterize genes and their products.

3 Conclusions

The *Vibrio*-squid system has proven to be a valuable study system for identifying principles of microbe-host interaction, continues to serve as a fertile field for discovery, and provides a useful road map for moving from patterns of intriguing phenotypes to discerning the molecular communication between microbe and host that is responsible for those patterns. By integrating approaches in genetics, genomics, molecular biology, imaging, physiology, evolutionary biology, and cell biology, each of the topic areas highlights an integrated and mechanistic view of how symbiotic partners functionally communicate in a model microbiome. In this manner, the *Vibrio*-squid system provides

a durable example for how to move from fascinating observations to molecular understanding of the processes by which very different organisms communicate and establish a productive partnership.

4 Conflict of Interest

The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

5 Author Contributions

MJM and AKD wrote the manuscript.

6 Funding

Research in the authors' laboratories is supported by National Science Foundation awards IOS-1456963 (MJM) and MCB-1050687 (AKD), and National Institutes of Health Awards R35GM119627 (MJM) and R21AI117262 (MJM). The content is solely the responsibility of the authors and does not necessarily represent the official views of the funding agencies.

7 Acknowledgments

We thank Ella Rotman and Denise Tarnowski for comments on the manuscript.

8 References

- Alegado, R. A., Brown, L. W., Cao, S., Dermenjian, R. K., Zuzow, R., Fairclough, S. R., et al. (2012). A bacterial sulfonolipid triggers multicellular development in the closest living relatives of animals. *Elife* 1, e00013. doi:[10.7554/eLife.00013](https://doi.org/10.7554/eLife.00013).
- Altura, M. A., Heath-Heckman, E. A. C., Gillette, A., Kremer, N., Krachler, A. M., Brennan, C., et al. (2013). The first engagement of partners in the *Euprymna scolopes*-*Vibrio fischeri* symbiosis is a two-step process initiated by a few environmental symbiont cells. *Environ Microbiol* 15, 2937–2950. doi:[10.1111/1462-2920.12179](https://doi.org/10.1111/1462-2920.12179).
- Altura, M. A., Stabb, E., Goldman, W., Apicella, M., and McFall-Ngai, M. J. (2011). Attenuation of host NO production by MAMPs potentiates development of the host in the squid-vibrio symbiosis. *Cell Microbiol* 13, 527–537. doi:[10.1111/j.1462-5822.2010.01552.x](https://doi.org/10.1111/j.1462-5822.2010.01552.x).
- Aschtgen, M.-S., Lynch, J. B., Koch, E., Schwartzman, J., McFall-Ngai, M., and Ruby, E. (2016). Rotation of *Vibrio fischeri* Flagella Produces Outer Membrane Vesicles That Induce Host Development. *J. Bacteriol.* 198, 2156–2165. doi:[10.1128/JB.00101-16](https://doi.org/10.1128/JB.00101-16).
- Beeby, M., Ribardo, D. A., Brennan, C. A., Ruby, E. G., Jensen, G. J., and Hendrixson, D. R. (2016). Diverse high-torque bacterial flagellar motors assemble wider stator rings using a conserved protein scaffold. *Proc. Natl. Acad. Sci. U.S.A.* 113, E1917–1926. doi:[10.1073/pnas.1518952113](https://doi.org/10.1073/pnas.1518952113).
- Bose, J. L., Rosenberg, C. S., and Stabb, E. V. (2008). Effects of luxCDABEG induction in *Vibrio fischeri*: Enhancement of symbiotic colonization and conditional attenuation of growth in culture. *Arch Microbiol* 190, 169–183. doi:[10.1007/s00203-008-0387-1](https://doi.org/10.1007/s00203-008-0387-1).

- 468 Brennan, C. A., Hunt, J. R., Kremer, N., Krasity, B. C., Apicella, M. A., McFall-Ngai, M. J., et al.
469 (2014). A model symbiosis reveals a role for sheathed-flagellum rotation in the release of
470 immunogenic lipopolysaccharide. *Elife* 3, e01579. doi:[10.7554/eLife.01579.019](https://doi.org/10.7554/eLife.01579.019).
- 471 Brennan, C. A., Mandel, M. J., Gyllborg, M. C., Thomasgard, K. A., and Ruby, E. G. (2013). Genetic
472 determinants of swimming motility in the squid light-organ symbiont *Vibrio fischeri*.
473 *MicrobiologyOpen* 2, 576–594. doi:[10.1002/mbo3.96](https://doi.org/10.1002/mbo3.96).
- 474 Brooks, J. F., and Mandel, M. J. (2016). The histidine kinase BinK is a negative regulator of biofilm
475 formation and squid colonization. *Journal of Bacteriology*, JB.00037–16. doi:[10.1128/JB.00037-16](https://doi.org/10.1128/JB.00037-16).
- 476 Cao, X., Studer, S. V., Wassarman, K., Zhang, Y., Ruby, E. G., and Miyashiro, T. (2012). The Novel
477 Sigma Factor-Like Regulator RpoQ Controls Luminescence, Chitinase Activity, and Motility in
478 *Vibrio fischeri*. *mBio* 3. doi:[10.1128/mBio.00285-11](https://doi.org/10.1128/mBio.00285-11).
- 479 Cowles, C. E., and Goodrich-Blair, H. (2008). The *Xenorhabdus nematophila* *niABC* genes confer
480 the ability of *Xenorhabdus* spp. to colonize *Steinernema carpocapsae* nematodes. *J Bacteriol* 190,
481 4121–4128. doi:[10.1128/JB.00123-08](https://doi.org/10.1128/JB.00123-08).
- 482 Crookes, W. J., Ding, L.-L., Huang, Q. L., Kimbell, J. R., Horwitz, J., and McFall-Ngai, M. J.
483 (2004). Reflectins: The unusual proteins of squid reflective tissues. *Science* 303, 235–238.
484 doi:[10.1126/science.1091288](https://doi.org/10.1126/science.1091288).
- 485 Damiani, I., Pauly, N., Puppo, A., Brouquisse, R., and Boscari, A. (2016). Reactive Oxygen Species
486 and Nitric Oxide Control Early Steps of the Legume - Rhizobium Symbiotic Interaction. *Front Plant*
487 *Sci* 7, 454. doi:[10.3389/fpls.2016.00454](https://doi.org/10.3389/fpls.2016.00454).
- 488 Davidson, S. K., Koropatnick, T. A., Kossmehl, R., Sycuro, L., and McFall-Ngai, M. J. (2004). NO
489 means 'yes' in the squid-vibrio symbiosis: Nitric oxide (NO) during the initial stages of a beneficial
490 association. *Cell Microbiol* 6, 1139–1151. doi:[10.1111/j.1462-5822.2004.00429.x](https://doi.org/10.1111/j.1462-5822.2004.00429.x).
- 491 DeLoney-Marino, C. R., and Visick, K. L. (2012). Role for *cheR* of *Vibrio fischeri* in the *Vibrio*-
492 squid symbiosis. *Can J Microbiol* 58, 29–38. doi:[10.1139/w11-107](https://doi.org/10.1139/w11-107).
- 493 Dunn, A. K., Karr, E. A., Wang, Y., Batton, A. R., Ruby, E. G., and Stabb, E. V. (2010). The
494 alternative oxidase (AOX) gene in *Vibrio fischeri* is controlled by *NsrR* and upregulated in response
495 to nitric oxide. *Mol. Microbiol.* 77, 44–55. doi:[10.1111/j.1365-2958.2010.07194.x](https://doi.org/10.1111/j.1365-2958.2010.07194.x).
- 496 Fang, F. C. (2004). Antimicrobial reactive oxygen and nitrogen species: Concepts and controversies.
497 *Nat. Rev. Microbiol.* 2, 820–832. doi:[10.1038/nrmicro1004](https://doi.org/10.1038/nrmicro1004).
- 498 Ghou, M., Griffin, A. S., and West, S. A. (2014). Toward an evolutionary definition of cheating.
499 *Evolution* 68, 318–331. doi:[10.1111/evo.12266](https://doi.org/10.1111/evo.12266).
- 500 Graf, J., and Ruby, E. G. (2000). Novel effects of a transposon insertion in the *Vibrio fischeri* *glnD*
501 gene: Defects in iron uptake and symbiotic persistence in addition to nitrogen utilization. *Mol*
502 *Microbiol* 37, 168–179. Available at: [http://onlinelibrary.wiley.com/doi/10.1046/j.1365-](http://onlinelibrary.wiley.com/doi/10.1046/j.1365-2958.2000.01984.x/abstract)
503 [2958.2000.01984.x/abstract](http://onlinelibrary.wiley.com/doi/10.1046/j.1365-2958.2000.01984.x/abstract).

- 504 Graf, J., Dunlap, P. V., and Ruby, E. G. (1994). Effect of transposon-induced motility mutations on
505 colonization of the host light organ by *Vibrio fischeri*. *J Bacteriol* 176, 6986–6991. Available at:
506 <http://jb.asm.org/cgi/reprint/176/22/6986?view=long&pmid=7961462>.
- 507 Heath-Heckman, E. A. C., and McFall-Ngai, M. J. (2011). The occurrence of chitin in the hemocytes
508 of invertebrates. *Zoology (Jena)* 114, 191–198. doi:[10.1016/j.zool.2011.02.002](https://doi.org/10.1016/j.zool.2011.02.002).
- 509 Heath-Heckman, E. A. C., Peyer, S. M., Whistler, C. A., Apicella, M. A., Goldman, W. E., and
510 McFall-Ngai, M. J. (2013). Bacterial bioluminescence regulates expression of a host cryptochrome
511 gene in the squid-*Vibrio* symbiosis. *MBio* 4. doi:[10.1128/mBio.00167-13](https://doi.org/10.1128/mBio.00167-13).
- 512 Hooper, L. V., and Gordon, J. I. (2001). Glycans as legislators of host-microbial interactions:
513 Spanning the spectrum from symbiosis to pathogenicity. *Glycobiology* 11, 1R–10R. Available at:
514 <http://glycob.oxfordjournals.org/cgi/content/full/11/2/1R?view=long&pmid=11287395>.
- 515 Human Microbiome Project Consortium (2012). Structure, function and diversity of the healthy
516 human microbiome. *Nature* 486, 207–214. doi:[10.1038/nature11234](https://doi.org/10.1038/nature11234).
- 517 Hutchison, C. A., Chuang, R.-Y., Noskov, V. N., Assad-Garcia, N., Deerinck, T. J., Ellisman, M. H.,
518 et al. (2016). Design and synthesis of a minimal bacterial genome. *Science* 351, aad6253.
519 doi:[10.1126/science.aad6253](https://doi.org/10.1126/science.aad6253).
- 520 Jones, B., and Nishiguchi, M. (2004). Counterillumination in the Hawaiian bobtail squid, *Euprymna*
521 *scolopes* Berry (Mollusca: Cephalopoda). *Marine Biol* 144, 1151–1155. Available at:
522 <http://www.springerlink.com/index/DFVKDCWEXPMC28DC.pdf>.
- 523 Kiers, E. T., Rousseau, R. A., West, S. A., and Denison, R. F. (2003). Host sanctions and the legume-
524 rhizobium mutualism. *Nature* 425, 78–81. doi:[10.1038/nature01931](https://doi.org/10.1038/nature01931).
- 525 Koch, E. J., Miyashiro, T., McFall-Ngai, M. J., and Ruby, E. G. (2013). Features governing symbiont
526 persistence in the squid-vibrio association. *Mol Ecol* 23, 1624–1634. doi:[10.1111/mec.12474](https://doi.org/10.1111/mec.12474).
- 527 Koropatnick, T. A., Engle, J. T., Apicella, M. A., Stabb, E. V., Goldman, W. E., and McFall-Ngai,
528 M. J. (2004). Microbial factor-mediated development in a host-bacterial mutualism. *Science* 306,
529 1186–1188. doi:[10.1126/science.1102218](https://doi.org/10.1126/science.1102218).
- 530 Kremer, N., Philipp, E. E. R., Carpentier, M.-C., Brennan, C. A., Kraemer, L., Altura, M. A., et al.
531 (2013). Initial Symbiont Contact Orchestrates Host-Organ-wide Transcriptional Changes that Prime
532 Tissue Colonization. *Cell Host Microbe* 14, 183–194. doi:[10.1016/j.chom.2013.07.006](https://doi.org/10.1016/j.chom.2013.07.006).
- 533 Mandel, M. J. (2010). Models and approaches to dissect host-symbiont specificity. *Trends Microbiol*
534 18, 504–511. doi:[10.1016/j.tim.2010.07.005](https://doi.org/10.1016/j.tim.2010.07.005).
- 535 Mandel, M. J., Schaefer, A. L., Brennan, C. A., Heath-Heckman, E. A. C., DeLoney-Marino, C. R.,
536 McFall-Ngai, M. J., et al. (2012). Squid-derived chitin oligosaccharides are a chemotactic signal
537 during colonization by *Vibrio fischeri*. *Appl Environ Microbiol* 78, 4620–4626.
538 doi:[10.1128/AEM.00377-12](https://doi.org/10.1128/AEM.00377-12).

- 539 Mandel, M. J., Wollenberg, M. S., Stabb, E. V., Visick, K. L., and Ruby, E. G. (2009). A single
540 regulatory gene is sufficient to alter bacterial host range. *Nature* 458, 215–218.
541 doi:[10.1038/nature07660](https://doi.org/10.1038/nature07660).
- 542 Mazmanian, S. K., Round, J. L., and Kasper, D. L. (2008). A microbial symbiosis factor prevents
543 intestinal inflammatory disease. *Nature* 453, 620–625. doi:[10.1038/nature07008](https://doi.org/10.1038/nature07008).
- 544 McDonald, A. E., and Vanlerberghe, G. C. (2005). Alternative oxidase and plastoquinol terminal
545 oxidase in marine prokaryotes of the Sargasso Sea. *Gene* 349, 15–24.
546 doi:[10.1016/j.gene.2004.12.049](https://doi.org/10.1016/j.gene.2004.12.049).
- 547 McFall-Ngai, M. J., and Ruby, E. G. (1991). Symbiont recognition and subsequent morphogenesis as
548 early events in an animal-bacterial mutualism. *Science* 254, 1491–1494. Available at:
549 [http://eutils.ncbi.nlm.nih.gov/entrez/eutils/elink.fcgi?dbfrom=pubmed&id=1962208&retmode=ref&c](http://eutils.ncbi.nlm.nih.gov/entrez/eutils/elink.fcgi?dbfrom=pubmed&id=1962208&retmode=ref&cmd=prlinks)
550 [md=prlinks](http://eutils.ncbi.nlm.nih.gov/entrez/eutils/elink.fcgi?dbfrom=pubmed&id=1962208&retmode=ref&cmd=prlinks).
- 551 McFall-Ngai, M., Hadfield, M. G., Bosch, T. C. G., Carey, H. V., Domazet-Loso, T., Douglas, A. E.,
552 et al. (2013). Animals in a bacterial world, a new imperative for the life sciences. *Proc Natl Acad Sci*
553 *USA*. doi:[10.1073/pnas.1218525110](https://doi.org/10.1073/pnas.1218525110).
- 554 Millikan, D. S., and Ruby, E. G. (2003). FlrA, a σ ⁵⁴-dependent transcriptional activator in *Vibrio*
555 *fischeri*, is required for motility and symbiotic light-organ colonization. *J Bacteriol* 185, 3547–3557.
556 Available at:
557 [http://eutils.ncbi.nlm.nih.gov/entrez/eutils/elink.fcgi?dbfrom=pubmed&id=12775692&retmode=ref&](http://eutils.ncbi.nlm.nih.gov/entrez/eutils/elink.fcgi?dbfrom=pubmed&id=12775692&retmode=ref&cmd=prlinks)
558 [cmd=prlinks](http://eutils.ncbi.nlm.nih.gov/entrez/eutils/elink.fcgi?dbfrom=pubmed&id=12775692&retmode=ref&cmd=prlinks).
- 559 Millikan, D. S., and Ruby, E. G. (2004). *Vibrio fischeri* flagellin A is essential for normal motility
560 and for symbiotic competence during initial squid light organ colonization. *J Bacteriol* 186, 4315–
561 4325. doi:[10.1128/JB.186.13.4315-4325.2004](https://doi.org/10.1128/JB.186.13.4315-4325.2004).
- 562 Montgomery, M. K., and McFall-Ngai, M. (1994). Bacterial symbionts induce host organ
563 morphogenesis during early postembryonic development of the squid *Euprymna scolopes*.
564 *Development* 120, 1719–1729.
- 565 Montgomery, M. K., and McFall-Ngai, M. J. (1998). Late postembryonic development of the
566 symbiotic light organ of *Euprymna scolopes* (Cephalopoda: Sepiolidae). *Biol. Bull.* 195, 326–336.
- 567 Morris, A. R., and Visick, K. L. (2013). The response regulator SypE controls biofilm formation and
568 colonization through phosphorylation of the syp-encoded regulator SypA in *Vibrio fischeri*. *Mol*
569 *Microbiol* 87, 509–525. doi:[10.1111/mmi.12109](https://doi.org/10.1111/mmi.12109).
- 570 Nikolakakis, K., Monfils, K., Moriano-Gutierrez, S., Brennan, C. A., and Ruby, E. G. (2016).
571 Characterization of the *Vibrio fischeri* Fatty Acid Chemoreceptors, VfcB and VfcB2. *Applied and*
572 *Environmental Microbiology* 82, 696–704. doi:[10.1128/AEM.02856-15](https://doi.org/10.1128/AEM.02856-15).
- 573 Norsworthy, A. N., and Visick, K. L. (2015). Signaling between two interacting sensor kinases
574 promotes biofilms and colonization by a bacterial symbiont. *Mol Microbiol* 96, 233–248.
575 doi:[10.1111/mmi.12932](https://doi.org/10.1111/mmi.12932).

- 576 Nyholm, S. V., and McFall-Ngai, M. J. (2003). Dominance of *Vibrio fischeri* in secreted mucus
577 outside the light organ of *Euprymna scolopes*: The first site of symbiont specificity. *Appl Environ*
578 *Microbiol* 69, 3932–3937. Available at:
579 <http://aem.asm.org/cgi/content/full/69/7/3932?view=long&pmid=12839763>.
- 580 Nyholm, S. V., Stabb, E. V., Ruby, E. G., and McFall-Ngai, M. J. (2000). Establishment of an
581 animal-bacterial association: Recruiting symbiotic vibrios from the environment. *Proc Natl Acad Sci*
582 *USA* 97, 10231–10235. Available at:
583 [http://eutils.ncbi.nlm.nih.gov/entrez/eutils/elink.fcgi?dbfrom=pubmed&id=10963683&retmode=ref&](http://eutils.ncbi.nlm.nih.gov/entrez/eutils/elink.fcgi?dbfrom=pubmed&id=10963683&retmode=ref&cmd=prlinks)
584 [cmd=prlinks](http://eutils.ncbi.nlm.nih.gov/entrez/eutils/elink.fcgi?dbfrom=pubmed&id=10963683&retmode=ref&cmd=prlinks).
- 585 O’Shea, T. M., DeLoney-Marino, C. R., Shibata, S., Aizawa, S.-I., Wolfe, A. J., and Visick, K. L.
586 (2005). Magnesium promotes flagellation of *Vibrio fischeri*. *J Bacteriol* 187, 2058–2065.
587 doi:[10.1128/JB.187.6.2058-2065.2005](https://doi.org/10.1128/JB.187.6.2058-2065.2005).
- 588 Peyer, S. M., Pankey, M. S., Oakley, T. H., and McFall-Ngai, M. J. (2014). Eye-specification genes
589 in the bacterial light organ of the bobtail squid *Euprymna scolopes*, and their expression in response
590 to symbiont cues. *Mech. Dev.* 131, 111–126. doi:[10.1016/j.mod.2013.09.004](https://doi.org/10.1016/j.mod.2013.09.004).
- 591 Poole, R. K., and Hughes, M. N. (2000). New functions for the ancient globin family: Bacterial
592 responses to nitric oxide and nitrosative stress. *Mol. Microbiol.* 36, 775–783.
- 593 Post, D. M. B., Yu, L., Krasity, B. C., Choudhury, B., Mandel, M. J., Brennan, C. A., et al. (2012).
594 O-antigen and core carbohydrate of *Vibrio fischeri* lipopolysaccharide: Composition and analysis of
595 their role in *Euprymna scolopes* light organ colonization. *J Biol Chem* 287, 8515–8530.
596 doi:[10.1074/jbc.M111.324012](https://doi.org/10.1074/jbc.M111.324012).
- 597 Rader, B. A., Kremer, N., Apicella, M. A., Goldman, W. E., and McFall-Ngai, M. J. (2012).
598 Modulation of symbiont lipid a signaling by host alkaline phosphatases in the squid-*Vibrio*
599 symbiosis. *mBio* 3. doi:[10.1128/mBio.00093-12](https://doi.org/10.1128/mBio.00093-12).
- 600 Ray, V. A., Driks, A., and Visick, K. L. (2015). Identification of a novel matrix protein that promotes
601 biofilm maturation in *Vibrio fischeri*. *J Bacteriol* 197, 518–528. doi:[10.1128/JB.02292-14](https://doi.org/10.1128/JB.02292-14).
- 602 Rodionov, D. A., Dubchak, I. L., Arkin, A. P., Alm, E. J., and Gelfand, M. S. (2005). Dissimilatory
603 metabolism of nitrogen oxides in bacteria: Comparative reconstruction of transcriptional networks.
604 *PLoS Comput. Biol.* 1, e55. doi:[10.1371/journal.pcbi.0010055](https://doi.org/10.1371/journal.pcbi.0010055).
- 605 Ruby, E. G. (2008). Symbiotic conversations are revealed under genetic interrogation. *Nat Rev*
606 *Microbiol* 6, 752–762. doi:[10.1038/nrmicro1958](https://doi.org/10.1038/nrmicro1958).
- 607 Ruby, E. G., and Asato, L. M. (1993). Growth and flagellation of *Vibrio fischeri* during initiation of
608 the sepiolid squid light organ symbiosis. *Arch Microbiol* 159, 160–167.
- 609 Ruby, E. G., and Lee, K.-H. (1998). The *Vibrio fischeri*-*Euprymna scolopes* light organ association:
610 Current ecological paradigms. *Appl Environ Microbiol* 64, 805–812. Available at:
611 <http://aem.asm.org/cgi/content/full/64/3/805?view=long&pmid=16349524>.

- 612 Ruby, E. G., and McFall-Ngai, M. J. (1992). A squid that glows in the night: Development of an
613 animal-bacterial mutualism. *J Bacteriol* 174, 4865–4870. Available at:
614 <http://jb.asm.org/cgi/reprint/174/15/4865?view=long&pmid=1629148>.
- 615 Ruby, E. G., Urbanowski, M., Campbell, J., Dunn, A., Faini, M., Gunsalus, R., et al. (2005).
616 Complete genome sequence of *Vibrio fischeri*: A symbiotic bacterium with pathogenic congeners.
617 *Proc Natl Acad Sci USA* 102, 3004–3009. doi:[10.1073/pnas.0409900102](https://doi.org/10.1073/pnas.0409900102).
- 618 Septer, A. N., and Stabb, E. V. (2012). Coordination of the arc regulatory system and pheromone-
619 mediated positive feedback in controlling the *Vibrio fischeri* lux operon. *PLoS ONE* 7, e49590.
620 doi:[10.1371/journal.pone.0049590](https://doi.org/10.1371/journal.pone.0049590).
- 621 Septer, A. N., Wang, Y., Ruby, E. G., Stabb, E. V., and Dunn, A. K. (2011). The haem-uptake gene
622 cluster in *Vibrio fischeri* is regulated by Fur and contributes to symbiotic colonization. *Environ*
623 *Microbiol* 13, 2855–2864. doi:[10.1111/j.1462-2920.2011.02558.x](https://doi.org/10.1111/j.1462-2920.2011.02558.x).
- 624 Shikuma, N. J., Pilhofer, M., Weiss, G. L., Hadfield, M. G., Jensen, G. J., and Newman, D. K.
625 (2014). Marine tubeworm metamorphosis induced by arrays of bacterial phage tail-like structures.
626 *Science* 343, 529–533. doi:[10.1126/science.1246794](https://doi.org/10.1126/science.1246794).
- 627 Stabb, E. V., and Visick, K. L. (2013). *Vibrio fischeri*: Squid symbiosis. *The Prokaryotes*.
628 doi:[10.1007/978-3-642-30194-0_118](https://doi.org/10.1007/978-3-642-30194-0_118).
- 629 Stenmark, P., and Nordlund, P. (2003). A prokaryotic alternative oxidase present in the bacterium
630 *Novosphingobium aromaticivorans*. *FEBS Lett.* 552, 189–192.
- 631 Sun, Y.-C., Hinnebusch, B. J., and Darby, C. (2008). Experimental evidence for negative selection in
632 the evolution of a *Yersinia pestis* pseudogene. *Proc Natl Acad Sci USA* 105, 8097–8101.
633 doi:[10.1073/pnas.0803525105](https://doi.org/10.1073/pnas.0803525105).
- 634 Sun, Y.-C., Jarrett, C. O., Bosio, C. F., and Hinnebusch, B. J. (2014). Retracing the evolutionary path
635 that led to flea-borne transmission of *Yersinia pestis*. *Cell Host Microbe* 15, 578–586.
636 doi:[10.1016/j.chom.2014.04.003](https://doi.org/10.1016/j.chom.2014.04.003).
- 637 Troll, J. V., Bent, E. H., Pacquette, N., Wier, A. M., Goldman, W. E., Silverman, N., et al. (2010).
638 Taming the symbiont for coexistence: A host PGRP neutralizes a bacterial symbiont toxin. *Environ*
639 *Microbiol* 12, 2190–2203. doi:[10.1111/j.1462-2920.2009.02121.x](https://doi.org/10.1111/j.1462-2920.2009.02121.x).
- 640 Tucker, N. P., Le Brun, N. E., Dixon, R., and Hutchings, M. I. (2010). There's NO stopping NsrR, a
641 global regulator of the bacterial NO stress response. *Trends Microbiol.* 18, 149–156.
642 doi:[10.1016/j.tim.2009.12.009](https://doi.org/10.1016/j.tim.2009.12.009).
- 643 Vanlerberghe, G. C. (2013). Alternative oxidase: A mitochondrial respiratory pathway to maintain
644 metabolic and signaling homeostasis during abiotic and biotic stress in plants. *Int J Mol Sci* 14,
645 6805–6847. doi:[10.3390/ijms14046805](https://doi.org/10.3390/ijms14046805).
- 646 Vanlerberghe, G. C., and McIntosh, L. (1997). ALTERNATIVE OXIDASE: From Gene to Function.
647 *Annu. Rev. Plant Physiol. Plant Mol. Biol.* 48, 703–734. doi:[10.1146/annurev.arplant.48.1.703](https://doi.org/10.1146/annurev.arplant.48.1.703).

- 648 Vine, C. E., and Cole, J. A. (2011). Unresolved sources, sinks, and pathways for the recovery of
649 enteric bacteria from nitrosative stress. *FEMS Microbiol. Lett.* 325, 99–107. doi:[10.1111/j.1574-
650 6968.2011.02425.x](https://doi.org/10.1111/j.1574-6968.2011.02425.x).
- 651 Visick, K. L. (2009). An intricate network of regulators controls biofilm formation and colonization
652 by *Vibrio fischeri*. *Mol Microbiol* 74, 782–789. doi:[10.1111/j.1365-2958.2009.06899.x](https://doi.org/10.1111/j.1365-2958.2009.06899.x).
- 653 Visick, K. L., and Skoufos, L. M. (2001). Two-component sensor required for normal symbiotic
654 colonization of *Euprymna scolopes* by *Vibrio fischeri*. *J Bacteriol* 183, 835–842.
655 doi:[10.1128/JB.183.3.835-842.2001](https://doi.org/10.1128/JB.183.3.835-842.2001).
- 656 Visick, K. L., Foster, J., Doino, J., McFall-Ngai, M., and Ruby, E. G. (2000). *Vibrio fischeri* lux
657 genes play an important role in colonization and development of the host light organ. *J Bacteriol* 182,
658 4578–4586. Available at:
659 <http://jb.asm.org/cgi/content/full/182/16/4578?view=long&pmid=10913092>.
- 660 Wang, Y., and Ruby, E. G. (2011). The roles of NO in microbial symbioses. *Cell. Microbiol.* 13,
661 518–526. doi:[10.1111/j.1462-5822.2011.01576.x](https://doi.org/10.1111/j.1462-5822.2011.01576.x).
- 662 Wang, Y., Dufour, Y. S., Carlson, H. K., Donohue, T. J., Marletta, M. A., and Ruby, E. G. (2010a).
663 H-NOX-mediated nitric oxide sensing modulates symbiotic colonization by *Vibrio fischeri*. *Proc*
664 *Natl Acad Sci USA* 107, 8375–8380. doi:[10.1073/pnas.1003571107](https://doi.org/10.1073/pnas.1003571107).
- 665 Wang, Y., Dunn, A. K., Wilneff, J., McFall-Ngai, M. J., Spiro, S., and Ruby, E. G. (2010b). *Vibrio*
666 *fischeri* flavohaemoglobin protects against nitric oxide during initiation of the squid-*Vibrio*
667 symbiosis. *Mol Microbiol* 78, 903–915. doi:[10.1111/j.1365-2958.2010.07376.x](https://doi.org/10.1111/j.1365-2958.2010.07376.x).
- 668 Wier, A. M., Nyholm, S. V., Mandel, M. J., Massengo-Tiassé, R. P., Schaefer, A. L., Koroleva, I., et
669 al. (2010). Transcriptional patterns in both host and bacterium underlie a daily rhythm of anatomical
670 and metabolic change in a beneficial symbiosis. *Proc. Natl. Acad. Sci. U.S.A* 107, 2259–2264.
671 doi:[10.1073/pnas.0909712107](https://doi.org/10.1073/pnas.0909712107).
- 672 Yip, E. S., Geszvain, K., DeLoney-Marino, C. R., and Visick, K. L. (2006). The symbiosis regulator
673 RscS controls the *Syp* gene locus, biofilm formation and symbiotic aggregation by *Vibrio Fischeri*.
674 *Mol Microbiol* 62, 1586–1600. doi:[10.1111/j.1365-2958.2006.05475.x](https://doi.org/10.1111/j.1365-2958.2006.05475.x).
- 675 Yip, E. S., Grublesky, B. T., Hussa, E. A., and Visick, K. L. (2005). A novel, conserved cluster of
676 genes promotes symbiotic colonization and σ^{54} -dependent biofilm formation by *Vibrio fischeri*. *Mol*
677 *Microbiol* 57, 1485–1498. doi:[10.1111/j.1365-2958.2005.04784.x](https://doi.org/10.1111/j.1365-2958.2005.04784.x).
- 678 Zimblér, D. L., Schroeder, J. A., Eddy, J. L., and Lathem, W. W. (2015). Early emergence of
679 *Yersinia pestis* as a severe respiratory pathogen. *Nat Commun* 6, 1–10. doi:[10.1038/ncomms8487](https://doi.org/10.1038/ncomms8487).