

The Ages of Stars and Other Stories: Redefining the Standard for 1D Stellar Modeling

Smith College Research Talk
¶ 15 December, 2022


 @MeridithJoyceGR
www.meridithjoyce.com
github.com/mjoyceGR
meridith.joyce@csfk.org

Dr Meridith Joyce
Marie Curie Widening Fellow: MATISSE
CSFK Konkoly Observatory, Budapest
 MESA Developers

About Me:

Aspiring international sensation

I have enjoyed work and life all over the world, and now I'm working on a Marie Curie (WF) in Europe!



About Me:

Astronomy software developer

Modules for Experiments in Stellar Astrophysics
(MESA)



Josiah Schwab



Adam Jermyn



Meridith Joyce



Evan Bauer




Earl Bellinger



Anne Thoul



Radek Smolec



Rob Farmer



Bill Wolf



Pablo Marchant



Warrick Ball



Aaron Dotter




Rich Townsend



Frank Timmes




Bill Paxton



Lars Bildsten



Matteo Cantiello




MESA

GYRE



About Me:

Part-time D-list science celebrity



SKY & TELESCOPE
THE ESSENTIAL GUIDE TO ASTRONOMY

WING TOOLS COMMUNITY

STELLAR SCIENCE **HOW BIG IS BETELGEUSE REALLY?**

BY: MONICA YOUNG | NOVEMBER 6, 2020 | 12

Knowing Betelgeuse's size is crucial to understanding its recent bizarre behavior — and predicting when it will go supernova. But it's harder to figure out than you might think.

AN **Astronomy Now**
The UK's best astronomy magazine

HOME THE MAGAZINE NEWS OBSERVING REVIEWS SPACEFLIGHT NOW S

LATEST NEWS [25 October 2020] OSIRIS-REx captures treasure trove of samples from asteroid Bennu

Red supergiant Betelgeuse not so bright, not so giant, not so far away

CNN World Africa Americas Asia Australia China Europe India Middle East United Kingdom Edition


SPACE + SCIENCE

Betelgeuse, the unusually dimming star, is smaller and closer than scientists thought

By Ashley Strickland, CNN Updated 1640 GMT (0040 HKT) October 19, 2020

About Me:

...and dubious Wikipedia source!



The screenshot shows the Wikipedia article for Betelgeuse. The page title is "Betelgeuse". The top navigation bar includes links for "Article", "Talk", "Read", "Edit", "View history", and "Search Wikipedia". The main content starts with a brief summary: "From Wikipedia, the free encyclopedia". Below this is a note about the article being about the star, not the film. The text describes Betelgeuse as the tenth-brightest star in the night sky, after Rigel, and notes it is a semiregular variable star. To the right, there is a small map of the Orion constellation showing the position of Betelgeuse relative to other stars like Aldebaran and Rigel. Coordinates for Betelgeuse are listed as 05^h 55^m 10.3053^s, +07° 24' 25.426".

In 2020, new observational data from the space-based *Solar Mass Ejection Imager* aboard the [Coriolis satellite](#) and three different modeling techniques produced a refined parallax of $5.95^{+0.58}_{-0.85}$ mas, a radius of $764^{+116}_{-62} R_{\odot}$, and a distance of $168.1^{+27.5}_{-14.4}$ pc or 548^{+90}_{-49} ly, which, if accurate, would mean Betelgeuse is nearly 25% smaller and 25% closer to Earth than previously thought.^[11]

All of astronomy and
cosmology hinges on results
from stellar models.

Exoplanets? Stellar models.

Exoplanets? Stellar models.

Exoplanet radii are determined with respect to the host star; most radii can't be measured directly → stellar models required

Exoplanets? Stellar models.

Exoplanet radii are determined with respect to the host star; most radii can't be measured directly → stellar models required

Cosmic chemistry? Stellar models.

Exoplanets? Stellar models.

Exoplanet radii are determined with respect to the host star; most radii can't be measured directly → stellar models required

Cosmic chemistry? Stellar models.

Intermediate and high mass stars chemically enrich the Galaxy, and our understanding of the **production of heavy elements** is closely tied to the quality of our models for nuclear burning and other interior processes

Exoplanets? Stellar models.

Exoplanet radii are determined with respect to the host star; most radii can't be measured directly → stellar models required

Cosmic chemistry? Stellar models.

Intermediate and high mass stars chemically enrich the Galaxy, and our understanding of the **production of heavy elements** is closely tied to the quality of our models for nuclear burning and other interior processes

Galaxy formation? Stellar models.

Exoplanets? Stellar models.

Exoplanet radii are determined with respect to the host star; most radii can't be measured directly → stellar models required

Cosmic chemistry? Stellar models.

Intermediate and high mass stars chemically enrich the Galaxy, and our understanding of the **production of heavy elements** is closely tied to the quality of our models for nuclear burning and other interior processes

Galaxy formation? Stellar models.

The formation history of the Galaxy is understood through maps of stellar abundances and stellar ages as a function of distance. → stellar models required

Exoplanets? Stellar models.

Exoplanet radii are determined with respect to the host star; most radii can't be measured directly → stellar models required

Cosmic chemistry? Stellar models.

Intermediate and high mass stars chemically enrich the Galaxy, and our understanding of the **production of heavy elements** is closely tied to the quality of our models for nuclear burning and other interior processes

Galaxy formation? Stellar models.

The formation history of the Galaxy is understood through maps of stellar abundances and stellar ages as a function of distance. → stellar models required

Further, the model of our Galaxy's history and structure sets the template against which all others are compared

Exoplanets? Stellar models.

Exoplanet radii are determined with respect to the host star; most radii can't be measured directly → stellar models required

Cosmic chemistry? Stellar models.

Intermediate and high mass stars chemically enrich the Galaxy, and our understanding of the **production of heavy elements** is closely tied to the quality of our models for nuclear burning and other interior processes

Galaxy formation? Stellar models.

The formation history of the Galaxy is understood through maps of stellar abundances and stellar ages as a function of distance. → stellar models required

Further, the model of our Galaxy's history and structure sets the template against which all others are compared

Cosmological questions? Stellar models.

Exoplanets? Stellar models.

Exoplanet radii are determined with respect to the host star; most radii can't be measured directly → stellar models required

Cosmic chemistry? Stellar models.

Intermediate and high mass stars chemically enrich the Galaxy, and our understanding of the **production of heavy elements** is closely tied to the quality of our models for nuclear burning and other interior processes

Galaxy formation? Stellar models.

The formation history of the Galaxy is understood through maps of stellar abundances and stellar ages as a function of distance. → stellar models required

Further, the model of our Galaxy's history and structure sets the template against which all others are compared

Cosmological questions? Stellar models.

Ancient, metal-poor stellar populations are still best fit by models with ages older than the accepted age of the universe, as are some peculiar individual stars...what if the resolution to this tension is right under our noses?

Exoplanets? Stellar models.

Exoplanet radii are determined with respect to the host star; most radii can't be measured directly → stellar models required

Cosmic chemistry? Stellar models.

Intermediate and high mass stars chemically enrich the Galaxy, and our understanding of the **production of heavy elements** is closely tied to the quality of our models for nuclear burning and other interior processes

Galaxy formation? Stellar models.

The formation history of the Galaxy is understood through maps of stellar abundances and stellar ages as a function of distance. → stellar models required

Further, the model of our Galaxy's history and structure sets the template against which all others are compared

Cosmological questions? Stellar models.

Ancient, metal-poor stellar populations are still best fit by models with ages older than the accepted age of the universe, as are some peculiar individual stars...what if the resolution to this tension is right under our noses?

No matter what you study, your work likely relies on results from stellar models


But this should concern you...

But this should concern you...

...because stellar models have many flaws.

But this should concern you...

...because stellar models have many flaws.



Together, we can fix them.


The power of stellar models and statistical inference in practice: determining the fundamental properties of

Betelgeuse

Standing on the shoulders of giants: New mass and distance estimates for Betelgeuse through a combination of evolutionary, asteroseismic, and hydrodynamical simulations with MESA

Meridith Joyce, Shing-Chi Leung, László Molnár,
Michael J. Ireland, Chiaki Kobayashi, and Ken'ichi Nomoto
ApJ, 2020

**From November 2019 to January 2020,
Betelgeuse underwent a huge brightness
drop—The “Great Dimming”**




Such severe brightness changes are thought to precede supernovae



Dr. Nash Rafeeq @nashrafeeg · Feb 26

GO SUPER NOVA!!!!!! twitter.com/IsBetelgeuseOK...




Betelgeuse
@IsBetelgeuseOK

You first.

Like everyone, I wanted to know what's going on.
But being a stellar evolution expert, I had some
tools at my disposal that might actually answer
this question

Like everyone, I wanted to know what's going on.
But being a stellar evolution expert, I had some
tools at my disposal that might actually answer
this question

My approach: reproduce this lightcurve
via simulation to understand why
Betelgeuse became dim or rule out causes



Like everyone, I wanted to know what's going on.
But being a stellar evolution expert, I had some
tools at my disposal that might actually answer
this question

My approach: reproduce this lightcurve
via simulation to understand why
Betelgeuse became dim or rule out causes

...but in order to understand why this dimming
event was “unprecedented,” we must first
understand Betelgeuse’s normal periodic
variations

Key tools:



Modules for Experiments in Stellar Astrophysics

Generates structural, evolutionary, hydrodynamic models




Solves the oscillation equations to predict a body's response to perturbations, generating a theoretical frequency spectrum

→ in this case, the “body” is a model of **Betelgeuse**

Part 1: Observational analysis

Betelgeuse's brightness variability over 100 years




Time →

Fig by László Molnár

The ‘unprecedented’ dimming of Betelgeuse




-First question: How unprecedented, *really*?



Actually, very. This is
the dimmest it's been in
nearly 100 years

Key Concept: Stars pulsate!




They “ring” like bells, in response to physical mechanisms causing waves inside them



Key Concept: Stars pulsate!

They “ring” like bells, in response to physical mechanisms causing waves inside them

The frequencies (pitches) at which they ring can tell us what they’re made out of






Key Concept: Stars pulsate!


They “ring” like bells, in response to physical mechanisms causing waves inside them

The frequencies (pitches) at which they ring can tell us what they’re made out of

The use of these pulsations to learn about stellar structure is called **asteroseismology**




Fourier Analysis: What patterns do we detect in the frequency spectrum?



Fourier Analysis: What patterns do we detect in the frequency spectrum?

From new and archival brightness measurements, we find strong periods at:

- **416 days**
- **185 days**
- **5.6 years**




Fourier Analysis: What patterns do we detect in the frequency spectrum?

From new and archival brightness measurements, we find strong periods at:

- 416 days
- 185 days
- 5.6 years

Determining the drivers of different frequencies tells us about the structure of the star



Fourier Analysis: What patterns do we detect in the frequency spectrum?


From new and archival brightness measurements, we find strong periods at:

- 416 days
- 185 days
- 5.6 years

Determining the drivers of different frequencies tells us about the structure of the star

What we want to know...

Is the 416-day periodicity a type of frequency called the ***fundamental mode***?



Fourier Analysis: What patterns do we detect in the frequency spectrum?

From new and archival brightness measurements, we find strong periods at:


- 416 days
- 185 days
- 5.6 years

Determining the drivers of different frequencies tells us about the structure of the star

What we want to know...

Is the 416-day periodicity a type of frequency called the ***fundamental mode***?

Is the 185-day periodicity the ***first overtone***?



Fourier Analysis: What patterns do we detect in the frequency spectrum?

From new and archival brightness measurements, we find strong periods at:

- 416 days
- 185 days
- 5.6 years


Determining the drivers of different frequencies tells us about the structure of the star

What we want to know...


Is the 416-day periodicity a type of frequency called the ***fundamental mode***?

Is the 185-day periodicity the ***first overtone***?


We can answer these questions by determining whether the 416-day periodicity is **driven by the kappa mechanism**



Step 2: Build evolutionary models that agree with observations




Part 3: Add in asteroseismology




Use GYRE to perform linear seismic analysis on observationally consistent tracks:
those which intersect the temperature constraints and their uncertainties


Determining which models are seismically compatible



Determining which models are seismically compatible





It turns out that this **theoretical and statistical method**
constrains Betelgeuse's physical radius **better than traditional
observational methods**



Key insight from evolutionary models and statistical methods:

Statistically, it is most likely that Betelgeuse is in its core helium burning phase

→ we demonstrate this using a **Monte Carlo interpolation** over a grid of tracks spanning the masses, metallicities, and radii compatible with the observations



Classical & Seismic results:

General finding: on the (initial) mass of Betelgeuse, our results are consistent with other modeling efforts; not particularly more precise: 18-24 Msolar

Classical & Seismic results:

General finding: on the (initial) mass of Betelgeuse, our results are consistent with other modeling efforts; not particularly more precise: 18-24 Msolar

...but our models permit only a **very small range** for the physical radius: a 3σ band of 150 Rsolar

Classical & Seismic results:

General finding: on the (initial) mass of Betelgeuse, our results are consistent with other modeling efforts; not particularly more precise: 18-24 Msolar

...but our models permit only a **very small range** for the physical radius: a 3σ band of 150 Rsolar


In fact, this range is considerably smaller than predictions for the physical radius provided by traditional observational methods!

Inversion of the distance inference:

Angular diameter

$$R = \frac{1}{2}\theta D$$

Fig by Tim White




Parallax ($1/D$) is a tricky trigonometric distance inference that can be combined with a measurement of the apparent size, or angular diameter (theta), of Betelgeuse on the sky to make a prediction for Betelgeuse's physical radius

Inversion of the distance inference:

Angular diameter

$$R = \frac{1}{2}\theta D$$

Fig by Tim White




But in our case, thanks to statistics and asteroseismology, we already have a theoretical value for Betelgeuse's physical radius from our models...

Inversion of the distance inference:

Angular diameter

Fig by Tim White

$$R = \frac{1}{2}\theta D$$



We may then invert the argument by combining
our theoretical radius (from calculations)
& the observed angular diameter (theta)
→ parallax ($1/D$)

So, we put the radius in and get the distance out, instead of
the other way around

Classical & Seismic results:


General finding: on the (initial) mass of Betelgeuse, our results are consistent with other modeling efforts; not particularly more precise: 18-24 Msolar

...but our models permit only a **very small range** for the physical radius: a 3σ band of 150 Rsolar

In fact, this range is considerably smaller than predictions for the physical radius provided by traditional observational methods!

Unanticipated Bonus:

precision modelled radius + measured angular diameter
= **new parallax distance estimate**



Revised distance from seismic parallax





Fig by László Molnár

Part 4: Hydrodynamical Simulations


We can reproduce GYRE's linear mode classifications with hydrodynamics

Self-consistent mode classifications: From short-timescale hydro simulations, we can extract cycle lengths as a secondary means of estimating oscillation periods






With hydrodynamic calculations, can we...



- (1) activate a mode consistent with the FM? → yes
- (2) conclusively determine the driving physics of that mode? → ??




Examine the envelope structure at critical points in the pulsation cycle



Yes, it's the κ mechanism




Key results from four analysis methods:



Radius: $764 +116, -62$ Rsolar
Distance: $168 +27, -15$ pc
Parallax: $5.95 +0.58, -0.85$ mas

Key results from four analysis methods:

We used hydrodynamics to confirm that the kappa mechanism drives the 416 d period, which means the 416d period is the FM




Radius: $764 +116, -62$ Rsolar
Distance: $168 +27, -15$ pc
Parallax: $5.95 +0.58, -0.85$ mas

Key results from four analysis methods:

We used hydrodynamics to confirm that the kappa mechanism drives the 416 d period, which means the 416d period is the FM

We discovered that Betelgeuse is a **double-mode pulsator**, with overtone period 185 ± 13 d




Radius: $764 +116, -62$ Rsolar
Distance: $168 +27, -15$ pc
Parallax: $5.95 +0.58, -0.85$ mas

Key results from four analysis methods:

We used hydrodynamics to confirm that the kappa mechanism drives the 416 d period, which means the 416d period is the FM

We discovered that Betelgeuse is a **double-mode pulsator**, with overtone period 185 ± 13 d

We accidentally invented a new way to measure distance!



PC: ALMA

Radius: $764 +116, -62$ Rsolar
Distance: $168 +27, -15$ pc
Parallax: $5.95 +0.58, -0.85$ mas


Key results from four analysis methods:

We used hydrodynamics to confirm that the kappa mechanism drives the 416 d period, which means the 416d period is the FM

We discovered that Betelgeuse is a **double-mode pulsator**, with overtone period 185 ± 13 d

We accidentally invented a new way to measure distance!

We found that Betelgeuse is **less massive, smaller and closer** than we thought



Radius: $764 \pm 116, -62$ Rsolar
Distance: $168 \pm 27, -15$ pc
Parallax: $5.95 \pm 0.58, -0.85$ mas

Key results from four analysis methods:


We used hydrodynamics to confirm that the kappa mechanism drives the 416 d period, which means the 416d period is the FM

We discovered that Betelgeuse is a **double-mode pulsator**, with overtone period 185 ± 13 d

We accidentally invented a new way to measure distance!

We found that Betelgeuse is **less massive, smaller and closer** than we thought

We inferred Betelgeuse's current nuclear burning state statistically, and that suggests it will not "go supernova" for ~100,000 years



PC: ALMA

Radius: $764 +116, -62$ Rsolar
Distance: $168 +27, -15$ pc
Parallax: $5.95 +0.58, -0.85$ mas

Key results from four analysis methods:

We used hydrodynamics to confirm that the kappa mechanism drives the 416 d period, which means the 416d period is the FM


We discovered that Betelgeuse is a **double-mode pulsator**, with overtone period 185 ± 13 d

We accidentally invented a new way to measure distance!

We found that Betelgeuse is **less massive, smaller and closer** than we thought

We inferred Betelgeuse's current nuclear burning state statistically, and that suggests it will not "go supernova" for $\sim 100,000$ years

Seismic parameters were the key to the modeling problem



Radius: $764 \pm 116, -62$ Rsolar
Distance: $168 \pm 27, -15$ pc
Parallax: $5.95 \pm 0.58, -0.85$ mas

Where did this project use data science methods?

Where did this project use data science methods?

Everywhere!

Where did this project use data science methods?

Everywhere!

- We performed a Fourier decomposition of the variability observations to determine the spectrum's composite frequencies

Where did this project use data science methods?

Everywhere!

- We performed a Fourier decomposition of the variability observations to determine the spectrum's composite frequencies
- We used Monte Carlo simulations constrained by physical information (priors) to find the most probable evolutionary stage of Betelgeuse

Where did this project use data science methods?

Everywhere!

- We performed a Fourier decomposition of the variability observations to determine the spectrum's composite frequencies
- We used Monte Carlo simulations constrained by physical information (priors) to find the most probable evolutionary stage of Betelgeuse
- behind the scenes: we used **grid modeling, multi-dimensional parameter optimization, error propagation techniques, and high performance computing** to generate the different bases of models used to fit to Betelgeuse, and then again to determine the *best fit*

How can students get involved in research like this?

How can students get involved in research like this?

- learn how to use observational catalogs and SQL querying to find and assemble the best observational information to use as priors/constraints for our models

How can students get involved in research like this?

- learn how to use observational catalogs and SQL querying to find and assemble the best observational information to use as priors/constraints for our models
- create the models we use to fit the stars! Create better models than the ones I made, using data-driven methods in place of out-of-date formulations from before we had good observational information

How can students get involved in research like this?

- learn how to use observational catalogs and SQL querying to find and assemble the best observational information to use as priors/constraints for our models
- create the models we use to fit the stars! Create better models than the ones I made, using data-driven methods in place of out-of-date formulations from before we had good observational information
- build open-source databases for these models so students can share their work and we can build a large project collaboratively and with minimal redundancy

How can students get involved in research like this?

- learn how to use observational catalogs and SQL querying to find and assemble the best observational information to use as priors/constraints for our models
- create the models we use to fit the stars! Create better models than the ones I made, using data-driven methods in place of out-of-date formulations from before we had good observational information
- build open-source databases for these models so students can share their work and we can build a large project collaboratively and with minimal redundancy
- write software that automatically analyzes asteroseismic information

How can students get involved in research like this?

- learn how to use observational catalogs and SQL querying to find and assemble the best observational information to use as priors/constraints for our models
- create the models we use to fit the stars! Create better models than the ones I made, using data-driven methods in place of out-of-date formulations from before we had good observational information
- build open-source databases for these models so students can share their work and we can build a large project collaboratively and with minimal redundancy
- write software that automatically analyzes asteroseismic information
- find better ways to create the models. Make running the stellar structure and evolution software obsolete. Replace ancient partial differential equation solvers—which is what stellar structure and evolution programs are, by and large—with machine learning parameter generators trained on stellar models instead

Is it really a “fundamental tension,”
or is it a matter of error bars?

The Age of the Galactic Bulge

The Ages of Galactic Bulge Stars with Realistic Uncertainties
Meridith Joyce, Christian I Johnson, Tommaso Marchetti,
R. Michael Rich, Iulia Simion, and John Bourke
ApJ (in revision), 2022

In 2017, a catalog of ages for 91 micro-lensed subdwarfs was put forth by T. Bensby and collaborators:

**Chemical evolution of the Galactic bulge as traced
by microlensed dwarf and subgiant stars**

**VI. Age and abundance structure of the stellar populations
in the central sub-kpc of the Milky Way^{*,**}**

T. Bensby¹, S. Feltzing¹, A. Gould^{2,3,4}, J. C. Yee⁵, J. A. Johnson⁴, M. Asplund⁶, J. Meléndez⁷, S. Lucatello⁸, L. M. Howes¹, A. McWilliam⁹, A. Udalski^{10,***}, M. K. Szymański^{10,****}, I. Soszyński^{10,****}, R. Poleski^{10,4,***}, Ł. Wyrzykowski^{10,***}, K. Ulaczyk^{10,11,***}, S. Kozłowski^{10,***}, P. Pietrukowicz^{10,***}, J. Skowron^{10,***}, P. Mróz^{10,***}, M. Pawlak^{10,***}, F. Abe^{12,*****}, Y. Asakura^{12,*****}, A. Bhattacharya^{13,*****}, I. A. Bond^{14,*****}, D. P. Bennett^{15,*****}, Y. Hirao^{16,*****}, M. Nagakane^{16,*****}, N. Koshimoto^{16,*****}, T. Sumi^{16,*****}, D. Suzuki^{15,*****}, and P. J. Tristram^{17,****}

In 2017, a catalog of ages for 91 micro-lensed subdwarfs was put forth by T. Bensby and collaborators:

**Chemical evolution of the Galactic bulge as traced
by microlensed dwarf and subgiant stars**

**VI. Age and abundance structure of the stellar populations
in the central sub-kpc of the Milky Way^{*,**}**


T. Bensby¹, S. Feltzing¹, A. Gould^{2,3,4}, J. C. Yee⁵, J. A. Johnson⁴, M. Asplund⁶, J. Meléndez⁷, S. Lucatello⁸, L. M. Howes¹, A. McWilliam⁹, A. Udalski^{10,***}, M. K. Szymański^{10,****}, I. Soszyński^{10,****}, R. Poleski^{10,4,***}, Ł. Wyrzykowski^{10,***}, K. Ulaczyk^{10,11,***}, S. Kozłowski^{10,***}, P. Pietrukowicz^{10,***}, J. Skowron^{10,***}, P. Mróz^{10,***}, M. Pawlak^{10,***}, F. Abe^{12,*****}, Y. Asakura^{12,*****}, A. Bhattacharya^{13,*****}, I. A. Bond^{14,*****}, D. P. Bennett^{15,*****}, Y. Hirao^{16,*****}, M. Nagakane^{16,*****}, N. Koshimoto^{16,*****}, T. Sumi^{16,*****}, D. Suzuki^{15,*****}, and P. J. Tristram^{17,****}



We played with this data
set in our mock class

In 2017, a catalog of ages for 91 micro-lensed subdwarfs was put forth by T. Bensby and collaborators

...that suggested the presence of a significant population of young stars in the Galactic Bulge



Young stars in the Galactic bulge?

Why is this contentious?

Young stars in the Galactic bulge?

Why is this contentious?

- Galactic bulge long been thought to be old

Young stars in the Galactic bulge?

Why is this contentious?

- Galactic bulge long been thought to be old
- presence of young stars implies prolonged star formation, and our understanding of the Galaxy is not consistent with recent star formation episodes in this region

Young stars in the Galactic bulge?

Why is this contentious?

- Galactic bulge long been thought to be old
- presence of young stars implies prolonged star formation, and our understanding of the Galaxy is not consistent with recent star formation episodes in this region
- an overabundance of young stars in this region thus calls into question the formation history of the Galaxy and galaxy evolution mechanisms more generally

Young stars in the Galactic bulge?

Big picture:

Young stars in the Galactic bulge?

Big picture:

There is tension in the literature between ages derived from **photometry**, which claims a **uniformly old bulge**

Young stars in the Galactic bulge?

Big picture:

There is tension in the literature between ages derived from **photometry**, which claims a **uniformly old bulge**

...and ages derived from **microlensing** (**spectroscopy**), which claims a **broad age distribution**

Young stars in the Galactic bulge?

Big picture:

There is tension in the literature between ages derived from **photometry**, which claims a **uniformly old bulge**

...and ages derived from **microlensing** (**spectroscopy**), which claims a **broad age distribution**

...but in order for something to be true, **it must be true regardless of inference method**

A rare and powerful dataset

Microlensing permits the direct inference of physical, spectroscopic coordinates (Teff, logg) of faint, cool stars due to the 10-1000x brightness magnification they experience during these events







Fig 6, Bensby et al. 2017: 91 stars on Yale isochrones




Resulting age distribution (Bensby et al.; Yale)




Reproduction with MIST isochrones





Resulting age distribution (Joyce et al.; MIST)



Distributions overlaid





One could reasonably ask...

Is it because the MIST and Yale models use wildly different physical assumptions and therefore yield different ages?

→ No

Is it because we use more sophisticated assumptions about the chemical distribution of the Galaxy in our isochrones and the other age determinations do not?

→ Also no

*see full paper (arXiv: 2205.07964) for rigorous demonstration of this using math & histograms

One could reasonably ask...

Is it because the MIST and Yale models use wildly different physical assumptions and therefore yield different ages?

→ No

Is it because we use more sophisticated assumptions about the chemical distribution of the Galaxy in our isochrones and the other age determinations do not?


→ Also no

*see full paper (arXiv: 2205.07964) for rigorous demonstration of this using math & histograms

A huge part of my research is the art of *ruling out* explanations for discrepancies, using statistics, probability, and computer modeling

Research is not just about finding a model that works, it's about demonstrating mathematically why other explanations are not viable

Punchline: we (Joyce + MIST) do not find an abundance of young stars at high metallicities



Bensby+ : 42 stars < 7 Gyr, all $[Fe/H] > -0.6$
13 stars > 7, < 10 Gyr
36 stars > 10 Gyr

Joyce+ : 15 stars < 7 Gyr, all $[Fe/H] > -0.3$
18 stars > 7, < 10 Gyr
58 stars > 10 Gyr

Why should you believe Joyce+MIST over the previous result?

Why should you believe Joyce+MIST over the previous result?



Carefully considered
hybrid statistical techniques

Age determination algorithm

Challenges:



Though it is straightforward to fit an isochrone to an observation “by eye,” it is much more difficult to construct a mathematically rigorous definition of a best-fitting model. This is especially true in a situation where:


1. many observations are plausibly consistent with a large number of isochrones (i.e., the isochrone falls within the star’s 1σ uncertainties);
2. the isochrones are discretely spaced in age and metallicity and thus limited in resolution; and
3. the models contain their own intrinsic uncertainties that are both difficult to quantify and highly variant over different parameter regimes—a point to which we will return in later discussion.

Age determination algorithm

Challenges:

Though it is straightforward to fit an isochrone to an observation “by eye,” it is much more difficult to construct a mathematically rigorous definition of a best-fitting model. This is especially true in a situation where:

- 
- 
1. many observations are plausibly consistent with a large number of isochrones (i.e., the isochrone falls within the star’s 1σ uncertainties);
 2. the isochrones are discretely spaced in age and metallicity and thus limited in resolution; and
 3. the models contain their own intrinsic uncertainties that are both difficult to quantify and highly variant over different parameter regimes—a point to which we will return in later discussion.



For the enthusiasts:

To compute the *weighted-average age* of a given star, we apply a combination of a **frequentist approach, goodness-of-fit, and maximum likelihood analysis**

For the enthusiasts:

To compute the *weighted-average age* of a given star, we apply a combination of a **frequentist approach**, **goodness-of-fit**, and **maximum likelihood analysis**

$$\chi^2_{\text{for } i \text{ DoF}} = \sum_i \frac{(o_i - t_i)^2}{\sigma_i^2}$$

(1) For a given star, compute the (3 DoF) chisq score for the fit of that star's observational parameters to **every** point along a candidate model (e.g. isochrone)

$$\chi^2_{\text{B17}} = \frac{(\log g_o - \log g_t)^2}{\sigma_{\log g,o}^2} + \frac{(T_{\text{eff},o} - T_{\text{eff},t})^2}{\sigma_{T_{\text{eff}},o}^2} + \frac{(Z_o - Z_t)^2}{\sigma_{Z,o}^2},$$

For the enthusiasts:

To compute the *weighted-average age* of a given star, we apply a combination of a **frequentist approach**, **goodness-of-fit**, and **maximum likelihood analysis**

$$\chi^2_{\text{for } i \text{ DoF}} = \sum_i \frac{(o_i - t_i)^2}{\sigma_i^2}$$

(1) For a given star, compute the (3 DoF) chisq score for the fit of that star's observational parameters to **every** point along a candidate model (e.g. isochrone)

$$\chi^2_{\text{B17}} = \frac{(\log g_o - \log g_t)^2}{\sigma_{\log g,o}^2} + \frac{(T_{\text{eff},o} - T_{\text{eff},t})^2}{\sigma_{T_{\text{eff}},o}^2} + \frac{(Z_o - Z_t)^2}{\sigma_{Z,o}^2},$$

(2) Each chisq score corresponds to a relative likelihood, p_n , given by the density function of the chi square distribution with 3 DoF at that score.

For the enthusiasts:

To compute the *weighted-average age* of a given star, we apply a combination of a **frequentist approach**, **goodness-of-fit**, and **maximum likelihood analysis**

$$\chi^2_{\text{for } i \text{ DoF}} = \sum_i \frac{(o_i - t_i)^2}{\sigma_i^2}$$

(1) For a given star, compute the (3 DoF) chisq score for the fit of that star's observational parameters to **every** point along a candidate model (e.g. isochrone)

$$\chi^2_{\text{B17}} = \frac{(\log g_o - \log g_t)^2}{\sigma_{\log g, o}^2} + \frac{(T_{\text{eff}, o} - T_{\text{eff}, t})^2}{\sigma_{T_{\text{eff}}, o}^2} + \frac{(Z_o - Z_t)^2}{\sigma_{Z, o}^2},$$

- (3) - Compute a weighted average over all candidate hypotheses (each point on each isochrone)
- A point with age t_n is weighted by its likelihood, p_n , of being an appropriate fit to the star
- The final weighted average, t_S , is our estimate for the age of the star


(2) Each chisq score corresponds to a relative likelihood, p_n , given by the density function of the chi square distribution with 3 DoF at that score.

$$t_S = \frac{\sum_n t_n p_n}{\sum_n p_n},$$


Visually...

Each model is
uniquely defined by
 $\{\text{age}, \text{metallicity}\}$

But consider all points
on all isochrones
agnostically



Error bars: Monte Carlo resampling



Construct three independent normal distributions with densities

$$f(x) = \frac{1}{\sigma\sqrt{2\pi}} e^{-(x-\mu)^2/2\sigma^2}$$

defined by

$$\mu = T_{\text{eff},S}, \quad \sigma = \sigma_{T_{\text{eff},S}},$$

$$\mu = \log g_S, \quad \sigma = \sigma_{\log g_S},$$

$$\mu = Z_S, \quad \sigma = \sigma_{Z_S}$$

Observational vs global uncertainties

There is an entire, second set of systematics we must understand in order to perform fully correct age determinations:

Observational vs global uncertainties

There is an entire, second set of systematics we must understand in order to perform fully correct age determinations:



...the intrinsic uncertainty of the stellar models themselves

There are **many** physical choices in stellar models one should worry about...

There are **many** physical choices in stellar models one should worry about...

Convective & energy transport parameters:

- mixing efficiency → convective mixing length
- convective overshoot
- convective boundaries; how the Schwarzschild/Ledoux criterion is evaluated

Heavy element diffusion:

- is it included, and where?
- how is it implemented?
- gravitational settling?
- are all isotopes treated the same?

Atmospheric boundary conditions:

- Is it a T-tau relation, & what kind of integration? Eddington vs Krishna-Swamy
- if instead using a table-based treatment from external simulations (e.g. PHOENIX, Kurucz), what solar scale and other physics were used in those simulations? Are they self-consistent with the assumptions in the stellar models?

There are **many** physical choices in stellar models one should worry about...

Convective & energy transport parameters:

- mixing efficiency → convective mixing length
- convective overshoot
- convective boundaries; how the Schwarzschild criterion is treated

Heavy element distribution


- is it a simple adiabatic gradient?
- is it a more complex treatment?
- is it treated the same?

Atmospheric boundary conditions:

- Is it a T-tau relation, & what kind of integration? Eddington vs Krishna-Swamy
- if instead using a table-based treatment from external simulations (e.g. PHOENIX, Kurucz), what solar scale and other physics were used in those simulations? Are they self-consistent with the assumptions in the stellar models?

Like all of these!

We choose to use the convective mixing length, α_{MLT} , for demonstration




We choose to use the convective mixing length, α_{MLT} , for demonstration



While far from the only source of modeling uncertainty, varying α_{MLT} provides a sharp demonstration of the danger of failing to account for theoretical uncertainties in age determinations—and hardly anyone does!



The impact of α_{MLT}





Incorporating the isochrones' "shift" in our error simulations

Uniformly shift the isochrones according to the sampling of normal distributions given by

$$\mu = 0, \quad \sigma = \sigma_{T_{\text{eff}, \text{th}}}$$

$$\mu = 0, \quad \sigma = \sigma_{\log g_{\text{th}}},$$



What happens when we incorporate variation in the isochrones' position in logg-Teff in the MC simulations?

	Name	.. α -enhanced	...model err
1	MOA-2009-BLG-174S	9.3 ± 2.0	9.3 ± 4.0
2	MOA-2009-BLG-259S	7.8 ± 2.4	7.8 ± 3.8
3	MOA-2010-BLG-167S ★	15.4 ± 1.7	15.4 ± 3.8
4	MOA-2010-BLG-311S	7.7 ± 2.7	7.7 ± 3.3
5	MOA-2010-BLG-446S	4.9 ± 1.2	4.9 ± 2.0
6	MOA-2010-BLG-523S ★	7.5 ± 2.5	7.5 ± 3.8
7	OGLE-2011-BLG-0950S ★ †	3.3 ± 1.1	3.3 ± 1.6
8	OGLE-2011-BLG-0969S ★	13.4 ± 1.1	13.4 ± 1.5
9	MOA-2011-BLG-034S	14.6 ± 1.9	14.6 ± 2.8
10	MOA-2011-BLG-058S	13.8 ± 1.9	13.8 ± 2.8
11	OGLE-2011-BLG-1072S	8.2 ± 2.3	8.2 ± 3.4
12	MOA-2011-BLG-090S	17.4 ± 1.1	17.4 ± 2.7
13	MOA-2011-BLG-104S	13.3 ± 1.2	13.3 ± 1.8
14	OGLE-2011-BLG-1105S	9.9 ± 1.9	9.9 ± 2.9
15	MOA-2011-BLG-174S	5.0 ± 1.0	5.0 ± 2.5
16	MOA-2011-BLG-191S	8.1 ± 3.1	8.1 ± 4.1
17	MOA-2011-BLG-234S †	8.3 ± 1.8	8.3 ± 3.0
18	MOA-2011-BLG-278S	12.7 ± 1.5	12.7 ± 1.8
19	OGLE-2011-BLG-1410S	10.8 ± 1.6	10.8 ± 2.4
20	MOA-2011-BLG-445S	11.1 ± 1.8	11.1 ± 2.8
21	MOA-2012-BLG-022S	5.0 ± 1.3	5.0 ± 2.2
22	OGLE-2012-BLG-0026S	11.1 ± 1.5	11.1 ± 2.5

What happens when we incorporate variation in the isochrones' position in logg-Teff in the MC simulations?

	Name	.. α -enhanced	...model err
1	MOA-2009-BLG-174S	9.3 ± 2.0	9.3 ± 4.0
2	MOA-2009-BLG-259S	7.8 ± 2.4	7.8 ± 3.8
3	MOA-2010-BLG-167S ★	15.4 ± 1.7	15.4 ± 3.8
4	MOA-2010-BLG-311S	7.7 ± 2.7	7.7 ± 3.3
5	MOA-2010-BLG-446S	4.9 ± 1.2	2.0
6	MOA-2010-BLG-523S ★	7.5 ± 1.5	3.8
7	OGLE-2011-BLG-0950S ★†	10.5 ± 1.5	6
8	OGLE-2011-BLG-1105S	17.4 ± 1.1	5
9	MOA-2011-BLG-174S	13.3 ± 1.2	3.8 ± 2.8
10	OGLE-2011-BLG-1105S	9.9 ± 1.9	8.2 ± 3.4
11	MOA-2011-BLG-174S	5.0 ± 1.0	5.0 ± 2.5
12	OGLE-2011-BLG-1105S	8.1 ± 3.1	8.1 ± 4.1
13	MOA-2011-BLG-234S †	8.3 ± 1.8	8.3 ± 3.0
14	MOA-2011-BLG-278S	12.7 ± 1.5	12.7 ± 1.8
15	OGLE-2011-BLG-1410S	10.8 ± 1.6	10.8 ± 2.4
16	MOA-2011-BLG-445S	11.1 ± 1.8	11.1 ± 2.8
17	MOA-2012-BLG-022S	5.0 ± 1.3	5.0 ± 2.2
18	OGLE-2012-BLG-0026S	11.1 ± 1.5	11.1 ± 2.5

uncertainties increase
by a factor of 1.5 - 2x

Tension between
inference methods?

Conclusions

Conclusions

Bensby+ 2017 (Yale isochrones) find a large population of metal-rich, young stars in the bulge, suggesting prolonged star formation in the region, which is in conflict with previous/other understanding of the formation history of the Galaxy

Conclusions

Bensby+ 2017 (Yale isochrones) find a large population of metal-rich, young stars in the bulge, suggesting prolonged star formation in the region, which is in conflict with previous/other understanding of the formation history of the Galaxy

Joyce (me)+ 2022 (MIST isochrones) do not find a large constituency of metal-rich young stars in this region, despite using Bensby+2017's parameters verbatim

Conclusions

Bensby+ 2017 (Yale isochrones) find a large population of metal-rich, young stars in the bulge, suggesting prolonged star formation in the region, which is in conflict with previous/other understanding of the formation history of the Galaxy

Joyce (me)+ 2022 (MIST isochrones) do not find a large constituency of metal-rich young stars in this region, despite using Bensby+2017's parameters verbatim

There is no significant discrepancy between the physical assumptions adopted between both isochrone databases, nor can differences in alpha-abundance scale explain the striking difference in derived age distributions

Conclusions

Bensby+ 2017 (Yale isochrones) find a large population of metal-rich, young stars in the bulge, suggesting prolonged star formation in the region, which is in conflict with previous/other understanding of the formation history of the Galaxy

Joyce (me)+ 2022 (MIST isochrones) do not find a large constituency of metal-rich young stars in this region, despite using Bensby+2017's parameters verbatim

There is no significant discrepancy between the physical assumptions adopted between both isochrone databases, nor can differences in alpha-abundance scale explain the striking difference in derived age distributions

Bensby+2017's age distribution is statistically consistent with a uniform distribution across 1 to 15 Gyr, whereas Joyce+2022 finds a clear peak at 13 Gyr and a median of 10.8 Gyr.

Conclusions

Bensby+ 2017 (Yale isochrones) find a large population of metal-rich, young stars in the bulge, suggesting prolonged star formation in the region, which is in conflict with previous/other understanding of the formation history of the Galaxy

Joyce (me)+ 2022 (MIST isochrones) do not find a large constituency of metal-rich young stars in this region, despite using Bensby+2017's parameters verbatim

There is no significant discrepancy between the physical assumptions adopted between both isochrone databases, nor can differences in alpha-abundance scale explain the striking difference in derived age distributions

Bensby+2017's age distribution is statistically consistent with a uniform distribution across 1 to 15 Gyr, whereas Joyce+2022 finds a clear peak at 13 Gyr and a median of 10.8 Gyr.

While still showing some slight age spread, Joyce+2022 results are **more consistent with photometric analyses of this region** despite being **based on the same spectroscopic, microlensed sample** analyzed in Bensby+2017

Conclusions

Bensby+ 2017 (Yale isochrones) find a large population of metal-rich, young stars in the bulge, suggesting prolonged star formation in the region, which is in conflict with previous/other understanding of the formation history of the Galaxy

Joyce (me)+ 2022 (MIST isochrones) do not find a large constituency of metal-rich young stars in this region, despite using Bensby+2017's parameters verbatim

There is no significant discrepancy between the physical assumptions adopted between both isochrone databases, nor can differences in alpha-abundance scale explain the striking difference in derived age distributions

Bensby+2017's age distribution is statistically consistent with a uniform distribution across 1 to 15 Gyr, whereas Joyce+2022 finds a clear peak at 13 Gyr and a median of 10.8 Gyr.

While still showing some slight age spread, Joyce+2022 results are **more consistent with photometric analyses of this region** despite being **based on the same spectroscopic, microlensed sample** analyzed in Bensby+2017

Have we resolved the spectroscopic/photometric tension? Not entirely, but careful application of statistics puts the picture in better focus

Conclusions

Bensby+ 2017 (Yale isochrones) find a large population of metal-rich, young stars in the bulge, suggesting prolonged star formation in the region, which is in conflict with previous/other understanding of the formation history of the Galaxy

Joyce (me)+ 2022 (MIST isochrones) do not find a large constituency of metal-rich young stars in this region, despite using Bensby+2017's parameters verbatim

There is no significant discrepancy between the physical assumptions adopted between both isochrone databases, nor can differences in alpha-abundance scale explain the striking difference in derived age distributions

Bensby+2017's age distribution is statistically consistent with a uniform distribution across 1 to 15 Gyr, whereas Joyce+2022 finds a clear peak at 13 Gyr and a median of 10.8 Gyr.

While still showing some slight age spread, Joyce+2022 results are **more consistent with photometric analyses of this region** despite being **based on the same spectroscopic, microlensed sample** analyzed in Bensby+2017


Have we resolved the spectroscopic/photometric tension? Not entirely, but careful application of statistics puts the picture in better focus

Ages are hard! Be careful with math.

How can students get involved in research like this?

How can students get involved in research like this?

You already were, in our mock class!



The necessity of stellar structure and evolution (SSE) programs as tools of inference

Even if you are not a modeller,
your work relies on results
from stellar models

But like any instrument, **stellar structure and evolution codes** are subject to calibration errors, biases, and “black-box” treatment

Barriers to the effective use of stellar structure and evolution programs



Barriers to the effective use of stellar structure and evolution programs



- Hard to develop a deep understanding of software you did not write yourself

Barriers to the effective use of stellar structure and evolution programs



- Hard to develop a deep understanding of software you did not write yourself
- users give up because of non-science technical problems (installation, SDKs, segfaults...)

Barriers to the effective use of stellar structure and evolution programs



- Hard to develop a deep understanding of software you did not write yourself
- users give up because of non-science technical problems (installation, SDKs, segfaults...)
- overinterpretation of results

Barriers to the effective use of stellar structure and evolution programs



- Hard to develop a deep understanding of software you did not write yourself
- users give up because of non-science technical problems (installation, SDKs, segfaults...)
- overinterpretation of results
- Not enough attention given to modeling uncertainties and parameter choices
“If I don’t know what this parameter means, I ignore it...”

Barriers to the effective use of stellar structure and evolution programs



- Hard to develop a deep understanding of software you did not write yourself
- users give up because of non-science technical problems (installation, SDKs, segfaults...)
- overinterpretation of results
- Not enough attention given to modeling uncertainties and parameter choices
“If I don’t know what this parameter means, I ignore it...”

These are powerful tools, but only if we use them properly and understand their limitations!