

# Consequences of ignoring dispersal variation in network models for landscape connectivity

DOI:10.1111/COBI.13640

Lauren L. Sullivan<sup>1,2</sup>, Matthew J. Michalska-Smith<sup>3,4</sup>, Katie P. Sperry<sup>2,5</sup>, David A. Moeller<sup>6</sup>, and Allison K. Shaw<sup>2</sup>

<sup>1</sup>Division of Biological Sciences, University of Missouri, Columbia, MO, U.S.A.

<sup>2</sup>Department of Ecology, Evolution, and Behavior, University of Minnesota, Saint Paul, MN, U.S.A.

<sup>3</sup>Department of Veterinary Population Medicine, University of Minnesota, Saint Paul, MN, U.S.A.

<sup>4</sup>Department of Plant Pathology, University of Minnesota, Saint Paul, MN, U.S.A.

<sup>5</sup>Marine and Environmental Sciences, Northeastern University, Boston, MA, U.S.A.

<sup>6</sup>Department of Plant and Microbial Biology, University of Minnesota, Saint Paul, MN, U.S.A.

First published: 25 September 2020

This is the peer reviewed version of the following article:

LL Sullivan, MJ Michalska-Smith, KP Sperry, DA Moeller, and AK Shaw. "Consequences of ignoring dispersal variation in network models for landscape connectivity." *Conservation Biology* 35, no. 3 (2021): 944-954.

which has been published in final form at <https://doi.org/10.1111/cobi.13640>. This article may be used for non-commercial purposes in accordance with Wiley Terms and Conditions for Use of Self-Archived Versions. This article may not be enhanced, enriched or otherwise transformed into a derivative work, without express permission from Wiley or by statutory rights under applicable legislation. Copyright notices must not be removed, obscured or modified. The article must be linked to Wiley's version of record on Wiley Online Library and any embedding, framing or otherwise making available the article or pages thereof by third parties from platforms, services and websites other than Wiley Online Library must be prohibited.

**Abstract**

Habitat loss and fragmentation can negatively impact population persistence and biodiversity, but these effects can be mitigated if species successfully disperse between isolated habitat patches. Network models are the primary tool for quantifying landscape connectivity, yet as practiced, they take an overly simplistic view of species dispersal. These models often ignore individual variation in dispersal ability, assuming all individuals move the same fixed distance with equal probability. Here we develop a modeling approach to address this problem. We incorporate dispersal kernels into network models to determine how individual variation in dispersal alters our understanding of landscape-level connectivity, and test our approach on a fragmented grassland landscape in Minnesota. We show that ignoring dispersal variation consistently overestimates a population’s robustness to local extinctions while simultaneously underestimating its robustness to local habitat loss. Furthermore, a simplified view of dispersal underestimates the amount of habitat sub-structure for small populations but overestimates habitat sub-structure for large populations. Our results demonstrate that considering biologically realistic dispersal alters our understanding of landscape connectivity for ecological theory and conservation practice.

**Introduction**

Loss of habitat due to land-use alteration is one of the largest anthropogenic threats to Earth’s planetary systems (Rockström 2009), contributing to major declines in biodiversity (Newbold et al. 2016) and other ecosystem services (Haddad et al. 2015). Habitat loss fundamentally alters landscapes by simultaneously decreasing the overall amount of native habitat, and changing how the remaining habitat patches are arranged with respect to each other