

Relative K -theory for C^* -algebras and factor groupoids

Groups, Operators, and Banach Algebras Seminar

Mitch Haslehurst

Department of Mathematics & Statistics
University of Victoria

June 7, 2021

Let A be a C^* -algebra.

Let A be a C^* -algebra.

Two projections are *Murray-von Neumann* equivalent if there is a partial isometry v with $v^*v = p$ and $vv^* = q$.

Let A be a C^* -algebra.

Two projections are *Murray-von Neumann* equivalent if there is a partial isometry v with $v^*v = p$ and $vv^* = q$.

The abelian group $K_0(A)$ consists of stable Murray-von Neumann equivalence classes of projections p in $M_\infty(\tilde{A}) = \bigcup_n M_n(\tilde{A})$, with group operation diagonal sum: $[p] + [q]$ is the class of

$$p \oplus q = \begin{bmatrix} p & 0 \\ 0 & q \end{bmatrix}.$$

The abelian group $K_1(A)$ consists of stable homotopy equivalence classes of unitaries u in $M_\infty(\tilde{A}) = \bigcup_n M_n(\tilde{A})$, with group operation diagonal sum: $[u] + [v]$ is the class of $u \oplus v = \begin{bmatrix} u & 0 \\ 0 & v \end{bmatrix}$.

The abelian group $K_1(A)$ consists of stable homotopy equivalence classes of unitaries u in $M_\infty(\tilde{A}) = \bigcup_n M_n(\tilde{A})$, with group operation diagonal sum: $[u] + [v]$ is the class of $u \oplus v = \begin{bmatrix} u & 0 \\ 0 & v \end{bmatrix}$.

K_0 and K_1 are functors: if $\phi : A \rightarrow B$ is a $*$ -homomorphism, then there are induced group homomorphisms $\phi_* : K_j(A) \rightarrow K_j(B)$ that satisfy

$$\phi_*([a]) = [\phi(a)]$$

If the sequence

$$0 \longrightarrow I \hookrightarrow A \xrightarrow{\pi} A/I \longrightarrow 0$$

of C^* -algebras is exact,

If the sequence

$$0 \longrightarrow I \hookrightarrow A \xrightarrow{\pi} A/I \longrightarrow 0$$

of C^* -algebras is exact, there are group homomorphisms δ_0 and δ_1 such that the sequence

$$\begin{array}{ccccc} K_0(I) & \xrightarrow{\iota_*} & K_0(A) & \xrightarrow{\pi_*} & K_0(A/I) \\ \delta_1 \uparrow & & & & \downarrow \delta_0 \\ K_1(A/I) & \xleftarrow{\pi_*} & K_1(A) & \xleftarrow{\iota_*} & K_1(I) \end{array}$$

is exact.

Let $\phi : A \rightarrow B$ be a $*$ -homomorphism. Let $\Gamma_0(\phi)$ be all triples (p, q, v) where $p, q \in M_\infty(\tilde{A})$ are projections and $v \in M_\infty(\tilde{B})$ is a partial isometry with $v^*v = \phi(p)$ and $vv^* = \phi(q)$.

Let $\phi : A \rightarrow B$ be a $*$ -homomorphism. Let $\Gamma_0(\phi)$ be all triples (p, q, v) where $p, q \in M_\infty(\tilde{A})$ are projections and $v \in M_\infty(\tilde{B})$ is a partial isometry with $v^*v = \phi(p)$ and $vv^* = \phi(q)$.

We can add them:

$$(p, q, v) \oplus (p', q', v') = \left(\begin{bmatrix} p & 0 \\ 0 & p' \end{bmatrix}, \begin{bmatrix} q & 0 \\ 0 & q' \end{bmatrix}, \begin{bmatrix} v & 0 \\ 0 & v' \end{bmatrix} \right)$$

Let $\phi : A \rightarrow B$ be a $*$ -homomorphism. Let $\Gamma_0(\phi)$ be all triples (p, q, v) where $p, q \in M_\infty(\tilde{A})$ are projections and $v \in M_\infty(\tilde{B})$ is a partial isometry with $v^*v = \phi(p)$ and $vv^* = \phi(q)$.

We can add them:

$$(p, q, v) \oplus (p', q', v') = \left(\begin{bmatrix} p & 0 \\ 0 & p' \end{bmatrix}, \begin{bmatrix} q & 0 \\ 0 & q' \end{bmatrix}, \begin{bmatrix} v & 0 \\ 0 & v' \end{bmatrix} \right)$$

Say $(p, q, v) \cong (p', q', v')$ if there are partial isometries $c, d \in M_\infty(\tilde{A})$ with $c^*c = p$, $cc^* = p'$, $d^*d = q$, $dd^* = q'$, and $\phi(d)v = v'\phi(c)$.

Say (p, p, v) is elementary if there is a homotopy v_t with $v_1 = v$, $v_0 = \phi(p)$, and $v_t^* v_t = v_t v_t^* = \phi(p)$ for all t .

Say (p, p, v) is elementary if there is a homotopy v_t with $v_1 = v$, $v_0 = \phi(p)$, and $v_t^* v_t = v_t v_t^* = \phi(p)$ for all t . Equivalently, v is a unitary homotopic to the identity in $\phi(p)M_\infty(\tilde{B})\phi(p)$.

Say (p, p, v) is elementary if there is a homotopy v_t with $v_1 = v$, $v_0 = \phi(p)$, and $v_t^* v_t = v_t v_t^* = \phi(p)$ for all t . Equivalently, v is a unitary homotopic to the identity in $\phi(p)M_\infty(\tilde{B})\phi(p)$.

Say $(p, q, v) \sim (p', q', v')$ if there are elementary triples (r, r, c) and (s, s, d) such that

$$(p, q, v) \oplus (r, r, c) \cong (p', q', v') \oplus (s, s, d)$$

Say (p, p, v) is elementary if there is a homotopy v_t with $v_1 = v$, $v_0 = \phi(p)$, and $v_t^* v_t = v_t v_t^* = \phi(p)$ for all t . Equivalently, v is a unitary homotopic to the identity in $\phi(p)M_\infty(\tilde{B})\phi(p)$.

Say $(p, q, v) \sim (p', q', v')$ if there are elementary triples (r, r, c) and (s, s, d) such that

$$(p, q, v) \oplus (r, r, c) \cong (p', q', v') \oplus (s, s, d)$$

Definition

$$K_0(\phi) := \Gamma_0(\phi) / \sim$$

Properties of $K_0(\phi)$:

- 1 It is an abelian group with addition being direct sum,

Properties of $K_0(\phi)$:

- 1 It is an abelian group with addition being direct sum,
- 2 $[0, 0, 0] = 0$,

Properties of $K_0(\phi)$:

- ① It is an abelian group with addition being direct sum,
- ② $[0, 0, 0] = 0$,
- ③ $-[p, q, v] = [q, p, v^*]$,

Properties of $K_0(\phi)$:

- ① It is an abelian group with addition being direct sum,
- ② $[0, 0, 0] = 0$,
- ③ $-[p, q, v] = [q, p, v^*]$,
- ④ If $pp' = qq' = 0$, then
$$[p, q, v] + [p', q', v'] = [p + p', q + q', v + v'],$$

Properties of $K_0(\phi)$:

- ① It is an abelian group with addition being direct sum,
- ② $[0, 0, 0] = 0$,
- ③ $-[p, q, v] = [q, p, v^*]$,
- ④ If $pp' = qq' = 0$, then
$$[p, q, v] + [p', q', v'] = [p + p', q + q', v + v'],$$
- ⑤ $[v^*v, vv^*, \phi(v)] = 0$ for v a partial isometry in $M_\infty(\tilde{A})$,

Properties of $K_0(\phi)$:

- ① It is an abelian group with addition being direct sum,
- ② $[0, 0, 0] = 0$,
- ③ $-[p, q, v] = [q, p, v^*]$,
- ④ If $pp' = qq' = 0$, then
$$[p, q, v] + [p', q', v'] = [p + p', q + q', v + v'],$$
- ⑤ $[v^*v, vv^*, \phi(v)] = 0$ for v a partial isometry in $M_\infty(\tilde{A})$,
- ⑥ $[p, q, v] = [p, q, v']$ if v and v' are homotopic,

Properties of $K_0(\phi)$:

- ① It is an abelian group with addition being direct sum,
- ② $[0, 0, 0] = 0$,
- ③ $-[p, q, v] = [q, p, v^*]$,
- ④ If $pp' = qq' = 0$, then
$$[p, q, v] + [p', q', v'] = [p + p', q + q', v + v'],$$
- ⑤ $[v^*v, vv^*, \phi(v)] = 0$ for v a partial isometry in $M_\infty(\tilde{A})$,
- ⑥ $[p, q, v] = [p, q, v']$ if v and v' are homotopic,
- ⑦ $[p, q, v] + [q, r, v'] = [p, r, v'v]$.

Let $\Gamma_1(\phi)$ be all triples (p, u, g) where $p \in M_\infty(\tilde{A})$ is a projection, u is a unitary in $pM_\infty(\tilde{A})p$, and g is a unitary in $C[0, 1] \otimes \phi(p)M_\infty(\tilde{B})\phi(p)$ with $g(0) = \phi(p)$ and $g(1) = \phi(u)$.

Let $\Gamma_1(\phi)$ be all triples (p, u, g) where $p \in M_\infty(\tilde{A})$ is a projection, u is a unitary in $pM_\infty(\tilde{A})p$, and g is a unitary in $C[0, 1] \otimes \phi(p)M_\infty(\tilde{B})\phi(p)$ with $g(0) = \phi(p)$ and $g(1) = \phi(u)$.

We can add them by diagonal sum, as with $\Gamma_0(\phi)$.

Let $\Gamma_1(\phi)$ be all triples (p, u, g) where $p \in M_\infty(\tilde{A})$ is a projection, u is a unitary in $pM_\infty(\tilde{A})p$, and g is a unitary in $C[0, 1] \otimes \phi(p)M_\infty(\tilde{B})\phi(p)$ with $g(0) = \phi(p)$ and $g(1) = \phi(u)$.

We can add them by diagonal sum, as with $\Gamma_0(\phi)$.

Say $(p, u, g) \cong (p', u', g')$ if there is a partial isometry $v \in M_\infty(\tilde{A})$ with $v^*v = p$, $vv^* = p'$, $vu = u'v$, and $vg(s) = g'(s)v$ for all $0 \leq s \leq 1$.

Say (p, u, g) is elementary if there are homotopies u_t and g_t with $u_1 = u$, $g_1 = g$, $u_0 = p$, $g_0(s) = \phi(p)$ for all $0 \leq s \leq 1$, and $g_t(1) = \phi(u_t)$ for all $0 \leq t \leq 1$.

Say (p, u, g) is elementary if there are homotopies u_t and g_t with $u_1 = u$, $g_1 = g$, $u_0 = p$, $g_0(s) = \phi(p)$ for all $0 \leq s \leq 1$, and $g_t(1) = \phi(u_t)$ for all $0 \leq t \leq 1$.

Say $(p, u, g) \sim (p', u', g')$ if there are elementary triples (p'', u'', g'') and (p''', u''', g''') such that

$$(p, u, g) \oplus (p'', u'', g'') \cong (p', u', g') \oplus (p''', u''', g''')$$

Say (p, u, g) is elementary if there are homotopies u_t and g_t with $u_1 = u$, $g_1 = g$, $u_0 = p$, $g_0(s) = \phi(p)$ for all $0 \leq s \leq 1$, and $g_t(1) = \phi(u_t)$ for all $0 \leq t \leq 1$.

Say $(p, u, g) \sim (p', u', g')$ if there are elementary triples (p'', u'', g'') and (p''', u''', g''') such that

$$(p, u, g) \oplus (p'', u'', g'') \cong (p', u', g') \oplus (p''', u''', g''')$$

Definition

$$K_1(\phi) := \Gamma_1(\phi) / \sim$$

Every element of $K_1(\phi)$ is represented by $(1_n, u, g)$, for some n .

Properties of $K_1(\phi)$:

- 1 It is an abelian group with addition being direct sum,

Properties of $K_1(\phi)$:

- 1 It is an abelian group with addition being direct sum,
- 2 $[0, 0, 0] = 0$,

Properties of $K_1(\phi)$:

- ① It is an abelian group with addition being direct sum,
- ② $[0, 0, 0] = 0$,
- ③ $-[p, u, g] = [p, u^*, g^*]$,

Properties of $K_1(\phi)$:

- 1 It is an abelian group with addition being direct sum,
- 2 $[0, 0, 0] = 0$,
- 3 $-[p, u, g] = [p, u^*, g^*]$,
- 4 If $pp' = 0$, then $[p, u, g] + [p', u', g'] = [p + p', u + u', g + g']$,

Properties of $K_1(\phi)$:

- ① It is an abelian group with addition being direct sum,
- ② $[0, 0, 0] = 0$,
- ③ $-[p, u, g] = [p, u^*, g^*]$,
- ④ If $pp' = 0$, then $[p, u, g] + [p', u', g'] = [p + p', u + u', g + g']$,
- ⑤ $[p, u, g] = 0$ if $g(s) = \phi(u_s)$ for a path u_s in $M_\infty(\tilde{A})$ with $u_1 = u$ and $u_0 = p$,

Properties of $K_1(\phi)$:

- ① It is an abelian group with addition being direct sum,
- ② $[0, 0, 0] = 0$,
- ③ $-[p, u, g] = [p, u^*, g^*]$,
- ④ If $pp' = 0$, then $[p, u, g] + [p', u', g'] = [p + p', u + u', g + g']$,
- ⑤ $[p, u, g] = 0$ if $g(s) = \phi(u_s)$ for a path u_s in $M_\infty(\tilde{A})$ with $u_1 = u$ and $u_0 = p$,
- ⑥ $[p, u, g] = [p, u', g']$ if u and u' are homotopic via u_t , g and g' are homotopic via g_t , and $g_t(1) = \phi(u_t)$,

Properties of $K_1(\phi)$:

- ① It is an abelian group with addition being direct sum,
- ② $[0, 0, 0] = 0$,
- ③ $-[p, u, g] = [p, u^*, g^*]$,
- ④ If $pp' = 0$, then $[p, u, g] + [p', u', g'] = [p + p', u + u', g + g']$,
- ⑤ $[p, u, g] = 0$ if $g(s) = \phi(u_s)$ for a path u_s in $M_\infty(\tilde{A})$ with $u_1 = u$ and $u_0 = p$,
- ⑥ $[p, u, g] = [p, u', g']$ if u and u' are homotopic via u_t , g and g' are homotopic via g_t , and $g_t(1) = \phi(u_t)$,
- ⑦ $[p, u, g] + [p, u', g'] = [p, uu', gg']$.

Relative K -theory

Define the maps

$$\nu_0 : K_0(\phi) \rightarrow K_0(A) : [p, q, v] \mapsto [p] - [q]$$

$$\nu_1 : K_1(\phi) \rightarrow K_1(A) : [1_n, u, g] \mapsto [u]$$

Relative K -theory

Define the maps

$$\nu_0 : K_0(\phi) \rightarrow K_0(A) : [p, q, v] \mapsto [p] - [q]$$

$$\nu_1 : K_1(\phi) \rightarrow K_1(A) : [1_n, u, g] \mapsto [u]$$

$$\mu_0 : K_1(B) \rightarrow K_0(\phi) : [u] \mapsto [1_n, 1_n, u]$$

$$\mu_1 : K_0(B) \rightarrow K_1(\phi) : [p] - [q] \mapsto [1_n, 1_n, f_p f_q^*]$$

where $f_p(t) = e^{2\pi i t p}$.

Define the maps

$$\nu_0 : K_0(\phi) \rightarrow K_0(A) : [p, q, v] \mapsto [p] - [q]$$

$$\nu_1 : K_1(\phi) \rightarrow K_1(A) : [1_n, u, g] \mapsto [u]$$

$$\mu_0 : K_1(B) \rightarrow K_0(\phi) : [u] \mapsto [1_n, 1_n, u]$$

$$\mu_1 : K_0(B) \rightarrow K_1(\phi) : [p] - [q] \mapsto [1_n, 1_n, f_p f_q^*]$$

where $f_p(t) = e^{2\pi i t p}$. Then we have the six-term exact sequence

$$\begin{array}{ccccc} K_1(B) & \xrightarrow{\mu_0} & K_0(\phi) & \xrightarrow{\nu_0} & K_0(A) \\ \uparrow \phi_* & & & & \downarrow \phi_* \\ K_1(A) & \xleftarrow{\nu_1} & K_1(\phi) & \xleftarrow{\mu_1} & K_0(B) \end{array}$$

Example 1.

- 1 If $\pi : A \rightarrow A/I$ is a quotient map, then $K_j(\pi) \cong K_j(I)$.

Example 1.

- 1 If $\pi : A \rightarrow A/I$ is a quotient map, then $K_j(\pi) \cong K_j(I)$.

This is a noncommutative version of a standard excision result:
If $Y \subseteq X$, then the restriction map $\pi : C(X) \rightarrow C(Y)$ is a quotient map with kernel $C_0(X - Y)$, and

Example 1.

- ❶ If $\pi : A \rightarrow A/I$ is a quotient map, then $K_j(\pi) \cong K_j(I)$.

This is a noncommutative version of a standard excision result:
If $Y \subseteq X$, then the restriction map $\pi : C(X) \rightarrow C(Y)$ is a quotient map with kernel $C_0(X - Y)$, and

$$K_j(\pi) \cong K_j(C_0(X - Y)) \cong K^j(X, Y)$$

Example 1.

- ❶ If $\pi : A \rightarrow A/I$ is a quotient map, then $K_j(\pi) \cong K_j(I)$.

This is a noncommutative version of a standard excision result:
If $Y \subseteq X$, then the restriction map $\pi : C(X) \rightarrow C(Y)$ is a quotient map with kernel $C_0(X - Y)$, and

$$K_j(\pi) \cong K_j(C_0(X - Y)) \cong K^j(X, Y)$$

- ❷ If $K_*(B) = 0$, then $K_*(\phi) = K_*(A)$. If $K_*(A) = 0$, then $K_j(\phi) = K_{1-j}(B)$.

Example 1.

- 1 If $\pi : A \rightarrow A/I$ is a quotient map, then $K_j(\pi) \cong K_j(I)$.

This is a noncommutative version of a standard excision result:
If $Y \subseteq X$, then the restriction map $\pi : C(X) \rightarrow C(Y)$ is a quotient map with kernel $C_0(X - Y)$, and

$$K_j(\pi) \cong K_j(C_0(X - Y)) \cong K^j(X, Y)$$

- 2 If $K_*(B) = 0$, then $K_*(\phi) = K_*(A)$. If $K_*(A) = 0$, then $K_j(\phi) = K_{1-j}(B)$.
- 3 If ϕ is a $*$ -isomorphism, then $K_*(\phi) = 0$.

Example 2. Let D be a C^* -algebra, $B = M_2(D)$, and $A \subseteq B$ the subalgebra of diagonal matrices. With $\phi : A \rightarrow B$ being the inclusion map, denote $K_j(A; B) := K_j(\phi)$.

Example 2. Let D be a C^* -algebra, $B = M_2(D)$, and $A \subseteq B$ the subalgebra of diagonal matrices. With $\phi : A \rightarrow B$ being the inclusion map, denote $K_j(A; B) := K_j(\phi)$.

$$\begin{array}{ccccc}
 K_1(D) & \xrightarrow{\mu_0} & K_0(A; B) & \xrightarrow{\nu_0} & K_0(D) \oplus K_0(D) \\
 \uparrow \phi_* & & & & \downarrow \phi_* \\
 K_1(D) \oplus K_1(D) & \xleftarrow{\nu_1} & K_1(A; B) & \xleftarrow{\mu_1} & K_0(D)
 \end{array}$$

Example 2. Let D be a C^* -algebra, $B = M_2(D)$, and $A \subseteq B$ the subalgebra of diagonal matrices. With $\phi : A \rightarrow B$ being the inclusion map, denote $K_j(A; B) := K_j(\phi)$.

$$\begin{array}{ccccc}
 K_1(D) & \xrightarrow{\mu_0} & K_0(A; B) & \xrightarrow{\nu_0} & K_0(D) \oplus K_0(D) \\
 \uparrow \phi_* & & & & \downarrow \phi_* \\
 K_1(D) \oplus K_1(D) & \xleftarrow{\nu_1} & K_1(A; B) & \xleftarrow{\mu_1} & K_0(D)
 \end{array}$$

The vertical maps are surjective, so $K_j(A; B) \cong \ker \phi_* \cong K_j(D)$.

Let \mathcal{H} be a separable Hilbert space and \mathcal{M} a proper nonzero subspace. Regard $A = \mathcal{K}(\mathcal{M}) \oplus \mathcal{K}(\mathcal{M}^\perp)$ as a subalgebra of $B = \mathcal{K}(\mathcal{H})$ via $(a, b) \mapsto \begin{bmatrix} a & 0 \\ 0 & b \end{bmatrix}$.

Let \mathcal{H} be a separable Hilbert space and \mathcal{M} a proper nonzero subspace. Regard $A = \mathcal{K}(\mathcal{M}) \oplus \mathcal{K}(\mathcal{M}^\perp)$ as a subalgebra of

$$B = \mathcal{K}(\mathcal{H}) \text{ via } (a, b) \mapsto \begin{bmatrix} a & 0 \\ 0 & b \end{bmatrix}.$$

Let $\xi \in \mathcal{M}$ and $\eta \in \mathcal{M}^\perp$ be nonzero vectors, and p_ξ, p_η be the projections onto $\text{span}\{\xi\}$ and $\text{span}\{\eta\}$.

Let \mathcal{H} be a separable Hilbert space and \mathcal{M} a proper nonzero subspace. Regard $A = \mathcal{K}(\mathcal{M}) \oplus \mathcal{K}(\mathcal{M}^\perp)$ as a subalgebra of $B = \mathcal{K}(\mathcal{H})$ via $(a, b) \mapsto \begin{bmatrix} a & 0 \\ 0 & b \end{bmatrix}$.

Let $\xi \in \mathcal{M}$ and $\eta \in \mathcal{M}^\perp$ be nonzero vectors, and p_ξ, p_η be the projections onto $\text{span}\{\xi\}$ and $\text{span}\{\eta\}$.

Then $K_0(A; B) \cong \mathbb{Z}$ is generated by the triple (p_ξ, p_η, v) , where v is a partial isometry in $\mathcal{K}(\mathcal{H})$ with $v^*v = p_\xi$ and $vv^* = p_\eta$.

Example 3. Let D be a C^* -algebra, $B = D \oplus D$, and $A = \{(a, a) \mid a \in D\} \subseteq B$, which we identify with D . This time the vertical maps are injective, and we have $K_j(A; B) \cong K_{1-j}(D)$.

Example 3. Let D be a C^* -algebra, $B = D \oplus D$, and $A = \{(a, a) \mid a \in D\} \subseteq B$, which we identify with D . This time the vertical maps are injective, and we have $K_j(A; B) \cong K_{1-j}(D)$.

In the case that $D = \mathcal{K}$, $K_1(A; B) \cong \mathbb{Z}$ is generated by the triple (p, p, g) , where p is any rank one projection in \mathcal{K} and $g(t) = (e^{2\pi it} p, p)$.

Suppose we have a commutative diagram

$$\begin{array}{ccc} A & \xrightarrow{\alpha} & C \\ \downarrow \phi & & \downarrow \psi \\ B & \xrightarrow{\beta} & D \end{array}$$

Suppose we have a commutative diagram

$$\begin{array}{ccc} A & \xrightarrow{\alpha} & C \\ \downarrow \phi & & \downarrow \psi \\ B & \xrightarrow{\beta} & D \end{array}$$

Then there is a group homomorphism $\alpha_* : K_0(\phi) \rightarrow K_0(\psi)$ such that

$$\alpha_*([p, q, v]) = [\alpha(p), \alpha(q), \beta(v)]$$

and similarly for $K_1(\phi) \rightarrow K_1(\psi)$.

If

$$\begin{array}{ccccccccc} 0 & \longrightarrow & I & \xrightarrow{\iota_A} & A & \xrightarrow{\pi_A} & A/I & \longrightarrow & 0 \\ & & \downarrow \psi & & \downarrow \phi & & \downarrow \gamma & & \\ 0 & \longrightarrow & J & \xrightarrow{\iota_B} & B & \xrightarrow{\pi_B} & B/J & \longrightarrow & 0 \end{array}$$

is commutative with exact rows,

If

$$\begin{array}{ccccccccc}
 0 & \longrightarrow & I & \xrightarrow{\iota_A} & A & \xrightarrow{\pi_A} & A/I & \longrightarrow & 0 \\
 & & \downarrow \psi & & \downarrow \phi & & \downarrow \gamma & & \\
 0 & \longrightarrow & J & \xrightarrow{\iota_B} & B & \xrightarrow{\pi_B} & B/J & \longrightarrow & 0
 \end{array}$$

is commutative with exact rows, we have a six-term exact sequence

$$\begin{array}{ccccc}
 K_0(\psi) & \xrightarrow{\iota_*} & K_0(\phi) & \xrightarrow{\pi_*} & K_0(\gamma) \\
 \uparrow \partial_1 & & & & \downarrow \partial_0 \\
 K_1(\gamma) & \xleftarrow{\pi_*} & K_1(\phi) & \xleftarrow{\iota_*} & K_1(\psi)
 \end{array}$$

Consider the commutative diagram

$$\begin{array}{ccccccc} 0 & \longrightarrow & C_0(\mathbb{R}) & \hookrightarrow & C(S^1) & \xrightarrow{\text{ev}_1} & \mathbb{C} \longrightarrow 0 \\ & & \parallel & & \downarrow \phi & & \downarrow \gamma \\ 0 & \longrightarrow & C_0(\mathbb{R}) & \hookrightarrow & C[0,1] & \xrightarrow{\text{ev}_{0,1}} & \mathbb{C} \oplus \mathbb{C} \longrightarrow 0 \end{array}$$

where $\phi(f)(t) = f(e^{2\pi it})$ and $\gamma(z) = (z, z)$.

Consider the commutative diagram

$$\begin{array}{ccccccc}
 0 & \longrightarrow & C_0(\mathbb{R}) & \hookrightarrow & C(S^1) & \xrightarrow{\text{ev}_1} & \mathbb{C} \longrightarrow 0 \\
 & & \parallel & & \downarrow \phi & & \downarrow \gamma \\
 0 & \longrightarrow & C_0(\mathbb{R}) & \hookrightarrow & C[0,1] & \xrightarrow{\text{ev}_{0,1}} & \mathbb{C} \oplus \mathbb{C} \longrightarrow 0
 \end{array}$$

where $\phi(f)(t) = f(e^{2\pi it})$ and $\gamma(z) = (z, z)$.

$$\begin{array}{ccccc}
 0 & \longrightarrow & K_0(C(S^1); C[0,1]) & \xrightarrow{\text{ev}_*} & K_0(\mathbb{C}; \mathbb{C} \oplus \mathbb{C}) \\
 \uparrow & & & & \downarrow \\
 K_1(\mathbb{C}; \mathbb{C} \oplus \mathbb{C}) & \xleftarrow{\text{ev}_*} & K_1(C(S^1); C[0,1]) & \xleftarrow{\quad} & 0
 \end{array}$$

Consider the commutative diagram

$$\begin{array}{ccccccc}
 0 & \longrightarrow & C_0(\mathbb{R}) & \hookrightarrow & C(S^1) & \xrightarrow{\text{ev}_1} & \mathbb{C} \longrightarrow 0 \\
 & & \parallel & & \downarrow \phi & & \downarrow \gamma \\
 0 & \longrightarrow & C_0(\mathbb{R}) & \hookrightarrow & C[0,1] & \xrightarrow{\text{ev}_{0,1}} & \mathbb{C} \oplus \mathbb{C} \longrightarrow 0
 \end{array}$$

where $\phi(f)(t) = f(e^{2\pi i t})$ and $\gamma(z) = (z, z)$.

$$\begin{array}{ccccc}
 0 & \longrightarrow & K_0(C(S^1); C[0,1]) & \xrightarrow{\text{ev}_*} & K_0(\mathbb{C}; \mathbb{C} \oplus \mathbb{C}) \\
 \uparrow & & & & \downarrow \\
 K_1(\mathbb{C}; \mathbb{C} \oplus \mathbb{C}) & \xleftarrow{\text{ev}_*} & K_1(C(S^1); C[0,1]) & \xleftarrow{\quad} & 0
 \end{array}$$

So $K_*(C(S^1); C[0,1]) \cong K_*(\mathbb{C}; \mathbb{C} \oplus \mathbb{C})$.

Consider the diagram

$$\begin{array}{ccccccccc} 0 & \longrightarrow & C_0(\mathbb{R}^2) & \hookrightarrow & C(\mathbb{D}) & \longrightarrow & C(S^1) & \longrightarrow & 0 \\ & & \downarrow & & \downarrow \phi & & \downarrow & & \\ 0 & \longrightarrow & 0 & \longrightarrow & C[0,1] & \xlongequal{\quad} & C[0,1] & \longrightarrow & 0 \end{array}$$

where ϕ is the composition of the restriction to S^1 and the inclusion to $[0,1]$.

Consider the diagram

$$\begin{array}{ccccccccc}
 0 & \longrightarrow & C_0(\mathbb{R}^2) & \hookrightarrow & C(\mathbb{D}) & \longrightarrow & C(S^1) & \longrightarrow & 0 \\
 & & \downarrow & & \downarrow \phi & & \downarrow & & \\
 0 & \longrightarrow & 0 & \longrightarrow & C[0, 1] & \xlongequal{\quad} & C[0, 1] & \longrightarrow & 0
 \end{array}$$

where ϕ is the composition of the restriction to S^1 and the inclusion to $[0, 1]$.

$$\begin{array}{ccccccc}
 K_0(C_0(\mathbb{R}^2)) & \longrightarrow & 0 & \longrightarrow & K_0(C(S^1); C[0, 1]) \\
 \uparrow & & & & \downarrow \\
 K_1(C(S^1); C[0, 1]) & \longleftarrow & 0 & \longleftarrow & K_1(C_0(\mathbb{R}^2))
 \end{array}$$

The map $\partial_1 : K_1(C(S^1); C[0, 1]) \rightarrow K_0(C_0(\mathbb{R}^2))$ is an isomorphism.

The map $\partial_1 : K_1(C(S^1); C[0, 1]) \rightarrow K_0(C_0(\mathbb{R}^2))$ is an isomorphism.

$K_1(C(S^1); C[0, 1])$ is generated by $(1, e^{2\pi it}, e^{2\pi ist})$, where $0 \leq s, t \leq 1$.

The map $\partial_1 : K_1(C(S^1); C[0, 1]) \rightarrow K_0(C_0(\mathbb{R}^2))$ is an isomorphism.

$K_1(C(S^1); C[0, 1])$ is generated by $(1, e^{2\pi it}, e^{2\pi ist})$, where $0 \leq s, t \leq 1$.

The map ∂_1 takes this element to $[p] - [1 \oplus 0]$ in $K_0(C_0(\mathbb{R}^2))$, where p is the Bott projection:

$$p(z) = \begin{bmatrix} |z|^2 & z(1 - |z|^2)^{1/2} \\ \bar{z}(1 - |z|^2)^{1/2} & 1 - |z|^2 \end{bmatrix}$$

for z in the unit disk $\mathbb{D} \subseteq \mathbb{C}$

Relative K -theory

The mapping cone C_ϕ gives an alternative portrait of relative K -theory.

Relative K -theory

The mapping cone C_ϕ gives an alternative portrait of relative K -theory.

$$C_\phi = \{(a, f) \mid f(1) = \phi(a)\} \subseteq A \oplus (C_0(0, 1] \otimes B)$$

Relative K -theory

The mapping cone C_ϕ gives an alternative portrait of relative K -theory.

$$C_\phi = \{(a, f) \mid f(1) = \phi(a)\} \subseteq A \oplus (C_0(0, 1] \otimes B)$$

There are natural isomorphisms $\Delta_j : K_j(\phi) \rightarrow K_j(C_\phi)$ for $j = 0, 1$:

Relative K -theory

The mapping cone C_ϕ gives an alternative portrait of relative K -theory.

$$C_\phi = \{(a, f) \mid f(1) = \phi(a)\} \subseteq A \oplus (C_0(0, 1] \otimes B)$$

There are natural isomorphisms $\Delta_j : K_j(\phi) \rightarrow K_j(C_\phi)$ for $j = 0, 1$:

$$\begin{array}{ccc} A & \xrightarrow{\alpha} & C \\ \downarrow \phi & & \downarrow \psi \\ B & \xrightarrow{\beta} & D \end{array}$$

commutes, then so does

Relative K -theory

The mapping cone C_ϕ gives an alternative portrait of relative K -theory.

$$C_\phi = \{(a, f) \mid f(1) = \phi(a)\} \subseteq A \oplus (C_0(0, 1] \otimes B)$$

There are natural isomorphisms $\Delta_j : K_j(\phi) \rightarrow K_j(C_\phi)$ for $j = 0, 1$:

$$\begin{array}{ccc} A & \xrightarrow{\alpha} & C \\ \downarrow \phi & & \downarrow \psi \\ B & \xrightarrow{\beta} & D \end{array}$$

commutes, then so does

$$\begin{array}{ccc} K_j(\phi) & \xrightarrow{\alpha_*} & K_j(\psi) \\ \downarrow \Delta_j & & \downarrow \Delta_j \\ K_j(C_\phi) & \xrightarrow{\alpha_*} & K_j(C_\psi) \end{array}$$

Problem. Given some K -theory data, find a groupoid G such that the data is $K_*(C_r^*(G))$.

Problem. Given some K -theory data, find a groupoid G such that the data is $K_*(C_r^*(G))$.

Notable references:

Li, X. "Every classifiable simple C^* -algebra has a Cartan subalgebra". *Invent. math.* 219, 653–699 (2020).

Putnam, I.F. "Some classifiable groupoid C^* -algebras with prescribed K -theory". *Math. Ann.* 370, 1361–1387 (2018).

G' is a *factor groupoid* of G if $\pi : G \rightarrow G'$ is a surjective groupoid morphism.

G' is a *factor groupoid* of G if $\pi : G \rightarrow G'$ is a surjective groupoid morphism.

Assume:

- 1 G and G' are locally compact Hausdorff and étale,
- 2 π is continuous and proper,
- 3 $\pi|_{G^u} : G^u \rightarrow (G')^{\pi(u)}$ is bijective for all u in $G^{(0)}$.

G' is a *factor groupoid* of G if $\pi : G \rightarrow G'$ is a surjective groupoid morphism.

Assume:


- 1 G and G' are locally compact Hausdorff and étale,
- 2 π is continuous and proper,
- 3 $\pi|_{G^u} : G^u \rightarrow (G')^{\pi(u)}$ is bijective for all u in $G^{(0)}$.

Obtain an inclusion $C_r^*(G') \subseteq C_r^*(G)$ via $b \mapsto b \circ \pi$ (b in $C_c(G')$)

Let (V, E) be a Bratteli diagram.


Factor groupoids

Let (V, E) be a Bratteli diagram.



Factor groupoids


Let (V, E) be a Bratteli diagram.



The *infinite path space* X_E of (V, E) is a totally disconnected compact metric space.

Factor groupoids

Let (V, E) be a Bratteli diagram.




The *infinite path space* X_E of (V, E) is a totally disconnected compact metric space.

Tail-equivalence $R_E \subseteq X_E \times X_E$ has an étale topology in which $C_r^*(R_E)$ is an AF-algebra.

Factor groupoids

Let (V, E) be a Bratteli diagram.



The *infinite path space* X_E of (V, E) is a totally disconnected compact metric space.

Tail-equivalence $R_E \subseteq X_E \times X_E$ has an étale topology in which $C_r^*(R_E)$ is an AF-algebra.

Goal: make a factor groupoid of R_E .

Let (V, E) and (W, F) be two Bratteli diagrams.

Let (V, E) and (W, F) be two Bratteli diagrams.

Two *graph embeddings* $\xi^0, \xi^1 : (W, F) \rightarrow (V, E)$ with $\xi^0|_W = \xi^1|_W$ and $\xi^0(F) \cap \xi^1(F) = \emptyset$.

Let (V, E) and (W, F) be two Bratteli diagrams.

Two *graph embeddings* $\xi^0, \xi^1 : (W, F) \rightarrow (V, E)$ with $\xi^0|_W = \xi^1|_W$ and $\xi^0(F) \cap \xi^1(F) = \emptyset$.

Equivalence relation \sim_ξ on X_E :

$$(x_1, x_2, \dots, x_{n_0-1}, x_{n_0}, \xi^0(z_{n_0+1}), \xi^0(z_{n_0+2}), \dots) \quad (1)$$

$$\sim_\xi (x_1, x_2, \dots, x_{n_0-1}, x'_{n_0}, \xi^1(z_{n_0+1}), \xi^1(z_{n_0+2}), \dots) \quad (2)$$

Denote $X_\xi := X_E / \sim_\xi$ and $\rho : X_E \rightarrow X_\xi$ the quotient map.

Denote $X_\xi := X_E / \sim_\xi$ and $\rho : X_E \rightarrow X_\xi$ the quotient map.

Facts:

- ① X_ξ is a second-countable compact Hausdorff space,
- ② the covering dimension of X_ξ is 1,
- ③ each connected component is either a single point or homeomorphic to S^1 .

Example 4. We let (V, E) be the Bratteli diagram with one vertex at each level and two edges at each level. Identify X_E with $\{0, 1\}^\omega$.

(W, F) is a single path, and for f in F , $\xi^j(f) = j$ for $j = 0, 1$.

Example 4. We let (V, E) be the Bratteli diagram with one vertex at each level and two edges at each level. Identify X_E with $\{0, 1\}^\omega$.

(W, F) is a single path, and for f in F , $\xi^j(f) = j$ for $j = 0, 1$.

$$(x_1, x_2, \dots, x_n, 1, 0, 0, 0, 0, \dots) \quad (3)$$

$$\sim_\xi (x_1, x_2, \dots, x_n, 0, 1, 1, 1, 1, \dots) \quad (4)$$

Example 4. We let (V, E) be the Bratteli diagram with one vertex at each level and two edges at each level. Identify X_E with $\{0, 1\}^\omega$.

(W, F) is a single path, and for f in F , $\xi^j(f) = j$ for $j = 0, 1$.

$$(x_1, x_2, \dots, x_n, 1, 0, 0, 0, 0, \dots) \quad (3)$$

$$\sim_\xi (x_1, x_2, \dots, x_n, 0, 1, 1, 1, 1, \dots) \quad (4)$$

$$\{0, 1\}^\omega \rightarrow S^1 : (x_n) \mapsto \exp \left(2\pi i \sum_{n=1}^{\infty} x_n 2^{-n} \right)$$

Example 4. We let (V, E) be the Bratteli diagram with one vertex at each level and two edges at each level. Identify X_E with $\{0, 1\}^\omega$.

(W, F) is a single path, and for f in F , $\xi^j(f) = j$ for $j = 0, 1$.

$$(x_1, x_2, \dots, x_n, 1, 0, 0, 0, 0, \dots) \quad (3)$$

$$\sim_\xi (x_1, x_2, \dots, x_n, 0, 1, 1, 1, 1, \dots) \quad (4)$$

$$\{0, 1\}^\omega \rightarrow S^1 : (x_n) \mapsto \exp \left(2\pi i \sum_{n=1}^{\infty} x_n 2^{-n} \right)$$

The fibres are precisely the \sim_ξ equivalence classes, so X_ξ is homeomorphic to S^1 .

Example 5. Let (V, E) have one vertex and three edges at each level. Identify X_E with $\{0, 1, 2\}^\omega$.

(W, F) is again a single path, and for f in F , $\xi^0(f) = 0$ and $\xi^1(f) = 2$.


Example 5. Let (V, E) have one vertex and three edges at each level. Identify X_E with $\{0, 1, 2\}^\omega$.

(W, F) is again a single path, and for f in F , $\xi^0(f) = 0$ and $\xi^1(f) = 2$.

There is a nested sequence $X_1 \subseteq X_2 \subseteq X_3 \subseteq \cdots \subseteq X_E$ such that


$$X_E = \overline{\bigcup_{n=1}^{\infty} X_n}$$

and each $\rho(X_n)$ is a disjoint union of finitely many circles.




$$\rho(X_1)$$

Factor groupoids




$$\rho(X_2)$$

Factor groupoids




$$\rho(X_3)$$

Factor groupoids




$$\rho(X_4)$$

Factor groupoids



$$\rho(X_5)$$

Factor groupoids



$$\rho(X_6)$$

Let $R_\xi = \rho \times \rho(R_E)$.

With the quotient topology, R_ξ is a second-countable locally compact Hausdorff étale groupoid, and a factor of R_E via $\rho \times \rho : R_E \rightarrow R_\xi$.

Let $R_\xi = \rho \times \rho(R_E)$.

With the quotient topology, R_ξ is a second-countable locally compact Hausdorff étale groupoid, and a factor of R_E via $\rho \times \rho : R_E \rightarrow R_\xi$.

We want to analyze the K -theory of $C_r^*(R_\xi) \subseteq C_r^*(R_E)$.

Factor groupoids

$$\begin{array}{ccccc}
 K_1(C_r^*(R_E)) & \longrightarrow & K_0(C_r^*(R_\xi); C_r^*(R_E)) & \longrightarrow & K_0(C_r^*(R_\xi)) \\
 \uparrow \scriptstyle L_* & & & & \downarrow \scriptstyle L_* \\
 K_1(C_r^*(R_\xi)) & \longleftarrow & K_1(C_r^*(R_\xi); C_r^*(R_E)) & \longleftarrow & K_0(C_r^*(R_E))
 \end{array}$$

Factor groupoids

$$\begin{array}{ccccc}
 K_1(C_r^*(R_E)) & \longrightarrow & K_0(C_r^*(R_\xi); C_r^*(R_E)) & \longrightarrow & K_0(C_r^*(R_\xi)) \\
 \uparrow \scriptstyle L_* & & & & \downarrow \scriptstyle L_* \\
 K_1(C_r^*(R_\xi)) & \longleftarrow & K_1(C_r^*(R_\xi); C_r^*(R_E)) & \longleftarrow & K_0(C_r^*(R_E))
 \end{array}$$

Let

$$H' = \{(x, y) \in R_\xi \mid \#(\rho \times \rho)^{-1}(x, y) > 1\}$$

and $H = (\rho \times \rho)^{-1}(H')$. Then we have *excision*: (Putnam, 2020)

$$K_*(C_r^*(R_\xi); C_r^*(R_E)) \cong K_*(C_r^*(H'); C_r^*(H))$$

Factor groupoids

It turns out that $C_r^*(H')$ is stably isomorphic to $C_r^*(R_F)$, and $C_r^*(H)$ may be identified with $C_r^*(H') \oplus C_r^*(H')$. Moreover,

$$\begin{array}{ccc} C_r^*(H') & \hookrightarrow & C_r^*(H) \\ \parallel & & \downarrow \cong \\ C_r^*(H') & \xrightarrow{a \oplus a} & C_r^*(H') \oplus C_r^*(H') \end{array}$$

commutes.

It turns out that $C_r^*(H')$ is stably isomorphic to $C_r^*(R_F)$, and $C_r^*(H)$ may be identified with $C_r^*(H') \oplus C_r^*(H')$. Moreover,

$$\begin{array}{ccc} C_r^*(H') & \hookrightarrow & C_r^*(H) \\ \parallel & & \downarrow \cong \\ C_r^*(H') & \xrightarrow{a \oplus a} & C_r^*(H') \oplus C_r^*(H') \end{array}$$

commutes.

By Example 3, we have

$$K_j(C_r^*(H'); C_r^*(H)) \cong K_{1-j}(C_r^*(H')) \cong K_{1-j}(C_r^*(R_F))$$

Factor groupoids

$$\begin{array}{ccccc}
 K_1(C_r^*(R_E)) & \longrightarrow & K_0(C_r^*(R_\xi); C_r^*(R_E)) & \longrightarrow & K_0(C_r^*(R_\xi)) \\
 \uparrow \scriptstyle L_* & & & & \downarrow \scriptstyle L_* \\
 K_1(C_r^*(R_\xi)) & \longleftarrow & K_1(C_r^*(R_\xi); C_r^*(R_E)) & \longleftarrow & K_0(C_r^*(R_E))
 \end{array}$$

Factor groupoids

$$\begin{array}{ccccc}
 0 & \longrightarrow & 0 & \longrightarrow & K_0(C_r^*(R_\xi)) \\
 \uparrow & & & & \downarrow \\
 K_1(C_r^*(R_\xi)) & \longleftarrow & K_0(C_r^*(R_F)) & \longleftarrow & K_0(C_r^*(R_E))
 \end{array}$$

Factor groupoids

$$\begin{array}{ccccc} 0 & \longrightarrow & 0 & \longrightarrow & K_0(C_r^*(R_\xi)) \\ \uparrow & & & & \downarrow \\ K_1(C_r^*(R_\xi)) & \longleftarrow & K_0(C_r^*(R_F)) & \longleftarrow & K_0(C_r^*(R_E)) \end{array}$$

If one can show that the lower right map is zero...

Factor groupoids

$$\begin{array}{ccccccc}
 0 & \longrightarrow & 0 & \longrightarrow & K_0(C_r^*(R_\xi)) \\
 \uparrow & & & & \downarrow \cong \\
 K_1(C_r^*(R_\xi)) & \xleftarrow{\cong} & K_0(C_r^*(R_F)) & \xleftarrow{0} & K_0(C_r^*(R_E))
 \end{array}$$

If one can show that the lower right map is zero...

Thus we obtain

$$K_0(C_r^*(R_\xi)) \cong K_0(C_r^*(R_E)) \quad K_1(C_r^*(R_\xi)) \cong K_0(C_r^*(R_F))$$

Thus we obtain

$$K_0(C_r^*(R_\xi)) \cong K_0(C_r^*(R_E)) \quad K_1(C_r^*(R_\xi)) \cong K_0(C_r^*(R_F))$$

Through the set-up $\xi^0, \xi^1 : (W, F) \rightarrow (V, E)$, we can prescribe $K_*(C_r^*(R_\xi))$.

Thank you!

- ① Deeley, R.J.; Putnam, I.F.; Strung, K.R. "Non-homogeneous extensions of Cantor minimal systems". to appear, Proc. A.M.S.
- ② Haslehurst, M.J. "Relative K -theory for C^* -algebras". arXiv: <https://arxiv.org/abs/2106.02620>
- ③ Haslehurst, M.J. "Some examples of factor groupoids". (in preparation)
- ④ Karoubi, M. *K-Theory: An Introduction*. Springer-Verlag, 1978. ISBN: 0-387-08090-2.
- ⑤ Li, X. "Every classifiable simple C^* -algebra has a Cartan subalgebra". Invent. math. 219, 653–699 (2020).

- ⑥ Rørdam, M.; Larsen, F.; Lausten, N.J. *An Introduction to K-theory for C^* -Algebras*. London Mathematical Society Student Texts. Cambridge University Press, 2000. ISBN: 0521 78944 3.
- ⑥ Putnam, I.F. "An excision theorem for the K -theory of C^* -algebras, with applications to groupoid C^* -algebras". Munster Mathematics Journal (to appear).