

# $C^*$ -algebras, groupoids, and symmetry

Tutte Institute for Mathematics and Computing  
Communications Security Establishment

Mitch Haslehurst

Department of Mathematics & Statistics  
University of Victoria

March 1, 2022

# Outline

## 1 C\*-algebras

- Definition(s) and examples
- Main theorems and structure
- Motivation for study

## 2 Groupoids

- Definition and examples
- Interlude:  $K$ -theory
- Recent results in the Elliott classification program

## 3 Applications: dynamics and fractals

- Iterated function systems
- Data interpolation with fractals
- Closing remarks

# C\*-algebras

What is a C\*-algebra?

What is a C\*-algebra?

**Abstract definition:** A complete normed complex algebra with a conjugate linear involution  $*$  such that, if  $a$  and  $b$  are in  $A$ ,

$$(ab)^* = b^* a^* \quad \|ab\| \leq \|a\| \|b\| \quad \|a^* a\| = \|a\|^2.$$

What is a C\*-algebra?

**Abstract definition:** A complete normed complex algebra with a conjugate linear involution  $*$  such that, if  $a$  and  $b$  are in  $A$ ,

$$(ab)^* = b^* a^* \quad \|ab\| \leq \|a\| \|b\| \quad \|a^* a\| = \|a\|^2.$$

**Concrete definition:** Any subalgebra of  $\mathcal{B}(\mathcal{H})$ , the bounded linear operators on a Hilbert space  $\mathcal{H}$ , that is closed in the uniform norm, and closed under the Hilbert space adjoint operation.

**Example.** The complex numbers  $\mathbb{C}$ , with its usual addition, multiplication,  $z^* = \bar{z}$ , and norm  $|z|$ .

**Example.** The complex numbers  $\mathbb{C}$ , with its usual addition, multiplication,  $z^* = \bar{z}$ , and norm  $|z|$ .

**Example.** If  $X$  is a locally compact Hausdorff space, then

$$C_0(X) = \{f : X \rightarrow \mathbb{C} \mid f \text{ is continuous and vanishes at infinity}\}$$

with  $f^*(x) = \overline{f(x)}$  and norm  $\|f\| = \sup_{x \in X} |f(x)|$  is a C\*-algebra.



**Example.** The complex numbers  $\mathbb{C}$ , with its usual addition, multiplication,  $z^* = \bar{z}$ , and norm  $|z|$ .

**Example.** If  $X$  is a locally compact Hausdorff space, then

$$C_0(X) = \{f : X \rightarrow \mathbb{C} \mid f \text{ is continuous and vanishes at infinity}\}$$

with  $f^*(x) = \overline{f(x)}$  and norm  $\|f\| = \sup_{x \in X} |f(x)|$  is a C\*-algebra.

**Example.**  $\mathcal{B}(\mathcal{H})$  with the Hilbert space operator adjoint and uniform norm. When  $\mathcal{H}$  is finite-dimensional, we may identify  $\mathcal{B}(\mathcal{H})$  with  $M_n(\mathbb{C})$  for some  $n$ .

## Theorem (Gelfand-Naimark)

If  $A$  is a commutative  $C^*$ -algebra, then there is a locally compact Hausdorff space  $X$  such that  $A \cong C_0(X)$ .

### Theorem (Gelfand-Naimark)

If  $A$  is a commutative  $C^*$ -algebra, then there is a locally compact Hausdorff space  $X$  such that  $A \cong C_0(X)$ .

### Theorem (Gelfand-Naimark)

If  $A$  is a  $C^*$ -algebra, then there exists a Hilbert space  $\mathcal{H}$  and an injective  $*$ -homomorphism  $\varphi : A \rightarrow \mathcal{B}(\mathcal{H})$ .

### Theorem (Gelfand-Naimark)

If  $A$  is a commutative  $C^*$ -algebra, then there is a locally compact Hausdorff space  $X$  such that  $A \cong C_0(X)$ .

### Theorem (Gelfand-Naimark)

If  $A$  is a  $C^*$ -algebra, then there exists a Hilbert space  $\mathcal{H}$  and an injective  $*$ -homomorphism  $\varphi : A \rightarrow \mathcal{B}(\mathcal{H})$ .

All commutative  $C^*$ -algebras look like continuous functions.

### Theorem (Gelfand-Naimark)

If  $A$  is a commutative C\*-algebra, then there is a locally compact Hausdorff space  $X$  such that  $A \cong C_0(X)$ .

### Theorem (Gelfand-Naimark)

If  $A$  is a C\*-algebra, then there exists a Hilbert space  $\mathcal{H}$  and an injective \*-homomorphism  $\varphi : A \rightarrow \mathcal{B}(\mathcal{H})$ .

All commutative C\*-algebras look like continuous functions.

All C\*-algebras (commutative or not) look like a subalgebra of bounded linear operators.

The pairing  $X \leftrightarrow C_0(X)$  is a complete categorical equivalence between locally compact Hausdorff spaces and commutative C\*-algebras.

The pairing  $X \leftrightarrow C_0(X)$  is a complete categorical equivalence between locally compact Hausdorff spaces and commutative C\*-algebras.

C\*-algebras are often referred to as "noncommutative spaces"

The pairing  $X \leftrightarrow C_0(X)$  is a complete categorical equivalence between locally compact Hausdorff spaces and commutative C\*-algebras.

C\*-algebras are often referred to as "noncommutative spaces"

Classical mechanics  $\longleftrightarrow$  Calculus and topology



The pairing  $X \leftrightarrow C_0(X)$  is a complete categorical equivalence between locally compact Hausdorff spaces and commutative C\*-algebras.


C\*-algebras are often referred to as "noncommutative spaces"


Classical mechanics  $\longleftrightarrow$  Calculus and topology


Quantum mechanics  $\longleftrightarrow$  Operator algebras

"...Heisenberg postulated that the mathematics describing quantum physics should be the mathematics, **not of functions on a space**, but of **linear operators on a Hilbert space**, which, taken as an algebra, behaves, algebraically, much like the algebra of continuous functions on a space, but is not commutative..."

-Heath Emerson, *An introduction to C\*-algebras and Noncommutative Geometry*.







C\*-algebras have found interactions with:

- 1 Group theory
- 2 Harmonic analysis
- 3 Dynamical systems
- 4 Probability
- 5 Logic
- 6 Number theory
- 7 Graph theory
- 8 Geometry
- 9 Knot theory
- 10 Quantum information theory

# Groupoids

Groups encode "global symmetry".



Groups encode "global symmetry".

### Theorem

Every group  $\Gamma$  is a subgroup of a permutation group.

Groups encode "global symmetry".

### Theorem

Every group  $\Gamma$  is a subgroup of a permutation group.

Groupoids, on the other hand, encode "local symmetry".

Groups encode "global symmetry".


### Theorem

Every group  $\Gamma$  is a subgroup of a permutation group.

Groupoids, on the other hand, encode "local symmetry".

They are useful for dynamics, fractal geometry, quasicrystals, tilings.

Groupoids are collections of "partial symmetries".



A groupoid  $G$  is like a group, but not every pair of elements can be multiplied.

A groupoid  $G$  is like a group, but not every pair of elements can be multiplied.

**Example.** Let  $X$  be a nonempty set and  $R \subseteq X \times X$  an equivalence relation. Then  $R$  is a groupoid:

$$(x, y)(y', z) = (x, z) \quad (x, y)^{-1} = (y, x)$$

the product being defined only when  $y = y'$ .

Given a groupoid with some nice topological properties, we can make a  $C^*$ -algebra out of it.

Given a groupoid with some nice topological properties, we can make a C\*-algebra out of it.

$C_c(G)$  = all continuous, compactly supported functions  $f : G \rightarrow \mathbb{C}$ .

$$(f \star g)(x) = \sum_{yy^{-1}=xx^{-1}} f(y)g(y^{-1}x) \quad f^*(x) = \overline{f(x^{-1})}$$



Given a groupoid with some nice topological properties, we can make a C\*-algebra out of it.

$C_c(G)$  = all continuous, compactly supported functions  $f : G \rightarrow \mathbb{C}$ .

$$(f \star g)(x) = \sum_{yy^{-1}=xx^{-1}} f(y)g(y^{-1}x) \quad f^*(x) = \overline{f(x^{-1})}$$

To get a complete norm, represent  $C_c(G)$  on a Hilbert space and take the closure to get the *reduced* C\*-algebra of  $G$ , called  $C_r^*(G)$ .

**Example.** Let  $X = \{1, 2, \dots, n\}$  and  $R = X \times X$ .

**Example.** Let  $X = \{1, 2, \dots, n\}$  and  $R = X \times X$ . Then if  $f$  and  $g$  are in  $C_c(R)$ , we have

$$(f \star g)(i, k) = \sum_{j=1}^n f(i, j)g(j, k) \quad f^*(i, k) = \overline{f(k, i)}$$

**Example.** Let  $X = \{1, 2, \dots, n\}$  and  $R = X \times X$ . Then if  $f$  and  $g$  are in  $C_c(R)$ , we have


$$(f \star g)(i, k) = \sum_{j=1}^n f(i, j)g(j, k) \quad f^*(i, k) = \overline{f(k, i)}$$

and  $C_c(R) \cong M_n(\mathbb{C})$ .

**Example.** If  $X$  is a locally compact Hausdorff space and  $R = \{(x, x) \mid x \in X\}$ , then  $C_r^*(R) \cong C_0(X)$ .

**Example.** If  $X$  is a locally compact Hausdorff space and  $R = \{(x, x) \mid x \in X\}$ , then  $C_r^*(R) \cong C_0(X)$ .

**Example.** If  $\Gamma$  is a group acting on a space  $X$  (a dynamical system), then  $G = X \rtimes \Gamma$  is a groupoid and  $C_r^*(G)$  is the *crossed product*  $C_0(X) \rtimes \Gamma$ .



**Question.** If  $A$  is a  $C^*$ -algebra, is there a groupoid  $G$  such that  $A \cong C_r^*(G)$ ?



**Question.** If  $A$  is a  $C^*$ -algebra, is there a groupoid  $G$  such that  $A \cong C_r^*(G)$ ?

To answer this question, we need some homological tools.

**Question.** If  $A$  is a  $C^*$ -algebra, is there a groupoid  $G$  such that  $A \cong C_r^*(G)$ ?

To answer this question, we need some homological tools.

$C^*$ -algebras have a useful homology theory: to each  $C^*$ -algebra  $A$  there are two abelian groups called  $K_0(A)$  and  $K_1(A)$ .

**Question.** If  $A$  is a  $C^*$ -algebra, is there a groupoid  $G$  such that  $A \cong C_r^*(G)$ ?

To answer this question, we need some homological tools.

$C^*$ -algebras have a useful homology theory: to each  $C^*$ -algebra  $A$  there are two abelian groups called  $K_0(A)$  and  $K_1(A)$ .

$K_0(A)$  consists of equivalence classes of projections  $p$  in  $\bigcup_n M_n(A)$  that "have the same rank".  $K_1(A)$  consists of equivalence classes of unitaries  $u$  in  $\bigcup_n M_n(A)$  that are "stably homotopic".

If the sequence

$$0 \longrightarrow I \xrightarrow{\iota} A \xrightarrow{\pi} A/I \longrightarrow 0$$

is exact,

If the sequence

$$0 \longrightarrow I \xrightarrow{\iota} A \xrightarrow{\pi} A/I \longrightarrow 0$$

is exact, there are group homomorphisms  $\delta_0$  and  $\delta_1$  such that the sequence

$$\begin{array}{ccccc} K_0(I) & \xrightarrow{\iota_*} & K_0(A) & \xrightarrow{\pi_*} & K_0(A/I) \\ \delta_1 \uparrow & & & & \downarrow \delta_0 \\ K_1(A/I) & \xleftarrow{\pi_*} & K_1(A) & \xleftarrow{\iota_*} & K_1(I) \end{array}$$

is exact.

**Examples.**  $K_0(\mathbb{C}) \cong \mathbb{Z}$  (associate a projection in  $\bigcup_n M_n(\mathbb{C})$  to its rank).  $K_0(M_n(\mathbb{C})) \cong \mathbb{Z}$  for the same reason.

**Examples.**  $K_0(\mathbb{C}) \cong \mathbb{Z}$  (associate a projection in  $\bigcup_n M_n(\mathbb{C})$  to its rank).  $K_0(M_n(\mathbb{C})) \cong \mathbb{Z}$  for the same reason.

$K_1(\mathbb{C}) = 0$  because the unitary group of  $M_n(\mathbb{C})$  is connected.

**Examples.**  $K_0(\mathbb{C}) \cong \mathbb{Z}$  (associate a projection in  $\bigcup_n M_n(\mathbb{C})$  to its rank).  $K_0(M_n(\mathbb{C})) \cong \mathbb{Z}$  for the same reason.

$K_1(\mathbb{C}) = 0$  because the unitary group of  $M_n(\mathbb{C})$  is connected.

$K_0(C(X)) \cong \mathbb{Z}$  and  $K_1(C(X)) = 0$  if  $X$  is contractible ( $K$ -theory is homotopy invariant).



**Examples.**  $K_0(\mathbb{C}) \cong \mathbb{Z}$  (associate a projection in  $\bigcup_n M_n(\mathbb{C})$  to its rank).  $K_0(M_n(\mathbb{C})) \cong \mathbb{Z}$  for the same reason.

$K_1(\mathbb{C}) = 0$  because the unitary group of  $M_n(\mathbb{C})$  is connected.

$K_0(C(X)) \cong \mathbb{Z}$  and  $K_1(C(X)) = 0$  if  $X$  is contractible ( $K$ -theory is homotopy invariant).

$K_0(\mathcal{B}(\mathcal{H})) = K_1(\mathcal{B}(\mathcal{H})) = 0$  if  $\mathcal{H}$  is infinite dimensional.

**Examples.**  $K_0(\mathbb{C}) \cong \mathbb{Z}$  (associate a projection in  $\bigcup_n M_n(\mathbb{C})$  to its rank).  $K_0(M_n(\mathbb{C})) \cong \mathbb{Z}$  for the same reason.

$K_1(\mathbb{C}) = 0$  because the unitary group of  $M_n(\mathbb{C})$  is connected.

$K_0(C(X)) \cong \mathbb{Z}$  and  $K_1(C(X)) = 0$  if  $X$  is contractible ( $K$ -theory is homotopy invariant).

$K_0(\mathcal{B}(\mathcal{H})) = K_1(\mathcal{B}(\mathcal{H})) = 0$  if  $\mathcal{H}$  is infinite dimensional.

$K_1(C(S^1)) \cong \mathbb{Z}$  where  $S^1 = \{z \in \mathbb{C} \mid |z| = 1\}$  (associate a unitary to its winding number).

There is an immense amount of research being done into what  $K$ -theory can tell us about C\*-algebras.

There is an immense amount of research being done into what  $K$ -theory can tell us about C\*-algebras.

**The Elliott classification program.** Suppose  $A$  and  $B$  are two unital, simple, separable, nuclear,  $\mathcal{Z}$ -stable C\*-algebras that satisfy the Universal Coefficient Theorem. If  $A$  and  $B$  have the same  $K$ -theory, then they are isomorphic.

**Question.** If  $A$  is a  $C^*$ -algebra, is there a groupoid  $G$  such that  $A \cong C_r^*(G)$ ?

**Question.** If  $A$  is a  $C^*$ -algebra, is there a groupoid  $G$  such that  $A \cong C_r^*(G)$ ?

**Answer.** Yes, if  $A$  is classifiable in the Elliott program.

**Question.** If  $A$  is a  $C^*$ -algebra, is there a groupoid  $G$  such that  $A \cong C_r^*(G)$ ?

**Answer.** Yes, if  $A$  is classifiable in the Elliott program.

**Theorem (Li, 2019)**

Given some  $K$ -theory data  $D$ , there is a groupoid  $G$  such that  $K_*(C_r^*(G)) \cong D$ .

**Question.** If  $A$  is a  $C^*$ -algebra, is there a groupoid  $G$  such that  $A \cong C_r^*(G)$ ?

**Answer.** Yes, if  $A$  is classifiable in the Elliott program.

**Theorem (Li, 2019)**

Given some  $K$ -theory data  $D$ , there is a groupoid  $G$  such that  $K_*(C_r^*(G)) \cong D$ .

**Theorem (Putnam, 2017)**

Given some torsion-free  $K$ -theory data  $D$ , there is an groupoid  $G$  on a Cantor set such that  $K_*(C_r^*(G)) \cong D$ .



## Theorem (H., 2021)

Given some torsion-free  $K$ -theory data  $D$ , there is a quotient space  $X$  of a Cantor set and a factor groupoid  $G$  on  $X$  such that  $K_*(C_r^*(G)) \cong D$ .

## Theorem (H., 2021)

Given some torsion-free  $K$ -theory data  $D$ , there is a quotient space  $X$  of a Cantor set and a factor groupoid  $G$  on  $X$  such that  $K_*(C_r^*(G)) \cong D$ .

The spaces  $X$  are constructed using a "generalized" Cantor ternary function.

## Theorem (H., 2021)

Given some torsion-free  $K$ -theory data  $D$ , there is a quotient space  $X$  of a Cantor set and a factor groupoid  $G$  on  $X$  such that  $K_*(C_r^*(G)) \cong D$ .

The spaces  $X$  are constructed using a "generalized" Cantor ternary function.

$$\{0, 1\}^{\mathbb{N}} \rightarrow S^1 \quad (x_n) \mapsto \exp \left( 2\pi i \sum_{n=1}^{\infty} x_n 2^{-n} \right)$$

## Theorem (H., 2021)

Given some torsion-free  $K$ -theory data  $D$ , there is a quotient space  $X$  of a Cantor set and a factor groupoid  $G$  on  $X$  such that  $K_*(C_r^*(G)) \cong D$ .

The spaces  $X$  are constructed using a "generalized" Cantor ternary function.

$$\{0, 1\}^{\mathbb{N}} \rightarrow S^1 \quad (x_n) \mapsto \exp \left( 2\pi i \sum_{n=1}^{\infty} x_n 2^{-n} \right)$$

In general, the spaces have connected components either a single point or homeomorphic to  $S^1 = \{z \in \mathbb{C} \mid |z| = 1\}$ .

## Theorem (H., 2021)


Given some torsion-free  $K$ -theory data  $D$ , there is a quotient space  $X$  of a Cantor set and a factor groupoid  $G$  on  $X$  such that  $K_*(C_r^*(G)) \cong D$ .

The spaces  $X$  are constructed using a "generalized" Cantor ternary function.

$$\{0, 1\}^{\mathbb{N}} \rightarrow S^1 \quad (x_n) \mapsto \exp \left( 2\pi i \sum_{n=1}^{\infty} x_n 2^{-n} \right)$$

In general, the spaces have connected components either a single point or homeomorphic to  $S^1 = \{z \in \mathbb{C} \mid |z| = 1\}$ .

They also often have some self-similar fractal structure.



# Applications: dynamics and fractals

An *iterated function system*  $(X, \{f_j\}_{j=1}^n)$  (abbreviated IFS) is a complete metric space  $X$  with a finite set of functions  $f_j : X \rightarrow X$  for  $j = 1, 2, \dots, n$ .



An *iterated function system*  $(X, \{f_j\}_{j=1}^n)$  (abbreviated IFS) is a complete metric space  $X$  with a finite set of functions  $f_j : X \rightarrow X$  for  $j = 1, 2, \dots, n$ .

If the IFS is hyperbolic (every  $f_j$  is a contraction) then there is a unique compact subset  $K \subseteq X$  such that

$$K = \bigcup_{j=1}^n f_j(K)$$

$K$  is called the *attractor* of the IFS.


**Example.** Take  $X = \mathbb{R}$  and  $f_1(x) = \frac{1}{2}x$  and  $f_2(x) = \frac{1}{2}x + \frac{1}{2}$ . The attractor is  $K = [0, 1]$ .

**Example.** Take  $X = \mathbb{R}$  and  $f_1(x) = \frac{1}{2}x$  and  $f_2(x) = \frac{1}{2}x + \frac{1}{2}$ . The attractor is  $K = [0, 1]$ .

**Example.** Take  $X = \mathbb{R}$  and  $f_1(x) = \frac{1}{3}x$  and  $f_2(x) = \frac{1}{3}x + \frac{2}{3}$ . The attractor  $K$  is the Cantor set.

**Example.** Take  $X = \mathbb{R}$  and  $f_1(x) = \frac{1}{2}x$  and  $f_2(x) = \frac{1}{2}x + \frac{1}{2}$ . The attractor is  $K = [0, 1]$ .

**Example.** Take  $X = \mathbb{R}$  and  $f_1(x) = \frac{1}{3}x$  and  $f_2(x) = \frac{1}{3}x + \frac{2}{3}$ . The attractor  $K$  is the Cantor set.



**Example.** Take  $X = \mathbb{R}^2$  and

$$f_1(x) = \frac{1}{2}x \quad f_2(x) = \frac{1}{2}x + \begin{bmatrix} 1/2 \\ 0 \end{bmatrix} \quad f_3(x) = \frac{1}{2}x + \begin{bmatrix} 1/4 \\ \sqrt{3}/4 \end{bmatrix}$$

**Example.** Take  $X = \mathbb{R}^2$  and

$$f_1(x) = \frac{1}{2}x \quad f_2(x) = \frac{1}{2}x + \begin{bmatrix} 1/2 \\ 0 \end{bmatrix} \quad f_3(x) = \frac{1}{2}x + \begin{bmatrix} 1/4 \\ \sqrt{3}/4 \end{bmatrix}$$

The attractor  $K$  is the Sierpiński triangle.



In 2020, E.R. Korfanty constructed a groupoid on the attractor  $K$  of an arbitrary IFS that captures the self-similarity of  $K$  through partial symmetry.

In 2020, E.R. Korfanty constructed a groupoid on the attractor  $K$  of an arbitrary IFS that captures the self-similarity of  $K$  through partial symmetry.

### Theorem (Korfanty, 2020)

Suppose  $(K, \{f_j\}_{j=1}^n)$  and  $(K', \{f'_j\}_{j=1}^n)$  are two compact hyperbolic IFS's with respective groupoids  $G$  and  $G'$ . If  $(K, \{f_j\}_{j=1}^n)$  and  $(K', \{f'_j\}_{j=1}^n)$  are topologically conjugate, then  $C_r^*(G)$  and  $C_r^*(G')$  are isomorphic.



In 2020, E.R. Korfanty constructed a groupoid on the attractor  $K$  of an arbitrary IFS that captures the self-similarity of  $K$  through partial symmetry.

### Theorem (Korfanty, 2020)

Suppose  $(K, \{f_j\}_{j=1}^n)$  and  $(K', \{f'_j\}_{j=1}^n)$  are two compact hyperbolic IFS's with respective groupoids  $G$  and  $G'$ . If  $(K, \{f_j\}_{j=1}^n)$  and  $(K', \{f'_j\}_{j=1}^n)$  are topologically conjugate, then  $C_r^*(G)$  and  $C_r^*(G')$  are isomorphic.

**The converse problem.** If  $C_r^*(G)$  and  $C_r^*(G')$  are isomorphic, are the two systems topologically conjugate?

In 2020, E.R. Korfanty constructed a groupoid on the attractor  $K$  of an arbitrary IFS that captures the self-similarity of  $K$  through partial symmetry.

### Theorem (Korfanty, 2020)

Suppose  $(K, \{f_j\}_{j=1}^n)$  and  $(K', \{f'_j\}_{j=1}^n)$  are two compact hyperbolic IFS's with respective groupoids  $G$  and  $G'$ . If  $(K, \{f_j\}_{j=1}^n)$  and  $(K', \{f'_j\}_{j=1}^n)$  are topologically conjugate, then  $C_r^*(G)$  and  $C_r^*(G')$  are isomorphic.

**The converse problem.** If  $C_r^*(G)$  and  $C_r^*(G')$  are isomorphic, are the two systems topologically conjugate?

**Conjecture.** Most likely not.

However...

However...

### Theorem (Giordano, Putnam, Skau, 1995)

If  $K$  is a Cantor set and  $\varphi, \psi : K \rightarrow K$  are two minimal homeomorphisms (every orbit is dense) and  $C(K) \rtimes_{\varphi} \mathbb{Z}$  and  $C(K) \rtimes_{\psi} \mathbb{Z}$  have the same  $K$ -theory, then  $(K, \varphi)$  and  $(K, \psi)$  are orbit-equivalent as dynamical systems.

However...

### Theorem (Giordano, Putnam, Skau, 1995)

If  $K$  is a Cantor set and  $\varphi, \psi : K \rightarrow K$  are two minimal homeomorphisms (every orbit is dense) and  $C(K) \rtimes_{\varphi} \mathbb{Z}$  and  $C(K) \rtimes_{\psi} \mathbb{Z}$  have the same  $K$ -theory, then  $(K, \varphi)$  and  $(K, \psi)$  are orbit-equivalent as dynamical systems.

If C\*-algebras are to give an invariant for IFS's:

However...

### Theorem (Giordano, Putnam, Skau, 1995)

If  $K$  is a Cantor set and  $\varphi, \psi : K \rightarrow K$  are two minimal homeomorphisms (every orbit is dense) and  $C(K) \rtimes_{\varphi} \mathbb{Z}$  and  $C(K) \rtimes_{\psi} \mathbb{Z}$  have the same  $K$ -theory, then  $(K, \varphi)$  and  $(K, \psi)$  are orbit-equivalent as dynamical systems.

If C\*-algebras are to give an invariant for IFS's:

- 1  $K$ -theory will likely be crucial,

However...

### Theorem (Giordano, Putnam, Skau, 1995)

If  $K$  is a Cantor set and  $\varphi, \psi : K \rightarrow K$  are two minimal homeomorphisms (every orbit is dense) and  $C(K) \rtimes_{\varphi} \mathbb{Z}$  and  $C(K) \rtimes_{\psi} \mathbb{Z}$  have the same  $K$ -theory, then  $(K, \varphi)$  and  $(K, \psi)$  are orbit-equivalent as dynamical systems.

If C\*-algebras are to give an invariant for IFS's:

- 1  $K$ -theory will likely be crucial,
- 2 "topological conjugacy" may need to be weakened.

## Theorem (Barnsley, 1986)

Let  $x_1 < x_2 < \cdots < x_n$  be real numbers and  $\{(x_j, y_j) \mid j = 1, 2, \dots, n\} \subseteq \mathbb{R}^2$  be a data set. Then there is an IFS  $(\mathbb{R}^2, \{f_j\}_{j=1}^n)$  such that the attractor  $K$  is the graph of a continuous function  $F : [x_1, x_n] \rightarrow \mathbb{R}$  with  $F(x_j) = y_j$  for all  $j = 1, 2, \dots, n$ .



## Theorem (Barnsley, 1986)


Let  $x_1 < x_2 < \cdots < x_n$  be real numbers and  $\{(x_j, y_j) \mid j = 1, 2, \dots, n\} \subseteq \mathbb{R}^2$  be a data set. Then there is an IFS  $(\mathbb{R}^2, \{f_j\}_{j=1}^n)$  such that the attractor  $K$  is the graph of a continuous function  $F : [x_1, x_n] \rightarrow \mathbb{R}$  with  $F(x_j) = y_j$  for all  $j = 1, 2, \dots, n$ .

Data interpolation using fractal interpolation functions:

- 1 Like smooth functions such as polynomials and trigonometric functions, can be approximated recurrently using formulae.
- 2 The fractal structure of the curve captures "irregularities" well as opposed to smooth functions.

## Closing remarks

- 1 Groupoids provide the algebraic groundwork for studying local symmetry, while C\*-algebras provide an immense amount of structure and powerful tools.
- 2 Interplay between the dynamics of iterated function systems and C\*-algebras: only scratched the surface.



Thank you!