

Examples of factor groupoids: Cantor functions and iterated function systems

Glasgow Analysis Seminar

Mitch Haslehurst

Department of Mathematics & Statistics
University of Victoria

April 7, 2022

Groupoids

A groupoid G is like a group, but not every pair of elements can be multiplied.

Groupoids

A groupoid G is like a group, but not every pair of elements can be multiplied.

Every element x of G has an inverse x^{-1} , but there are many "units". The set of units is denoted $G^{(0)}$ and they are of the form $x^{-1}x$. Define the maps $r(x) = xx^{-1}$ and $s(x) = x^{-1}x$.

Groupoids

A groupoid G is like a group, but not every pair of elements can be multiplied.

Every element x of G has an inverse x^{-1} , but there are many "units". The set of units is denoted $G^{(0)}$ and they are of the form $x^{-1}x$. Define the maps $r(x) = xx^{-1}$ and $s(x) = x^{-1}x$.

Example. Let X be a nonempty set and $R \subseteq X \times X$ an equivalence relation. Then R is a groupoid:

$$(x, y)(y', z) = (x, z) \quad (x, y)^{-1} = (y, x)$$

only when $y = y'$.

Given a groupoid with some nice topological properties, we can make a C^* -algebra out of it. Assume G is locally compact Hausdorff and étale (the map r is a local homeomorphism).

Given a groupoid with some nice topological properties, we can make a C^* -algebra out of it. Assume G is locally compact Hausdorff and étale (the map r is a local homeomorphism).

$C_c(G)$ = all continuous, compactly supported functions $f : G \rightarrow \mathbb{C}$.

$$(f \star g)(x) = \sum_{r(y)=r(x)} f(y)g(y^{-1}x) \quad f^*(x) = \overline{f(x^{-1})}$$

Given a groupoid with some nice topological properties, we can make a C^* -algebra out of it. Assume G is locally compact Hausdorff and étale (the map r is a local homeomorphism).

$C_c(G)$ = all continuous, compactly supported functions $f : G \rightarrow \mathbb{C}$.

$$(f \star g)(x) = \sum_{r(y)=r(x)} f(y)g(y^{-1}x) \quad f^*(x) = \overline{f(x^{-1})}$$

To get a complete norm, represent $C_c(G)$ on a Hilbert space and take the closure to get the *reduced* C^* -algebra of G , called $C_r^*(G)$.

Example. Let $X = \{1, 2, \dots, n\}$ and $R = X \times X$.

Example. Let $X = \{1, 2, \dots, n\}$ and $R = X \times X$. Then if f and g are in $C_c(R)$, we have

$$(f \star g)(i, k) = \sum_{j=1}^n f(i, j)g(j, k) \quad f^*(i, k) = \overline{f(k, i)}$$

Example. Let $X = \{1, 2, \dots, n\}$ and $R = X \times X$. Then if f and g are in $C_c(R)$, we have

$$(f \star g)(i, k) = \sum_{j=1}^n f(i, j)g(j, k) \quad f^*(i, k) = \overline{f(k, i)}$$

Hopefully these formulae look familiar. It's matrix multiplication and the conjugate transpose, so $C_c(R) \cong M_n(\mathbb{C})$.

Example. Let $X = \{1, 2, \dots, n\}$ and $R = X \times X$. Then if f and g are in $C_c(R)$, we have

$$(f \star g)(i, k) = \sum_{j=1}^n f(i, j)g(j, k) \quad f^*(i, k) = \overline{f(k, i)}$$

Hopefully these formulae look familiar. It's matrix multiplication and the conjugate transpose, so $C_c(R) \cong M_n(\mathbb{C})$.

If X is a locally compact Hausdorff space and $R = \{(x, x) \mid x \in X\}$, then $C_r^*(R) \cong C_0(X)$.

Let X be a compact Hausdorff space and Γ a discrete group acting on X by homeomorphisms. Then $X \times \Gamma$ is a locally compact Hausdorff étale groupoid:

$$(x, g)(y, h) = (x, gh) \quad (x, g)^{-1} = (g \cdot x, g^{-1})$$

only when $g \cdot x = y$. Moreover, $C_r^*(X \times \Gamma)$ is isomorphic to the (reduced) crossed product $C(X) \rtimes \Gamma$.

Problem. Given a C^* -algebra A , find a groupoid G such that $A \cong C_r^*(G)$.

Problem. Given a C^* -algebra A , find a groupoid G such that $A \cong C_r^*(G)$.

If A is classifiable, then this becomes

Problem. Given a C^* -algebra A , find a groupoid G such that $A \cong C_r^*(G)$.

If A is classifiable, then this becomes

Problem. Given some K -theory data, find a groupoid G such that the data is $K_*(C_r^*(G))$.

Problem. Given a C^* -algebra A , find a groupoid G such that $A \cong C_r^*(G)$.

If A is classifiable, then this becomes

Problem. Given some K -theory data, find a groupoid G such that the data is $K_*(C_r^*(G))$.

Notable references:

Li, X. "Every classifiable simple C^* -algebra has a Cartan subalgebra". *Invent. math.* 219, 653–699 (2020).

Putnam, I.F. "Some classifiable groupoid C^* -algebras with prescribed K -theory". *Math. Ann.* 370, 1361–1387 (2018).

Factor groupoids

G' is a *factor groupoid* of G if $\pi : G \rightarrow G'$ is a surjective groupoid homomorphism.

Factor groupoids

G' is a *factor groupoid* of G if $\pi : G \rightarrow G'$ is a surjective groupoid homomorphism.

Assume:

- 1 G and G' are locally compact Hausdorff and étale,
- 2 π is continuous and proper,
- 3 $\pi|_{G^u} : G^u \rightarrow (G')^{\pi(u)}$ is bijective for all u in $G^{(0)}$.

Factor groupoids

G' is a *factor groupoid* of G if $\pi : G \rightarrow G'$ is a surjective groupoid homomorphism.

Assume:

- ① G and G' are locally compact Hausdorff and étale,
- ② π is continuous and proper,
- ③ $\pi|_{G^u} : G^u \rightarrow (G')^{\pi(u)}$ is bijective for all u in $G^{(0)}$.

Obtain an inclusion $C_r^*(G') \subseteq C_r^*(G)$ via $b \mapsto b \circ \pi$ (b in $C_c(G')$)

The Cantor function

Define the map $\varphi : \{0, 1\}^{\mathbb{N}} \rightarrow S^1$ by

$$\varphi(\{x_n\}) = \exp \left(2\pi i \sum_{n=1}^{\infty} x_n 2^{-n} \right)$$

The Cantor function

Define the map $\varphi : \{0, 1\}^{\mathbb{N}} \rightarrow S^1$ by

$$\varphi(\{x_n\}) = \exp \left(2\pi i \sum_{n=1}^{\infty} x_n 2^{-n} \right)$$

Let S be *tail-equivalence* on $\{0, 1\}^{\mathbb{N}}$, that is, $(\{x_n\}, \{y_n\}) \in S$ if and only if $x_n = y_n$ for sufficiently large n .

The Cantor function

Define the map $\varphi : \{0, 1\}^{\mathbb{N}} \rightarrow S^1$ by

$$\varphi(\{x_n\}) = \exp \left(2\pi i \sum_{n=1}^{\infty} x_n 2^{-n} \right)$$

Let S be *tail-equivalence* on $\{0, 1\}^{\mathbb{N}}$, that is, $(\{x_n\}, \{y_n\}) \in S$ if and only if $x_n = y_n$ for sufficiently large n .

S has a natural étale topology generated by the basic sets

$$\gamma(p, q) = \left\{ (\{x_n\}, \{y_n\}) \mid \begin{array}{l} (x_1, x_2, \dots, x_n) = (p_1, p_2, \dots, p_n) \\ (y_1, y_2, \dots, y_n) = (q_1, q_2, \dots, q_n) \\ x_k = y_k \text{ for all } k \geq n+1 \end{array} \right\}$$

where $p, q \in \{0, 1\}^n$, and $C_r^*(S) \cong M_{2^\infty}$.

The Cantor function

What happens when we let $T = (\varphi \times \varphi)(S)$ and give T the quotient topology?

The Cantor function

What happens when we let $T = (\varphi \times \varphi)(S)$ and give T the quotient topology? We get

$$T = \{(w, z) \in S^1 \times S^1 \mid w = e^{2\pi i \theta} z \text{ for some } \theta \in \mathbb{Z}[\frac{1}{2}]\}$$

with basic open sets $U_{W, \theta} = \{(z, e^{2\pi i \theta} z) \mid z \in W\}$ where $W \subseteq S^1$ is open and $\theta \in \mathbb{Z}[\frac{1}{2}]$.

The Cantor function

What happens when we let $T = (\varphi \times \varphi)(S)$ and give T the quotient topology? We get

$$T = \{(w, z) \in S^1 \times S^1 \mid w = e^{2\pi i \theta} z \text{ for some } \theta \in \mathbb{Z}[\frac{1}{2}]\}$$

with basic open sets $U_{W, \theta} = \{(z, e^{2\pi i \theta} z) \mid z \in W\}$ where $W \subseteq S^1$ is open and $\theta \in \mathbb{Z}[\frac{1}{2}]$.

T is a second-countable locally compact Hausdorff étale groupoid in the quotient topology, and $C_r^*(T) \cong B$, where B is the Bunce-Deddens algebra of type 2^∞ .

The Cantor function

What happens when we let $T = (\varphi \times \varphi)(S)$ and give T the quotient topology? We get

$$T = \{(w, z) \in S^1 \times S^1 \mid w = e^{2\pi i \theta} z \text{ for some } \theta \in \mathbb{Z}[\frac{1}{2}]\}$$

with basic open sets $U_{W, \theta} = \{(z, e^{2\pi i \theta} z) \mid z \in W\}$ where $W \subseteq S^1$ is open and $\theta \in \mathbb{Z}[\frac{1}{2}]$.

T is a second-countable locally compact Hausdorff étale groupoid in the quotient topology, and $C_r^*(T) \cong B$, where B is the Bunce-Deddens algebra of type 2^∞ .

$$K_0(C_r^*(T)) \cong \mathbb{Z}[\frac{1}{2}] \quad K_1(C_r^*(T)) \cong \mathbb{Z}$$


and $C_r^*(T)$ has a unique tracial state.

Bratteli diagrams

Let (V, E) be a Bratteli diagram.


Bratteli diagrams

Let (V, E) be a Bratteli diagram.



Bratteli diagrams


Let (V, E) be a Bratteli diagram.



The *infinite path space* X_E of (V, E) is a totally disconnected compact metric space.

Bratteli diagrams

Let (V, E) be a Bratteli diagram.



The *infinite path space* X_E of (V, E) is a totally disconnected compact metric space.

Tail-equivalence $R_E \subseteq X_E \times X_E$ has an étale topology in which $C_r^*(R_E)$ is an AF-algebra.

Let (V, E) and (W, F) be two Bratteli diagrams.

The space X_ξ

Let (V, E) and (W, F) be two Bratteli diagrams.

Two *graph embeddings* $\xi^0, \xi^1 : (W, F) \rightarrow (V, E)$ with $\xi^0|_W = \xi^1|_W$ and $\xi^0(F) \cap \xi^1(F) = \emptyset$.

The space X_ξ

Let (V, E) and (W, F) be two Bratteli diagrams.

Two *graph embeddings* $\xi^0, \xi^1 : (W, F) \rightarrow (V, E)$ with $\xi^0|_W = \xi^1|_W$ and $\xi^0(F) \cap \xi^1(F) = \emptyset$.

Equivalence relation \sim_ξ on X_E :

$$(x_1, x_2, \dots, x_{n_0-1}, x_{n_0}, \xi^0(z_{n_0+1}), \xi^0(z_{n_0+2}), \dots) \quad (1)$$

$$\sim_\xi (x_1, x_2, \dots, x_{n_0-1}, x'_{n_0}, \xi^1(z_{n_0+1}), \xi^1(z_{n_0+2}), \dots) \quad (2)$$

where $x_{n_0} = x'_{n_0}$ if $x_{n_0} \notin \xi^0(F) \cup \xi^1(F)$, and $x_{n_0} = \xi^1(f)$ if $x'_{n_0} = \xi^0(f)$.

The space X_ξ

Denote $X_\xi := X_E / \sim_\xi$ and $\rho : X_E \rightarrow X_\xi$ the quotient map.

The space X_ξ

Denote $X_\xi := X_E / \sim_\xi$ and $\rho : X_E \rightarrow X_\xi$ the quotient map.

Facts:

- 1 X_ξ is a second-countable compact metrizable space,
- 2 the covering dimension of X_ξ is 1,
- 3 each connected component is either a single point or homeomorphic to S^1 .

Proposition. If $E = \xi^0(F) \cup \xi^1(F)$, then X_ξ is homeomorphic to $X_F \times S^1$.

Examples

Proposition. If $E = \xi^0(F) \cup \xi^1(F)$, then X_ξ is homeomorphic to $X_F \times S^1$.

Example. Let (V, E) have one vertex and three edges at each level. Identify X_E with $\{0, 1, 2\}^{\mathbb{N}}$.

Let (W, F) consist of a single path, and for f in F , $\xi^0(f) = 0$ and $\xi^1(f) = 1$.

Examples

Proposition. If $E = \xi^0(F) \cup \xi^1(F)$, then X_ξ is homeomorphic to $X_F \times S^1$.


Example. Let (V, E) have one vertex and three edges at each level. Identify X_E with $\{0, 1, 2\}^\mathbb{N}$.

Let (W, F) consist of a single path, and for f in F , $\xi^0(f) = 0$ and $\xi^1(f) = 1$.

There is a nested sequence $X_1 \subseteq X_2 \subseteq X_3 \subseteq \cdots \subseteq X_E$ such that


$$X_E = \overline{\bigcup_{n=1}^{\infty} X_n}$$

and each $\rho(X_n)$ is a disjoint union of finitely many circles.




$$\rho(X_1)$$

Examples




$$\rho(X_2)$$

Examples




$$\rho(X_3)$$

Examples




$$\rho(X_4)$$

Examples



$$\rho(X_5)$$

Examples



$$\rho(X_6)$$

The groupoid R_ξ

Let $R_\xi = \rho \times \rho(R_E)$.

With the quotient topology, R_ξ is a second-countable locally compact Hausdorff étale groupoid, and a factor of R_E via $\rho \times \rho : R_E \rightarrow R_\xi$.

The groupoid R_ξ

Let $R_\xi = \rho \times \rho(R_E)$.

With the quotient topology, R_ξ is a second-countable locally compact Hausdorff étale groupoid, and a factor of R_E via $\rho \times \rho : R_E \rightarrow R_\xi$.

We want to analyze the K -theory of $C_r^*(R_\xi) \subseteq C_r^*(R_E)$.

If $A' \subseteq A$, there is a six-term exact sequence

If $A' \subseteq A$, there is a six-term exact sequence

$$\begin{array}{ccccc}
 K_1(A) & \longrightarrow & K_0(A'; A) & \longrightarrow & K_0(A') \\
 \uparrow \scriptstyle \iota_* & & & & \downarrow \scriptstyle \iota_* \\
 K_1(A') & \longleftarrow & K_1(A'; A) & \longleftarrow & K_0(A)
 \end{array}$$

If $A' \subseteq A$, there is a six-term exact sequence

$$\begin{array}{ccccc} K_1(A) & \longrightarrow & K_0(A'; A) & \longrightarrow & K_0(A') \\ \uparrow \scriptstyle \iota_* & & & & \downarrow \scriptstyle \iota_* \\ K_1(A') & \longleftarrow & K_1(A'; A) & \longleftarrow & K_0(A) \end{array}$$

$K_*(A'; A)$ is the *relative* K-theory of the inclusion $A' \subseteq A$.

If $A' \subseteq A$, there is a six-term exact sequence

$$\begin{array}{ccccc}
 K_1(A) & \longrightarrow & K_0(A'; A) & \longrightarrow & K_0(A') \\
 \uparrow \scriptstyle \iota_* & & & & \downarrow \scriptstyle \iota_* \\
 K_1(A') & \longleftarrow & K_1(A'; A) & \longleftarrow & K_0(A)
 \end{array}$$

$K_*(A'; A)$ is the *relative* K-theory of the inclusion $A' \subseteq A$.

Excision: (Putnam, 2020)

$$K_*(C_r^*(G'); C_r^*(G)) \cong K_*(C_r^*(H'); C_r^*(H))$$

where $H \subseteq G$ and $H' \subseteq G'$ are where π is not one-to-one.

Theorem (H.)

We have $K_0(C_r^(R_\xi)) \cong K_0(C_r^*(R_E))$ and $K_1(C_r^*(R_\xi)) \cong K_0(C_r^*(R_F))$. If R_E is minimal, then $C_r^*(R_\xi)$ is classifiable.*

Theorem (H.)

We have $K_0(C_r^(R_\xi)) \cong K_0(C_r^*(R_E))$ and $K_1(C_r^*(R_\xi)) \cong K_0(C_r^*(R_F))$. If R_E is minimal, then $C_r^*(R_\xi)$ is classifiable.*

Through the set-up $\xi^0, \xi^1 : (W, F) \rightarrow (V, E)$, we can prescribe $K_*(C_r^*(R_\xi))$.

Corollary

If G_0 is a simple acyclic dimension group and G_1 is a countable torsion free abelian group, we can find R_ξ such that $K_0(C_r^(R_\xi)) \cong G_0$ (with order) and $K_1(C_r^*(R_\xi)) \cong G_1$.*

Iterated function systems

An *iterated function system* (X, \mathcal{F}) (abbreviated IFS) is a complete metric space X with a finite set \mathcal{F} of functions $f_j : X \rightarrow X$ for $j = 1, 2, \dots, n$.

Iterated function systems

An *iterated function system* (X, \mathcal{F}) (abbreviated IFS) is a complete metric space X with a finite set \mathcal{F} of functions $f_j : X \rightarrow X$ for $j = 1, 2, \dots, n$.

If the IFS is hyperbolic (every f_j is a contraction) then there is a unique compact subset $C \subseteq X$ such that

$$C = \bigcup_{j=1}^n f_j(C)$$

C is called the *attractor* of the IFS.

Iterated function systems

An *iterated function system* (X, \mathcal{F}) (abbreviated IFS) is a complete metric space X with a finite set \mathcal{F} of functions $f_j : X \rightarrow X$ for $j = 1, 2, \dots, n$.

If the IFS is hyperbolic (every f_j is a contraction) then there is a unique compact subset $C \subseteq X$ such that

$$C = \bigcup_{j=1}^n f_j(C)$$

C is called the *attractor* of the IFS.

We will also assume every function f_j is injective.

Example. Take $X = \mathbb{R}$ and $f_1(x) = \frac{1}{2}x$ and $f_2(x) = \frac{1}{2}x + \frac{1}{2}$. The attractor is $C = [0, 1]$.


Example. Take $X = \mathbb{R}$ and $f_1(x) = \frac{1}{2}x$ and $f_2(x) = \frac{1}{2}x + \frac{1}{2}$. The attractor is $C = [0, 1]$.

Example. Take $X = \mathbb{R}$ and $f_1(x) = \frac{1}{3}x$ and $f_2(x) = \frac{1}{3}x + \frac{2}{3}$. The attractor C is the Cantor set.

Iterated function systems

Example. Take $X = \mathbb{R}$ and $f_1(x) = \frac{1}{2}x$ and $f_2(x) = \frac{1}{2}x + \frac{1}{2}$. The attractor is $C = [0, 1]$.

Example. Take $X = \mathbb{R}$ and $f_1(x) = \frac{1}{3}x$ and $f_2(x) = \frac{1}{3}x + \frac{2}{3}$. The attractor C is the Cantor set.



Example. Take $X = \mathbb{R}^2$ and

$$f_1(x) = \frac{1}{2}x \quad f_2(x) = \frac{1}{2}x + \begin{bmatrix} 1/2 \\ 0 \end{bmatrix} \quad f_3(x) = \frac{1}{2}x + \begin{bmatrix} 1/4 \\ \sqrt{3}/4 \end{bmatrix}$$

Example. Take $X = \mathbb{R}^2$ and

$$f_1(x) = \frac{1}{2}x \quad f_2(x) = \frac{1}{2}x + \begin{bmatrix} 1/2 \\ 0 \end{bmatrix} \quad f_3(x) = \frac{1}{2}x + \begin{bmatrix} 1/4 \\ \sqrt{3}/4 \end{bmatrix}$$

The attractor C is the Sierpiński triangle.



Nonhomogeneous extensions

Theorem (Deeley, Putnam, Strung. 2019)

Let (C, \mathcal{F}) be a compact hyperbolic IFS and (X, φ) a Cantor minimal system. There exists a minimal extension $(\tilde{X}, \tilde{\varphi})$ of (X, φ) with factor map $\tilde{\pi} : \tilde{X} \rightarrow X$ such that $\tilde{\pi}^{-1}(x)$ is either a single point or homeomorphic to C (both possibilities always occur).

Nonhomogeneous extensions

Theorem (Deeley, Putnam, Strung. 2019)

Let (C, \mathcal{F}) be a compact hyperbolic IFS and (X, φ) a Cantor minimal system. There exists a minimal extension $(\tilde{X}, \tilde{\varphi})$ of (X, φ) with factor map $\tilde{\pi} : \tilde{X} \rightarrow X$ such that $\tilde{\pi}^{-1}(x)$ is either a single point or homeomorphic to C (both possibilities always occur).

Idea: (X, φ) is topologically conjugate to a Bratteli-Vershik system on a Bratteli diagram (V, E) . To each edge e in E , assign a function f_e in $\mathcal{F} \cup \{\text{id}_C\}$.

$$\tilde{X}_n = \{(x, c) \in X_E \times C \mid c \in f_{x_1} \circ f_{x_2} \circ \cdots \circ f_{x_n}(C)\}$$

Set $\tilde{X} = \bigcap_{n=1}^{\infty} \tilde{X}_n$ and $\tilde{\pi}(x, c) = x$.

Nonhomogeneous extensions

We get a surjective groupoid homomorphism
 $\tilde{\pi} \times \text{id}_{\mathbb{Z}} : \tilde{X} \times \mathbb{Z} \rightarrow X_E \times \mathbb{Z}.$

Nonhomogeneous extensions

We get a surjective groupoid homomorphism

$\tilde{\pi} \times \text{id}_{\mathbb{Z}} : \tilde{X} \times \mathbb{Z} \rightarrow X_E \times \mathbb{Z}$. Using the excision theorem and the Pimsner-Voiculescu exact sequence, we get an exact sequence

$$0 \rightarrow K_0(C_r^*(R_E)) \rightarrow K_0(C_r^*(\tilde{X} \times \mathbb{Z})) \rightarrow K_0(C_r^*(R_E^C)) \otimes (K^0(C)/\mathbb{Z}) \rightarrow 0$$

and an exact sequence

$$0 \rightarrow \mathbb{Z} \rightarrow K_1(C_r^*(\tilde{X} \times \mathbb{Z})) \rightarrow K_0(C_r^*(R_E^C)) \otimes K^{-1}(C) \rightarrow 0$$

$R_E^C \subseteq R_E$ are the tail-equivalent paths that are "eventually id_C ".

Nonhomogeneous extensions

If there is only one path in X_E with $f_{x_n} = \text{id}_C$ for all n , then we get

$$K_0(C_r^*(\tilde{X} \times \mathbb{Z}))/K_0(C_r^*(R_E)) \cong K^0(C)/\mathbb{Z}$$

and

$$K_1(C_r^*(\tilde{X} \times \mathbb{Z}))/\mathbb{Z} \cong K^{-1}(C)$$

Nonhomogeneous extensions

If there is only one path in X_E with $f_{x_n} = \text{id}_C$ for all n , then we get

$$K_0(C_r^*(\tilde{X} \times \mathbb{Z}))/K_0(C_r^*(R_E)) \cong K^0(C)/\mathbb{Z}$$

and

$$K_1(C_r^*(\tilde{X} \times \mathbb{Z}))/\mathbb{Z} \cong K^{-1}(C)$$

- 1 Possibilities for $K^*(C)$?
- 2 When do the sequences split?

Thank you!

- ① Deeley, R.J.; Putnam, I.F.; Strung, K.R. "Non-homogeneous extensions of Cantor minimal systems". to appear, Proc. A.M.S.
- ② Haslehurst, M.J. "Relative K -theory for C^* -algebras", preprint.
- ③ Haslehurst, M.J. "Some examples of factor groupoids". (in preparation)
- ④ Li, X. "Every classifiable simple C^* -algebra has a Cartan subalgebra". Invent. math. 219, 653–699 (2020).
- ⑤ Putnam, I.F. "Some classifiable groupoid C^* -algebras with prescribed K -theory". Math. Ann. 370, 1361–1387 (2018).
- ⑥ Putnam, I.F. "An excision theorem for the K -theory of C^* -algebras, with applications to groupoid C^* -algebras". to appear, Munster Mathematics Journal.