

# Factor groupoids and prescribed $K$ -theory

## GPOTS 2021

Mitch Haslehurst

Department of Mathematics & Statistics  
University of Victoria

May 14, 2021

**Problem.** Given some  $K$ -theory data, find a groupoid  $G$  such that the data is  $K_*(C_r^*(G))$ .

**Problem.** Given some  $K$ -theory data, find a groupoid  $G$  such that the data is  $K_*(C_r^*(G))$ .

Notable references:

Li, X. "Every classifiable simple  $C^*$ -algebra has a Cartan subalgebra". *Invent. math.* 219, 653–699 (2020).

Putnam, I.F. "Some classifiable groupoid  $C^*$ -algebras with prescribed  $K$ -theory". *Math. Ann.* 370, 1361–1387 (2018).

$G'$  is a *factor groupoid* of  $G$  if  $\pi : G \rightarrow G'$  is a surjective groupoid morphism.

$G'$  is a *factor groupoid* of  $G$  if  $\pi : G \rightarrow G'$  is a surjective groupoid morphism.

Assume:

- 1  $G$  and  $G'$  are locally compact Hausdorff and étale,
- 2  $\pi$  is continuous and proper,
- 3  $\pi|_{G^u} : G^u \rightarrow (G')^{\pi(u)}$  is bijective for all  $u$  in  $G^{(0)}$ .

$G'$  is a *factor groupoid* of  $G$  if  $\pi : G \rightarrow G'$  is a surjective groupoid morphism.

Assume:

- 1  $G$  and  $G'$  are locally compact Hausdorff and étale,
- 2  $\pi$  is continuous and proper,
- 3  $\pi|_{G^u} : G^u \rightarrow (G')^{\pi(u)}$  is bijective for all  $u$  in  $G^{(0)}$ .


Obtain an inclusion  $C_r^*(G') \subseteq C_r^*(G)$  via  $b \mapsto b \circ \pi$  ( $b$  in  $C_c(G')$ )

# Bratteli diagrams

Let  $(V, E)$  be a Bratteli diagram.

# Bratteli diagrams


Let  $(V, E)$  be a Brattelli diagram.





# Bratteli diagrams


Let  $(V, E)$  be a Bratteli diagram.



The *infinite path space*  $X_E$  of  $(V, E)$  is a totally disconnected compact metric space.

# Bratteli diagrams

Let  $(V, E)$  be a Bratteli diagram.




The *infinite path space*  $X_E$  of  $(V, E)$  is a totally disconnected compact metric space.

Tail-equivalence  $R_E \subseteq X_E \times X_E$  has an étale topology in which  $C_r^*(R_E)$  is an AF-algebra.

# Bratteli diagrams

Let  $(V, E)$  be a Bratteli diagram.



The *infinite path space*  $X_E$  of  $(V, E)$  is a totally disconnected compact metric space.

Tail-equivalence  $R_E \subseteq X_E \times X_E$  has an étale topology in which  $C_r^*(R_E)$  is an AF-algebra.

**Goal:** make a factor groupoid of  $R_E$ .

Let  $(V, E)$  and  $(W, F)$  be two Bratteli diagrams.

Let  $(V, E)$  and  $(W, F)$  be two Bratteli diagrams.

Two *graph embeddings*  $\xi^0, \xi^1 : (W, F) \rightarrow (V, E)$  with  $\xi^0|_W = \xi^1|_W$  and  $\xi^0(F) \cap \xi^1(F) = \emptyset$ .

# The space $X_\xi$

Let  $(V, E)$  and  $(W, F)$  be two Bratteli diagrams.

Two *graph embeddings*  $\xi^0, \xi^1 : (W, F) \rightarrow (V, E)$  with  $\xi^0|_W = \xi^1|_W$  and  $\xi^0(F) \cap \xi^1(F) = \emptyset$ .

Equivalence relation  $\sim_\xi$  on  $X_E$ :

$$(x_1, x_2, \dots, x_{n_0-1}, x_{n_0}, \xi^0(z_{n_0+1}), \xi^0(z_{n_0+2}), \dots) \quad (1)$$

$$\sim_\xi (x_1, x_2, \dots, x_{n_0-1}, x'_{n_0}, \xi^1(z_{n_0+1}), \xi^1(z_{n_0+2}), \dots) \quad (2)$$

Denote  $X_\xi := X_E / \sim_\xi$  and  $\rho : X_E \rightarrow X_\xi$  the quotient map.

# The space $X_\xi$

Denote  $X_\xi := X_E / \sim_\xi$  and  $\rho : X_E \rightarrow X_\xi$  the quotient map.

Facts:

- 1  $X_\xi$  is a second-countable compact Hausdorff space,
- 2 the covering dimension of  $X_\xi$  is 1,
- 3 each connected component is either a single point or homeomorphic to  $S^1$ .



**Example 1.** We let  $(V, E)$  be the Bratteli diagram with one vertex at each level and two edges at each level. Identify  $X_E$  with  $\{0, 1\}^\omega$ .

$(W, F)$  is a single path, and for  $f$  in  $F$ ,  $\xi^j(f) = j$  for  $j = 0, 1$ .

**Example 1.** We let  $(V, E)$  be the Bratteli diagram with one vertex at each level and two edges at each level. Identify  $X_E$  with  $\{0, 1\}^\omega$ .

$(W, F)$  is a single path, and for  $f$  in  $F$ ,  $\xi^j(f) = j$  for  $j = 0, 1$ .

$$(x_1, x_2, \dots, x_n, 1, 0, 0, 0, 0, \dots) \quad (3)$$

$$\sim_\xi (x_1, x_2, \dots, x_n, 0, 1, 1, 1, 1, \dots) \quad (4)$$

**Example 1.** We let  $(V, E)$  be the Bratteli diagram with one vertex at each level and two edges at each level. Identify  $X_E$  with  $\{0, 1\}^\omega$ .

$(W, F)$  is a single path, and for  $f$  in  $F$ ,  $\xi^j(f) = j$  for  $j = 0, 1$ .

$$(x_1, x_2, \dots, x_n, 1, 0, 0, 0, 0, \dots) \quad (3)$$

$$\sim_\xi (x_1, x_2, \dots, x_n, 0, 1, 1, 1, 1, \dots) \quad (4)$$

$$\{0, 1\}^\omega \rightarrow S^1 : (x_n) \mapsto \exp \left( 2\pi i \sum_{n=1}^{\infty} x_n 2^{-n} \right)$$

**Example 1.** We let  $(V, E)$  be the Bratteli diagram with one vertex at each level and two edges at each level. Identify  $X_E$  with  $\{0, 1\}^\omega$ .

$(W, F)$  is a single path, and for  $f$  in  $F$ ,  $\xi^j(f) = j$  for  $j = 0, 1$ .

$$(x_1, x_2, \dots, x_n, 1, 0, 0, 0, 0, \dots) \quad (3)$$

$$\sim_\xi (x_1, x_2, \dots, x_n, 0, 1, 1, 1, 1, \dots) \quad (4)$$

$$\{0, 1\}^\omega \rightarrow S^1 : (x_n) \mapsto \exp \left( 2\pi i \sum_{n=1}^{\infty} x_n 2^{-n} \right)$$

The fibres are precisely the  $\sim_\xi$  equivalence classes, so  $X_\xi$  is homeomorphic to  $S^1$ .

**Example 2.** Let  $(V, E)$  have one vertex and three edges at each level. Identify  $X_E$  with  $\{0, 1, 2\}^\omega$ .

$(W, F)$  is again a single path, and for  $f$  in  $F$ ,  $\xi^0(f) = 0$  and  $\xi^1(f) = 2$ .


**Example 2.** Let  $(V, E)$  have one vertex and three edges at each level. Identify  $X_E$  with  $\{0, 1, 2\}^\omega$ .

$(W, F)$  is again a single path, and for  $f$  in  $F$ ,  $\xi^0(f) = 0$  and  $\xi^1(f) = 2$ .


There is a nested sequence  $X_1 \subseteq X_2 \subseteq X_3 \subseteq \cdots \subseteq X_E$  such that

$$X_E = \overline{\bigcup_{n=1}^{\infty} X_n}$$

and each  $\rho(X_n)$  is a disjoint union of finitely many circles.




$$\rho(X_1)$$



$$\rho(X_2)$$




# Examples




$\rho(X_3)$

# Examples




$$\rho(X_4)$$

# Examples



$$\rho(X_5)$$



$$\rho(X_6)$$

# The groupoid $R_\xi$

Let  $R_\xi = \rho \times \rho(R_E)$ .

With the quotient topology,  $R_\xi$  is a second-countable locally compact Hausdorff étale groupoid, and a factor of  $R_E$  via  $\rho \times \rho : R_E \rightarrow R_\xi$ .

# The groupoid $R_\xi$

Let  $R_\xi = \rho \times \rho(R_E)$ .

With the quotient topology,  $R_\xi$  is a second-countable locally compact Hausdorff étale groupoid, and a factor of  $R_E$  via  $\rho \times \rho : R_E \rightarrow R_\xi$ .

We want to analyze the  $K$ -theory of  $C_r^*(R_\xi) \subseteq C_r^*(R_E)$ .

If  $A' \subseteq A$ , there is a six-term exact sequence

If  $A' \subseteq A$ , there is a six-term exact sequence

$$\begin{array}{ccccc} K_1(A) & \longrightarrow & K_0(A'; A) & \longrightarrow & K_0(A') \\ \uparrow \scriptstyle \iota_* & & & & \downarrow \scriptstyle \iota_* \\ K_1(A') & \longleftarrow & K_1(A'; A) & \longleftarrow & K_0(A) \end{array}$$



If  $A' \subseteq A$ , there is a six-term exact sequence

$$\begin{array}{ccccc} K_1(A) & \longrightarrow & K_0(A'; A) & \longrightarrow & K_0(A') \\ \uparrow \scriptstyle \iota_* & & & & \downarrow \scriptstyle \iota_* \\ K_1(A') & \longleftarrow & K_1(A'; A) & \longleftarrow & K_0(A) \end{array}$$

$K_*(A'; A)$  is the *relative*  $K$ -theory of the inclusion  $A' \subseteq A$ .

If  $A' \subseteq A$ , there is a six-term exact sequence

$$\begin{array}{ccccc}
 K_1(A) & \longrightarrow & K_0(A'; A) & \longrightarrow & K_0(A') \\
 \uparrow \iota_* & & & & \downarrow \iota_* \\
 K_1(A') & \longleftarrow & K_1(A'; A) & \longleftarrow & K_0(A)
 \end{array}$$

$K_*(A'; A)$  is the *relative* K-theory of the inclusion  $A' \subseteq A$ .

Excision: (Putnam, 2020)

$$K_*(C_r^*(G'); C_r^*(G)) \cong K_*(C_r^*(H'); C_r^*(H))$$

where  $H \subseteq G$  and  $H' \subseteq G'$  are where  $\pi$  is not one-to-one.

Through the results on the previous slide, we obtain

$$K_0(C_r^*(R_\xi)) \cong K_0(C_r^*(R_E)) \quad K_1(C_r^*(R_\xi)) \cong K_0(C_r^*(R_F))$$

Through the results on the previous slide, we obtain

$$K_0(C_r^*(R_\xi)) \cong K_0(C_r^*(R_E)) \quad K_1(C_r^*(R_\xi)) \cong K_0(C_r^*(R_F))$$

Through the set-up  $\xi^0, \xi^1 : (W, F) \rightarrow (V, E)$ , we can prescribe  $K_*(C_r^*(R_\xi))$ .

- ① Increasing the number of embeddings  $\xi^j$ :
  - ① More complicated connected components
  - ② Still one-dimensional
  - ③  $K_0$  the same,  $K_1$  more direct summands
- ② (Based on work of Deeley, Putnam, Strung)  $\pi : \tilde{X} \rightarrow X_E$   
where  $\pi^{-1}(x)$  is either a single point or homeomorphic to a fixed point set of an iterated function system

Thank you!

- ① Deeley, R.J.; Putnam, I.F.; Strung, K.R. "Non-homogeneous extensions of Cantor minimal systems". to appear, Proc. A.M.S.
- ② Haslehurst, M.J. "Relative  $K$ -theory for  $C^*$ -algebras". (in preparation)
- ③ Haslehurst, M.J. "Some examples of factor groupoids". (in preparation)
- ④ Li, X. "Every classifiable simple  $C^*$ -algebra has a Cartan subalgebra". Invent. math. 219, 653–699 (2020).
- ⑤ Putnam, I.F. "Some classifiable groupoid  $C^*$ -algebras with prescribed  $K$ -theory". Math. Ann. 370, 1361–1387 (2018).
- ⑥ Putnam, I.F. "An excision theorem for the  $K$ -theory of  $C^*$ -algebras, with applications to groupoid  $C^*$ -algebras". to appear, Munster Mathematics Journal.