

HYPOTHESIS TESTING



PRESENTED BY:
DR. KALPESH M. POPAT
ASSISTANT PROFESSOR
ATMIYA UNIVERSITY
RAJKOT, GUJARAT(INDIA)

Types of Hypothesis

Types of Hypothesis

Null & Alternative Hypothesis

Types of Hypothesis

Null & Alternative Hypothesis

Types of Hypothesis

Null & Alternative Hypothesis

- ▶ The null hypothesis states that a population parameter (such as the mean, the standard deviation, and so on) is equal to a hypothesized value.

Types of Hypothesis

Null & Alternative Hypothesis

- ▶ The null hypothesis states that a population parameter (such as the mean, the standard deviation, and so on) is equal to a hypothesized value.
- ▶ The null hypothesis is often an initial claim that is based on previous analyses or specialized knowledge.

Types of Hypothesis

Null & Alternative Hypothesis

- ▶ The null hypothesis states that a population parameter (such as the mean, the standard deviation, and so on) is equal to a hypothesized value.
- ▶ The null hypothesis is often an initial claim that is based on previous analyses or specialized knowledge.
- ▶ The alternative hypothesis states that a population parameter is smaller, greater, or different than the hypothesized value in the null hypothesis.

Types of Hypothesis

Null & Alternative Hypothesis

- ▶ The null hypothesis states that a population parameter (such as the mean, the standard deviation, and so on) is equal to a hypothesized value.
- ▶ The null hypothesis is often an initial claim that is based on previous analyses or specialized knowledge.
- ▶ The alternative hypothesis states that a population parameter is smaller, greater, or different than the hypothesized value in the null hypothesis.
- ▶ The alternative hypothesis is what you might believe to be true or hope to prove true.

Null Hypothesis & Alternative Hypothesis

Null Hypothesis & Alternative Hypothesis

- ▶ As an example suppose flour packaged by a manufacturer want to test to determine whether their packaging process is out of control as to determine by weight of flour package is 40 ounce or not.

Null Hypothesis & Alternative Hypothesis

- ▶ As an example suppose flour packaged by a manufacturer want to test to determine whether their packaging process is out of control as to determine by weight of flour package is 40 ounce or not.
- ▶ In this case null and alternate hypothesis stated as follow

Null Hypothesis & Alternative Hypothesis

- ▶ As an example suppose flour packaged by a manufacturer want to test to determine whether their packaging process is out of control as to determine by weight of flour package is 40 ounce or not.
- ▶ In this case null and alternate hypothesis stated as follow

$$H_o : \mu = 40,$$

Null Hypothesis & Alternative Hypothesis

- ▶ As an example suppose flour packaged by a manufacturer want to test to determine whether their packaging process is out of control as to determine by weight of flour package is 40 ounce or not.
- ▶ In this case null and alternate hypothesis stated as follow

$$H_o : \mu = 40, H_a : \mu \neq 40$$

Rejection & Non Rejection Areas

Rejection & Non Rejection Areas

- ▶ Area of statistical outcomes that result in the rejection of null hypothesis is called rejection area.

Rejection & Non Rejection Areas

- ▶ Area of statistical outcomes that result in the rejection of null hypothesis is called rejection area.
- ▶ Area of Statistical outcomes that fail to reject null hypothesis is called non rejection area.

Rejection & Non Rejection Areas

- ▶ Area of statistical outcomes that result in the rejection of null hypothesis is called rejection area.
- ▶ Area of Statistical outcomes that fail to reject null hypothesis is called non rejection area.

Rejection & Non Rejection Areas

Rejection & Non Rejection Areas

- For an example consider flour package manufacturing process example.

Rejection & Non Rejection Areas

- For an example consider flour package manufacturing process example. The null hypothesis is that the average fill for the population packages is 40 ounce.

Rejection & Non Rejection Areas

- ▶ For an example consider flour package manufacturing process example. The null hypothesis is that the average fill for the population packages is 40 ounce.
- ▶ Suppose a sample of 100 such packages is randomly selected

Rejection & Non Rejection Areas

- ▶ For an example consider flour package manufacturing process example. The null hypothesis is that the average fill for the population packages is 40 ounce.
- ▶ Suppose a sample of 100 such packages is randomly selected and a sample mean of 40.01 ounce is obtained.

Rejection & Non Rejection Areas

- ▶ For an example consider flour package manufacturing process example. The null hypothesis is that the average fill for the population packages is 40 ounce.
- ▶ Suppose a sample of 100 such packages is randomly selected and a sample mean of 40.01 ounce is obtained.
- ▶ Because the mean is 40 ounce,

Rejection & Non Rejection Areas

- ▶ For an example consider flour package manufacturing process example. The null hypothesis is that the average fill for the population packages is 40 ounce.
- ▶ Suppose a sample of 100 such packages is randomly selected and a sample mean of 40.01 ounce is obtained.
- ▶ Because the mean is 40 ounce, should the business research decides to reject the null hypothesis?

Rejection & Non Rejection Areas

- ▶ For an example consider flour package manufacturing process example. The null hypothesis is that the average fill for the population packages is 40 ounce.
- ▶ Suppose a sample of 100 such packages is randomly selected and a sample mean of 40.01 ounce is obtained.
- ▶ Because the mean is 40 ounce, should the business research decides to reject the null hypothesis?

Rejection & Non Rejection Areas

Rejection & Non Rejection Areas

- ▶ It make sense that in making random sample from a population with a mean of 40 ounces not all sample means will equal to 40 ounces.

Rejection & Non Rejection Areas

- ▶ It make sense that in making random sample from a population with a mean of 40 ounces not all sample means will equal to 40 ounces.
- ▶ In fact central limit theorem states that for large sample sizes,

Rejection & Non Rejection Areas

- ▶ It make sense that in making random sample from a population with a mean of 40 ounces not all sample means will equal to 40 ounces.
- ▶ In fact central limit theorem states that for large sample sizes, sample means are normally distributed around population mean.

Rejection & Non Rejection Areas

- ▶ It make sense that in making random sample from a population with a mean of 40 ounces not all sample means will equal to 40 ounces.
- ▶ In fact central limit theorem states that for large sample sizes, sample means are normally distributed around population mean.
- ▶ Thus even when the population mean is 40 ounces,

Rejection & Non Rejection Areas

- ▶ It make sense that in making random sample from a population with a mean of 40 ounces not all sample means will equal to 40 ounces.
- ▶ In fact central limit theorem states that for large sample sizes, sample means are normally distributed around population mean.
- ▶ Thus even when the population mean is 40 ounces, a sample mean might still be 40.1, 38.6, or even 44.2

Rejection & Non Rejection Areas

- ▶ It make sense that in making random sample from a population with a mean of 40 ounces not all sample means will equal to 40 ounces.
- ▶ In fact central limit theorem states that for large sample sizes, sample means are normally distributed around population mean.
- ▶ Thus even when the population mean is 40 ounces, a sample mean might still be 40.1, 38.6, or even 44.2
- ▶ How ever a sample mean of 50 ounces is obtained for 100 packages.

Rejection & Non Rejection Areas


- ▶ It make sense that in making random sample from a population with a mean of 40 ounces not all sample means will equal to 40 ounces.
- ▶ In fact central limit theorem states that for large sample sizes, sample means are normally distributed around population mean.
- ▶ Thus even when the population mean is 40 ounces, a sample mean might still be 40.1, 38.6, or even 44.2
- ▶ How ever a sample mean of 50 ounces is obtained for 100 packages.
- ▶ This sample mean is so far away what is reasonable to expect for a population mean is 40 ounces.

Rejection & Non Rejection Areas

- ▶ It make sense that in making random sample from a population with a mean of 40 ounces not all sample means will equal to 40 ounces.
- ▶ In fact central limit theorem states that for large sample sizes, sample means are normally distributed around population mean.
- ▶ Thus even when the population mean is 40 ounces, a sample mean might still be 40.1, 38.6, or even 44.2
- ▶ How ever a sample mean of 50 ounces is obtained for 100 packages.
- ▶ This sample mean is so far away what is reasonable to expect for a population mean is 40 ounces. That the decision is made to reject null hypothesis.

Rejection & Non Rejection Areas

- ▶ It make sense that in making random sample from a population with a mean of 40 ounces not all sample means will equal to 40 ounces.
- ▶ In fact central limit theorem states that for large sample sizes, sample means are normally distributed around population mean.
- ▶ Thus even when the population mean is 40 ounces, a sample mean might still be 40.1, 38.6, or even 44.2
- ▶ How ever a sample mean of 50 ounces is obtained for 100 packages.
- ▶ This sample mean is so far away what is reasonable to expect for a population mean is 40 ounces. That the decision is made to reject null hypothesis.



Rejection & Non Rejection Areas

Rejection & Non Rejection Areas

- Note that critical value in each end(tail) of the distribution.

Rejection & Non Rejection Areas

- ▶ Note that critical value in each end(tail) of the distribution.
- ▶ In each direction beyond the critical values lie the rejection area

Rejection & Non Rejection Areas

- ▶ Note that critical value in each end(tail) of the distribution.
- ▶ In each direction beyond the critical values lie the rejection area
- ▶ Any sample mean that falls in that region will lead the business researcher to reject the null hypothesis.

Rejection & Non Rejection Areas

- ▶ Note that critical value in each end(tail) of the distribution.
- ▶ In each direction beyond the critical values lie the rejection area
- ▶ Any sample mean that falls in that region will lead the business researcher to reject the null hypothesis.
- ▶ Sample means that fall between two critical value are close enough to the population mean that business researcher will decide not to reject null hypothesis.

Rejection & Non Rejection Areas

- ▶ Note that critical value in each end(tail) of the distribution.
- ▶ In each direction beyond the critical values lie the rejection area
- ▶ Any sample mean that falls in that region will lead the business researcher to reject the null hypothesis.
- ▶ Sample means that fall between two critical value are close enough to the population mean that business researcher will decide not to reject null hypothesis.

Type-I Error

Type-I Error

Type-I Error

Type-I Error

Type-I Error

- ▶ A Type-I error is committed by rejecting true null hypothesis.

Type-I Error

Type-I Error

- ▶ A Type-I error is committed by rejecting true null hypothesis.
- ▶ With a Type-I error a null hypothesis is true but the business researcher decided that it is not.

Type-I Error

Type-I Error

- ▶ A Type-I error is committed by rejecting true null hypothesis.
- ▶ With a Type-I error a null hypothesis is true but the business researcher decided that it is not.

Type-I Error

- ▶ As an example of flour packaging process actually in control and is averaging 40 ounce per flour package.

Type-I Error

- ▶ As an example of flour packaging process actually in control and is averaging 40 ounce per flour package.
- ▶ Suppose also that a business researcher randomly select 100 packages,

Type-I Error

- ▶ As an example of flour packaging process actually in control and is averaging 40 ounce per flour package.
- ▶ Suppose also that a business researcher randomly select 100 packages, weight the content of each and compute the sample mean.

Type-I Error

- ▶ As an example of flour packaging process actually in control and is averaging 40 ounce per flour package.
- ▶ Suppose also that a business researcher randomly select 100 packages, weight the content of each and compute the sample mean.
- ▶ It is possible by chance to randomly select 100 of more extreme packages result in a mean that falls in the rejection area.

Type-I Error

- ▶ As an example of flour packaging process actually in control and is averaging 40 ounce per flour package.
- ▶ Suppose also that a business researcher randomly select 100 packages, weight the content of each and compute the sample mean.
- ▶ It is possible by chance to randomly select 100 of more extreme packages result in a mean that falls in the rejection area.
- ▶ The decision is to reject null hypothesis even though the population mean is actually 40 ounce

Type-I Error

- ▶ As an example of flour packaging process actually in control and is averaging 40 ounce per flour package.
- ▶ Suppose also that a business researcher randomly select 100 packages, weight the content of each and compute the sample mean.
- ▶ It is possible by chance to randomly select 100 of more extreme packages result in a mean that falls in the rejection area.
- ▶ The decision is to reject null hypothesis even though the population mean is actually 40 ounce in that case the business researcher has committed Type-I error.

Level of Significance

Level of Significance

- ▶ The probability of committing Type-I error is called α or Level of Significance.

Level of Significance

- ▶ The probability of committing Type-I error is called α or Level of Significance.
- ▶ The value of α set before the experiment or study is undertaken.

Level of Significance

- ▶ The probability of committing Type-I error is called α or Level of Significance.
- ▶ The value of α set before the experiment or study is undertaken.
- ▶ Standard values of α are 0.05 and 0.01.

Level of Significance

- ▶ The probability of committing Type-I error is called α or Level of Significance.
- ▶ The value of α set before the experiment or study is undertaken.
- ▶ Standard values of α are 0.05 and 0.01.

Type-II Error

Type-II Error

Type-II Error

Type-II Error

Type-II Error

- ▶ Type-II error is committed when a business researcher fails to reject false hypothesis.

Type-II Error

Type-II Error

- ▶ Type-II error is committed when a business researcher fails to reject false hypothesis.
- ▶ In this case null hypothesis is false but the decision is to made not to reject it.

Type-II Error

Type-II Error

- ▶ Type-II error is committed when a business researcher fails to reject false hypothesis.
- ▶ In this case null hypothesis is false but the decision is to made not to reject it.

Type-II Error

Type-II Error

- ▶ Suppose in the case of flour packaging problem that the packaging process is actually producing a population mean of 41 ounce even though the null hypothesis is 40 ounce.

Type-II Error

- ▶ Suppose in the case of flour packaging problem that the packaging process is actually producing a population mean of 41 ounce even though the null hypothesis is 40 ounce.
- ▶ A sample of 100 packages yields a sample mean of 40.2 ounces which falls in the non-rejection area.

Type-II Error

- ▶ Suppose in the case of flour packaging problem that the packaging process is actually producing a population mean of 41 ounce even though the null hypothesis is 40 ounce.
- ▶ A sample of 100 packages yields a sample mean of 40.2 ounces which falls in the non-rejection area.
- ▶ The business decision maker decided not to reject null hypothesis.

Type-II Error

- ▶ Suppose in the case of flour packaging problem that the packaging process is actually producing a population mean of 41 ounce even though the null hypothesis is 40 ounce.
- ▶ A sample of 100 packages yields a sample mean of 40.2 ounces which falls in the non-rejection area.
- ▶ The business decision maker decided not to reject null hypothesis.
- ▶ In this case a Type-II error has committed

Type-II Error

- ▶ Suppose in the case of flour packaging problem that the packaging process is actually producing a population mean of 41 ounce even though the null hypothesis is 40 ounce.
- ▶ A sample of 100 packages yields a sample mean of 40.2 ounces which falls in the non-rejection area.
- ▶ The business decision maker decided not to reject null hypothesis.
- ▶ In this case a Type-II error has committed

Table

Accept Null

Table

Accept Null

Reject Null

Table

	Accept Null	Reject Null
Null True		

Table

	Accept Null	Reject Null
Null True		
Null False		

Table

	Accept Null	Reject Null
Null True	Correct Decision	
Null False		

Table

	Accept Null	Reject Null
Null True	Correct Decision	Type-I Error
Null False		

Table

	Accept Null	Reject Null
Null True	Correct Decision	Type-I Error
Null False	Type-II Error	

Table

	Accept Null	Reject Null
Null True	Correct Decision	Type-I Error
Null False	Type-II Error	Correct Decision

Method-I Testing a hypothesis for single mean(large samples)(z-test)

Method-I Testing a hypothesis for single mean (large samples) (z-test)

Necessary Conditions

Method-I Testing a hypothesis for single mean(large samples)(z-test)

Necessary Conditions

- 1 Sample size must be large($n \geq 30$).

Method-I Testing a hypothesis for single mean(large samples)(z-test)

Necessary Conditions

- 1 Sample size must be large($n \geq 30$).
- 2 Population standard deviation must be given(σ -known)

Method-I Testing a hypothesis for single mean(large samples)(z-test)

Necessary Conditions

- 1 Sample size must be large($n \geq 30$).
- 2 Population standard deviation must be given(σ -known)

$$z = \frac{\bar{x} - \mu}{\frac{\sigma}{\sqrt{n}}}$$

Method-I Testing a hypothesis for single mean(large samples)(z-test)

Necessary Conditions

- 1 Sample size must be large($n \geq 30$).
- 2 Population standard deviation must be given(σ -known)

$$z = \frac{\bar{x} - \mu}{\frac{\sigma}{\sqrt{n}}}$$

Where

- \bar{x} =Sample Mean.

Method-I Testing a hypothesis for single mean(large samples)(z-test)

Necessary Conditions

- 1 Sample size must be large($n \geq 30$).
- 2 Population standard deviation must be given(σ -known)

$$z = \frac{\bar{x} - \mu}{\frac{\sigma}{\sqrt{n}}}$$

Where

- ▶ \bar{x} =Sample Mean.
- ▶ μ =Population Mean.

Method-I Testing a hypothesis for single mean(large samples)(z-test)

Necessary Conditions

- 1 Sample size must be large($n \geq 30$).
- 2 Population standard deviation must be given(σ -known)

$$z = \frac{\bar{x} - \mu}{\frac{\sigma}{\sqrt{n}}}$$

Where

- ▶ \bar{x} =Sample Mean.
- ▶ μ =Population Mean.
- ▶ σ =Population standard deviation.

Method-I Testing a hypothesis for single mean(large samples)(z-test)

Necessary Conditions

- 1 Sample size must be large($n \geq 30$).
- 2 Population standard deviation must be given(σ -known)

$$z = \frac{\bar{x} - \mu}{\frac{\sigma}{\sqrt{n}}}$$

Where

- ▶ \bar{x} =Sample Mean.
- ▶ μ =Population Mean.
- ▶ σ =Population standard deviation.
- ▶ n =Sample size(must be ≥ 30)

Method-I Testing a hypothesis for single mean(large samples)(z-test)

Necessary Conditions

- 1 Sample size must be large($n \geq 30$).
- 2 Population standard deviation must be given(σ -known)

$$z = \frac{\bar{x} - \mu}{\frac{\sigma}{\sqrt{n}}}$$

Where

- ▶ \bar{x} =Sample Mean.
- ▶ μ =Population Mean.
- ▶ σ =Population standard deviation.
- ▶ n =Sample size(must be ≥ 30)

Confidence Interval

Confidence Interval

Confidence Interval

Confidence Interval

Confidence Interval

$$\bar{x} - z_{\alpha/2} \frac{\sigma}{\sqrt{n}} \leq \mu \leq \bar{x} + z_{\alpha/2} \frac{\sigma}{\sqrt{n}}$$

Confidence Interval

Confidence Interval

$$\bar{x} - z_{\alpha/2} \frac{\sigma}{\sqrt{n}} \leq \mu \leq \bar{x} + z_{\alpha/2} \frac{\sigma}{\sqrt{n}}$$

Example:

Confidence Interval

Confidence Interval

$$\bar{x} - z_{\alpha/2} \frac{\sigma}{\sqrt{n}} \leq \mu \leq \bar{x} + z_{\alpha/2} \frac{\sigma}{\sqrt{n}}$$

Example: The mean height obtained from a random sample of size 100 is 64 inches.

Confidence Interval

Confidence Interval

$$\bar{x} - z_{\alpha/2} \frac{\sigma}{\sqrt{n}} \leq \mu \leq \bar{x} + z_{\alpha/2} \frac{\sigma}{\sqrt{n}}$$

Example: The mean height obtained from a random sample of size 100 is 64 inches. The standard deviation of distribution of height is known to be 3 inches.

Confidence Interval

Confidence Interval

$$\bar{x} - z_{\alpha/2} \frac{\sigma}{\sqrt{n}} \leq \mu \leq \bar{x} + z_{\alpha/2} \frac{\sigma}{\sqrt{n}}$$

Example: The mean height obtained from a random sample of size 100 is 64 inches. The standard deviation of distribution of height is known to be 3 inches. Test the statement that mean height of population is 67 inches at 5% level of significance.

Confidence Interval

Confidence Interval

$$\bar{x} - z_{\alpha/2} \frac{\sigma}{\sqrt{n}} \leq \mu \leq \bar{x} + z_{\alpha/2} \frac{\sigma}{\sqrt{n}}$$

Example: The mean height obtained from a random sample of size 100 is 64 inches. The standard deviation of distribution of height is known to be 3 inches. Test the statement that mean height of population is 67 inches at 5% level of significance.

Examples

Examples

Example:

Examples

Example: A sample of 100 tires are taken from a lot. The mean life of tires is found to be 39,350 kms

Examples

Example: A sample of 100 tires are taken from a lot. The mean life of tires is found to be 39,350 kms with population standard deviation of 3260 kms.

Examples

Example: A sample of 100 tires are taken from a lot. The mean life of tires is found to be 39,350 kms with population standard deviation of 3260 kms. Could the sample come from a population with mean of 40,000 kms?

Examples

Example: A sample of 100 tires are taken from a lot. The mean life of tires is found to be 39,350 kms with population standard deviation of 3260 kms. Could the sample come from a population with mean of 40,000 kms? (Take $\alpha = 0.05$)

Examples

Example: A sample of 100 tires are taken from a lot. The mean life of tires is found to be 39,350 kms with population standard deviation of 3260 kms. Could the sample come from a population with mean of 40,000 kms? (Take $\alpha = 0.05$) Also Establish 99% confidence limit in which mean life of tires expected to be lie.

Examples

Example: A sample of 100 tires are taken from a lot. The mean life of tires is found to be 39,350 kms with population standard deviation of 3260 kms. Could the sample come from a population with mean of 40,000 kms? (Take $\alpha = 0.05$) Also Establish 99% confidence limit in which mean life of tires expected to be lie.

Example:

Examples

Example: A sample of 100 tires are taken from a lot. The mean life of tires is found to be 39,350 kms with population standard deviation of 3260 kms. Could the sample come from a population with mean of 40,000 kms? (Take $\alpha = 0.05$) Also Establish 99% confidence limit in which mean life of tires expected to be lie.

Example: The mean life of a sample of 400 florescent light bulbs produced by the company is found to be 1570 hrs

Examples

Example: A sample of 100 tires are taken from a lot. The mean life of tires is found to be 39,350 kms with population standard deviation of 3260 kms. Could the sample come from a population with mean of 40,000 kms? (Take $\alpha = 0.05$) Also Establish 99% confidence limit in which mean life of tires expected to be lie.

Example: The mean life of a sample of 400 florescent light bulbs produced by the company is found to be 1570 hrs with a standard deviation of 150 hrs.

Examples

Example: A sample of 100 tires are taken from a lot. The mean life of tires is found to be 39,350 kms with population standard deviation of 3260 kms. Could the sample come from a population with mean of 40,000 kms? (Take $\alpha = 0.05$) Also Establish 99% confidence limit in which mean life of tires expected to be lie.

Example: The mean life of a sample of 400 florescent light bulbs produced by the company is found to be 1570 hrs with a standard deviation of 150 hrs. Test the hypothesis that the mean life of light bulbs produced by the company is 1600 hrs

Examples

Example: A sample of 100 tires are taken from a lot. The mean life of tires is found to be 39,350 kms with population standard deviation of 3260 kms. Could the sample come from a population with mean of 40,000 kms? (Take $\alpha = 0.05$) Also Establish 99% confidence limit in which mean life of tires expected to be lie.

Example: The mean life of a sample of 400 florescent light bulbs produced by the company is found to be 1570 hrs with a standard deviation of 150 hrs. Test the hypothesis that the mean life of light bulbs produced by the company is 1600 hrs against alternative hypothesis that is mean life is less than 1600 hrs at 1% level of significance.

Examples

Example: A sample of 100 tires are taken from a lot. The mean life of tires is found to be 39,350 kms with population standard deviation of 3260 kms. Could the sample come from a population with mean of 40,000 kms? (Take $\alpha = 0.05$) Also Establish 99% confidence limit in which mean life of tires expected to be lie.

Example: The mean life of a sample of 400 florescent light bulbs produced by the company is found to be 1570 hrs with a standard deviation of 150 hrs. Test the hypothesis that the mean life of light bulbs produced by the company is 1600 hrs against alternative hypothesis that is mean life is less than 1600 hrs at 1% level of significance.

Method-II Testing a hypothesis for a single mean (small sample) (t-test)

Method-II Testing a hypothesis for a single mean (small sample) (t-test)

Conditions

Method-II Testing a hypothesis for a single mean (small sample) (t-test)

Conditions

One of the following cases occur.

Method-II Testing a hypothesis for a single mean (small sample) (t-test)

Conditions

One of the following cases occur.

Method-II Testing a hypothesis for a single mean(small sample)(t-test)

Conditions

One of the following case occur.

- 1 Sample size is small($n < 30$).

Method-II Testing a hypothesis for a single mean(small sample)(t-test)

Conditions

One of the following case occur.

- 1 Sample size is small($n < 30$).
- 2 Population standard deviation is not given (σ -unknown)

Method-II Testing a hypothesis for a single mean(small sample)(t-test)

Conditions

One of the following case occur.

- 1 Sample size is small($n < 30$).
- 2 Population standard deviation is not given (σ -unknown)

Method-II Testing a hypothesis for a single mean(small sample)(t-test)

Conditions

One of the following case occur.

- 1 Sample size is small($n < 30$).
- 2 Population standard deviation is not given (σ -unknown)

$$t = \frac{\bar{x} - \mu}{\frac{s}{\sqrt{n}}}, d.f = n - 1$$

Method-II Testing a hypothesis for a single mean(small sample)(t-test)

Conditions

One of the following case occur.

- 1 Sample size is small($n < 30$).
- 2 Population standard deviation is not given (σ -unknown)

$$t = \frac{\bar{x} - \mu}{\frac{s}{\sqrt{n}}}, d.f = n - 1$$

Where

Method-II Testing a hypothesis for a single mean(small sample)(t-test)

Conditions

One of the following case occur.

- 1 Sample size is small($n < 30$).
- 2 Population standard deviation is not given (σ -unknown)

$$t = \frac{\bar{x} - \mu}{\frac{s}{\sqrt{n}}}, d.f = n - 1$$

Where

- \bar{x} =Sample Mean.

Method-II Testing a hypothesis for a single mean(small sample)(t-test)

Conditions

One of the following case occur.

- 1 Sample size is small($n < 30$).
- 2 Population standard deviation is not given (σ -unknown)

$$t = \frac{\bar{x} - \mu}{\frac{s}{\sqrt{n}}}, d.f = n - 1$$

Where

- ▶ \bar{x} =Sample Mean.
- ▶ μ =Population Mean.

Method-II Testing a hypothesis for a single mean(small sample)(t-test)

Conditions

One of the following case occur.

- 1 Sample size is small($n < 30$).
- 2 Population standard deviation is not given (σ -unknown)

$$t = \frac{\bar{x} - \mu}{\frac{s}{\sqrt{n}}}, d.f = n - 1$$

Where

- ▶ \bar{x} =Sample Mean.
- ▶ μ =Population Mean.
- ▶ s =Sample standard deviation.

Method-II Testing a hypothesis for a single mean(small sample)(t-test)

Conditions

One of the following case occur.

- 1 Sample size is small($n < 30$).
- 2 Population standard deviation is not given (σ -unknown)

$$t = \frac{\bar{x} - \mu}{\frac{s}{\sqrt{n}}}, d.f = n - 1$$

Where

- ▶ \bar{x} =Sample Mean.
- ▶ μ =Population Mean.
- ▶ s =Sample standard deviation.
- ▶ n =Sample size(may be < 30)

Method-II Testing a hypothesis for a single mean(small sample)(t-test)

Conditions

One of the following case occur.

- 1 Sample size is small($n < 30$).
- 2 Population standard deviation is not given (σ -unknown)

$$t = \frac{\bar{x} - \mu}{\frac{s}{\sqrt{n}}}, d.f = n - 1$$

Where

- ▶ \bar{x} =Sample Mean.
- ▶ μ =Population Mean.
- ▶ s =Sample standard deviation.
- ▶ n =Sample size(may be < 30)

Confidence Interval

Confidence Interval

Confidence interval

Confidence Interval

Confidence interval

$$\bar{x} - t_{\alpha/2} \frac{s}{\sqrt{n}} \leq \mu \leq \bar{x} + t_{\alpha/2} \frac{s}{\sqrt{n}}$$

Confidence Interval

Confidence interval

$$\bar{x} - t_{\alpha/2} \frac{s}{\sqrt{n}} \leq \mu \leq \bar{x} + t_{\alpha/2} \frac{s}{\sqrt{n}}$$

Example:

Confidence Interval

Confidence interval

$$\bar{x} - t_{\alpha/2} \frac{s}{\sqrt{n}} \leq \mu \leq \bar{x} + t_{\alpha/2} \frac{s}{\sqrt{n}}$$

Example: A random sample of size 16 has 53 as mean.

Confidence Interval

Confidence interval

$$\bar{x} - t_{\alpha/2} \frac{s}{\sqrt{n}} \leq \mu \leq \bar{x} + t_{\alpha/2} \frac{s}{\sqrt{n}}$$

Example: A random sample of size 16 has 53 as mean. The sum of the square of deviation taken from mean is 135.

Confidence Interval

Confidence interval

$$\bar{x} - t_{\alpha/2} \frac{s}{\sqrt{n}} \leq \mu \leq \bar{x} + t_{\alpha/2} \frac{s}{\sqrt{n}}$$

Example: A random sample of size 16 has 53 as mean. The sum of the square of deviation taken from mean is 135. Can this sample be regarded as taken from the population having mean as 56?

Confidence Interval

Confidence interval

$$\bar{x} - t_{\alpha/2} \frac{s}{\sqrt{n}} \leq \mu \leq \bar{x} + t_{\alpha/2} \frac{s}{\sqrt{n}}$$

Example: A random sample of size 16 has 53 as mean. The sum of the square of deviation taken from mean is 135. Can this sample be regarded as taken from the population having mean as 56? (Take $\alpha = 0.05$) Also obtain 95% and 99% confidence limit of the population mean.

Confidence Interval

Confidence interval

$$\bar{x} - t_{\alpha/2} \frac{s}{\sqrt{n}} \leq \mu \leq \bar{x} + t_{\alpha/2} \frac{s}{\sqrt{n}}$$

Example: A random sample of size 16 has 53 as mean. The sum of the square of deviation taken from mean is 135. Can this sample be regarded as taken from the population having mean as 56? (Take $\alpha = 0.05$) Also obtain 95% and 99% confidence limit of the population mean.

Examples

Examples

Example:

Examples

Example: The manufacturer of a certain types of bulb claims that this bulb have a mean of 25 months with a standard deviation of 5 months.

Examples

Example: The manufacturer of a certain types of bulb claims that this bulb have a mean of 25 months with a standard deviation of 5 months. A random sample of 6 such bulb gave following values

Examples

Example: The manufacturer of a certain types of bulb claims that this bulb have a mean of 25 months with a standard deviation of 5 months. A random sample of 6 such bulb gave following values
24,

Examples

Example: The manufacturer of a certain types of bulb claims that this bulb have a mean of 25 months with a standard deviation of 5 months. A random sample of 6 such bulb gave following values
24, 26,

Examples

Example: The manufacturer of a certain types of bulb claims that this bulb have a mean of 25 months with a standard deviation of 5 months. A random sample of 6 such bulb gave following values
24, 26, 30,

Examples

Example: The manufacturer of a certain types of bulb claims that this bulb have a mean of 25 months with a standard deviation of 5 months. A random sample of 6 such bulb gave following values
24, 26, 30, 20,

Examples

Example: The manufacturer of a certain types of bulb claims that this bulb have a mean of 25 months with a standard deviation of 5 months. A random sample of 6 such bulb gave following values
24, 26, 30, 20, 20,

Examples

Example: The manufacturer of a certain types of bulb claims that this bulb have a mean of 25 months with a standard deviation of 5 months. A random sample of 6 such bulb gave following values
24, 26, 30, 20, 20, 18.

Examples

Example: The manufacturer of a certain types of bulb claims that this bulb have a mean of 25 months with a standard deviation of 5 months. A random sample of 6 such bulb gave following values 24, 26, 30, 20, 20, 18. Can you regards the manufacturer's claim to be valid at 1% level of significance?

Examples

Example: The manufacturer of a certain types of bulb claims that this bulb have a mean of 25 months with a standard deviation of 5 months. A random sample of 6 such bulb gave following values 24, 26, 30, 20, 20, 18. Can you regards the manufacturer's claim to be valid at 1% level of significance?

Example:

Examples

Example: The manufacturer of a certain types of bulb claims that this bulb have a mean of 25 months with a standard deviation of 5 months. A random sample of 6 such bulb gave following values 24, 26, 30, 20, 20, 18. Can you regards the manufacturer's claim to be valid at 1% level of significance?

Example: The life time of electric bulbs for a random sample of 10 from large consignment gave the following data:

Examples

Example: The manufacturer of a certain types of bulb claims that this bulb have a mean of 25 months with a standard deviation of 5 months. A random sample of 6 such bulb gave following values 24,26,30,20,20,18. Can you regards the manufacturer's claim to be valid at 1% level of significance?

Example: The life time of electric bulbs for a random sample of 10 from large consignment gave the following data:

Item	1	2	3	4	5	6	7	8	9	10
Life ('000 hrs)	4.2	4.6	3.9	4.1	5.2	3.8	3.9	4.3	4.4	5.0

Examples

Example: The manufacturer of a certain types of bulb claims that this bulb have a mean of 25 months with a standard deviation of 5 months. A random sample of 6 such bulb gave following values 24, 26, 30, 20, 20, 18. Can you regards the manufacturer's claim to be valid at 1% level of significance?

Example: The life time of electric bulbs for a random sample of 10 from large consignment gave the following data:

Item	1	2	3	4	5	6	7	8	9	10
Life ('000 hrs)	4.2	4.6	3.9	4.1	5.2	3.8	3.9	4.3	4.4	5.1

Can you regard the mean life of light bulbs is atleast 4000 hrs?

Examples

Example: The manufacturer of a certain types of bulb claims that this bulb have a mean of 25 months with a standard deviation of 5 months. A random sample of 6 such bulb gave following values 24, 26, 30, 20, 20, 18. Can you regards the manufacturer's claim to be valid at 1% level of significance?

Example: The life time of electric bulbs for a random sample of 10 from large consignment gave the following data:

Item	1	2	3	4	5	6	7	8	9	10
Life ('000 hrs)	4.2	4.6	3.9	4.1	5.2	3.8	3.9	4.3	4.4	5.1

Can you regard the mean life of light bulbs is atleast 4000 hrs? (Take $\alpha = 0.05$)

Method-III Testing a hypothesis for single proportion(z-test)

Method-III Testing a hypothesis for single proportion(z-test)

Necessary Conditions

Method-III Testing a hypothesis for single proportion(z-test)

Necessary Conditions

Method-III Testing a hypothesis for single proportion(z-test)

Necessary Conditions

1 $n.p \geq 5$

Method-III Testing a hypothesis for single proportion(z-test)

Necessary Conditions

- 1 $n.p \geq 5$
- 2 $n.(1-p) \geq 5$

Method-III Testing a hypothesis for single proportion(z-test)

Necessary Conditions

- 1 $n.p \geq 5$
- 2 $n.(1-p) \geq 5$

Method-III Testing a hypothesis for single proportion(z-test)

Necessary Conditions

1 $n.p \geq 5$

2 $n.(1-p) \geq 5$

$$z = \frac{\bar{p} - p}{\sqrt{\frac{p.(1-p)}{n}}}$$

Method-III Testing a hypothesis for single proportion(z-test)

Necessary Conditions

1 $n.p \geq 5$

2 $n.(1-p) \geq 5$

$$z = \frac{\bar{p} - p}{\sqrt{\frac{p.(1-p)}{n}}}$$

Where

Method-III Testing a hypothesis for single proportion(z-test)

Necessary Conditions

1 $n.p \geq 5$

2 $n.(1-p) \geq 5$

$$z = \frac{\bar{p} - p}{\sqrt{\frac{p.(1-p)}{n}}}$$

Where

Method-III Testing a hypothesis for single proportion(z-test)

Necessary Conditions

1 $n.p \geq 5$

2 $n.(1-p) \geq 5$

$$z = \frac{\bar{p} - p}{\sqrt{\frac{p.(1-p)}{n}}}$$

Where

► \bar{p} =Sample proportion.

Method-III Testing a hypothesis for single proportion(z-test)

Necessary Conditions

1 $n.p \geq 5$

2 $n.(1-p) \geq 5$

$$z = \frac{\bar{p} - p}{\sqrt{\frac{p.(1-p)}{n}}}$$

Where

- ▶ \bar{p} =Sample proportion.
- ▶ p =Population proportion.

Method-III Testing a hypothesis for single proportion(z-test)

Necessary Conditions

1 $n.p \geq 5$

2 $n.(1-p) \geq 5$

$$z = \frac{\bar{p} - p}{\sqrt{\frac{p.(1-p)}{n}}}$$

Where

- ▶ \bar{p} =Sample proportion.
- ▶ p =Population proportion.
- ▶ n =Sample size.

Method-III Testing a hypothesis for single proportion(z-test)

Necessary Conditions

1 $n.p \geq 5$

2 $n.(1-p) \geq 5$

$$z = \frac{\bar{p} - p}{\sqrt{\frac{p.(1-p)}{n}}}$$

Where

- ▶ \bar{p} =Sample proportion.
- ▶ p =Population proportion.
- ▶ n =Sample size.

Method-III Testing a hypothesis for single proportion(z-test)

Necessary Conditions

1 $n.p \geq 5$

2 $n.(1-p) \geq 5$

$$z = \frac{\bar{p} - p}{\sqrt{\frac{p.(1-p)}{n}}}$$

Where

- ▶ \bar{p} =Sample proportion.
- ▶ p =Population proportion.
- ▶ n =Sample size.

Confidence Interval

Confidence Interval

Confidence interval

Confidence Interval

Confidence interval

$$\bar{p} - z_{\alpha/2} \sqrt{\frac{\bar{p}(1 - \bar{p})}{n}} \leq p \leq \bar{p} + z_{\alpha/2} \sqrt{\frac{\bar{p}(1 - \bar{p})}{n}}$$

Confidence Interval

Confidence interval

$$\bar{p} - z_{\alpha/2} \sqrt{\frac{\bar{p}(1 - \bar{p})}{n}} \leq p \leq \bar{p} + z_{\alpha/2} \sqrt{\frac{\bar{p}(1 - \bar{p})}{n}}$$

Example:

Confidence Interval

Confidence interval

$$\bar{p} - z_{\alpha/2} \sqrt{\frac{\bar{p}(1 - \bar{p})}{n}} \leq p \leq \bar{p} + z_{\alpha/2} \sqrt{\frac{\bar{p}(1 - \bar{p})}{n}}$$

Example: A coin tossed 400 times and head turned up 216 times.

Confidence Interval

Confidence interval

$$\bar{p} - z_{\alpha/2} \sqrt{\frac{\bar{p}(1 - \bar{p})}{n}} \leq p \leq \bar{p} + z_{\alpha/2} \sqrt{\frac{\bar{p}(1 - \bar{p})}{n}}$$

Example: A coin tossed 400 times and head turned up 216 times. Test the hypothesis that the coin is unbiased.

Confidence Interval

Confidence interval

$$\bar{p} - z_{\alpha/2} \sqrt{\frac{\bar{p}(1 - \bar{p})}{n}} \leq p \leq \bar{p} + z_{\alpha/2} \sqrt{\frac{\bar{p}(1 - \bar{p})}{n}}$$

Example: A coin tossed 400 times and head turned up 216 times. Test the hypothesis that the coin is unbiased. (Take $\alpha = 0.05$)

Confidence Interval

Confidence interval

$$\bar{p} - z_{\alpha/2} \sqrt{\frac{\bar{p}(1 - \bar{p})}{n}} \leq p \leq \bar{p} + z_{\alpha/2} \sqrt{\frac{\bar{p}(1 - \bar{p})}{n}}$$

Example: A coin tossed 400 times and head turned up 216 times. Test the hypothesis that the coin is unbiased. (Take $\alpha = 0.05$)

Example:

Confidence Interval

Confidence interval

$$\bar{p} - z_{\alpha/2} \sqrt{\frac{\bar{p}(1 - \bar{p})}{n}} \leq p \leq \bar{p} + z_{\alpha/2} \sqrt{\frac{\bar{p}(1 - \bar{p})}{n}}$$

Example: A coin tossed 400 times and head turned up 216 times. Test the hypothesis that the coin is unbiased. (Take $\alpha = 0.05$)

Example: In a hospital 480 female and 520 male babies were born in a week.

Confidence Interval

Confidence interval

$$\bar{p} - z_{\alpha/2} \sqrt{\frac{\bar{p}(1 - \bar{p})}{n}} \leq p \leq \bar{p} + z_{\alpha/2} \sqrt{\frac{\bar{p}(1 - \bar{p})}{n}}$$

Example: A coin tossed 400 times and head turned up 216 times. Test the hypothesis that the coin is unbiased. (Take $\alpha = 0.05$)

Example: In a hospital 480 female and 520 male babies were born in a week. Do this figure confirm the hypothesis that male and female were born in equal numbers?

Confidence Interval

Confidence interval

$$\bar{p} - z_{\alpha/2} \sqrt{\frac{\bar{p}(1 - \bar{p})}{n}} \leq p \leq \bar{p} + z_{\alpha/2} \sqrt{\frac{\bar{p}(1 - \bar{p})}{n}}$$

Example: A coin tossed 400 times and head turned up 216 times. Test the hypothesis that the coin is unbiased. (Take $\alpha = 0.05$)

Example: In a hospital 480 female and 520 male babies were born in a week. Do this figure confirm the hypothesis that male and female were born in equal numbers? (Take $\alpha = 0.05$)

Confidence Interval

Confidence interval

$$\bar{p} - z_{\alpha/2} \sqrt{\frac{\bar{p}(1 - \bar{p})}{n}} \leq p \leq \bar{p} + z_{\alpha/2} \sqrt{\frac{\bar{p}(1 - \bar{p})}{n}}$$

Example: A coin tossed 400 times and head turned up 216 times. Test the hypothesis that the coin is unbiased. (Take $\alpha = 0.05$)

Example: In a hospital 480 female and 520 male babies were born in a week. Do this figure confirm the hypothesis that male and female were born in equal numbers? (Take $\alpha = 0.05$)

Examples

Examples

Example: Out of 20,000 customer ledger accounts a sample of 600 were taken to test the accuracy of positing balancing

Examples

Example: Out of 20,000 customer ledger accounts a sample of 600 were taken to test the accuracy of positing balancing and 45 mistakes were found.

Examples

Example: Out of 20,000 customer ledger accounts a sample of 600 were taken to test the accuracy of positing balancing and 45 mistakes were found. Assign the limit in which the no of mistakes can be expected to be lie at 95% confidence.

Examples

Example: Out of 20,000 customer ledger accounts a sample of 600 were taken to test the accuracy of positing balancing and 45 mistakes were found. Assign the limit in which the no of mistakes can be expected to be lie at 95% confidence.

Example:

Examples

Example: Out of 20,000 customer ledger accounts a sample of 600 were taken to test the accuracy of positing balancing and 45 mistakes were found. Assign the limit in which the no of mistakes can be expected to be lie at 95% confidence.

Example: A factory is producing 50,000 pair of shoes daily.

Examples

Example: Out of 20,000 customer ledger accounts a sample of 600 were taken to test the accuracy of positing balancing and 45 mistakes were found. Assign the limit in which the no of mistakes can be expected to be lie at 95% confidence.

Example: A factory is producing 50,000 pair of shoes daily. From a sample of 500 pairs 2% were found to be sub-standard quality.

Examples

Example: Out of 20,000 customer ledger accounts a sample of 600 were taken to test the accuracy of positing balancing and 45 mistakes were found. Assign the limit in which the no of mistakes can be expected to be lie at 95% confidence.

Example: A factory is producing 50,000 pair of shoes daily. From a sample of 500 pairs 2% were found to be sub-standard quality. Estimates the no of pairs to be spoiled in the daily production at 95% confidence.

Examples

Example: Out of 20,000 customer ledger accounts a sample of 600 were taken to test the accuracy of positing balancing and 45 mistakes were found. Assign the limit in which the no of mistakes can be expected to be lie at 95% confidence.

Example: A factory is producing 50,000 pair of shoes daily. From a sample of 500 pairs 2% were found to be sub-standard quality. Estimates the no of pairs to be spoiled in the daily production at 95% confidence.

Method-IV Testing a hypothesis for double population mean(large samples)(z-test)

Method-IV Testing a hypothesis for double population mean(large samples)(z-test)

Necessary Conditions

Method-IV Testing a hypothesis for double population mean(large samples)(z-test)

Necessary Conditions

- 1 Both sample sizes must be large($n_1 \geq 30, n_2 \geq 30$)

Method-IV Testing a hypothesis for double population mean(large samples)(z-test)

Necessary Conditions

- 1 Both sample sizes must be large($n_1 \geq 30, n_2 \geq 30$)
- 2 Both population standard deviations must be given. (σ_1 and σ_2 known)

Method-IV Testing a hypothesis for double population mean(large samples)(z-test)

Necessary Conditions

- 1 Both sample sizes must be large($n_1 \geq 30, n_2 \geq 30$)
- 2 Both population standard deviations must be given. (σ_1 and σ_2 known)

Types of hypothesis

Method-IV Testing a hypothesis for double population mean(large samples)(z-test)

Necessary Conditions

- 1 Both sample sizes must be large($n_1 \geq 30, n_2 \geq 30$)
- 2 Both population standard deviations must be given. (σ_1 and σ_2 known)

Types of hypothesis

- 1 $H_o : \mu_1 - \mu_2 = 0, H_a : \mu_1 - \mu_2 \neq 0$

Method-IV Testing a hypothesis for double population mean(large samples)(z-test)

Necessary Conditions

- 1 Both sample sizes must be large($n_1 \geq 30, n_2 \geq 30$)
- 2 Both population standard deviations must be given. (σ_1 and σ_2 known)

Types of hypothesis

- 1 $H_o : \mu_1 - \mu_2 = 0, H_a : \mu_1 - \mu_2 \neq 0$
- 2 $H_o : \mu_1 - \mu_2 \geq 0, H_a : \mu_1 - \mu_2 < 0$

Method-IV Testing a hypothesis for double population mean(large samples)(z-test)

Necessary Conditions

- 1 Both sample sizes must be large($n_1 \geq 30, n_2 \geq 30$)
- 2 Both population standard deviations must be given. (σ_1 and σ_2 known)

Types of hypothesis

- 1 $H_o : \mu_1 - \mu_2 = 0, H_a : \mu_1 - \mu_2 \neq 0$
- 2 $H_o : \mu_1 - \mu_2 \geq 0, H_a : \mu_1 - \mu_2 < 0$
- 3 $H_o : \mu_1 - \mu_2 \leq 0, H_a : \mu_1 - \mu_2 > 0$

Method-IV Testing a hypothesis for double population mean(large samples)(z-test)

Necessary Conditions

- 1 Both sample sizes must be large($n_1 \geq 30, n_2 \geq 30$)
- 2 Both population standard deviations must be given. (σ_1 and σ_2 known)

Types of hypothesis

- 1 $H_o : \mu_1 - \mu_2 = 0, H_a : \mu_1 - \mu_2 \neq 0$
- 2 $H_o : \mu_1 - \mu_2 \geq 0, H_a : \mu_1 - \mu_2 < 0$
- 3 $H_o : \mu_1 - \mu_2 \leq 0, H_a : \mu_1 - \mu_2 > 0$

z-formula

z-formula

$$z = \frac{(\bar{x}_1 - \bar{x}_2) - (\mu_1 - \mu_2)}{\sqrt{\frac{\sigma_1^2}{n_1} + \frac{\sigma_2^2}{n_2}}}$$

z-formula

$$z = \frac{(\bar{x}_1 - \bar{x}_2) - (\mu_1 - \mu_2)}{\sqrt{\frac{\sigma_1^2}{n_1} + \frac{\sigma_2^2}{n_2}}}$$

Where

z-formula

$$z = \frac{(\bar{x}_1 - \bar{x}_2) - (\mu_1 - \mu_2)}{\sqrt{\frac{\sigma_1^2}{n_1} + \frac{\sigma_2^2}{n_2}}}$$

Where

► \bar{x}_1 = First sample mean.

z-formula

$$z = \frac{(\bar{x}_1 - \bar{x}_2) - (\mu_1 - \mu_2)}{\sqrt{\frac{\sigma_1^2}{n_1} + \frac{\sigma_2^2}{n_2}}}$$

Where

- ▶ \bar{x}_1 =First sample mean.
- ▶ \bar{x}_2 =Second sample mean.

z-formula

$$z = \frac{(\bar{x}_1 - \bar{x}_2) - (\mu_1 - \mu_2)}{\sqrt{\frac{\sigma_1^2}{n_1} + \frac{\sigma_2^2}{n_2}}}$$

Where

- ▶ \bar{x}_1 =First sample mean.
- ▶ \bar{x}_2 =Second sample mean.
- ▶ μ_1 =First population mean.

z-formula

$$z = \frac{(\bar{x}_1 - \bar{x}_2) - (\mu_1 - \mu_2)}{\sqrt{\frac{\sigma_1^2}{n_1} + \frac{\sigma_2^2}{n_2}}}$$

Where

- ▶ \bar{x}_1 =First sample mean.
- ▶ \bar{x}_2 =Second sample mean.
- ▶ μ_1 =First population mean.
- ▶ μ_2 =Second population mean.

z-formula

$$z = \frac{(\bar{x}_1 - \bar{x}_2) - (\mu_1 - \mu_2)}{\sqrt{\frac{\sigma_1^2}{n_1} + \frac{\sigma_2^2}{n_2}}}$$

Where

- ▶ \bar{x}_1 =First sample mean.
- ▶ \bar{x}_2 =Second sample mean.
- ▶ μ_1 =First population mean.
- ▶ μ_2 =Second population mean.
- ▶ σ_1 =First population standard deviation.

z-formula

$$z = \frac{(\bar{x}_1 - \bar{x}_2) - (\mu_1 - \mu_2)}{\sqrt{\frac{\sigma_1^2}{n_1} + \frac{\sigma_2^2}{n_2}}}$$

Where

- ▶ \bar{x}_1 =First sample mean.
- ▶ \bar{x}_2 =Second sample mean.
- ▶ μ_1 =First population mean.
- ▶ μ_2 =Second population mean.
- ▶ σ_1 =First population standard deviation.
- ▶ σ_2 =Second population standard deviation.

z-formula

$$z = \frac{(\bar{x}_1 - \bar{x}_2) - (\mu_1 - \mu_2)}{\sqrt{\frac{\sigma_1^2}{n_1} + \frac{\sigma_2^2}{n_2}}}$$

Where

- ▶ \bar{x}_1 =First sample mean.
- ▶ \bar{x}_2 =Second sample mean.
- ▶ μ_1 =First population mean.
- ▶ μ_2 =Second population mean.
- ▶ σ_1 =First population standard deviation.
- ▶ σ_2 =Second population standard deviation.
- ▶ n_1 =First sample size.(≥ 30)

z-formula

$$z = \frac{(\bar{x}_1 - \bar{x}_2) - (\mu_1 - \mu_2)}{\sqrt{\frac{\sigma_1^2}{n_1} + \frac{\sigma_2^2}{n_2}}}$$

Where

- ▶ \bar{x}_1 =First sample mean.
- ▶ \bar{x}_2 =Second sample mean.
- ▶ μ_1 =First population mean.
- ▶ μ_2 =Second population mean.
- ▶ σ_1 =First population standard deviation.
- ▶ σ_2 =Second population standard deviation.
- ▶ n_1 =First sample size.(≥ 30)
- ▶ n_2 =Second sample size.(≥ 30)

z-formula

$$z = \frac{(\bar{x}_1 - \bar{x}_2) - (\mu_1 - \mu_2)}{\sqrt{\frac{\sigma_1^2}{n_1} + \frac{\sigma_2^2}{n_2}}}$$

Where

- ▶ \bar{x}_1 =First sample mean.
- ▶ \bar{x}_2 =Second sample mean.
- ▶ μ_1 =First population mean.
- ▶ μ_2 =Second population mean.
- ▶ σ_1 =First population standard deviation.
- ▶ σ_2 =Second population standard deviation.
- ▶ n_1 =First sample size.(≥ 30)
- ▶ n_2 =Second sample size.(≥ 30)

Confidence Interval

Confidence Interval

Confidence interval

Confidence Interval

Confidence interval

$$(\bar{x}_1 - \bar{x}_2) - z_{\alpha/2} \sqrt{\frac{\sigma_1^2}{n_1} + \frac{\sigma_2^2}{n_2}} \leq (\mu_1 - \mu_2) \leq (\bar{x}_1 - \bar{x}_2) + z_{\alpha/2} \sqrt{\frac{\sigma_1^2}{n_1} + \frac{\sigma_2^2}{n_2}}$$

Confidence Interval

Confidence interval

$$(\bar{x}_1 - \bar{x}_2) - z_{\alpha/2} \sqrt{\frac{\sigma_1^2}{n_1} + \frac{\sigma_2^2}{n_2}} \leq (\mu_1 - \mu_2) \leq (\bar{x}_1 - \bar{x}_2) + z_{\alpha/2} \sqrt{\frac{\sigma_1^2}{n_1} + \frac{\sigma_2^2}{n_2}}$$

Example:

Confidence Interval

Confidence interval

$$(\bar{x}_1 - \bar{x}_2) - z_{\alpha/2} \sqrt{\frac{\sigma_1^2}{n_1} + \frac{\sigma_2^2}{n_2}} \leq (\mu_1 - \mu_2) \leq (\bar{x}_1 - \bar{x}_2) + z_{\alpha/2} \sqrt{\frac{\sigma_1^2}{n_1} + \frac{\sigma_2^2}{n_2}}$$

Example: A men buy 50 electric bulbs of Phillips and 50 electric bulbs of HMT.

Confidence Interval

Confidence interval

$$(\bar{x}_1 - \bar{x}_2) - z_{\alpha/2} \sqrt{\frac{\sigma_1^2}{n_1} + \frac{\sigma_2^2}{n_2}} \leq (\mu_1 - \mu_2) \leq (\bar{x}_1 - \bar{x}_2) + z_{\alpha/2} \sqrt{\frac{\sigma_1^2}{n_1} + \frac{\sigma_2^2}{n_2}}$$

Example: A men buy 50 electric bulbs of Phillips and 50 electric bulbs of HMT. He finds that Phillips bulb give an average life of 1500 hrs with a population standard deviation of 60 hrs

Confidence Interval

Confidence interval

$$(\bar{x}_1 - \bar{x}_2) - z_{\alpha/2} \sqrt{\frac{\sigma_1^2}{n_1} + \frac{\sigma_2^2}{n_2}} \leq (\mu_1 - \mu_2) \leq (\bar{x}_1 - \bar{x}_2) + z_{\alpha/2} \sqrt{\frac{\sigma_1^2}{n_1} + \frac{\sigma_2^2}{n_2}}$$

Example: A man buys 50 electric bulbs of Phillips and 50 electric bulbs of HMT. He finds that Phillips bulbs give an average life of 1500 hrs with a population standard deviation of 60 hrs and HMT bulbs give an average life of 1512 hrs with a standard deviation of 80 hrs.

Confidence Interval

Confidence interval

$$(\bar{x}_1 - \bar{x}_2) - z_{\alpha/2} \sqrt{\frac{\sigma_1^2}{n_1} + \frac{\sigma_2^2}{n_2}} \leq (\mu_1 - \mu_2) \leq (\bar{x}_1 - \bar{x}_2) + z_{\alpha/2} \sqrt{\frac{\sigma_1^2}{n_1} + \frac{\sigma_2^2}{n_2}}$$

Example: A men buy 50 electric bulbs of Phillips and 50 electric bulbs of HMT. He finds that Phillips bulb give an average life of 1500 hrs with a population standard deviation of 60 hrs and HMT bulbs give an average life of 1512 hrs with a standard deviation of 80 hrs. Is there any significant difference between the mean life of two types of bulbs.

Confidence Interval

Confidence interval

$$(\bar{x}_1 - \bar{x}_2) - z_{\alpha/2} \sqrt{\frac{\sigma_1^2}{n_1} + \frac{\sigma_2^2}{n_2}} \leq (\mu_1 - \mu_2) \leq (\bar{x}_1 - \bar{x}_2) + z_{\alpha/2} \sqrt{\frac{\sigma_1^2}{n_1} + \frac{\sigma_2^2}{n_2}}$$

Example: A men buy 50 electric bulbs of Phillips and 50 electric bulbs of HMT. He finds that Phillips bulb give an average life of 1500 hrs with a population standard deviation of 60 hrs and HMT bulbs give an average life of 1512 hrs with a standard deviation of 80 hrs. Is there any significant difference between the mean life of two types of bulbs. (Take $\alpha = 0.05$)

Confidence Interval

Confidence interval

$$(\bar{x}_1 - \bar{x}_2) - z_{\alpha/2} \sqrt{\frac{\sigma_1^2}{n_1} + \frac{\sigma_2^2}{n_2}} \leq (\mu_1 - \mu_2) \leq (\bar{x}_1 - \bar{x}_2) + z_{\alpha/2} \sqrt{\frac{\sigma_1^2}{n_1} + \frac{\sigma_2^2}{n_2}}$$

Example: A men buy 50 electric bulbs of Phillips and 50 electric bulbs of HMT. He finds that Phillips bulb give an average life of 1500 hrs with a population standard deviation of 60 hrs and HMT bulbs give an average life of 1512 hrs with a standard deviation of 80 hrs. Is there any significant difference between the mean life of two types of bulbs. (Take $\alpha = 0.05$) Also establish 95% and 99% confidence limit.

Examples

Examples

Example:

Examples

Example: A simple sample of height of 6,400 Englishmen has a mean of 67.85 inches

Examples

Example: A simple sample of height of 6,400 Englishmen has a mean of 67.85 inches with a population standard deviation of 2.52 inches.

Examples

Example: A simple sample of height of 6,400 Englishmen has a mean of 67.85 inches with a population standard deviation of 2.52 inches. While a simple sample of 1,600 Austrians has a mean height of 68.55 inches

Examples

Example: A simple sample of height of 6,400 Englishmen has a mean of 67.85 inches with a population standard deviation of 2.52 inches. While a simple sample of 1,600 Austrians has a mean height of 68.55 inches with a population standard deviation of 2.52 inches.

Examples

Example: A simple sample of height of 6,400 Englishmen has a mean of 67.85 inches with a population standard deviation of 2.52 inches. While a simple sample of 1,600 Austrians has a mean height of 68.55 inches with a population standard deviation of 2.52 inches. Do this data indicates that Austrians are on average taller than the Englishmen.

Examples

Example: A simple sample of height of 6,400 Englishmen has a mean of 67.85 inches with a population standard deviation of 2.52 inches. While a simple sample of 1,600 Austrians has a mean height of 68.55 inches with a population standard deviation of 2.52 inches. Do this data indicates that Austrians are on average taller than the Englishmen. (Take $\alpha = 0.01$)

Examples

Example: A simple sample of height of 6,400 Englishmen has a mean of 67.85 inches with a population standard deviation of 2.52 inches. While a simple sample of 1,600 Austrians has a mean height of 68.55 inches with a population standard deviation of 2.52 inches. Do this data indicates that Austrians are on average taller than the Englishmen. (Take $\alpha = 0.01$)

Example:

Examples

Example: A simple sample of height of 6,400 Englishmen has a mean of 67.85 inches with a population standard deviation of 2.52 inches. While a simple sample of 1,600 Austrians has a mean height of 68.55 inches with a population standard deviation of 2.52 inches. Do this data indicates that Austrians are on average taller than the Englishmen. (Take $\alpha = 0.01$)

Example: A random sample of 100 mill workers at Kanpur showed their mean wage to be Rs 350

Examples

Example: A simple sample of height of 6,400 Englishmen has a mean of 67.85 inches with a population standard deviation of 2.52 inches. While a simple sample of 1,600 Austrians has a mean height of 68.55 inches with a population standard deviation of 2.52 inches. Do this data indicates that Austrians are on average taller than the Englishmen. (Take $\alpha = 0.01$)

Example: A random sample of 100 mill workers at Kanpur showed their mean wage to be Rs 350 with a population standard deviation of Rs 28.

Examples

Example: A simple sample of height of 6,400 Englishmen has a mean of 67.85 inches with a population standard deviation of 2.52 inches. While a simple sample of 1,600 Austrians has a mean height of 68.55 inches with a population standard deviation of 2.52 inches. Do this data indicates that Austrians are on average taller than the Englishmen. (Take $\alpha = 0.01$)

Example: A random sample of 100 mill workers at Kanpur showed their mean wage to be Rs 350 with a population standard deviation of Rs 28. Another random sample of 150 mill workers in Nagpur showed wage to be Rs 390

Examples

Example: A simple sample of height of 6,400 Englishmen has a mean of 67.85 inches with a population standard deviation of 2.52 inches. While a simple sample of 1,600 Austrians has a mean height of 68.55 inches with a population standard deviation of 2.52 inches. Do this data indicates that Austrians are on average taller than the Englishmen. (Take $\alpha = 0.01$)

Example: A random sample of 100 mill workers at Kanpur showed their mean wage to be Rs 350 with a population standard deviation of Rs 28. Another random sample of 150 mill workers in Nagpur showed wage to be Rs 390 with a population standard deviation of Rs 40.

Examples

Example: A simple sample of height of 6,400 Englishmen has a mean of 67.85 inches with a population standard deviation of 2.52 inches. While a simple sample of 1,600 Austrians has a mean height of 68.55 inches with a population standard deviation of 2.52 inches. Do this data indicates that Austrians are on average taller than the Englishmen. (Take $\alpha = 0.01$)

Example: A random sample of 100 mill workers at Kanpur showed their mean wage to be Rs 350 with a population standard deviation of Rs 28. Another random sample of 150 mill workers in Nagpur showed wage to be Rs 390 with a population standard deviation of Rs 40. Do the mean wages of workers in Nagpur and Kanpur differ significantly?

Examples

Example: A simple sample of height of 6,400 Englishmen has a mean of 67.85 inches with a population standard deviation of 2.52 inches. While a simple sample of 1,600 Austrians has a mean height of 68.55 inches with a population standard deviation of 2.52 inches. Do this data indicates that Austrians are on average taller than the Englishmen. (Take $\alpha = 0.01$)

Example: A random sample of 100 mill workers at Kanpur showed their mean wage to be Rs 350 with a population standard deviation of Rs 28. Another random sample of 150 mill workers in Nagpur showed wage to be Rs 390 with a population standard deviation of Rs 40. Do the mean wages of workers in Nagpur and Kanpur differ significantly? (Take $\alpha = 0.05$)

Examples

Example: A simple sample of height of 6,400 Englishmen has a mean of 67.85 inches with a population standard deviation of 2.52 inches. While a simple sample of 1,600 Austrians has a mean height of 68.55 inches with a population standard deviation of 2.52 inches. Do this data indicates that Austrians are on average taller than the Englishmen. (Take $\alpha = 0.01$)

Example: A random sample of 100 mill workers at Kanpur showed their mean wage to be Rs 350 with a population standard deviation of Rs 28. Another random sample of 150 mill workers in Nagpur showed wage to be Rs 390 with a population standard deviation of Rs 40. Do the mean wages of workers in Nagpur and Kanpur differ significantly? (Take $\alpha = 0.05$)

Method-V Testing a hypothesis for double population mean (small samples) (t-test)

Method-V Testing a hypothesis for double population mean (small samples) (t-test)

One of the following case occur.

Method-V Testing a hypothesis for double population mean (small samples) (t-test)

One of the following cases occur.

- 1 One of the sample sizes is small ($n_1 \leq 30$ or $n_2 \leq 30$)

Method-V Testing a hypothesis for double population mean (small samples) (t-test)

One of the following cases occur.

- 1 One of the sample sizes is small ($n_1 \leq 30$ or $n_2 \leq 30$)
- 2 One of the population standard deviations is not given, (σ_1 is unknown or σ_2 is unknown)

Method-V Testing a hypothesis for double population mean (small samples) (t-test)

One of the following cases occur.

- 1 One of the sample sizes is small ($n_1 \leq 30$ or $n_2 \leq 30$)
- 2 One of the population standard deviations is not given, (σ_1 is unknown or σ_2 is unknown)

t-formula

t-formula

$$t = \frac{(\bar{x}_1 - \bar{x}_2) - (\mu_1 - \mu_2)}{\sqrt{\frac{s_1^2(n_1-1) + s_2^2(n_2-1)}{n_1+n_2-2}} \sqrt{\frac{1}{n_1} + \frac{1}{n_2}}}, d.f. = n_1 + n_2 - 2$$

t-formula

$$t = \frac{(\bar{x}_1 - \bar{x}_2) - (\mu_1 - \mu_2)}{\sqrt{\frac{s_1^2(n_1-1) + s_2^2(n_2-1)}{n_1+n_2-2}} \sqrt{\frac{1}{n_1} + \frac{1}{n_2}}}, d.f = n_1 + n_2 - 2$$

Where

► \bar{x}_1 = First sample mean.

t-formula

$$t = \frac{(\bar{x}_1 - \bar{x}_2) - (\mu_1 - \mu_2)}{\sqrt{\frac{s_1^2(n_1-1) + s_2^2(n_2-1)}{n_1+n_2-2}} \sqrt{\frac{1}{n_1} + \frac{1}{n_2}}}, d.f = n_1 + n_2 - 2$$

Where

- ▶ \bar{x}_1 =First sample mean.
- ▶ \bar{x}_2 =Second sample mean

t-formula

$$t = \frac{(\bar{x}_1 - \bar{x}_2) - (\mu_1 - \mu_2)}{\sqrt{\frac{s_1^2(n_1-1) + s_2^2(n_2-1)}{n_1+n_2-2}} \sqrt{\frac{1}{n_1} + \frac{1}{n_2}}}, d.f = n_1 + n_2 - 2$$

Where

- ▶ \bar{x}_1 =First sample mean.
- ▶ \bar{x}_2 =Second sample mean
- ▶ μ_1 =First population mean.

t-formula

$$t = \frac{(\bar{x}_1 - \bar{x}_2) - (\mu_1 - \mu_2)}{\sqrt{\frac{s_1^2(n_1-1) + s_2^2(n_2-1)}{n_1+n_2-2}} \sqrt{\frac{1}{n_1} + \frac{1}{n_2}}}, d.f = n_1 + n_2 - 2$$

Where

- ▶ \bar{x}_1 =First sample mean.
- ▶ \bar{x}_2 =Second sample mean
- ▶ μ_1 =First population mean.
- ▶ μ_2 =Second population mean.

t-formula

$$t = \frac{(\bar{x}_1 - \bar{x}_2) - (\mu_1 - \mu_2)}{\sqrt{\frac{s_1^2(n_1-1) + s_2^2(n_2-1)}{n_1+n_2-2}} \sqrt{\frac{1}{n_1} + \frac{1}{n_2}}}, d.f = n_1 + n_2 - 2$$

Where

- ▶ \bar{x}_1 =First sample mean.
- ▶ \bar{x}_2 =Second sample mean
- ▶ μ_1 =First population mean.
- ▶ μ_2 =Second population mean.
- ▶ s_1 =First sample standard deviation.

t-formula

$$t = \frac{(\bar{x}_1 - \bar{x}_2) - (\mu_1 - \mu_2)}{\sqrt{\frac{s_1^2(n_1-1) + s_2^2(n_2-1)}{n_1+n_2-2}} \sqrt{\frac{1}{n_1} + \frac{1}{n_2}}}, d.f = n_1 + n_2 - 2$$

Where

- ▶ \bar{x}_1 =First sample mean.
- ▶ \bar{x}_2 =Second sample mean
- ▶ μ_1 =First population mean.
- ▶ μ_2 =Second population mean.
- ▶ s_1 =First sample standard deviation.
- ▶ s_2 =Second sample standard deviation.

t-formula

$$t = \frac{(\bar{x}_1 - \bar{x}_2) - (\mu_1 - \mu_2)}{\sqrt{\frac{s_1^2(n_1-1) + s_2^2(n_2-1)}{n_1+n_2-2}} \sqrt{\frac{1}{n_1} + \frac{1}{n_2}}}, d.f = n_1 + n_2 - 2$$

Where

- ▶ \bar{x}_1 =First sample mean.
- ▶ \bar{x}_2 =Second sample mean
- ▶ μ_1 =First population mean.
- ▶ μ_2 =Second population mean.
- ▶ s_1 =First sample standard deviation.
- ▶ s_2 =Second sample standard deviation.
- ▶ n_1 =First sample size.

t-formula

$$t = \frac{(\bar{x}_1 - \bar{x}_2) - (\mu_1 - \mu_2)}{\sqrt{\frac{s_1^2(n_1-1) + s_2^2(n_2-1)}{n_1+n_2-2}} \sqrt{\frac{1}{n_1} + \frac{1}{n_2}}}, d.f = n_1 + n_2 - 2$$

Where

- ▶ \bar{x}_1 =First sample mean.
- ▶ \bar{x}_2 =Second sample mean
- ▶ μ_1 =First population mean.
- ▶ μ_2 =Second population mean.
- ▶ s_1 =First sample standard deviation.
- ▶ s_2 =Second sample standard deviation.
- ▶ n_1 =First sample size.
- ▶ n_2 =Second sample size.

t-formula

$$t = \frac{(\bar{x}_1 - \bar{x}_2) - (\mu_1 - \mu_2)}{\sqrt{\frac{s_1^2(n_1-1) + s_2^2(n_2-1)}{n_1+n_2-2}} \sqrt{\frac{1}{n_1} + \frac{1}{n_2}}}, d.f = n_1 + n_2 - 2$$

Where

- ▶ \bar{x}_1 =First sample mean.
- ▶ \bar{x}_2 =Second sample mean
- ▶ μ_1 =First population mean.
- ▶ μ_2 =Second population mean.
- ▶ s_1 =First sample standard deviation.
- ▶ s_2 =Second sample standard deviation.
- ▶ n_1 =First sample size.
- ▶ n_2 =Second sample size.

Confidence Interval

Confidence Interval

Confidence Interval

Confidence Interval

Confidence Interval

$$\begin{aligned} (\bar{x}_1 - \bar{x}_2) - t_{\alpha/2} \sqrt{\frac{s_1^2(n_1-1) + s_2^2(n_2-1)}{n_1+n_2-2}} \sqrt{\frac{1}{n_1} + \frac{1}{n_2}} \\ \leq (\mu_1 - \mu_2) \leq \\ (\bar{x}_1 - \bar{x}_2) + t_{\alpha/2} \sqrt{\frac{s_1^2(n_1-1) + s_2^2(n_2-1)}{n_1+n_2-2}} \sqrt{\frac{1}{n_1} + \frac{1}{n_2}} \end{aligned}$$

Confidence Interval

Confidence Interval

$$\begin{aligned} (\bar{x}_1 - \bar{x}_2) - t_{\alpha/2} \sqrt{\frac{s_1^2(n_1-1) + s_2^2(n_2-1)}{n_1+n_2-2}} \sqrt{\frac{1}{n_1} + \frac{1}{n_2}} \\ \leq (\mu_1 - \mu_2) \leq \\ (\bar{x}_1 - \bar{x}_2) + t_{\alpha/2} \sqrt{\frac{s_1^2(n_1-1) + s_2^2(n_2-1)}{n_1+n_2-2}} \sqrt{\frac{1}{n_1} + \frac{1}{n_2}} \end{aligned}$$

Example

Example

Example:

Example

Example: A company is interested in finding out if there is any difference in average salary received by managers of two divisions.

Example

Example: A company is interested in finding out if there is any difference in average salary received by managers of two divisions. Accordingly samples of 12 managers in the first division and 10 managers in the second were selected at random. The result are given below :

Example

Example: A company is interested in finding out if there is any difference in average salary received by managers of two divisions. Accordingly samples of 12 managers in the first division and 10 managers in the second were selected at random. The result are given below :

	I division	II division
Sample size	12	10
Average Monthly Salary	12500	11200
Standard Deviation	320	480

Example

Example: A company is interested in finding out if there is any difference in average salary received by managers of two divisions. Accordingly samples of 12 managers in the first division and 10 managers in the second were selected at random. The result are given below :

	I division	II division
Sample size	12	10
Average Monthly Salary	12500	11200
Standard Deviation	320	480

Is there any significant difference between salary received by managers of two types of division?

Example

Example: A company is interested in finding out if there is any difference in average salary received by managers of two divisions. Accordingly samples of 12 managers in the first division and 10 managers in the second were selected at random. The result are given below :

	I division	II division
Sample size	12	10
Average Monthly Salary	12500	11200
Standard Deviation	320	480

Is there any significant difference between salary received by managers of two types of division?

Example

Example

Example:

Example

Example: Two laboratories A and B carried out independent estimates of fat content in ice-creme made by a firm.

Example

Example: Two laboratories A and B carried out independent estimates of fat content in ice-cream made by a firm. A sample is taken from each batch. The fat content obtained by laboratories is recorded below:

Example

Example: Two laboratories A and B carried out independent estimates of fat content in ice-cream made by a firm. A sample is taken from each batch. The fat content obtained by laboratories is recorded below:

Batch No	1	2	3	4	5	6	7	8	9	10
Lab A	7	8	7	3	8	6	9	4	7	8
Lab B	9	8	8	4	7	7	9	6	6	6

Example

Example: Two laboratories A and B carried out independent estimates of fat content in ice-creme made by a firm. A sample is taken from each batch. The fat content obtained by laboratories is recorded below:

Batch No	1	2	3	4	5	6	7	8	9	10
Lab A	7	8	7	3	8	6	9	4	7	8
Lab B	9	8	8	4	7	7	9	6	6	6

Is there any significant difference between the mean fat content obtained by two laboratories.

Example

Example: Two laboratories A and B carried out independent estimates of fat content in ice-creme made by a firm. A sample is taken from each batch. The fat content obtained by laboratories is recorded below:

Batch No	1	2	3	4	5	6	7	8	9	10
Lab A	7	8	7	3	8	6	9	4	7	8
Lab B	9	8	8	4	7	7	9	6	6	6

Is there any significant difference between the mean fat content obtained by two laboratories. (Take $\alpha = 0.01$)

Example

Example: Two laboratories A and B carried out independent estimates of fat content in ice-cream made by a firm. A sample is taken from each batch. The fat content obtained by laboratories is recorded below:

Batch No	1	2	3	4	5	6	7	8	9	10
Lab A	7	8	7	3	8	6	9	4	7	8
Lab B	9	8	8	4	7	7	9	6	6	6

Is there any significant difference between the mean fat content obtained by two laboratories. (Take $\alpha = 0.01$) Also establish 90% confidence limit.

Method-VI Testing a hypothesis for dependent population(Difference-test)(t-test)

Method-VI Testing a hypothesis for dependent population(Difference-test)(t-test)

Condition

Method-VI Testing a hypothesis for dependent population(Difference-test)(t-test)

Condition

Method-VI Testing a hypothesis for dependent population(Difference-test)(t-test)

Condition

- ▶ Small sample sizes.

Method-VI Testing a hypothesis for dependent population(Difference-test)(t-test)

Condition

- ▶ Small sample sizes.
- ▶ Data in pairs.

Method-VI Testing a hypothesis for dependent population(Difference-test)(t-test)

Condition

- ▶ Small sample sizes.
- ▶ Data in pairs.

Method-VI Testing a hypothesis for dependent population(Difference-test)(t-test)

Condition

- ▶ Small sample sizes.
- ▶ Data in pairs.

Types of hypothesis

Method-VI Testing a hypothesis for dependent population(Difference-test)(t-test)

Condition

- ▶ Small sample sizes.
- ▶ Data in pairs.

Types of hypothesis

Method-VI Testing a hypothesis for dependent population(Difference-test)(t-test)

Condition

- ▶ Small sample sizes.
- ▶ Data in pairs.

Types of hypothesis

- ▶ $H_o : D = 0, H_a : D \neq 0$

Method-VI Testing a hypothesis for dependent population(Difference-test)(t-test)

Condition

- ▶ Small sample sizes.
- ▶ Data in pairs.

Types of hypothesis

- ▶ $H_o : D = 0, H_a : D \neq 0$
- ▶ $H_o : D \geq 0, H_a : D < 0$

Method-VI Testing a hypothesis for dependent population(Difference-test)(t-test)

Condition

- ▶ Small sample sizes.
- ▶ Data in pairs.

Types of hypothesis

- ▶ $H_o : D = 0, H_a : D \neq 0$
- ▶ $H_o : D \geq 0, H_a : D < 0$
- ▶ $H_o : D \leq 0, H_a : D > 0$

Method-VI Testing a hypothesis for dependent population(Difference-test)(t-test)

Condition

- ▶ Small sample sizes.
- ▶ Data in pairs.

Types of hypothesis

- ▶ $H_o : D = 0, H_a : D \neq 0$
- ▶ $H_o : D \geq 0, H_a : D < 0$
- ▶ $H_o : D \leq 0, H_a : D > 0$

Mid-Sem Mark(Statistical Methods) Before Assignment

Mid-Sem Mark(Statistical Methods)
Before Assignment

18

Mid-Sem Mark(Statistical Methods)
Before Assignment

18

7

Mid-Sem Mark(Statistical Methods)
Before Assignment

18

7

10

Mid-Sem Mark(Statistical Methods)
Before Assignment

18

7

10

22

Mid-Sem Mark(Statistical Methods)
Before Assignment

18

7

10

22

15

Mid-Sem Mark(Statistical Methods)
Before Assignment

18

7

10

22

15

0

Mid-Sem Mark(Statistical Methods)
Before Assignment

18

7

10

22

15

0

4

Mid-Sem Mark(Statistical Methods)
Before Assignment

18

7

10

22

15

0

4

25

Mid-Sem Mark(Statistical Methods)
Before Assignment

18

7

10

22

15

0

4

25

17

Mid-Sem Mark(Statistical Methods)
Before Assignment

18

7

10

22

15

0

4

25

17

5

Mid-Sem Mark(Statistical Methods)
Before Assignment After Assignments

18

7

10

22

15

0

4

25

17

5

Mid-Sem Mark(Statistical Methods)

Before Assignment	After Assignments
--------------------------	--------------------------

18	
----	--

	22
--	----

7	
---	--

10	
----	--

22	
----	--

15	
----	--

0	
---	--

4	
---	--

25	
----	--

17	
----	--

5	
---	--

Mid-Sem Mark(Statistical Methods)

Before Assignment	After Assignments
--------------------------	--------------------------

18	
----	--

	22
--	----

7	
---	--

	15
--	----

10	
----	--

22	
----	--

15	
----	--

0	
---	--

4	
---	--

25	
----	--

17	
----	--

5	
---	--

Mid-Sem Mark(Statistical Methods)

Before Assignment	After Assignments
--------------------------	--------------------------

18	
----	--

	22
--	----

7	
---	--

	15
--	----

10	
----	--

	12
--	----

22	
----	--

15	
----	--

0	
---	--

4	
---	--

25	
----	--

17	
----	--

5	
---	--

Mid-Sem Mark(Statistical Methods)

Before Assignment	After Assignments
--------------------------	--------------------------

18	
----	--

	22
--	----

7	
---	--

	15
--	----

10	
----	--

	12
--	----

22	
----	--

	28
--	----

15	
----	--

0	
---	--

4	
---	--

25	
----	--

17	
----	--

5	
---	--

Mid-Sem Mark(Statistical Methods)

Before Assignment	After Assignments
--------------------------	--------------------------

18	
----	--

	22
--	----

7	
---	--

	15
--	----

10	
----	--

	12
--	----

22	
----	--

	28
--	----

15	
----	--

	12
--	----

0	
---	--

4	
---	--

25	
----	--

17	
----	--

5	
---	--

Mid-Sem Mark(Statistical Methods)

Before Assignment	After Assignments
--------------------------	--------------------------

18	
----	--

	22
--	----

7	
---	--

	15
--	----

10	
----	--

	12
--	----

22	
----	--

	28
--	----

15	
----	--

	12
--	----

0	
---	--

	0
--	---

4	
---	--

25	
----	--

17	
----	--

5	
---	--

Mid-Sem Mark(Statistical Methods)

Before Assignment	After Assignments
--------------------------	--------------------------

18	
----	--

	22
--	----

7	
---	--

	15
--	----

10	
----	--

	12
--	----

22	
----	--

	28
--	----

15	
----	--

	12
--	----

0	
---	--

	0
--	---

4	
---	--

	2
--	---

25	
----	--

17	
----	--

5	
---	--

Mid-Sem Mark(Statistical Methods)

Before Assignment	After Assignments
--------------------------	--------------------------

18	
----	--

	22
--	----

7	
---	--

	15
--	----

10	
----	--

	12
--	----

22	
----	--

	28
--	----

15	
----	--

	12
--	----

0	
---	--

	0
--	---

4	
---	--

	2
--	---

25	
----	--

	14
--	----

17	
----	--

5	
---	--

Mid-Sem Mark(Statistical Methods)

Before Assignment	After Assignments
--------------------------	--------------------------

18	22
7	15
10	12
22	28
15	12
0	0
4	2
25	14
17	4
5	

Mid-Sem Mark(Statistical Methods)

Before Assignment	After Assignments
--------------------------	--------------------------

18	22
7	15
10	12
22	28
15	12
0	0
4	2
25	14
17	4
5	5

Mid-Sem Mark(Statistical Methods)

Before Assignment	After Assignments	d
18	22	
7	15	
10	12	
22	28	
15	12	
0	0	
4	2	
25	14	
17	4	
5	5	

Mid-Sem Mark(Statistical Methods)

Before Assignment	After Assignments	d
18	22	4
7	15	
10	12	
22	28	
15	12	
0	0	
4	2	
25	14	
17	4	
5	5	

Mid-Sem Mark(Statistical Methods)		
Before Assignment	After Assignments	d
18	22	4
7	15	8
10	12	
22	28	
15	12	
0	0	
4	2	
25	14	
17	4	
5	5	

Mid-Sem Mark(Statistical Methods)

Before Assignment	After Assignments	d
18	22	4
7	15	8
10	12	2
22	28	
15	12	
0	0	
4	2	
25	14	
17	4	
5	5	

Mid-Sem Mark(Statistical Methods)

Before Assignment	After Assignments	d
18	22	4
7	15	8
10	12	2
22	28	6
15	12	
0	0	
4	2	
25	14	
17	4	
5	5	

Mid-Sem Mark(Statistical Methods)

Before Assignment	After Assignments	d
18	22	4
7	15	8
10	12	2
22	28	6
15	12	-3
0	0	
4	2	
25	14	
17	4	
5	5	

Mid-Sem Mark(Statistical Methods)

Before Assignment	After Assignments	d
18	22	4
7	15	8
10	12	2
22	28	6
15	12	-3
0	0	0
4	2	
25	14	
17	4	
5	5	

Mid-Sem Mark(Statistical Methods)

Before Assignment	After Assignments	d
18	22	4
7	15	8
10	12	2
22	28	6
15	12	-3
0	0	0
4	2	-2
25	14	
17	4	
5	5	

Mid-Sem Mark(Statistical Methods)

Before Assignment	After Assignments	d
18	22	4
7	15	8
10	12	2
22	28	6
15	12	-3
0	0	0
4	2	-2
25	14	-11
17	4	
5	5	

Mid-Sem Mark(Statistical Methods)

Before Assignment	After Assignments	d
18	22	4
7	15	8
10	12	2
22	28	6
15	12	-3
0	0	0
4	2	-2
25	14	-11
17	4	-13
5	5	

Mid-Sem Mark(Statistical Methods)

Before Assignment	After Assignments	d
18	22	4
7	15	8
10	12	2
22	28	6
15	12	-3
0	0	0
4	2	-2
25	14	-11
17	4	-13
5	5	0

t-formula

t-formula

$$t = \frac{\bar{d} - D}{\frac{s_d}{\sqrt{n}}}, d.f = n - 1$$

t-formula

$$t = \frac{\bar{d} - D}{\frac{s_d}{\sqrt{n}}}, d.f = n - 1$$

Where

t-formula

$$t = \frac{\bar{d} - D}{\frac{s_d}{\sqrt{n}}}, d.f = n - 1$$

Where

t-formula

$$t = \frac{\bar{d} - D}{\frac{s_d}{\sqrt{n}}}, d.f = n - 1$$

Where

► \bar{d} = Sample mean of difference. = $\frac{\sum d}{n}$

t-formula

$$t = \frac{\bar{d} - D}{\frac{s_d}{\sqrt{n}}}, d.f = n - 1$$

Where

- ▶ \bar{d} =Sample mean of difference. $= \frac{\sum d}{n}$
- ▶ D =Population mean of difference.

t-formula

$$t = \frac{\bar{d} - D}{\frac{s_d}{\sqrt{n}}}, d.f = n - 1$$

Where

- ▶ \bar{d} =Sample mean of difference. $=\frac{\sum d}{n}$
- ▶ D =Population mean of difference.
- ▶ s_d =Sample standard deviation of difference. $=\sqrt{\frac{\sum (d-\bar{d})^2}{n-1}}$

t-formula

$$t = \frac{\bar{d} - D}{\frac{s_d}{\sqrt{n}}}, d.f = n - 1$$

Where

- ▶ \bar{d} =Sample mean of difference. $=\frac{\sum d}{n}$
- ▶ D =Population mean of difference.
- ▶ s_d =Sample standard deviation of difference. $=\sqrt{\frac{\sum (d-\bar{d})^2}{n-1}}$
- ▶ n =Sample size(no of pairs)

t-formula

$$t = \frac{\bar{d} - D}{\frac{s_d}{\sqrt{n}}}, d.f = n - 1$$

Where

- ▶ \bar{d} =Sample mean of difference. $=\frac{\sum d}{n}$
- ▶ D =Population mean of difference.
- ▶ s_d =Sample standard deviation of difference. $=\sqrt{\frac{\sum (d-\bar{d})^2}{n-1}}$
- ▶ n =Sample size(no of pairs)

t-formula

$$t = \frac{\bar{d} - D}{\frac{s_d}{\sqrt{n}}}, d.f = n - 1$$

Where

- ▶ \bar{d} =Sample mean of difference. $=\frac{\sum d}{n}$
- ▶ D =Population mean of difference.
- ▶ s_d =Sample standard deviation of difference. $=\sqrt{\frac{\sum (d-\bar{d})^2}{n-1}}$
- ▶ n =Sample size(no of pairs)

Confidence Interval

Confidence Interval

Confidence interval

Confidence Interval

Confidence interval

$$\bar{d} - t_{\alpha/2} \frac{s_d}{\sqrt{n}} \leq D \leq \bar{d} + t_{\alpha/2} \frac{s_d}{\sqrt{n}}$$

Confidence Interval

Confidence interval

$$\bar{d} - t_{\alpha/2} \frac{s_d}{\sqrt{n}} \leq D \leq \bar{d} + t_{\alpha/2} \frac{s_d}{\sqrt{n}}$$

Example:

Confidence Interval

Confidence interval

$$\bar{d} - t_{\alpha/2} \frac{s_d}{\sqrt{n}} \leq D \leq \bar{d} + t_{\alpha/2} \frac{s_d}{\sqrt{n}}$$

Example: To verify whether the course in accounting improved performance.

Confidence Interval

Confidence interval

$$\bar{d} - t_{\alpha/2} \frac{s_d}{\sqrt{n}} \leq D \leq \bar{d} + t_{\alpha/2} \frac{s_d}{\sqrt{n}}$$

Example: To verify whether the course in accounting improved performance. A similar test was given to 12 participants both before and after the course.

Confidence Interval

Confidence interval

$$\bar{d} - t_{\alpha/2} \frac{s_d}{\sqrt{n}} \leq D \leq \bar{d} + t_{\alpha/2} \frac{s_d}{\sqrt{n}}$$

Example: To verify whether the course in accounting improved performance. A similar test was given to 12 participants both before and after the course. The original mark recorded in alphabetical order of participants were

Confidence Interval

Confidence interval

$$\bar{d} - t_{\alpha/2} \frac{s_d}{\sqrt{n}} \leq D \leq \bar{d} + t_{\alpha/2} \frac{s_d}{\sqrt{n}}$$

Example: To verify whether the course in accounting improved performance. A similar test was given to 12 participants both before and after the course. The original mark recorded in alphabetical order of participants were 44,

Confidence Interval

Confidence interval

$$\bar{d} - t_{\alpha/2} \frac{s_d}{\sqrt{n}} \leq D \leq \bar{d} + t_{\alpha/2} \frac{s_d}{\sqrt{n}}$$

Example: To verify whether the course in accounting improved performance. A similar test was given to 12 participants both before and after the course. The original mark recorded in alphabetical order of participants were 44, 40,

Confidence Interval

Confidence interval

$$\bar{d} - t_{\alpha/2} \frac{s_d}{\sqrt{n}} \leq D \leq \bar{d} + t_{\alpha/2} \frac{s_d}{\sqrt{n}}$$

Example: To verify whether the course in accounting improved performance. A similar test was given to 12 participants both before and after the course. The original mark recorded in alphabetical order of participants were 44, 40, 61,

Confidence Interval

Confidence interval

$$\bar{d} - t_{\alpha/2} \frac{s_d}{\sqrt{n}} \leq D \leq \bar{d} + t_{\alpha/2} \frac{s_d}{\sqrt{n}}$$

Example: To verify whether the course in accounting improved performance. A similar test was given to 12 participants both before and after the course. The original mark recorded in alphabetical order of participants were 44, 40, 61, 52,

Confidence Interval

Confidence interval

$$\bar{d} - t_{\alpha/2} \frac{s_d}{\sqrt{n}} \leq D \leq \bar{d} + t_{\alpha/2} \frac{s_d}{\sqrt{n}}$$

Example: To verify whether the course in accounting improved performance. A similar test was given to 12 participants both before and after the course. The original mark recorded in alphabetical order of participants were 44, 40, 61, 52, 32,

Confidence Interval

Confidence interval

$$\bar{d} - t_{\alpha/2} \frac{s_d}{\sqrt{n}} \leq D \leq \bar{d} + t_{\alpha/2} \frac{s_d}{\sqrt{n}}$$

Example: To verify whether the course in accounting improved performance. A similar test was given to 12 participants both before and after the course. The original mark recorded in alphabetical order of participants were 44, 40, 61, 52, 32, 44,

Confidence Interval

Confidence interval

$$\bar{d} - t_{\alpha/2} \frac{s_d}{\sqrt{n}} \leq D \leq \bar{d} + t_{\alpha/2} \frac{s_d}{\sqrt{n}}$$

Example: To verify whether the course in accounting improved performance. A similar test was given to 12 participants both before and after the course. The original mark recorded in alphabetical order of participants were 44, 40, 61, 52, 32, 44, 70,

Confidence Interval

Confidence interval

$$\bar{d} - t_{\alpha/2} \frac{s_d}{\sqrt{n}} \leq D \leq \bar{d} + t_{\alpha/2} \frac{s_d}{\sqrt{n}}$$

Example: To verify whether the course in accounting improved performance. A similar test was given to 12 participants both before and after the course. The original mark recorded in alphabetical order of participants were 44, 40, 61, 52, 32, 44, 70, 41,

Confidence Interval

Confidence interval

$$\bar{d} - t_{\alpha/2} \frac{s_d}{\sqrt{n}} \leq D \leq \bar{d} + t_{\alpha/2} \frac{s_d}{\sqrt{n}}$$

Example: To verify whether the course in accounting improved performance. A similar test was given to 12 participants both before and after the course. The original mark recorded in alphabetical order of participants were 44, 40, 61, 52, 32, 44, 70, 41, 67,

Confidence Interval

Confidence interval

$$\bar{d} - t_{\alpha/2} \frac{s_d}{\sqrt{n}} \leq D \leq \bar{d} + t_{\alpha/2} \frac{s_d}{\sqrt{n}}$$

Example: To verify whether the course in accounting improved performance. A similar test was given to 12 participants both before and after the course. The original mark recorded in alphabetical order of participants were 44, 40, 61, 52, 32, 44, 70, 41, 67, 72,

Confidence Interval

Confidence interval

$$\bar{d} - t_{\alpha/2} \frac{s_d}{\sqrt{n}} \leq D \leq \bar{d} + t_{\alpha/2} \frac{s_d}{\sqrt{n}}$$

Example: To verify whether the course in accounting improved performance. A similar test was given to 12 participants both before and after the course. The original mark recorded in alphabetical order of participants were 44, 40, 61, 52, 32, 44, 70, 41, 67, 72, 53 and

Confidence Interval

Confidence interval

$$\bar{d} - t_{\alpha/2} \frac{s_d}{\sqrt{n}} \leq D \leq \bar{d} + t_{\alpha/2} \frac{s_d}{\sqrt{n}}$$

Example: To verify whether the course in accounting improved performance. A similar test was given to 12 participants both before and after the course. The original mark recorded in alphabetical order of participants were 44, 40, 61, 52, 32, 44, 70, 41, 67, 72, 53 and 72.

Confidence Interval

Confidence interval

$$\bar{d} - t_{\alpha/2} \frac{s_d}{\sqrt{n}} \leq D \leq \bar{d} + t_{\alpha/2} \frac{s_d}{\sqrt{n}}$$

Example: To verify whether the course in accounting improved performance. A similar test was given to 12 participants both before and after the course. The original mark recorded in alphabetical order of participants were 44, 40, 61, 52, 32, 44, 70, 41, 67, 72, 53 and 72. The marks obtained after the course in the same order were

Confidence Interval

Confidence interval

$$\bar{d} - t_{\alpha/2} \frac{s_d}{\sqrt{n}} \leq D \leq \bar{d} + t_{\alpha/2} \frac{s_d}{\sqrt{n}}$$

Example: To verify whether the course in accounting improved performance. A similar test was given to 12 participants both before and after the course. The original mark recorded in alphabetical order of participants were 44, 40, 61, 52, 32, 44, 70, 41, 67, 72, 53 and 72. The marks obtained after the course in the same order were 53,

Confidence Interval

Confidence interval

$$\bar{d} - t_{\alpha/2} \frac{s_d}{\sqrt{n}} \leq D \leq \bar{d} + t_{\alpha/2} \frac{s_d}{\sqrt{n}}$$

Example: To verify whether the course in accounting improved performance. A similar test was given to 12 participants both before and after the course. The original mark recorded in alphabetical order of participants were 44, 40, 61, 52, 32, 44, 70, 41, 67, 72, 53 and 72. The marks obtained after the course in the same order were 53, 38,

Confidence Interval

Confidence interval

$$\bar{d} - t_{\alpha/2} \frac{s_d}{\sqrt{n}} \leq D \leq \bar{d} + t_{\alpha/2} \frac{s_d}{\sqrt{n}}$$

Example: To verify whether the course in accounting improved performance. A similar test was given to 12 participants both before and after the course. The original mark recorded in alphabetical order of participants were 44, 40, 61, 52, 32, 44, 70, 41, 67, 72, 53 and 72. The marks obtained after the course in the same order were 53, 38, 69,

Confidence Interval

Confidence interval

$$\bar{d} - t_{\alpha/2} \frac{s_d}{\sqrt{n}} \leq D \leq \bar{d} + t_{\alpha/2} \frac{s_d}{\sqrt{n}}$$

Example: To verify whether the course in accounting improved performance. A similar test was given to 12 participants both before and after the course. The original mark recorded in alphabetical order of participants were 44, 40, 61, 52, 32, 44, 70, 41, 67, 72, 53 and 72. The marks obtained after the course in the same order were 53, 38, 69, 57,

Confidence Interval

Confidence interval

$$\bar{d} - t_{\alpha/2} \frac{s_d}{\sqrt{n}} \leq D \leq \bar{d} + t_{\alpha/2} \frac{s_d}{\sqrt{n}}$$

Example: To verify whether the course in accounting improved performance. A similar test was given to 12 participants both before and after the course. The original mark recorded in alphabetical order of participants were 44, 40, 61, 52, 32, 44, 70, 41, 67, 72, 53 and 72. The marks obtained after the course in the same order were 53, 38, 69, 57, 46,

Confidence Interval

Confidence interval

$$\bar{d} - t_{\alpha/2} \frac{s_d}{\sqrt{n}} \leq D \leq \bar{d} + t_{\alpha/2} \frac{s_d}{\sqrt{n}}$$

Example: To verify whether the course in accounting improved performance. A similar test was given to 12 participants both before and after the course. The original mark recorded in alphabetical order of participants were 44, 40, 61, 52, 32, 44, 70, 41, 67, 72, 53 and 72. The marks obtained after the course in the same order were 53, 38, 69, 57, 46, 39,

Confidence Interval

Confidence interval

$$\bar{d} - t_{\alpha/2} \frac{s_d}{\sqrt{n}} \leq D \leq \bar{d} + t_{\alpha/2} \frac{s_d}{\sqrt{n}}$$

Example: To verify whether the course in accounting improved performance. A similar test was given to 12 participants both before and after the course. The original mark recorded in alphabetical order of participants were 44, 40, 61, 52, 32, 44, 70, 41, 67, 72, 53 and 72. The marks obtained after the course in the same order were 53, 38, 69, 57, 46, 39, 73,

Confidence Interval

Confidence interval

$$\bar{d} - t_{\alpha/2} \frac{s_d}{\sqrt{n}} \leq D \leq \bar{d} + t_{\alpha/2} \frac{s_d}{\sqrt{n}}$$

Example: To verify whether the course in accounting improved performance. A similar test was given to 12 participants both before and after the course. The original mark recorded in alphabetical order of participants were 44, 40, 61, 52, 32, 44, 70, 41, 67, 72, 53 and 72. The marks obtained after the course in the same order were 53, 38, 69, 57, 46, 39, 73, 48,

Confidence Interval

Confidence interval

$$\bar{d} - t_{\alpha/2} \frac{s_d}{\sqrt{n}} \leq D \leq \bar{d} + t_{\alpha/2} \frac{s_d}{\sqrt{n}}$$

Example: To verify whether the course in accounting improved performance. A similar test was given to 12 participants both before and after the course. The original mark recorded in alphabetical order of participants were 44, 40, 61, 52, 32, 44, 70, 41, 67, 72, 53 and 72. The marks obtained after the course in the same order were 53, 38, 69, 57, 46, 39, 73, 48, 73,

Confidence Interval

Confidence interval

$$\bar{d} - t_{\alpha/2} \frac{s_d}{\sqrt{n}} \leq D \leq \bar{d} + t_{\alpha/2} \frac{s_d}{\sqrt{n}}$$

Example: To verify whether the course in accounting improved performance. A similar test was given to 12 participants both before and after the course. The original mark recorded in alphabetical order of participants were 44, 40, 61, 52, 32, 44, 70, 41, 67, 72, 53 and 72. The marks obtained after the course in the same order were 53, 38, 69, 57, 46, 39, 73, 48, 73, 74,

Confidence Interval

Confidence interval

$$\bar{d} - t_{\alpha/2} \frac{s_d}{\sqrt{n}} \leq D \leq \bar{d} + t_{\alpha/2} \frac{s_d}{\sqrt{n}}$$

Example: To verify whether the course in accounting improved performance. A similar test was given to 12 participants both before and after the course. The original mark recorded in alphabetical order of participants were 44, 40, 61, 52, 32, 44, 70, 41, 67, 72, 53 and 72. The marks obtained after the course in the same order were 53, 38, 69, 57, 46, 39, 73, 48, 73, 74, 60 and

Confidence Interval

Confidence interval

$$\bar{d} - t_{\alpha/2} \frac{s_d}{\sqrt{n}} \leq D \leq \bar{d} + t_{\alpha/2} \frac{s_d}{\sqrt{n}}$$

Example: To verify whether the course in accounting improved performance. A similar test was given to 12 participants both before and after the course. The original mark recorded in alphabetical order of participants were 44, 40, 61, 52, 32, 44, 70, 41, 67, 72, 53 and 72. The marks obtained after the course in the same order were 53, 38, 69, 57, 46, 39, 73, 48, 73, 74, 60 and 78.

Confidence Interval

Confidence interval

$$\bar{d} - t_{\alpha/2} \frac{s_d}{\sqrt{n}} \leq D \leq \bar{d} + t_{\alpha/2} \frac{s_d}{\sqrt{n}}$$

Example: To verify whether the course in accounting improved performance. A similar test was given to 12 participants both before and after the course. The original mark recorded in alphabetical order of participants were 44, 40, 61, 52, 32, 44, 70, 41, 67, 72, 53 and 72. The marks obtained after the course in the same order were 53, 38, 69, 57, 46, 39, 73, 48, 73, 74, 60 and 78. Determine whether the course is useful? (Take $\alpha = 0.05$).

Confidence Interval

Confidence interval

$$\bar{d} - t_{\alpha/2} \frac{s_d}{\sqrt{n}} \leq D \leq \bar{d} + t_{\alpha/2} \frac{s_d}{\sqrt{n}}$$

Example: To verify whether the course in accounting improved performance. A similar test was given to 12 participants both before and after the course. The original mark recorded in alphabetical order of participants were 44, 40, 61, 52, 32, 44, 70, 41, 67, 72, 53 and 72. The marks obtained after the course in the same order were 53, 38, 69, 57, 46, 39, 73, 48, 73, 74, 60 and 78. Determine whether the course is useful? (Take $\alpha = 0.05$).

Examples

Examples

Example:

Examples

Example: A drug was given to 10 patients and increments in their blood pressure were recorded to be

Examples

Example: A drug was given to 10 patients and increments in their blood pressure were recorded to be 3,

Examples

Example: A drug was given to 10 patients and increments in their blood pressure were recorded to be 3, 6,

Examples

Example: A drug was given to 10 patients and increments in their blood pressure were recorded to be 3, 6, -2,

Examples

Example: A drug was given to 10 patients and increments in their blood pressure were recorded to be 3, 6, -2, 4,

Examples

Example: A drug was given to 10 patients and increments in their blood pressure were recorded to be 3, 6, -2, 4, -3,

Examples

Example: A drug was given to 10 patients and increments in their blood pressure were recorded to be 3, 6, -2, 4, -3, 4,

Examples

Example: A drug was given to 10 patients and increments in their blood pressure were recorded to be 3, 6, -2, 4, -3, 4, 6,

Examples

Example: A drug was given to 10 patients and increments in their blood pressure were recorded to be 3, 6, -2, 4, -3, 4, 6, 0,

Examples

Example: A drug was given to 10 patients and increments in their blood pressure were recorded to be 3, 6, -2, 4, -3, 4, 6, 0, 0,

Examples

Example: A drug was given to 10 patients and increments in their blood pressure were recorded to be 3, 6, -2, 4, -3, 4, 6, 0, 0, 2.

Examples

Example: A drug was given to 10 patients and increments in their blood pressure were recorded to be 3, 6, -2, 4, -3, 4, 6, 0, 0, 2. Is it reasonable to believe that the drug has no effect on the change of blood pressure?

Examples

Example: A drug was given to 10 patients and increments in their blood pressure were recorded to be 3, 6, -2, 4, -3, 4, 6, 0, 0, 2. Is it reasonable to believe that the drug has no effect on the change of blood pressure? (Take $\alpha = 0.01$).

Examples

Example: A drug was given to 10 patients and increments in their blood pressure were recorded to be 3, 6, -2, 4, -3, 4, 6, 0, 0, 2. Is it reasonable to believe that the drug has no effect on the change of blood pressure? (Take $\alpha = 0.01$). Also establish 99% confidence limit.

Examples

Example: A drug was given to 10 patients and increments in their blood pressure were recorded to be 3, 6, -2, 4, -3, 4, 6, 0, 0, 2. Is it reasonable to believe that the drug has no effect on the change of blood pressure? (Take $\alpha = 0.01$). Also establish 99% confidence limit.

Example:

Examples

Example: A drug was given to 10 patients and increments in their blood pressure were recorded to be 3, 6, -2, 4, -3, 4, 6, 0, 0, 2. Is it reasonable to believe that the drug has no effect on the change of blood pressure? (Take $\alpha = 0.01$). Also establish 99% confidence limit.

Example: The sales of data of an item in six shops before and after a special promotional campaign are as under:

Examples

Example: A drug was given to 10 patients and increments in their blood pressure were recorded to be 3, 6, -2, 4, -3, 4, 6, 0, 0, 2. Is it reasonable to believe that the drug has no effect on the change of blood pressure? (Take $\alpha = 0.01$). Also establish 99% confidence limit.

Example: The sales of data of an item in six shops before and after a special promotional campaign are as under:

Shops	A	B	C	D	E	F
Before campaign	53	28	31	48	50	42
After campaign	58	29	30	55	56	45

Examples

Example: A drug was given to 10 patients and increments in their blood pressure were recorded to be 3, 6, -2, 4, -3, 4, 6, 0, 0, 2. Is it reasonable to believe that the drug has no effect on the change of blood pressure? (Take $\alpha = 0.01$). Also establish 99% confidence limit.

Example: The sales of data of an item in six shops before and after a special promotional campaign are as under:

Shops	A	B	C	D	E	F
Before campaign	53	28	31	48	50	42
After campaign	58	29	30	55	56	45

Did the campaign make any significant difference in sales at 5% Significant level?

Method-VII Testing a hypothesis for double proportion(z-test)

Method-VII Testing a hypothesis for double proportion(z-test)

Necessary condition

Method-VII Testing a hypothesis for double proportion(z-test)

Necessary condition

Method-VII Testing a hypothesis for double proportion(z-test)

Necessary condition

► $n_1 \cdot p_1 \geq 5$

Method-VII Testing a hypothesis for double proportion(z-test)

Necessary condition

- ▶ $n_1 \cdot p_1 \geq 5$
- ▶ $n_1 \cdot (1 - p_1) \geq 5$

Method-VII Testing a hypothesis for double proportion(z-test)

Necessary condition

- ▶ $n_1.p_1 \geq 5$
- ▶ $n_1.(1 - p_1) \geq 5$
- ▶ $n_2.p_2 \geq 5$

Method-VII Testing a hypothesis for double proportion(z-test)

Necessary condition

- ▶ $n_1.p_1 \geq 5$
- ▶ $n_1.(1 - p_1) \geq 5$
- ▶ $n_2.p_2 \geq 5$
- ▶ $n_2.(1 - p_2) \geq 5$

Method-VII Testing a hypothesis for double proportion(z-test)

Necessary condition

- ▶ $n_1.p_1 \geq 5$
- ▶ $n_1.(1 - p_1) \geq 5$
- ▶ $n_2.p_2 \geq 5$
- ▶ $n_2.(1 - p_2) \geq 5$

Method-VII Testing a hypothesis for double proportion(z-test)

Necessary condition

- ▶ $n_1.p_1 \geq 5$
- ▶ $n_1.(1 - p_1) \geq 5$
- ▶ $n_2.p_2 \geq 5$
- ▶ $n_2.(1 - p_2) \geq 5$

z-formula

z-formula

$$z = \frac{(\bar{p}_1 - \bar{p}_2) - (p_1 - p_2)}{\sqrt{\bar{p} \cdot \bar{q} \left(\frac{1}{n_1} + \frac{1}{n_2} \right)}}$$

z-formula

$$z = \frac{(\bar{p}_1 - \bar{p}_2) - (p_1 - p_2)}{\sqrt{\bar{p} \cdot \bar{q} \left(\frac{1}{n_1} + \frac{1}{n_2} \right)}}$$

$$\bar{p} = \frac{n_1 \cdot \bar{p}_1 + n_2 \cdot \bar{p}_2}{n_1 + n_2} = \frac{x_1 + x_2}{n_1 + n_2}$$

z-formula

$$z = \frac{(\bar{p}_1 - \bar{p}_2) - (p_1 - p_2)}{\sqrt{\bar{p} \cdot \bar{q} \left(\frac{1}{n_1} + \frac{1}{n_2} \right)}}$$

$$\bar{p} = \frac{n_1 \cdot \bar{p}_1 + n_2 \cdot \bar{p}_2}{n_1 + n_2} = \frac{x_1 + x_2}{n_1 + n_2}$$

$$\bar{q} = 1 - \bar{p}$$

z-formula

$$z = \frac{(\bar{p}_1 - \bar{p}_2) - (p_1 - p_2)}{\sqrt{\bar{p} \cdot \bar{q} \left(\frac{1}{n_1} + \frac{1}{n_2} \right)}}$$

$$\bar{p} = \frac{n_1 \cdot \bar{p}_1 + n_2 \cdot \bar{p}_2}{n_1 + n_2} = \frac{x_1 + x_2}{n_1 + n_2}$$

$$\bar{q} = 1 - \bar{p}$$

Where

z-formula

$$z = \frac{(\bar{p}_1 - \bar{p}_2) - (p_1 - p_2)}{\sqrt{\bar{p} \cdot \bar{q} \left(\frac{1}{n_1} + \frac{1}{n_2} \right)}}$$

$$\bar{p} = \frac{n_1 \cdot \bar{p}_1 + n_2 \cdot \bar{p}_2}{n_1 + n_2} = \frac{x_1 + x_2}{n_1 + n_2}$$

$$\bar{q} = 1 - \bar{p}$$

Where

z-formula

$$z = \frac{(\bar{p}_1 - \bar{p}_2) - (p_1 - p_2)}{\sqrt{\bar{p} \cdot \bar{q} \left(\frac{1}{n_1} + \frac{1}{n_2} \right)}}$$

$$\bar{p} = \frac{n_1 \cdot \bar{p}_1 + n_2 \cdot \bar{p}_2}{n_1 + n_2} = \frac{x_1 + x_2}{n_1 + n_2}$$

$$\bar{q} = 1 - \bar{p}$$

Where

► \bar{p}_1 = First sample proportion.

z-formula

$$z = \frac{(\bar{p}_1 - \bar{p}_2) - (p_1 - p_2)}{\sqrt{\bar{p} \cdot \bar{q} \left(\frac{1}{n_1} + \frac{1}{n_2} \right)}}$$

$$\bar{p} = \frac{n_1 \cdot \bar{p}_1 + n_2 \cdot \bar{p}_2}{n_1 + n_2} = \frac{x_1 + x_2}{n_1 + n_2}$$

$$\bar{q} = 1 - \bar{p}$$

Where

- ▶ \bar{p}_1 =First sample proportion.
- ▶ \bar{p}_2 =Second sample proportion.

z-formula

$$z = \frac{(\bar{p}_1 - \bar{p}_2) - (p_1 - p_2)}{\sqrt{\bar{p} \cdot \bar{q} \left(\frac{1}{n_1} + \frac{1}{n_2} \right)}}$$

$$\bar{p} = \frac{n_1 \cdot \bar{p}_1 + n_2 \cdot \bar{p}_2}{n_1 + n_2} = \frac{x_1 + x_2}{n_1 + n_2}$$

$$\bar{q} = 1 - \bar{p}$$

Where

- ▶ \bar{p}_1 = First sample proportion.
- ▶ \bar{p}_2 = Second sample proportion.
- ▶ p_1 = First population proportion.

z-formula

$$z = \frac{(\bar{p}_1 - \bar{p}_2) - (p_1 - p_2)}{\sqrt{\bar{p} \cdot \bar{q} \left(\frac{1}{n_1} + \frac{1}{n_2} \right)}}$$

$$\bar{p} = \frac{n_1 \cdot \bar{p}_1 + n_2 \cdot \bar{p}_2}{n_1 + n_2} = \frac{x_1 + x_2}{n_1 + n_2}$$

$$\bar{q} = 1 - \bar{p}$$

Where

- ▶ \bar{p}_1 =First sample proportion.
- ▶ \bar{p}_2 =Second sample proportion.
- ▶ p_1 =First population proportion.
- ▶ p_2 =Second population proportion.

z-formula

$$z = \frac{(\bar{p}_1 - \bar{p}_2) - (p_1 - p_2)}{\sqrt{\bar{p} \cdot \bar{q} \left(\frac{1}{n_1} + \frac{1}{n_2} \right)}}$$

$$\bar{p} = \frac{n_1 \cdot \bar{p}_1 + n_2 \cdot \bar{p}_2}{n_1 + n_2} = \frac{x_1 + x_2}{n_1 + n_2}$$

$$\bar{q} = 1 - \bar{p}$$

Where

- ▶ \bar{p}_1 =First sample proportion.
- ▶ \bar{p}_2 =Second sample proportion.
- ▶ p_1 =First population proportion.
- ▶ p_2 =Second population proportion.

z-formula

$$z = \frac{(\bar{p}_1 - \bar{p}_2) - (p_1 - p_2)}{\sqrt{\bar{p} \cdot \bar{q} \left(\frac{1}{n_1} + \frac{1}{n_2} \right)}}$$

$$\bar{p} = \frac{n_1 \cdot \bar{p}_1 + n_2 \cdot \bar{p}_2}{n_1 + n_2} = \frac{x_1 + x_2}{n_1 + n_2}$$

$$\bar{q} = 1 - \bar{p}$$

Where

- ▶ \bar{p}_1 =First sample proportion.
- ▶ \bar{p}_2 =Second sample proportion.
- ▶ p_1 =First population proportion.
- ▶ p_2 =Second population proportion.

Confidence Interval

Confidence Interval

Confidence interval

Confidence Interval

Confidence interval

$$\begin{aligned} (\bar{p}_1 - \bar{p}_2) - z_{\alpha/2} \sqrt{\frac{\bar{p}_1 \cdot (1 - \bar{p}_1)}{n_1} + \frac{\bar{p}_2 \cdot (1 - \bar{p}_2)}{n_2}} \\ \leq (p_1 - p_2) \leq \\ (\bar{p}_1 - \bar{p}_2) + z_{\alpha/2} \sqrt{\frac{\bar{p}_1 \cdot (1 - \bar{p}_1)}{n_1} + \frac{\bar{p}_2 \cdot (1 - \bar{p}_2)}{n_2}} \end{aligned}$$

Confidence Interval

Confidence interval

$$\begin{aligned} (\bar{p}_1 - \bar{p}_2) - z_{\alpha/2} \sqrt{\frac{\bar{p}_1 \cdot (1 - \bar{p}_1)}{n_1} + \frac{\bar{p}_2 \cdot (1 - \bar{p}_2)}{n_2}} \\ \leq (p_1 - p_2) \leq \\ (\bar{p}_1 - \bar{p}_2) + z_{\alpha/2} \sqrt{\frac{\bar{p}_1 \cdot (1 - \bar{p}_1)}{n_1} + \frac{\bar{p}_2 \cdot (1 - \bar{p}_2)}{n_2}} \end{aligned}$$

Examples

Examples

Example:

Examples

Example: A random sample of 500 persons belonging to urban area 200 are found to be commuters of public transport.

Examples

Example: A random sample of 500 persons belonging to urban area 200 are found to be commuters of public transport. In another sample of 400 persons belonging to rural area 200 are found to be commuter of public transport.

Examples

Example: A random sample of 500 persons belonging to urban area 200 are found to be commuters of public transport. In another sample of 400 persons belonging to ruler area 200 are found to be commuter of public transport. Discuss whether the data reveal that proportion of commuters of public transport is significantly higher for ruler area as compared to urban area?

Examples

Example: A random sample of 500 persons belonging to urban area 200 are found to be commuters of public transport. In another sample of 400 persons belonging to ruler area 200 are found to be commuter of public transport. Discuss whether the data reveal that proportion of commuters of public transport is significantly higher for ruler area as compared to urban area? (Take $\alpha = 0.01$)

Examples

Example: A random sample of 500 persons belonging to urban area 200 are found to be commuters of public transport. In another sample of 400 persons belonging to ruler area 200 are found to be commuter of public transport. Discuss whether the data reveal that proportion of commuters of public transport is significantly higher for ruler area as compared to urban area? (Take $\alpha = 0.01$)

Example:

Examples

Example: A random sample of 500 persons belonging to urban area 200 are found to be commuters of public transport. In another sample of 400 persons belonging to rural area 200 are found to be commuter of public transport. Discuss whether the data reveal that proportion of commuters of public transport is significantly higher for rural area as compared to urban area? (Take $\alpha = 0.01$)

Example: In a village 'A' out of a random sample of 1000 persons 100 were found to be vegetarians.

Examples

Example: A random sample of 500 persons belonging to urban area 200 are found to be commuters of public transport. In another sample of 400 persons belonging to rural area 200 are found to be commuter of public transport. Discuss whether the data reveal that proportion of commuters of public transport is significantly higher for rural area as compared to urban area? (Take $\alpha = 0.01$)

Example: In a village 'A' out of a random sample of 1000 persons 100 were found to be vegetarians. While in another village 'B' out of 1500 persons 180 were found to be vegetarians.

Examples

Example: A random sample of 500 persons belonging to urban area 200 are found to be commuters of public transport. In another sample of 400 persons belonging to rural area 200 are found to be commuter of public transport. Discuss whether the data reveal that proportion of commuters of public transport is significantly higher for rural area as compared to urban area? (Take $\alpha = 0.01$)

Example: In a village 'A' out of a random sample of 1000 persons 100 were found to be vegetarians. While in another village 'B' out of 1500 persons 180 were found to be vegetarians. Do you find a significant difference in the food habits of the people of the two village?

Examples

Example: A random sample of 500 persons belonging to urban area 200 are found to be commuters of public transport. In another sample of 400 persons belonging to rural area 200 are found to be commuter of public transport. Discuss whether the data reveal that proportion of commuters of public transport is significantly higher for rural area as compared to urban area? (Take $\alpha = 0.01$)

Example: In a village 'A' out of a random sample of 1000 persons 100 were found to be vegetarians. While in another village 'B' out of 1500 persons 180 were found to be vegetarians. Do you find a significant difference in the food habits of the people of the two village? (Take $\alpha=0.01$) Also establish 95% confidence limit.

Examples

Example: A random sample of 500 persons belonging to urban area 200 are found to be commuters of public transport. In another sample of 400 persons belonging to rural area 200 are found to be commuter of public transport. Discuss whether the data reveal that proportion of commuters of public transport is significantly higher for rural area as compared to urban area? (Take $\alpha = 0.01$)

Example: In a village 'A' out of a random sample of 1000 persons 100 were found to be vegetarians. While in another village 'B' out of 1500 persons 180 were found to be vegetarians. Do you find a significant difference in the food habits of the people of the two village? (Take $\alpha=0.01$) Also establish 95% confidence limit.

Method-VIII χ^2 Test(Contingency Table)

Method-VIII χ^2 Test(Contingency Table)

χ^2 formula:

Method-VIII χ^2 Test(Contingency Table)

χ^2 formula:

$$\chi^2 = \Sigma \Sigma \left(\frac{(O - E)^2}{E} \right)$$

Method-VIII χ^2 Test(Contingency Table)

χ^2 **formula:**

$$\chi^2 = \Sigma \Sigma \left(\frac{(O - E)^2}{E} \right)$$

$$df=(r-1)(c-1)$$

Method-VIII χ^2 Test(Contingency Table)

χ^2 **formula:**

$$\chi^2 = \Sigma \Sigma \left(\frac{(O - E)^2}{E} \right)$$

df=(r-1)(c-1)

where

Method-VIII χ^2 Test(Contingency Table)

χ^2 **formula:**

$$\chi^2 = \Sigma \Sigma \left(\frac{(O - E)^2}{E} \right)$$

$df = (r - 1)(c - 1)$

where

1 O=Observed frequency.

Method-VIII χ^2 Test(Contingency Table)

χ^2 formula:

$$\chi^2 = \Sigma \Sigma \left(\frac{(O - E)^2}{E} \right)$$

df=(r-1)(c-1)

where

1 O=Observed frequency.

2 E=Expected frequency.

Method-VIII χ^2 Test(Contingency Table)

χ^2 formula:

$$\chi^2 = \Sigma \Sigma \left(\frac{(O - E)^2}{E} \right)$$

df=(r-1)(c-1)

where

- 1** O=Observed frequency.
- 2** E=Expected frequency.
- 3** r=No of rows.

Method-VIII χ^2 Test(Contingency Table)

χ^2 formula:

$$\chi^2 = \Sigma \Sigma \left(\frac{(O - E)^2}{E} \right)$$

df=(r-1)(c-1)

where

- 1** O=Observed frequency.
- 2** E=Expected frequency.
- 3** r=No of rows.
- 4** c=No of columns.

Method-VIII χ^2 Test(Contingency Table)

χ^2 formula:

$$\chi^2 = \Sigma \Sigma \left(\frac{(O - E)^2}{E} \right)$$

df=(r-1)(c-1)

where

- 1** O=Observed frequency.
- 2** E=Expected frequency.
- 3** r=No of rows.
- 4** c=No of columns.

Method-VIII χ^2 Test(Contingency Table)

χ^2 formula:

$$\chi^2 = \Sigma \Sigma \left(\frac{(O - E)^2}{E} \right)$$

$df=(r-1)(c-1)$

where

- 1** O=Observed frequency.
- 2** E=Expected frequency.
- 3** r=No of rows.
- 4** c=No of columns.

Method-VIII χ^2 Test(Contingency Table)

Method-VIII χ^2 Test(Contingency Table)

Example:

GTU -May-2013

Method-VIII χ^2 Test(Contingency Table)

Example:

GTU -May-2013

The table below gives beverage preferences for random samples of teens and adults.

Method-VIII χ^2 Test(Contingency Table)

Example:

GTU -May-2013

The table below gives beverage preferences for random samples of teens and adults.

	Teens	Adults	Total
Coffee	50	200	250
Tea	100	150	250
Soft Drink	200	200	400
Other	50	50	100

Method-VIII χ^2 Test(Contingency Table)

Example:

GTU -May-2013

The table below gives beverage preferences for random samples of teens and adults.

	Teens	Adults	Total
Coffee	50	200	250
Tea	100	150	250
Soft Drink	200	200	400
Other	50	50	100

Test for independence between age (i.e., adult and teen) and drink preferences.(Take $\alpha = 0.05$).

Method-VIII χ^2 Test(Contingency Table)

Method-VIII χ^2 Test(Contingency Table)

Example:

GTU -Dec-2011

Method-VIII χ^2 Test(Contingency Table)

Example:

GTU -Dec-2011

A brand manager is concerned that her brand's share may be unevenly distributed throughout the country.

Method-VIII χ^2 Test(Contingency Table)

Example:

GTU -Dec-2011

A brand manager is concerned that her brand's share may be unevenly distributed throughout the country. In a survey in which the country was divided into four geographical regions, a random sampling of 100 consumers in each region was surveyed, with the following results:

Method-VIII χ^2 Test(Contingency Table)

Example:

GTU -Dec-2011

A brand manager is concerned that her brand's share may be unevenly distributed throughout the country. In a survey in which the country was divided into four geographical regions, a random sampling of 100 consumers in each region was surveyed, with the following results:

	Region				Total
	NE	NW	SE	SW	
Purchase the brand	40	55	45	50	190
Do not purchase	60	45	55	50	210
Total	100	100	100	100	400

Method-VIII χ^2 Test(Contingency Table)

Example:

GTU -Dec-2011

A brand manager is concerned that her brand's share may be unevenly distributed throughout the country. In a survey in which the country was divided into four geographical regions, a random sampling of 100 consumers in each region was surveyed, with the following results:

	Region				Total
	NE	NW	SE	SW	
Purchase the brand	40	55	45	50	190
Do not purchase	60	45	55	50	210
Total	100	100	100	100	400

State the null and alternative hypotheses and test whether brand share is the same across the four regions at $\alpha = 0.05$

Method-VIII χ^2 Test(Contingency Table)

Method-VIII χ^2 Test(Contingency Table)

Example:

GTU -Dec-2013

Method-VIII χ^2 Test(Contingency Table)

Example:

GTU -Dec-2013

The contingency table below summarizes the results obtained in a study conducted by a research organization, with respect to the performance of four competing brands of toothpaste among the users:

Method-VIII χ^2 Test(Contingency Table)

Example:

GTU -Dec-2013

The contingency table below summarizes the results obtained in a study conducted by a research organization, with respect to the performance of four competing brands of toothpaste among the users:

	Brand A	Brand B	Brand C	Brand D
No cavities	9	13	17	11
One to five cavities	63	70	85	82
More than five cavities	28	37	48	37
Total	100	120	150	130

Method-VIII χ^2 Test(Contingency Table)

Example:

GTU -Dec-2013

The contingency table below summarizes the results obtained in a study conducted by a research organization, with respect to the performance of four competing brands of toothpaste among the users:

	Brand A	Brand B	Brand C	Brand D
No cavities	9	13	17	11
One to five cavities	63	70	85	82
More than five cavities	28	37	48	37
Total	100	120	150	130

Using χ^2 test at 5% level of significance test the hypothesis that incidence of cavities is independent of the brand of the toothpaste used.

Method-VIII χ^2 Test(Contingency Table)

Example:

GTU -Dec-2013

The contingency table below summarizes the results obtained in a study conducted by a research organization, with respect to the performance of four competing brands of toothpaste among the users:

	Brand A	Brand B	Brand C	Brand D
No cavities	9	13	17	11
One to five cavities	63	70	85	82
More than five cavities	28	37	48	37
Total	100	120	150	130

Using χ^2 test at 5% level of significance test the hypothesis that incidence of cavities is independent of the brand of the toothpaste used.

Method-IX χ^2 Test(Uniform distribution)

Method-IX χ^2 Test(Uniform distribution)

χ^2 formula:

Method-IX χ^2 Test(Uniform distribution)

χ^2 formula:

$$\chi^2 = \Sigma \left(\frac{(O - E)^2}{E} \right)$$

df=n-1

Method-IX χ^2 Test(Uniform distribution)

χ^2 formula:

$$\chi^2 = \Sigma \left(\frac{(O - E)^2}{E} \right)$$

df=n-1

Example:

GTU -Jan-2011 & May-2014

Method-IX χ^2 Test(Uniform distribution)

χ^2 formula:

$$\chi^2 = \Sigma \left(\frac{(O - E)^2}{E} \right)$$

df=n-1

Example:

GTU -Jan-2011 & May-2014

Define the chi-square test.

Method-IX χ^2 Test(Uniform distribution)

χ^2 formula:

$$\chi^2 = \Sigma \left(\frac{(O - E)^2}{E} \right)$$

df=n-1

Example:

GTU -Jan-2011 & May-2014

Define the chi-square test. A die is thrown 150 times and the following results are obtained.

Method-IX χ^2 Test(Uniform distribution)

χ^2 formula:

$$\chi^2 = \Sigma \left(\frac{(O - E)^2}{E} \right)$$

df=n-1

Example:

GTU -Jan-2011 & May-2014

Define the chi-square test. A die is thrown 150 times and the following results are obtained.

Number turned up	1	2	3	4	5	6
Frequency	19	23	28	17	32	31

Method-IX χ^2 Test(Uniform distribution)

χ^2 formula:

$$\chi^2 = \Sigma \left(\frac{(O - E)^2}{E} \right)$$

df=n-1

Example:

GTU -Jan-2011 & May-2014

Define the chi-square test. A die is thrown 150 times and the following results are obtained.

Number turned up	1	2	3	4	5	6
Frequency	19	23	28	17	32	31

Test the hypothesis that the die is unbiased at 5 % level of significance.

Method-IX χ^2 Test(Uniform distribution)

χ^2 formula:

$$\chi^2 = \Sigma \left(\frac{(O - E)^2}{E} \right)$$

df=n-1

Example:

GTU -Jan-2011 & May-2014

Define the chi-square test. A die is thrown 150 times and the following results are obtained.

Number turned up	1	2	3	4	5	6
Frequency	19	23	28	17	32	31

Test the hypothesis that the die is unbiased at 5 % level of significance.

Method-IX χ^2 Test(Uniform distribution)

Method-IX χ^2 Test(Uniform distribution)

Example:

Method-IX χ^2 Test(Uniform distribution)

Example:

The demand for a particular spare part in a factory was found to vary from day to day. In a sample study the following information was obtained:

Method-IX χ^2 Test(Uniform distribution)

Example:

The demand for a particular spare part in a factory was found to vary from day to day. In a sample study the following information was obtained:

Day	Mon	Tue	Wed	Thu	Fri	Sat
Demand	1124	1125	1110	1120	1126	1115

Method-IX χ^2 Test(Uniform distribution)

Example:

The demand for a particular spare part in a factory was found to vary from day to day. In a sample study the following information was obtained:

Day	Mon	Tue	Wed	Thu	Fri	Sat
Demand	1124	1125	1110	1120	1126	1115

Test the hypothesis that the number of part demanded does not depend on the day of the week at 5% of significance.

Method-X χ^2 Test(Ratio test)

Method-X χ^2 Test(Ratio test)

$$\chi^2 = \Sigma \left(\frac{(O - E)^2}{E} \right)$$

df=n-1

Method-X χ^2 Test(Ratio test)

$$\chi^2 = \Sigma \left(\frac{(O - E)^2}{E} \right)$$

df=n-1

Example:

GTU -Dec-2012

Method-X χ^2 Test(Ratio test)

$$\chi^2 = \Sigma \left(\frac{(O - E)^2}{E} \right)$$

df=n-1

Example:

GTU -Dec-2012

A sample analysis of examination results of 500 students was made.

Method-X χ^2 Test(Ratio test)

$$\chi^2 = \Sigma \left(\frac{(O - E)^2}{E} \right)$$

df=n-1

Example:

GTU -Dec-2012

A sample analysis of examination results of 500 students was made. It was found that 220 students had failed,

Method-X χ^2 Test(Ratio test)

$$\chi^2 = \Sigma \left(\frac{(O - E)^2}{E} \right)$$

df=n-1

Example:

GTU -Dec-2012

A sample analysis of examination results of 500 students was made. It was found that 220 students had failed, 170 had secured a third class,

Method-X χ^2 Test(Ratio test)

$$\chi^2 = \Sigma \left(\frac{(O - E)^2}{E} \right)$$

df=n-1

Example:

GTU -Dec-2012

A sample analysis of examination results of 500 students was made. It was found that 220 students had failed, 170 had secured a third class, 90 were placed in second class and 20 got a first class.

Method-X χ^2 Test(Ratio test)

$$\chi^2 = \Sigma \left(\frac{(O - E)^2}{E} \right)$$

df=n-1

Example:

GTU -Dec-2012

A sample analysis of examination results of 500 students was made. It was found that 220 students had failed, 170 had secured a third class, 90 were placed in second class and 20 got a first class. Are these figures commensurate with the general examination result which is the ratio of 4 : 3 : 2 : 1 for the various categories respectively?

Method-X χ^2 Test(Ratio test)

$$\chi^2 = \Sigma \left(\frac{(O - E)^2}{E} \right)$$

df=n-1

Example:

GTU -Dec-2012

A sample analysis of examination results of 500 students was made. It was found that 220 students had failed, 170 had secured a third class, 90 were placed in second class and 20 got a first class. Are these figures commensurate with the general examination result which is the ratio of 4 : 3 : 2 : 1 for the various categories respectively?

Method-X χ^2 Test(Ratio test)

Method-X χ^2 Test(Ratio test)

Example:

Method-X χ^2 Test(Ratio test)

Example:

In a cross - breeding experiment with plants at certain species 240 offspring were classified into 4 classes with respect to the structure of their leaves as follow:

Method-X χ^2 Test(Ratio test)

Example:

In a cross - breeding experiment with plants at certain species 240 offspring were classified into 4 classes with respect to the structure of their leaves as follow:

Class	I	II	III	IV	Total
Frequency	21	127	40	52	240

Method-X χ^2 Test(Ratio test)

Example:

In a cross - breeding experiment with plants at certain species 240 offspring were classified into 4 classes with respect to the structure of their leaves as follow:

Class	I	II	III	IV	Total
Frequency	21	127	40	52	240

According to theory of heredity , the probabilities of the four class should be in ratio 1:9:3:3 . Are these data consistent with theory?

Method-X χ^2 Test(Ratio test)

Example:

In a cross - breeding experiment with plants at certain species 240 offspring were classified into 4 classes with respect to the structure of their leaves as follow:

Class	I	II	III	IV	Total
Frequency	21	127	40	52	240

According to theory of heredity , the probabilities of the four class should be in ratio 1:9:3:3 . Are these data consistent with theory?

Method-XI χ^2 Test(Poisson distribution)

Method-XI χ^2 Test(Poisson distribution)

$$\chi^2 = \Sigma \left(\frac{(O - E)^2}{E} \right)$$

,df=n-1

Method-XI χ^2 Test(Poisson distribution)

$$\chi^2 = \Sigma \left(\frac{(O - E)^2}{E} \right)$$

,df=n-1

Example:

GTU -May-2012

Method-XI χ^2 Test(Poisson distribution)

$$\chi^2 = \Sigma \left(\frac{(O - E)^2}{E} \right)$$

,df=n-1

Example:

GTU -May-2012

A typist in a company commits the following number of mistakes per page in typing 432 pages. Does this information verify that the mistakes are distributed according to Poisson law?

Method-XI χ^2 Test(Poisson distribution)

$$\chi^2 = \Sigma \left(\frac{(O - E)^2}{E} \right)$$

,df=n-1

Example:

GTU -May-2012

A typist in a company commits the following number of mistakes per page in typing 432 pages. Does this information verify that the mistakes are distributed according to Poisson law?

No. of mistakes per page	0	1	2	3	4	5
No. of pages	223	142	48	15	4	0

Method-XI χ^2 Test(Poisson distribution)

$$\chi^2 = \Sigma \left(\frac{(O - E)^2}{E} \right)$$

,df=n-1

Example:

GTU -May-2012

A typist in a company commits the following number of mistakes per page in typing 432 pages. Does this information verify that the mistakes are distributed according to Poisson law?

No. of mistakes per page	0	1	2	3	4	5
No. of pages	223	142	48	15	4	0

Method-XI χ^2 Test(Poisson distribution)

Method-XI χ^2 Test(Poisson distribution)

Example:

Method-XI χ^2 Test(Poisson distribution)

Example:

In the accounting department of a bank 100 accounts are selected at random and examined for errors. Suppose the following results have been obtained:

Method-XI χ^2 Test(Poisson distribution)

Example:

In the accounting department of a bank 100 accounts are selected at random and examined for errors. Suppose the following results have been obtained:

No of errors	0	1	2	3	4	5	6
No of account	35	40	19	2	0	2	2

Method-XI χ^2 Test(Poisson distribution)

Example:

In the accounting department of a bank 100 accounts are selected at random and examined for errors. Suppose the following results have been obtained:

No of errors	0	1	2	3	4	5	6
No of account	35	40	19	2	0	2	2

On the basis of this of this information can it be concluded that the errors are distributed according to the poisson probability law?

Method-XI χ^2 Test(Poisson distribution)

Example:

In the accounting department of a bank 100 accounts are selected at random and examined for errors. Suppose the following results have been obtained:

No of errors	0	1	2	3	4	5	6
No of account	35	40	19	2	0	2	2

On the basis of this of this information can it be concluded that the errors are distributed according to the poisson probability law?

References:

- 1** Anderson, Sweeney, Williams, “Statistics for business and economics”, 9th edition, Thompson Publication
- 2** S P Gupta, “Statistical Methods”, 30th edition, S Chand
- 3** J.Susan Milton & Jesse Arnold, “Introduction to Probability & Statistics: Principles & Applications for Engineering & Computing Sciences”
- 4** Bharat Jhunjhunwala, “Business Statistics”, first edition, S Chand, 2008
- 5** Richard Levin, David Rubin, “Statistics for Management”, 7th edition, PHI
- 6** Nabendu Pal, Sahadeb Sarkar, “Statistics-Concepts and Applications”, 2nd edition, PHI

Temporary page!

\LaTeX was unable to guess the total number of pages correctly. As there was some unprocessed data that should have been added to the final page this extra page has been added to receive it.

If you rerun the document (without altering it) this surplus page will go away, because \LaTeX now knows how many pages to expect for this document.