



US 20110000125A1

(19) United States

(12) Patent Application Publication

McDaniel et al.

(10) Pub. No.: US 2011/0000125 A1

(43) Pub. Date:

Jan. 6, 2011

(54) PRODUCTION OF FATTY ALCOHOLS WITH  
FATTY ALCOHOL FORMING ACYL-COA  
REDUCTASES (FAR)

(75) Inventors: Robert McDaniel, Palo Alto, CA  
(US); Behnaz Behrouzian,  
Sunnyvale, CA (US); Louis Clark,  
San Francisco, CA (US); Douglas  
Hattendorf, Mountain View, CA  
(US); Fernando Valle, Burlingame,  
CA (US)

Correspondence Address:

Codexis, Inc.  
200 Penobscot Drive  
Redwood City, CA 94063 (US)

(73) Assignee: CODEXIS, INC., Redwood City,  
CA (US)

(21) Appl. No.: 12/825,939

(22) Filed: Jun. 29, 2010

#### Related U.S. Application Data

(60) Provisional application No. 61/221,934, filed on Jun.  
30, 2009, provisional application No. 61/315,380,  
filed on Mar. 18, 2010.


#### Publication Classification

(51) Int. Cl.  
*C10L 1/19* (2006.01)  
*C12P 7/64* (2006.01)  
*C07H 21/00* (2006.01)  
*C12N 15/63* (2006.01)  
*C12N 1/21* (2006.01)  
*C12N 1/19* (2006.01)  
*C12N 9/02* (2006.01)  
*C12P 5/00* (2006.01)  
*C07C 9/00* (2006.01)  
*C09K 3/00* (2006.01)

(52) U.S. Cl. .... 44/388; 435/134; 536/23.2; 435/320.1;  
435/252.33; 435/254.21; 435/254.2; 435/189;  
435/166; 585/16; 252/182.12

#### ABSTRACT

The disclosure relates to methods of producing fatty alcohols from recombinant host cells comprising genes encoding heterologous fatty acyl-CoA reductase (FAR) enzymes. The disclosure further relates to FAR enzymes and functional fragments thereof derived from marine bacterium and particularly marine gamma proteobacterium such as *Marinobacter* and *Oceanobacter*; polynucleotides encoding the FAR enzymes and vectors and host cells comprising the same.






FIG. 1

## FIG.2

Codon optimized *M. aligicola* DG893 FAR DNA (SEQ ID NO: 1)

ATGGCTACTCAACAACAACAGAACGGTGCATCTGCATCCGGCGTCTTGGAAACAACCTCGT  
GGAAAGCACGTTCTTATCACAGGTACTACCGGATTTGGCAAAGTGGTTCTGGAAAAG  
TTGATTCGTACTGTTCCGGATATTGGAGGTATTCATCTGCTGATTCTGGCAATAAACGT  
CATCCAGCCGCTCGTGAACGTTCTGAACGAAATTGCGTCCTCCTCCGTCTCGAACGT  
TTGCGTCACGATGATAATGAAGCCTCGAGACCTTCTTGGAAAGAACGTGTTCACTGTATT  
ACCGGTGAGGTTACTGAATCCCGTTGGTTGACACCTGAACGTTCTGCTTGGCC  
GGTCAGGTTGACGCTTTATTAACAGCGCTGCAAGCGTGAACCTTCTGAGGAATTGGAT  
AAAGCCCTGAAAATCAACACCTTGTCTGAAAATGTTGCTGCTTGCAGAATTGAAC  
TCCGCTATGGCGGTATTCAAGGTTCCACTTGTACGTTAACGGTAAAAACTCCGGTCAA  
ATTACCGAATCCGTATTAAACCTGCTGGCGAATCCATTCCCCGTTCCACTGACGGTTAC  
TACGAGATCGAAGAATTGGTCCATCTGTTGCAAGACAAGATTCCGATGTTAAAGCTCGT  
TACTCCGGAAAAGTTCTGGAGAAAAATTGGTTGATTGGGTATTCTGAGGCCAATAAT  
TACGGATGGTCCGACACCTACACATTCAACAAATGGTTGGGTGAACAACGTGCTGATGAAG  
GCCTTGTCTGGTCGTTCTTGACTATTGTGCGTCCCTCTATTATTGAGTCCGCTTGGAA  
GAACCTTCCCCTGGTTGGATCGAAGGCCTAAAGTTGCCATGCCATTATCTGGCTTAT  
GCCCGTAAAAAGTTAGCCTGTTCCCTGGAAAACGTTCCGGCATTATTGATGTTATTCC  
GTCGATTGGTTGCGACTCCATCATCTTGTCTCTGGCTGAGGCCTGCTGGTTCTGGT  
CAACGTCGTATTATCAATGTTGCAGCGGTGGTTCTAATCCAATCTCCCTGGTAAGTTC  
ATTGATTATTGATGGCCGAGGCTAACGACCAACTATGCTGCCTACGATCAACTGTTTAT  
CGTCGTCCTACTAAACCTTCGTCGCCGTGAACCGTAAATTGTTGACGTTGGCTGGTCAAATCGT  
GGTATGCGTGGTCCCTTTCTATTGCCGGTAAAGCTATGCGTTGGCTGGTCAAATCGT  
GAGTTGAAAGTCTTAAGAACCTGATACGACCCGTTCCCTGCAACCATTGGCTTC  
TATACTGCTCCGACTATATCTTCCGTAACGATAGCTGATGCCCTGGCTCTCGTATG  
GGTGAATTGGATCGTGGTCTTCCAGTTGATGCTCGTCAAATTGATGGCAGTTGTAC  
TTGTGAAAATTCAATTGGGTGGTCTGAACCGTTACGCTTGAAGGAACGTAAACTGTAT  
TCTTGCGTGCTGCTGATACTCGTAAAAAGCTGCCTAA

## FIG. 3

Codon optimized *M. aligicola* DG893 FAR DNA (SEQ ID NO: 3)

```
ATGCCACCCAGCAGCAGCAGAACGGTGCATCCGCTCGGGCGTTCTGGAGCAGCTTAGA
GGCAAGCATGTCTGATTACCGGTACTACAGGATTCTGGAAAGGTGGTTCTGGAGAAG
CTGATCCGAACCGTGCTGACATCGGTGGTATTCATCTGCTGATTAGAGGCAACAAGAGA
CATCCTGCTGCCAGAGAAAGATTCTTGAACGAAATCGCCTTCCTCTGTGTTCGAGCGG
CTTAGACATGACGACAACGAAGCCTTGAGACTTCCTGGAGGAGCGTGTGCACTGCATC
ACCGGAGAAGTGACCGAGTCGAGATTGGCCTTACTCCTGAGCGGTTCCGAGCCCTTGCT
GGCCAAGTGGATGCCTTCATCAATTCCGCGCCTGTAACTTCAGAGAGGAGCTGGAC
AAGGCACATCAAGATCAACACCCCTGTGTTGGAGAACGTGGCTCTGGCCGAACCTAAC
TCCGCTATGGCAGTGATCCAAGTTCCACCTGTTACGTGAACGGCAAGAACTCTGGACAG
ATCACCGAGTCCGTTATCAAGCCCGCTGGCGAATCCATCCCCAGATCCACAGATGGCTAC
TACGAGATCGAGGAGCTGGTCCACCTCTGCAAGACAAGATCTCCGACGTGAAGGCTCGA
TACTCTGGCAAGGTGTTGGAGAAGAAGCTGGTGGACCTGGGATCCGAGAGGCGAACAAAC
TACGGCTGGTCTGACACCTACACCTCACCAAATGGCTCGGAGAGCAGCTCTGATGAAA
GCTCTGTCCGGAAGATCCCTGACTATCGTGCAGCCTTCATCATCGAGTCGGCTTTGAA
GAGCCTTCTCCAGGTTGGATCGAGGGCGTAAGGTTGCTGACGCCATCATCCTGCGTAC
GCCAGAGAGAAGGTTCTGTTCCCCGGCAAGCGATCTGGCATCATCGACGTTATCCCC
GTGGATCTGGTGGCCAACTCTATCATTCTCTCTGCTGAAGCCCTTCTGGATCTGGC
CAGCGTAGAATCTACCAATGTTGTTCTGGCGGTTCTAACCCGATTCTGGCAAGTTC
ATCGACTACCTTATGCCGAAGCCAAGACCAACTATGCTGCCTACGACCAGCTTTCTAC
CGACGACCCACCAAGCCCTCGTCGCTGTGAACCGAAAGCTGTTGATGTTGCGTGGGA
GGAATGCGAGTGCCTCTTCCATTGCTGGCAAGGCCATGAGATTGGCGGGTCAGAATCGA
GAATTGAAGGTTCTCAAGAACCTGACACTACTCGATCGCTCGCTACTATCTTGGATTG
TACACTGCTCCTGACTACATCTTCCGGAATGACTCTGATGGCTCTGCTTCCGAATG
GGAGAACTCGATCGTGTGCTGTTCCCTGTTGACGCTCGACAGATCGACTGGCAGCTAC
TTGTGTAAGATCCACCTGGCGGCTGAACCGATATGCTCTGAAAGAACGAAAGCTGTAC
AGCCTTAGAGCCGCTGATAACCCGAAAGAACGGCTGCT
```

## FIG. 4

Codon optimized *Oceanobacter sp.* RED65 FAR DNA (SEQ ID NO: 5)

```
ATGTCTCAGTACTCCCGTTCTCCGTTCTCAATCTTGAAGGGCAAACATATCTTCTTG  
ACAGGTGTTACGGTTCTGGCAAGGCATTCTGGAGAAACTGTTGACTCCGTTCCA  
CAATTGGCTCAGATTCATATCCTGGTCCGTGGTAAAGTTAGCGCTAAGAACGCTTC  
CAACATGACATCTGGGTTCTTCATCTTGAGCGTCTTAAGGAACAACATGGCGAACAT  
TTCGAAGAATGGGTTCAAAGCAAGATCAACCTGTCGAGGGCGAACTTACTCAACCTATG  
TTCGATTGCCTCTGCTGAGTCGCTGGCTGGCTAACCAATTGGATCTGATCATCAAT  
AGCGCTGCTTCTGTCAACTTCGTGAGAACTTGGAGAACGGCTCTGAACATCAATACCCTG  
TGTCTGAACACATCATTGCGCTGGCTCAGTACAACGTCGCTGCTCAAACCTGTTATG  
CAAATCTCACCTGCTATGTCAACGGCTCAACAAAGGCCAATCAACGAGGAAGTTGTT  
GGCCTGCTTCTGGTTGATTCCCTCAGTTGTCCTAAGACTGCTACGACATTGATTCCGTC  
TTCAAGCGTGTTCATTCCCAGATTGAACAAAGTCAAGAACGTAAGACCGACATTGAACAG  
CAAGAACAAAGCCTTGATCAAACCTGGGTATCAAGACCTCTCAACATTGCGCTGGAATGAT  
ACCTACACCTTCACCAAGTGGCTGGGTAAACAATTGTTGATTGAGAAGTTGGCAAGCAA  
TCTTGACAATTCTCGTCCGTCCATCATCGAATCTGCTGTTGACCTGCTCCTGGC  
TGGGTTGAAGGCCTGAAAGTTGCGGATGCCCTGATCTATGCTTACGCTAAAGGTCGTGTT  
AGCATCTTCCCGTGTGATGAAGGTATCTGGATGTCATCCCTGTTGATTGGTTGCT  
AATGCCGCCGATTGTCGCTGCTCAGTTATGGAATCTAACAGCAAACGGCTATCGT  
ATCTATCAATGCTGTTCTGGTCCCGTAATCCGATCAAGCTGAAAGAGTTCAATTGTCAC  
ATTCAAATGTTGCTCAAGCCCGTTACCAAGAACGGCTAAAGTTGTTGCTGACAAACCT  
CAGGAAGCCTTCAAGACCGTGTCTCCTAAACGTTCAAGCTGTACATGTCGGTTCACT  
GCTATCACATGGCTAAGACCATTGGTCGTCTTGGATCTAATGCTGCCCTCCAA  
CACATGTTGAAGGCTAAGACCACAGCATCTTGGCGAACATCTTGGTTCTACACTGCT  
CCTAACTACCGTTCTCCTCTCAGAAGTTGGAACAGTTGGTCAAACAGTTGATACTACC  
GAACAAACGTCTTACGACATTGCTGCTGACCATTGATGGAAGTACTACTTGAGGAA  
GTTCACATGGATGGTCTTCACAAGTATGCGTTGGCGATCGTCAAGAATTGAAGGCCAAA  
CATGTTGAAGAACGCTAAACGTGAAACGATTGCTCAAGCTGCTTAA
```

## FIG. 5

Codon optimized *Oceanobacter sp.* RED65 FAR DNA (SEQ ID NO: 7)

```
ATGTCCCAGTACTCGGCTTCCTGTTCTCAATCTGAAGGGCAAGCACATCTTCTC  
ACTGGTGTACTGGCTTCGGAAAGGCAATTCTGGAGAAGCTCTGTACTCGGTTCCC  
CAGCTGGCACAGATCCATATCCTGTGAGAGGCGCAAAGTTCTGCCAAGAACGGTTT  
CAGCACGACATCCTGGCTCTAGCATCTCGAGAGACTTAAGGAGAACACGGAGAGCAC  
TTTGAGGAATGGGTTAGTCCAAGATCAACCTGTGGAGGGAGAACTGACTCAACCAATG  
TTTGACCTCCCTCTGCTGAGTCGCTGGACTTGCTAACCGACTGGACCTGATCATCAAC  
TCTGCCGCTCGGTTAACCTTCGAGAGAACCTGGAGAACGGCTCTGAACATCAACACCCTG  
TGCCTGAACACATCATCGCTCTGGCTCAGTACAATGTCGCTGCCAGACTCCTGTGATG  
CAGATTCCACCTGCTACGTGAACGGCTCAACAAGGGCCAGATCAACGAAGAAGTTGTG  
GGACCTGCTCTGGACTGATCCCTCAGCTGTCTCAAGATTGCTACGACATCGACTCCGTC  
TTCAAGAGAGTGCATTCCAGATTGAGCAGGTGAAGAACAGAGAACAGACATTGAGCAG  
CAGGAACAAGCCATTATCAAGCTCGGTATCAAGACTTCCAGCACTTGGCTGGAATGAC  
ACTTACACCTCACCAAATGGCTCGCGAACAGCTGTTGATTAGCTGAGAGAGCCTGCCGGAA  
TCTCTCACCATTCTCGACCTCGATCATTGAGTCTGCTGTGAGAGAGCCTGCCGGAA  
TGGGTCGAAGGAGTGAAAGTCGCTGACGCCCTATCTACGCTTATGCTAACGGAAAGAGTC  
TCGATCTTCCTGGAAGAGACGAGGGCATCCTGATGTGATTCCCGTTGACCTGTTGCC  
AATGCTGCTGCTTTCTGCTGCACAACCATGGAGTCCAACCAACAGACTGGCTACCGA  
ATCTACCAATGCTGCTGGATCTCGAAACCCCATCAAGCTGAAGGAGTTCATCCGACAC  
ATCCAGAACGTGGCTCAGGCCGATAACCAAGGAATGCCCTAACGCTGTTGCCGATAAGCCT  
CAAGAGGCCCTCAAGACTGTCTCCCAAGAGATTCAAGCTGTACATGTCCGGCTTCACT  
GCCATCACCTGGCTAACGACATCATTGGCCGAGTGGCTGAGTCAACGCAGCCTCCAG  
CACATGCTGAAGGCAAAGACCACAGCTTCCCTGGCCAACATCTCGGCTTACACTGCA  
CCCCAACTACCGTTCTCTTCCCAGAAGTTGGAACAGCTGGTGAAGCAGTTGACACTACC  
GAGCAGCGACTGTACGACATTGAGCTGACCACCTCGACTGGAAGTACTACCTGCAAGAG  
GTTCACATGGATGGCCTTCACAAGTACGCTTGGCCATCGTCAGGAGCTGAAACCCAAG  
CACGTGAAGAAGAGAAAGCGGGAGACTATCCGACAAGCTGCG
```

FIG. 6

FAR peptide (SEQ ID NOs: 2 and 4) from codon optimized (SEQ ID NOs: 1 and 3) DNA

MATQQQNGASASGVLEQLRGKHVLITGTTGFLGVVLEKLIRTVPDIGGIHLLIRGNKR  
HPAARERFLNEIASSSVFERLRHDDNEAFETFLEERVHCITGEVTESRFGLTPERFRALA  
GQVDAFINSAASVNRFREELDKALKINTLCLENVAALAEELNSAMAVIQVSTCYVNGKNSQ  
ITESVIKPAGESIPRSTDGYYEIEELVHLLQDKISDVKARYSGKVLEKKLVLDLGIREANN  
YGWSDTYTFTKWLGEQLLMKALSGRSLTIVRPSIIESALEEPPSPGWIEGVKVADAIILAY  
AREKVSLFPGKRSGIIDVIPVDLVANSIILS LAEALSGSGQRRIYQCCSGGSNPISLGKF  
IDYLMAEAKTNYAAYDQLFYRRPTKPFVAVNRKLF DVVVGGMRVPLSIAGKAMRLAGQNR  
ELKVLKNLDTTRSLATIFGFYTAPDYIFRNDSL MALASRMGELDRVLF PVDARQIDWQLY  
LCKIHGLGLNRYALKERKLYSLRAADTRKAA

## FIG. 7

FAR peptide (SEQ ID NOs: 6 and 8) from codon optimized DNA (SEQ ID NOs: 5 and 7)

MSQYSAFSVSQSLKGKHIFLTGVTGFLGKAILEKLLYSVPQLAQIHIILVRGGKVSACKRF  
QHDILGSSIFERLKEQHGEHFEEWVQSKINLVEGELTQPMFDLPSAEFAGLANQLDLIIN  
SAASVNFRNLEKALNINTLCNNIIALAQNVAATPVMQISTCYVNGFNKGQINEEVV  
GPASGLIPQLSQDCYDIDSVFKRVHSQIEQVKKRKTDIEQQEQALIKLGIKTSQHFGWND  
TYTFTKWLGEQLLIQKLGKQSLTILRPSIIESAVREPAPGWEGVKVADALIYAYAKGRV  
SIFPPGRDEGILDVIPVDLVANAAALSAAQLMESNQQTGYRIYQCCSGSRNPIKLKEFIRH  
IQNVAQARYQEWPKLFADKPQEAFKTVSPKRFKLYMSGFTAITWAKTIIGRVFGSNAASQ  
HMLKAKTTASLANIFGFYTAPNYRFSSQKLEQLVKQFDTTEQRPLYDIRADHFDWKYYLQE  
VHMDGLHKYALADRQELPKHVKKRKRETIRQAA

## FIG. 8

*B. mori* FAR DNA (SEQ ID NO: 9)

ATGTCCCACAATGGTACTCTTGATGAGCATTATCAGACAGTCGTGAATTTATGACGGT  
AAGAGCGTTTATTACTGGCGCCACTGGCTTCTGGAAAAGCCTACGTTGAGAAACTG  
GCATATTCTTGTCCCGAATTGTCTCTATTATTTGATTGATTCGTATAAAAAGGGCTCC  
AACACGGAGGAGCGTATGCGTAAATATTGGACCAACCCATTTCTCTCGTATTAAATAT  
GAGCATCCAGAGTACTTCAAAAAGATTATCCCCATTCTGGCGATATTACGCCCTAAA  
CTTGGTCTTGCGACGAGGAGCGTAACATCCTGATTAATGAAGTGTCCATCGTTATCCAC  
TCTGCTGCCAGCGTTAAACTGAACGATCATCTTAAATTCACCTTGAACACCAACGTTGGT  
GGTACTATGAAAGTCTTGGAGCTTGTAAAGGAGATGAAAAATTGGCGATGTTGTGTAC  
GTTTCTACGGCTTATTCTAACACCAGCCAACGTATTTGGAAGAAAAATTGTACCCTCAG  
TCTCTGAATTGAACGAAATTCAAAGGCTGAAGAGCACTACATTCTGGGTAAAGGAT  
AACGACGAAATGATTAAATTCAATTGGAAACCACCTAACACTACGCTACACGAAAGCC  
CTGGCTGAGAATCTTGTGCTGAAGAACATGGAGAAATTCTACTATTATTATTCGTCT  
TCTATCATCACGCCTCCGCCAGGAACCCGTTCTGGATTGTTGATTCTGGAGCGGA  
GCCACGGCTATGGCTGCATTGCACTTAAAGGCTGGAACACATCATGTATTCCACCGGT  
GAAGAGAACATTGACTTGATCCCCCTGGATTACGTTGTCAACTTGACACTGGTGGCTATT  
GCTAAATATAAGCCAACGAAAGAGGTTACGGTGTACCATGTTACAACGAGCGACTTGAAC  
CCGATTAGCATCCGTCGTATTTCATCAAACGTGAGCGAGTTCGCCTCAAAAACCCAAC  
AGCAACGCTGCCCTTCGCTGCCACTACATTGCTGACCAACAGAAACCGCTTATTAAA  
CTGGTGACATTCTTATGCAGACCACACCAGCTTTGGCAGACTTGTGGATGAAACG  
CAGCGTAAAGAGGCCAAGTTCTGAAACACAGCACAACACTGGTCGTTCTCGTGTAC  
CTGGAGTTCTTACATCTCAGAGCTGGCTTCTGGCTGTGAGCGTGCACGTGTCCCTGTCC  
GCAGCCTTGTCGATTCCGACCGTGTCTCCGTTGCGATCCTCCACGATTGATTGG  
GATCAGTATCTCCTATCTACTTCGAAGGTATTAATAAACACCTGTTCAAAAATAAGTTG  
TAG

FIG. 9

*B. mori* FAR peptide (SEQ ID NO: 10)

MSHNGTLDEHYQTVREFYDGKSVFITGATGFLGKAYVEKLAYSCPGIVSIYILIRDKKGS  
NTEERMRKYLDQPIFSRIKYEHPEYFKKIIPISGDITAPKLGLCDEERNILINEVSIVIH  
SAASVKLNDHLKFTLNTNVGGTMKVLELVKEMKNLAMFVYVSTAYSNTSQRILEEKLYPQ  
SLNLNEIQKFAEEHYILGKDNDENMIKFIGNHPNTYAYTKALAENLVAEEHGEIPTIIIRP  
SIITASAEPPVRGFVDSWSGATAMAFAALKGWNNIMYSTGEENIDLIPLDYVVNLTVAI  
AKYKPTKEVTVYHVTTSDLNPISIRRIFIKLSEFASKNPTSNAAPFAATTLLTKQKPLIK  
LVTFLMQTTPAFLADLWMKTQRKEAKFVKQHNLVVRSRDQLEFFTSQSWLLRCERARVLS  
AALSDSDRAVFRCDPSTIDWDQYLPIYFEGINKHLFKNKL

FIG. 10

Codon optimized *B. mori* FAR DNA (SEQ ID NO: 11)

```
ATGTCTCACAAACGGAACCTTGACGAGCATTACCAAACGTGCGAGAGTTCTACGACGGCAAATC  
CGTGGTCATCACCGGAGCCACTGGATTCTGGTAAGGCATACGTGGAGAAGCTCGCTTACAGCT  
GTCCCGGTATCGTTCTATCTACATCCTGATTAGAGATAAGAAGGGCTCCAACACAGAAGAGCGG  
ATGCGAAAGTACCTGGACCAGCCCCTCTCTCCCGGATCAAGTACGAACACCCCTGAGTACTTCAA  
GAAGATCATCCCCATTAGCGGAGACATTACTGCCCAAAGCTGGGTTGTGACGAGGAGCGGA  
ACATCCTGATCAACGAGGTGTCCATCGTATTCTCGCCGCTCTGTTAACGTAACGACCAAC  
CTGAAGTTCACCTGAACACTAACGTGGGTGGAACGATGAAGGTCTGGAGCTGGTAAGGAGAT  
GAAGAACCTGGCATGTTCGTGTACGTGTACCGCTTACTCCAACACCTCTCAGAGAACCTGG  
AGGAGAACGCTGTACCCCTCAGTCCCTGAACCTGAACGAGATCCAGAACGTTCGCCGAGGAACACTAC  
ATCCTGGCAAGGACAACGACGAGATGATCAAGTTCATCGGAACCAACCCCCAACACCTACGCTTA  
CACCAAAGCCCTGGCTGAGAACCTTGTTGCCAAGAGACATGGAGAGATCCCCACCATCATCATCC  
GACCCCTCTATCATTACTGCATCCCGGAGGAGCCTGTGCGAGGCTCGTCGATTCTGGTCTGGC  
GCTACTGCTATGGCTGCTTCGCTCTCAAGGGATGGAACAACATCATGTATTCCACCGCGAAGA  
GAACATTGACCTGATCCCTCTCGACTACGTGGTAACCTTACCCCTGGCATCGCTAACGTACA  
AGCCTACCAAAGAGGTGACCGTGACCGACTACGACTTCGGATCTGAATCCATCTCCATCCGG  
CGAATCTTCATCAAGCTGTCTGAGTTGCTTCTAAGAACCCGACTTCTAATGCTGCTCCTTCGC  
TGCCACTACTCTGCTTACCAAGCAGAAACCCCTGATCAAGCTGGTTACGTTCTGATGCAAACCA  
CTCCTGCCTTCCTCGCTGACCTGTGGATGAGAACCCAGCGAAAGGAGGCCAAGTTCGTCAAACAG  
CATAACCTCGTCGTTAGATCGCGAGATCAGTTGGAGTTCTTACCTCCCAGTCCTGGCTGCTTCG  
TTGTGAAAGAGGCCAGAGTGCTGTCGCTGCTCTTAGCGACTCTGATCGTGGCTGTTAGATGTG  
ACCCCTCGACAATCGACTGGATCAGTACCTGCCATCTACTTCGAGGGCATCAACAAGCACCTG  
TTCAAGAACAAAGCTC
```

FIG. 11

Codon optimized *Marinobacter aquaeolei* FAR DNA (SEQ ID NO: 13)


```
ATGGCTATCCAGCAGGTTCATCACGCCACACATCCTCCTCTAAAGTCTGGGTCAACTT
CGTGGTAAACGTGTCCTGATTACCGGCACTACTGGATTCTGGTAAAGTCGTCTGGAA
CGTTGATTCGTGCGCTCCTGACATCGGTGCTATCTACCTGCTGATTGTAACAAG
CGTCACCCGGATGCTCGTCTCGTTCTGGAGGAGATTGCTACCTCCTGTCTTGT
CGTTGCGTGAAGCTGATTCCGAAGGTTCGATGCTTCCTGGAAGAACGTATTCACTGT
GTTACTGGTGAAGTTACTGAAGCTGGTTGGTATTGGTCAAGAGGACTATCGTAAGTTG
GCCACCGAATTGGACGCAGTCATCAATTCTGCTGCCTCCGTCAACTCCGTGAGGAGTTG
GATAAGGCTCTGCCATCAACACTCTGTGTTGCGTAACATCGCTGGTATGGTGGATCTT
AACCTAAGCTGGCGTCTTCAGTCTACGTGTTACGTCAACGGTATGAACACTCTGGT
CAAGTTACTGAATCCGTCAAAACCAGCTGGTGAAGCTGTTCCCTGTTCTCCTGATGGA
TTCTACGAGATCGAGGAATTGGTTCGTCTGCTGCAAGACAAGATTGAAGACGTTCAAGCA
CGTTACTCTGGTAAGGTGTTGGAGCGTAAGTGGTTGATTGGTATTGGTATTGGCTAAT
CGTTACGGTTGGTCTGATACATACACCTCACGAAATGGTGGGTAAACAACCTCTGATG
AAAGCCTTGAATGGTGTACCTGACTATTCTGCGTCTAGCATCATTGAATCTGTTG
GAAGAACCGCACCTGGTGGATTGAAGGCGTGAAGGTTACTACGTTCTGGTATCATTGATGTGATT
TATGCTCGTGAAGGTTACTTGTGTTCCGGTAAACGTTCTGGTATCATTGATGTGATT
CCTGGTACTGGTTGCCAATTCCATCATCTGTCTTGGCTGAGGCTCTGGCGAACCT
GGTCGTCGTATCTACCAATGTTGTTGGTGGTAATCCTATCTCCCTGGCGAACCT
TTCATTGATCACCTGATGGCTGAATCCAAAGCCAACATGCCGCATACGATCATCTGTTG
TACCGTCAACCCCTCCAAGCCTTCCTGCTGTCAACCGTGCTTGTGACTTGGTTATC
TCTGGTGTCCGTCTGCCCTTGTCTTGACCGACCGTGTCTGAAGCTGCTGGCAACTCC
CGTACCTGAAGATGCTGCGTAACCTGGATACTACGCAATCCCTGGCTACTATCTTGGC
TTCTACACAGCCCCCGACTACATCTCCGTAATGACGAGTTGATGGCCCTGGCTAACCGT
ATGGCGAGGTTGATAAGGGTTGTTCCCCGGTGTGCTCGTCTGATTGATTGGGAATTG
TACCTGCGTAAGATTCACCTGGCTGGTTGAACCGTACGCCTGAAGGAGCGTAAGGTT
TACTCTTGAAGACAGCCCGTCAGCGTAAGAAGGAGCAGCTTAA
```

FIG. 12

*Marinobacter aquaeolei* FAR peptide (SEQ ID NO: 14)

MAIQQVHHADTSSSKVLGQLRGKRVLITGTTGFLGVVLERLIRAVPDIGAIYLLIRGNK  
RHPDARSRFLEEIAATSSVFDRRLREADSEGFDAFLEERIHCVTGEVTEAGFGIGQEDYRKL  
ATELDAVINSAASVNFREELDKALAINTLCLRNIAGMVDLNPKLAVLQVSTCYVNGMNSG  
QVTESVIKPAGEAVPRSPDGFYEIEELVRLLQDKIEDVQARYSGKVLERKLVDLGIREAN  
RYGWSDTYTFTKWLGEQLLMKALNGRTLTILRPSIIESALEEPAPGWIEGVKVADAIILA  
YAREKVTLFPGKRSIIIDVIPVDLVANSIILS LAEALGEPEGRRRIYQCCSGGGNPISLGE  
FIDHLMAESKANYAAYDHLFYRQPSKPFLAVNRALFDLVISGVRLPLSLDRVLIKLLGNS  
RDLKMLRNLDTTQSLATIFGFYTAPDYIFRNDELMALANRMGEVDKGLFPVDARLIDWEL  
YLRKIHLAGLNRYALKERKVYSLKTAQRKKAA

FIG. 13



## PRODUCTION OF FATTY ALCOHOLS WITH FATTY ALCOHOL FORMING ACYL-COA REDUCTASES (FAR)

### CROSS REFERENCE TO RELATED APPLICATIONS

[0001] This application claims benefit of provisional application No. 61/221,934 filed Jun. 30, 2009 and provisional application No. 61/315,380 filed Mar. 18, 2010, the entire contents of which are incorporated herein by reference.

### REFERENCE TO SEQUENCE LISTING

[0002] The "Sequence Listing" submitted electronically concurrently herewith pursuant to 37 C.F.R. §1.821 in a computer readable form (CRF) via file name CX3-017US-ST25.txt is incorporated herein by reference. The electronic copy of the Sequence Listing was created Jun. 28, 2010, and the size on disk is 82 Kbytes.

### BACKGROUND

[0003] Diesel fuel is any fuel used in diesel engines and includes both petrodiesel and biodiesel. Petrodiesel is a specific fractional distillate of fossil fuel oil. It is comprised of about 75% saturated hydrocarbons and 25% aromatic hydrocarbons. Biodiesel is not derived from petroleum but from vegetable oil or animal fats and contains long chain alkyl esters. Biodiesel is made by the transesterification of lipids (e.g., spent vegetable oil from fryers or seed oils) with an alcohol and burns cleaner than petrodiesel. Biodiesel can be used alone or mixed with petrodiesel in any amount for use in modern engines.

[0004] Kerosene is a combustible hydrocarbon that is also a specific fractional distillate of fossil fuel and contains hydrocarbons having 6 to 16 carbon atoms. Kerosene has a heat of combustion comparable to that of petrodiesel and is widely used in jet fuel to power jet engines and for heating in certain countries. Kerosene-based fuels can also be burned with liquid oxygen and used as rocket fuel (e.g., RP-1).

[0005] Reliance on petroleum-derived fuels has depleted the supply of natural resources and has required increased reliance on imported gasoline and diesel products. In addition, the burning of petroleum-based fuels has increased the amount of greenhouse gasses (e.g., carbon dioxide and methane) in the atmosphere that is contributing to the gradual warming of the earth's climate.

[0006] Fuels, such as biodiesel, that are made from animal or vegetable products burn cleaner than petroleum-derived fuels and do not produce a net increase in greenhouse gases. Furthermore, they are a sustainable energy source and have the potential to reduce the United States' reliance on imported petroleum-based products. However, there is a concern that using land to produce fuel crops rather than food crops will contribute to world hunger.

[0007] Fatty acids are the principle component of cell membranes and are used by nearly all organisms as a primary source of energy storage. Fatty alcohols are the reduction products of fatty acids and, like fatty acids, can be produced enzymatically by cultured cells. Fatty alcohols can be reacted with acids to form ester compositions similar to those present in biodiesel fuel, or reduced to form kerosene-like compositions, or hydrocarbon compositions similar to petrodiesel. Enzymes that convert fatty acyl-thioester substrates (e.g., fatty acyl-CoA or fatty acyl-ACP) to fatty alcohols are com-

monly referred to as fatty alcohol forming acyl-CoA reductases or fatty acyl reductases ("FARs").

[0008] PCT Publication No. WO 2007/136762 discloses genetically engineered microorganisms for the production of fatty acid derivatives and methods of their use.

[0009] PCT Publication No. WO 2009/140695 discloses compositions comprising cyanobacterial genes encoding enzymes involved in hydrocarbon biosynthesis and methods of using them in the production of aldehydes and alcohols.

[0010] Steen et al., 2010, *Nature* 463:559-563 discloses the engineering of *E. coli* to produce specific fatty esters, fatty alcohols and waxes directly from simple sugars.

[0011] Schirmer, 2009, *Current Opinion in Microbiology* 12:274-281 describes the fermentative and nonfermentative metabolism of heterotrophic microorganisms used in the production of fuel-like molecules, biocatalysts that convert metabolic intermediates into fuel-like molecules, and the parameters that govern the cost effective production of such fuel-like molecules.

[0012] U.S. Pat. No. 5,370,996 and Metz et al. (2000) *Plant Physiology* 122:635-644 disclose isolation and characterization of a fatty acyl reductase (FAR) enzyme from the desert shrub *Simmondsia chinensis*, more commonly known as jojoba.

[0013] Moto et al. (2003) *Proc. Nat'l Acad. Sci. USA* 100 (16):9156-9161 discloses the isolation and characterization of a FAR enzyme from the silk moth *Bombyx mori*.

[0014] Reiser et al. (1997) *J. Bacteriol.* 179(9):2969-2975 discloses the isolation and characterization of a fatty acyl CoA reductase enzyme from the wax ester producing bacterium *Acinetobacter calcoaceticus* that reduces a fatty acyl-CoA substrate with chain lengths from C14 to C22 to the corresponding fatty aldehyde, requiring a dehydrogenase enzyme for conversion of the fatty aldehyde to the fatty alcohol.

[0015] In theory, these FAR enzymes could be expressed in heterologous hosts as a means of producing a non-petroleum-based, renewable source of fatty alcohols or derivative compositions for use in biofuels. However, when expressed in heterologous hosts such as *E. coli* and yeast, the yields of fatty alcohols obtained have been insufficient for certain applications. In addition, at most, only a small fraction of the produced fatty alcohols are secreted by the microorganisms, increasing substantially the cost of purification.

[0016] Accordingly, there remains a need in the art for enzymes such as FAR enzymes that can be used to efficiently produce fatty alcohols for use in industrial applications such as, but not limited to applications in the food industry, cosmetic industry, medical industry, and fuels industry.

### SUMMARY

[0017] The present disclosure provides isolated fatty acyl reductase (FAR) enzymes, nucleic acids encoding the FAR enzymes, recombinant microorganisms engineered to express the FAR enzymes, methods of producing fatty alcohols and other compositions utilizing the FAR enzymes and/or recombinant microorganisms, and compositions comprising and/or that are derived from the fatty alcohols. The various inventions described herein are based, in part, on the inventors' surprising discovery that certain genera of the class of gammaproteobacteria found in seawater contain FAR enzymes that are capable of generating high yields of total and secreted fatty alcohols when expressed in heterologous cells.

[0018] In particular, the present inventors have discovered that *Marinobacter* species *algicola* (strain DG893) ("FAR\_Maa"), *Marinobacter aquaeolei* VT8 ("FAR\_Maq"), and *Oceanobacter* species strain RED65 (recently reclassified as *Bermanella marisrubri* RED65) ("FAR\_Ocs") possess FAR enzymes capable of producing approximately 150- to 400-times more total fatty alcohol, respectively, than FAR enzymes from *B. mori*, when expressed in an *E. coli* host. Significantly greater yields of fatty alcohols are also achieved in yeast. Based on this surprising discovery, it is expected that other genera of marine bacteria and particularly marine gammaproteobacteria will likewise contain FAR enzymes that are especially suitable for the production of fatty alcohols, especially when expressed in heterologous host microorganisms. In certain embodiments, FAR enzymes that are especially suitable for the production of fatty alcohols may be identified using Hidden Markov Models ("HMMs"), which identify proteins by similarity to patterns found in pre-complied sets of protein sequences. The patterns inherent in each pre-complied set are encapsulated in HMMs. It is the HMM for a given sequence set that defines a protein domain or protein family. Using this technique it is possible to classify portions of a given protein sequence that share common domains with previously-identified suitable FAR enzymes. See, e.g., <http://pfam.sanger.ac.uk/>. In certain embodiments, the HMMs are used to identify NAD binding domains and/or sterile domains.

[0019] Accordingly, in one aspect the present disclosure provides isolated FAR enzymes derived from or obtained from marine bacteria, such as marine gammaproteobacteria, and/or functional fragments thereof having FAR activity. In some embodiments, the FAR enzyme is isolated directly from the marine gammaproteobacteria in which it occurs naturally. In some embodiments, the FAR enzyme and/or a function fragment is isolated from a heterologous host microorganism engineered to express the FAR enzyme or functional fragment.

[0020] In another aspect, the present invention relates to a FAR enzyme and functional fragment thereof derived or obtained from *M. algicola* strain DG893. In one embodiment, the FAR enzyme or functional fragment thereof has an amino acid sequence that is at least 30% identical, at least 75% identical, at least 80% identical and at least 90% identical to SEQ ID NO: 2 or a functional fragment thereof. In another specific embodiment, the isolated FAR enzyme has an amino acid sequence that is identical to SEQ ID NO: 2.

[0021] In a further aspect, the present invention relates to a FAR enzyme and functional fragment thereof derived or obtained from a strain of *Oceanobacter*. In one specific embodiment, the FAR enzyme is obtained or derived from *Oceanobacter* strain RED65 and has an amino acid sequence that is at least 30% identical, at least 75% identical, at least 80% identical, and at least 90% identical to SEQ ID NO: 6. In another specific embodiment, the isolated FAR enzyme has an amino acid sequence that is identical to SEQ ID NO: 6.

[0022] In an additional aspect, the present disclosure provides polynucleotides encoding a marine gammaproteobacterial FAR enzyme, or functional fragment thereof, as described above. The polynucleotide can be a DNA or RNA, and can be single-stranded or double-stranded. It can be isolated from a naturally occurring microorganism, or prepared wholly or partially via synthetic means. In one embodiment, the nucleic acid sequence encodes a fatty acyl-CoA reductase (FAR) enzyme comprising an amino acid sequence having at

least 30% sequence identity to SEQ ID NO: 2, SEQ ID NO: 6, SEQ ID NO: 14 or functional fragments thereof. In other embodiments, the nucleic acid encodes a FAR enzyme comprising at least 75% sequence identity (that is at least 75%, at least 80%, at least 90%, at least 95% and even 100% sequence identity) to SEQ ID NO: 2, SEQ ID NO: 6, SEQ ID NO: 14 or a functional fragment thereof. In some embodiments of this aspect, a vector comprises the nucleic acid encoding the FAR enzyme. In some embodiments, the polynucleotide encoding the FAR enzyme or functional fragment thereof is operably linked to a promoter, and optionally, to other control sequences and elements that control expression. In some embodiments, the vector is designed or selected so as to remain distinct from the genome of the host microorganism (for example, a plasmid) or to integrate into the genome of the host microorganism for example by either homologous recombination or site-specific integration. In a specific embodiment, the codons of the portion of the construct encoding the FAR enzyme or functional fragment are optimized for expression in a particular host microorganism.

[0023] In one more aspect, the present disclosure provides recombinant microorganisms engineered to express a heterologous marine gammaproteobacterial FAR gene, or a functional fragment thereof, as described herein. In some embodiments, recombinant host cells include bacteria, filamentous fungi and yeast. In particular embodiments, bacteria are *E. coli*. In other particular embodiments, the yeast is an oleaginous yeast. In certain embodiments, the oleaginous yeast is *Y. lipolytica*. In still other particular embodiments, the yeast is *S. cerevisiae*. In certain aspects, the microorganisms are wild-type microorganisms. In other aspects, the microorganisms are genetically modified.

[0024] In certain specific embodiments, the genetically modified microorganism further over expresses a gene encoding a protein that increases the rate at which the microorganism produces the substrate of a FAR enzyme, i.e., a fatty acyl-thioester substrate. In certain embodiments, the enzyme encoded by the over expressed gene is directly involved in fatty acid synthesis. In particular embodiments, the enzyme encoded by the over expressed gene is selected from a fatty acid synthase, an acyl-ACP thioesterase, a fatty acyl-CoA synthase and an acetyl-CoA carboxylase. In some embodiments, the protein encoded by the over expressed gene is endogenous to the microorganism. In other embodiments, the protein encoded by the over expressed gene is heterologous to the microorganism.

[0025] In yet another aspect, the present disclosure provides methods of producing compositions comprising fatty alcohols and/or acid esters and/or alkanes and/or alkenes derived from the fatty alcohols. In certain embodiments, the fatty alcohols are produced in a cell-free system where a FAR enzyme as described above is provided with a fatty acyl-thioester substrate and necessary cofactors under suitable conditions of temperature, pH and ionic strength for a time sufficient to produce fatty alcohols. In various embodiments, fatty alcohols are produced by a recombinant host cell comprising a heterologous gene encoding a FAR enzyme described above. In these embodiments, the host cell is cultured in an aqueous nutrient medium comprising an assimilable carbon source under conditions suitable for production of fatty alcohols. In certain specific embodiments, the recombinant host cell is a bacterium or yeast. In various embodi-

ments, the yeast is an oleaginous yeast. In particular embodiments, the bacterium is *E. coli* and the yeast is selected from *S. cerevisiae* and *Y. lipolytica*.

[0026] In some embodiments, the fatty alcohols are produced by a recombinant host cell comprising a heterologous gene encoding a FAR enzyme and one or more proteins that increase the rate at which the recombinant host cell produces the acyl-thioester substrate for FAR. In some embodiments, the one or more proteins are directly involved in fatty acid biosynthesis. In certain particular embodiments, the one or more proteins are selected from a fatty acid synthase, an acyl-ACP thioesterase, a fatty acyl-CoA synthase and an acetyl-CoA carboxylase. In some aspects, the fatty alcohols produced by the recombinant microorganisms are secreted from the cells into the aqueous nutrient medium. In certain embodiments, the methods described herein include a step of isolating the fatty alcohols.

[0027] In an additional aspect, the invention relates to a method of producing a fatty alcohol composition comprising culturing a recombinant microorganism in a suitable culture medium wherein the recombinant microorganism comprises a gene encoding a heterologous fatty acyl reductase (FAR) enzyme having at least 30% sequence identity to SEQ ID NO: 2, SEQ ID NO: 6, SEQ ID NO: 14 or a functional fragment thereof and allowing expression of said gene wherein said expression results in the production of a composition of fatty alcohols. In some embodiments of this aspect, the method further comprises isolating the produced fatty alcohols. In particular embodiments, the methods described herein produce at least about 0.5 g/L of fatty alcohols, also at least 5 g/L and even at least 15 g/L of fatty alcohols.

[0028] In yet another aspect, the invention relates to a method of producing a fatty alcohol composition by an acyl-CoA independent pathway comprising culturing a recombinant *E. coli* in a nutrient medium under suitable conditions, wherein said recombinant *E. coli* comprises a nucleic acid sequence encoding a heterologous fatty acyl-CoA reductase (FAR) enzyme comprising an amino acid sequence having at least 75% (such as at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 97% and even 100%) sequence identity to SEQ ID NO: 2, SEQ ID NO: 6, SEQ ID NO: 14 or functional fragments thereof; allowing expression of said FAR; and producing a fatty alcohol composition.

[0029] In yet a further aspect, the recombinant microorganism engineered to express a FAR encompassed by the invention is a bacterium and particularly an *E. coli* comprising an inactivated or silenced endogenous fatty acyl-CoA synthetase fadD gene. In other embodiments, the recombinant *E. coli* will comprise an inactivated or silenced endogenous short chain fatty acyl-CoA synthetase fadK gene. In still other embodiments, both genes will be inactivated or silenced.

[0030] In yet another aspect, the present disclosure provides for fatty alcohol compositions produced by a method described above. In some embodiments, the fatty alcohol composition is produced in a cell-free system. In other embodiments, the fatty alcohol composition is produced by a recombinant host cell. In particular embodiments, the compositions comprise fatty alcohols with saturated, unsaturated, linear, branched or cyclic hydrocarbon chains. In certain embodiments, the compositions comprise saturated fatty alcohols and/or monounsaturated fatty alcohols.

[0031] In still other aspects, the present disclosure provides for compositions comprising alkanes and/or alkenes derived by reducing the fatty alcohol compositions described herein.

In certain embodiments, the reduction reaction is carried out chemically. In other embodiments, the reduction reaction is carried out using a microorganism that has a biosynthetic pathway for reducing fatty alcohols. In certain embodiments, the compositions produced by reduction of fatty alcohols comprise substantially all alkanes and/or alkenes. In other embodiments, the compositions comprise a mixture of fatty alcohols and alkanes and/or alkenes derived there from.

[0032] In yet another aspect, the present disclosure provides for fuel compositions comprising fatty alcohols and/or acid esters and/or alkanes and/or alkenes derived there from. In certain embodiments, the fatty alcohol compositions produced by the methods described above are used directly in fuel compositions. In various embodiments, the fatty alcohols are reacted with a carboxylic acid to produce acid esters for use as biodiesel fuel compositions. In other embodiments, the fatty alcohols are reacted with a reducing agent to produce alkanes and/or alkenes. In certain embodiments, the fatty alcohols undergo esterification to form fatty esters. In some embodiments, the alkanes and/or alkenes derived from the fatty alcohol compositions are used as components of jet fuel compositions. In other embodiments, the alkanes and/or alkenes derived are used as components of rocket fuel. In still other embodiments, alkanes and/or alkenes derived from the fatty alcohol compositions are used as components of petro-diesel-like fuel compositions.

#### BRIEF DESCRIPTION OF THE FIGURES

[0033] FIG. 1 shows the replicative *Y. lipolytica* vector pCEN411 (approximately 9065 bp) containing cassettes encoding hygromycin (HygB) resistance and FAR expression. Ars18 is an autonomous replicating sequence isolated from *Y. lipolytica* chromosomal DNA.

[0034] FIG. 2 shows a codon optimized polynucleotide sequence (SEQ ID NO: 1) encoding a FAR enzyme of *Marinobacter algicola* strain DG893 having the amino acid sequence shown in FIG. 6 (SEQ ID NO: 2).

[0035] FIG. 3 shows a codon optimized polynucleotide sequence (SEQ ID NO: 3) encoding a FAR enzyme of *Marinobacter algicola* strain DG893 having the amino acid sequence shown in FIG. 6 (SEQ ID NO: 4) and optimized for expression in *Y. lipolytica*.

[0036] FIG. 4 shows a codon optimized polynucleotide sequence (SEQ ID NO: 5) encoding a FAR enzyme of *Oceanobacter* sp. strain RED65 having the amino acid sequence shown in FIG. 7 (SEQ ID NO: 6).

[0037] FIG. 5 shows a codon optimized polynucleotide sequence (SEQ ID NO: 7) encoding a FAR enzyme of *Oceanobacter* sp. strain RED65 having the amino acid sequence shown in FIG. 7 (SEQ ID NO: 8) and optimized for expression in *Y. lipolytica*.

[0038] FIG. 6 shows the amino acid sequence of a FAR polypeptide of *Marinobacter algicola* strain DG893. The amino acid sequence (SEQ ID NO: 2) encoded by the polynucleotide sequence of SEQ ID NO: 1 is identical to the amino acid sequence (SEQ ID NO: 4) encoded by the polynucleotide sequence of SEQ ID NO: 3. While it is understood that both SEQ ID NO: 2 and SEQ ID NO: 4 are identical and interchangeable, reference will generally be made to SEQ ID NO: 2 in this disclosure.

[0039] FIG. 7 shows the amino acid sequence of a FAR polypeptide of *Oceanobacter* sp. strain RED65. The amino acid sequence (SEQ ID NO: 6) encoded by the polynucleotide sequence of SEQ ID NO: 5 is identical to the amino acid

sequence (SEQ ID NO: 8) encoded by the polynucleotide sequence of SEQ ID NO: 7. While it is understood that both SEQ ID NO: 6 and SEQ ID NO: 8 are identical and interchangeable reference will generally be made to SEQ ID NO: 6 in this disclosure. The amino acid sequence of SEQ ID NO: 2 and the amino acid sequence of SEQ ID NO: 6 are about 47% identical.

[0040] FIG. 8 shows a codon optimized polynucleotide sequence (SEQ ID NO: 9) encoding a FAR polypeptide (SEQ ID NO: 10) from the silk moth *B. mori*.

[0041] FIG. 9 shows the amino acid sequence (SEQ ID NO: 10 and SEQ ID NO: 12) of a FAR polypeptide from the silk moth *B. mori*.

[0042] FIG. 10 shows a codon optimized polynucleotide sequence (SEQ ID NO: 11) encoding a FAR enzyme of *B. mori* having the amino acid sequence shown in SEQ ID NO: 12 and optimized for expression in *Y. lipolytica*.

[0043] FIG. 11 shows a codon optimized polynucleotide sequence (SEQ ID NO: 13) encoding a FAR enzyme of *Marinobacter aquaeolei* having the amino acid sequence shown in SEQ ID NO: 14.

[0044] FIG. 12 shows the amino acid sequence of a FAR polypeptide of *Marinobacter aquaeolei* (SEQ ID NO: 14). The amino acid sequence of *M. aquaeolei* (SEQ ID NO: 14) has about 78% sequence identity to the amino acid sequence of *M. algicola* DG893 (SEQ ID NO: 2).

[0045] FIG. 13 depicts biosynthetic pathways for fatty alcohol production via a) an acyl-CoA dependent pathway utilizing fatty acyl-ACP and fatty acyl-CoA intermediates and b) an acyl-CoA independent pathway utilizing fatty acyl-ACP intermediates but not a fatty acyl-CoA intermediate, wherein "R" as used in the compound formulas is a C8 to C24 saturated, unsaturated, linear, branched or cyclic hydrocarbon.

## DETAILED DESCRIPTION


[0046] The present disclosure provides FAR enzymes, recombinant host cells comprising a nucleic acid encoding a heterologous FAR enzyme, processes for the biosynthesis of fatty alcohols including the conversion of fatty acyl-thioester complexes (such as fatty acyl-CoA substrates and fatty acyl-ACP substrates) to fatty alcohols, and compositions comprising fatty alcohols produced by the biosynthetic methods.

## DEFINITIONS

[0047] Unless defined otherwise, all technical and scientific terms used herein generally have the same meaning as commonly understood by one of ordinary skill in the art to which this invention pertains. Generally, the nomenclature used herein and the laboratory procedures of cell culture, molecular genetics, organic chemistry, analytical chemistry and nucleic acid chemistry described below are those well known and commonly employed in the art.

[0048] As used herein, the following terms are intended to have the following meanings: The following abbreviations are used herein: "FAR" denotes fatty acyl reductase or fatty alcohol forming acyl-CoA reductase; "ACP" denotes acyl carrier protein; "CoA" denotes coenzyme A; "TE" denotes thioesterase; "FAS" denotes fatty acid synthase; "FACR" denotes fatty acyl-CoA reductase; "FACS" denotes fatty acyl-CoA synthase (synthetase) and acyl-CoA synthase (synthetase) as used interchangeably herein; and "ACC" denotes acetyl-CoA carboxylase.

[0049] "Fatty alcohol forming acyl-CoA reductase" or "fatty acyl reductase" as used interchangeably herein refers to an enzyme that catalyzes the reduction of a fatty acyl-CoA, a fatty acyl-ACP, or other fatty acyl thioester complex to a fatty alcohol, which is linked to the oxidation of NAD(P)H to NAD(P)<sup>+</sup>, as shown in Scheme I:



wherein "R" represents a C8 to C24 saturated, unsaturated, linear, branched or cyclic hydrocarbon, and "R<sub>1</sub>" represents CoA, ACP or other fatty acyl thioester substrates. CoA is a non-protein acyl carrier group factor (or moiety) involved in the synthesis and oxidation of fatty acids. "ACP" is a polypeptide or protein subunit of fatty acid synthase used in the synthesis of fatty acids. FARs are distinct from FACRs. FACRs reduce only fatty acyl-CoA intermediates to fatty aldehydes and require an additional oxidoreductase enzyme to generate the corresponding fatty alcohol. "Fatty aldehyde" as used herein refers to a saturated or unsaturated aliphatic aldehyde and reference is made to FIG. 13, wherein R is as defined above.

[0050] The term "fatty acid" as used herein refers to a compound of formula III:



wherein "R" is as defined above. Saturated or unsaturated fatty acids can be described as "Ca:b", wherein "a" is an integer that represents the total number of carbon atoms and "b" is an integer that refers to the number of double bonds in the carbon chain.

[0051] The term "fatty alcohol" as used herein refers to an aliphatic alcohol of the formula (II) wherein R is as defined above. Saturated or unsaturated fatty alcohols can be described as "Ca:b-OH", wherein "a" is an integer that represents the total number of carbon atoms in the fatty alcohol and "b" is an integer that refers to the number of double bonds in carbon chain.

[0052] Unsaturated fatty acids or fatty alcohols can be referred to as "cis Δ<sup>x</sup>" or "trans Δ<sup>x</sup>" wherein "cis" and "trans" refer to the carbon chain configuration around the double bond and "x" indicates the number of the first carbon of the double bond, wherein carbon 1 is the carboxylic acid carbon of the fatty acid or the carbon bound to the —OH group of the fatty alcohol.

[0053] The term "fatty acyl-thioester" or "fatty acyl-thioester complex" refers to a compound of formula (I), wherein a fatty acyl moiety is covalently linked via a thioester linkage to a carrier moiety. Fatty acyl-thioesters are substrates for the FAR enzymes described herein.

[0054] The term "fatty acyl-CoA" refers to a compound of formula (I) wherein R<sub>1</sub> is Coenzyme A.

[0055] The term "fatty acyl-ACP" refers to a compound of formula (I) wherein R<sub>1</sub> is acyl carrier protein.

[0056] The phrase "acyl-CoA independent pathway" refers to the production of fatty alcohols by the direct enzymatic conversion of fatty acyl-ACP substrates to fatty alcohols and does not involve the use of free fatty acids or fatty acyl-CoA intermediates. This biosynthetic pathway differs from a) the fatty acyl-CoA dependent pathway which converts fatty acyl-ACP directly to fatty acyl CoA via an acyl-transfer reaction, such as in yeast, and b) the fatty acyl-CoA dependent pathway which converts fatty acyl-ACP to fatty acyl-CoA via a free fatty acid intermediate, such as in bacteria and reference is made to FIG. 13.

[0057] The acyl-CoA independent pathway has the advantage of bypassing the step of form a fatty acyl-CoA substrate from free fatty acid, which requires the use of ATP. Therefore, the acyl-CoA independent pathway may use less energy than the acyl-CoA dependent pathway that utilizes a free fatty acid intermediate.

[0058] "Conversion" refers to the enzymatic conversion of the substrate to the corresponding product. "Percent conversion" refers to the percent of the substrate that is reduced to the product within a period of time under specified conditions. Thus, the "enzymatic activity" or "activity" of a polypeptide can be expressed as "percent conversion" of the substrate to the product.

[0059] "Percentage of sequence identity," "percent identity" and "percentage homology" are used interchangeably herein to refer to comparisons among polynucleotides and polypeptides, and are determined by comparing two optimally aligned sequences over a comparison window, wherein the portion of the polynucleotide or polypeptide sequence in the comparison window may comprise additions or deletions (i.e., gaps) as compared to the reference sequence (which may also contain gaps to optimize the alignment) for alignment of the two sequences. The percentage may be calculated by determining the number of positions at which the identical nucleic acid base or amino acid residue occurs in both sequences to yield the number of matched positions, dividing the number of matched positions by the total number of positions in the window of comparison (including positions where one of the sequences has a gap) and multiplying the result by 100 to yield the percentage of sequence identity. Alternatively, the percentage may be calculated by determining the number of positions at which either the identical nucleic acid base or amino acid residue occurs in both sequences or a nucleic acid base or amino acid residue is aligned with a gap to yield the number of matched positions, dividing the number of matched positions by the total number of positions in the window of comparison and multiplying the result by 100 to yield the percentage of sequence identity. Those of skill in the art appreciate that there are many established algorithms available to align two sequences and that different methods may give slightly different results.

[0060] Alignment of sequences for comparison can be conducted, e.g., by the local homology algorithm of Smith and Waterman, 1981, Adv. Appl. Math. 2:482, or the homology alignment algorithm of Needleman and Wunsch, 1970, J. Mol. Biol. 48:443, by the search for similarity method of Pearson and Lipman, 1988, Proc. Natl. Acad. Sci. USA 85:2444, by computerized implementations of these algorithms (GAP, BESTFIT, FASTA, and TFASTA in the GCG

Wisconsin Software Package), or by visual inspection (see generally, Current Protocols in Molecular Biology, F. M. Ausubel et al., eds., Current Protocols, a joint venture between Greene Publishing Associates, Inc. and John Wiley & Sons, Inc., (1995 Supplement) (Ausubel)). The Clustal (Chema R., Sugawara H., Koike T., Lopez R., Gibson T. J., Higgins D. G., Thompson J. D., (2003) Multiple sequence alignment with the Clustal series of programs, Nucleic Acids Res., 31, 3497-3500.) and T-Coffee (T-COFFEE: A novel method for multiple sequence alignments. Notredame, Higgins, Heringa, J M B 302 (205-217) 2000 software packages may also be used to align sequences.

[0061] Examples of algorithms that are suitable for determining percent sequence identity and sequence similarity are the BLAST and BLAST 2.0 algorithms, which are described in Altschul et al., 1990, J. Mol. Biol. 215: 403-410 and Altschul et al., 1977, Nucleic Acids Res. 3389-3402, respectively. Software for performing BLAST analyses is publicly available through the National Center for Biotechnology Information website. This algorithm involves first identifying high scoring sequence pairs (HSPs) by identifying short words of length W in the query sequence, which either match or satisfy some positive-valued threshold score T when aligned with a word of the same length in a database sequence. T is referred to as, the neighborhood word score threshold (Altschul et al, supra). These initial neighborhood word hits act as seeds for initiating searches to find longer HSPs containing them. The word hits are then extended in both directions along each sequence for as far as the cumulative alignment score can be increased. Cumulative scores are calculated using, for nucleotide sequences, the parameters M (reward score for a pair of matching residues; always >0) and N (penalty score for mismatching residues; always <0). For amino acid sequences, a scoring matrix is used to calculate the cumulative score. Extension of the word hits in each direction are halted when: the cumulative alignment score falls off by the quantity X from its maximum achieved value; the cumulative score goes to zero or below, due to the accumulation of one or more negative-scoring residue alignments; or the end of either sequence is reached. The BLAST algorithm parameters W, T, and X determine the sensitivity and speed of the alignment. The BLASTN program (for nucleotide sequences) uses as defaults a wordlength (W) of 11, an expectation (E) of 10, M=5, N=-4, and a comparison of both strands. For amino acid sequences, the BLASTP program uses as defaults a wordlength (W) of 3, an expectation (E) of 10, and the BLOSUM62 scoring matrix (see Henikoff and Henikoff, 1989, Proc Natl Acad Sci USA 89:10915). Exemplary determination of sequence alignment and % sequence identity can employ the BESTFIT or GAP programs in the GCG Wisconsin Software package (Accelrys, Madison Wis.), using default parameters provided.

[0062] "Reference sequence" refers to a defined sequence used as a basis for a sequence comparison. A reference sequence may be a subset of a larger sequence, for example, a segment of a full-length gene or polypeptide sequence. Generally, a reference sequence is at least 20 nucleotide or amino acid residues in length, at least 25 residues in length, at least 50 residues in length, or the full length of the nucleic acid or polypeptide. Since two polynucleotides or polypeptides may each (1) comprise a sequence (i.e., a portion of the complete sequence) that is similar between the two sequences, and (2) may further comprise a sequence that is divergent between the two sequences, sequence comparisons

between two (or more) polynucleotides or polypeptide are typically performed by comparing sequences of the two polynucleotides over a “comparison window” to identify and compare local regions of sequence similarity.

[0063] “Comparison window” refers to a conceptual segment of at least about 20 contiguous nucleotide positions or amino acids residues wherein a sequence may be compared to a reference sequence of at least 20 contiguous nucleotides or amino acids and wherein the portion of the sequence in the comparison window may comprise additions or deletions (i.e., gaps) of 20 percent or less as compared to the reference sequence (which does not comprise additions or deletions) for optimal alignment of the two sequences. The comparison window can be longer than 20 contiguous residues, and includes, optionally 30, 40, 50, 100, or longer windows.

[0064] “Conservative” amino acid substitutions or mutations refer to the interchangeability of residues having similar side chains, and thus typically involves substitution of the amino acid in the polypeptide with amino acids within the same or similar defined class of amino acids. However, as used herein, conservative mutations do not include substitutions from a hydrophilic to hydrophilic, hydrophobic to hydrophobic, hydroxyl-containing to hydroxyl-containing, or small to small residue, if the conservative mutation can instead be a substitution from an aliphatic to an aliphatic, non-polar to non-polar, polar to polar, acidic to acidic, basic to basic, aromatic to aromatic, or constrained to constrained residue. Further, as used herein, A, V, L, or I can be conservatively mutated to either another aliphatic residue or to another non-polar residue. Table 1 below shows exemplary conservative substitutions.

TABLE 1

Conservative Substitutions	
Residue	Possible Conservative Mutations
A, L, V, I	Other aliphatic (A, L, V, I) Other non-polar (A, L, V, I, G, M)
G, M	Other non-polar (A, L, V, I, G, M)
D, E	Other acidic (D, E)
K, R	Other basic (K, R)
P, H	Other constrained (P, H)
N, Q, S, T	Other polar (N, Q, S, T)
Y, W, F	Other aromatic (Y, W, F)
C	None

[0065] “Non-conservative substitution” refers to substitution or mutation of an amino acid in the polypeptide with an amino acid with significantly differing side chain properties. Non-conservative substitutions may use amino acids between, rather than within, the defined groups listed above. In one embodiment, a non-conservative mutation affects (a) the structure of the peptide backbone in the area of the substitution (e.g., proline for glycine) (b) the charge or hydrophobicity, or (c) the bulk of the side chain.

[0066] “Deletion” refers to modification to the polypeptide by removal of one or more amino acids from the reference polypeptide. Deletions can comprise removal of 1 or more amino acids, 2 or more amino acids, 5 or more amino acids, 10 or more amino acids, 15 or more amino acids, or 20 or more amino acids, up to 10% of the total number of amino acids, or up to 20% of the total number of amino acids making up the reference enzyme while retaining enzymatic activity and/or retaining the improved properties of an engineered enzyme. Deletions can be directed to the internal portions and/or ter-

minal portions of the polypeptide. In various embodiments, the deletion can comprise a continuous segment or can be discontinuous. The term “deletion” is also used to refer to a DNA modification in which one or more nucleotides or nucleotide base-pairs have been removed, as compared to the corresponding reference, parental or “wild type” DNA.

[0067] “Insertion” refers to modification to the polypeptide by addition of one or more amino acids to the reference polypeptide. In some embodiments, the modification comprises insertions of one or more amino acids to the naturally occurring polypeptide as well as insertions of one or more amino acids to other modified polypeptides. Insertions can be in the internal portions of the polypeptide, or to the carboxy or amino terminus. Insertions as used herein include fusion proteins as is known in the art. The insertion can be a contiguous segment of amino acids or separated by one or more of the amino acids in the naturally occurring polypeptide. The term “insertion” is also used to refer to a DNA modification in which one or more nucleotides or nucleotide base-pairs have been inserted, as compared to the corresponding reference, parental or “wild type” DNA.

[0068] “Different from” or “differs from” with respect to a designated reference sequence refers to difference of a given amino acid or polynucleotide sequence when aligned to the reference sequence. Generally, the differences can be determined when the two sequences are optimally aligned. Differences include insertions, deletions, or substitutions of amino acid residues in comparison to the reference sequence.

[0069] As used herein in the context of a polypeptide or polynucleotide, the phrase “derived from” a particular organism refers to a wild-type polynucleotide or polypeptide that originates in the organism and to mutant and variants thereof that either originate in the organism or are produced by human manipulation of the wild-type polynucleotide or polypeptide.

[0070] “Functional fragment” as used herein refers to a polypeptide that has an amino-terminal and/or carboxy-terminal deletion, but where the remaining amino acid sequence is identical to the corresponding positions in the sequence and that retains substantially all of the activity of the full-length polypeptide. Functional fragments can comprise up to 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, and 99% of the full-length polypeptide.

[0071] “Endogenous” polynucleotide, gene, promoter or polypeptide refers to any polynucleotide, gene, promoter or polypeptide that originates in a particular host cell. A polynucleotide, gene, promoter or polypeptide is not endogenous to a host cell if it has been removed from the host cell, subjected to laboratory manipulation, and then reintroduced into a host cell.

[0072] “Heterologous” polynucleotide, gene, promoter or polypeptide refers to any polynucleotide, gene, promoter or polypeptide that is introduced into a host cell that is not normally present in that cell, and includes any polynucleotide, gene, promoter or polypeptide that is removed from the host cell and then reintroduced into the host cell.

[0073] “Inactive” or “inactivated” in reference to a gene refers to a gene having at least one function that is impaired. Genes can be inactivated in a variety of ways known in the art, including but not limited to insertion of a mobile genetic element (e.g., a transposon); deletion of all or part of the gene, such that the gene product is not made, or is truncated and is non-functional; mutation of the gene such that the gene product is not made, or is truncated and is non-functional; deletion

or mutation of one or more control elements that control expression of the gene such that the gene product is not made; and the like. In certain embodiments genes can be inactivated by methods other than genetic modification, for example, by gene silencing at the transcriptional level or at the post-transcriptional level using for example RNAi.

[0074] “Recombinant host cell” refers to a cell into which has been introduced a heterologous polynucleotide, gene, promoter, e.g., an expression vector, or to a cell having a heterologous polynucleotide or gene integrated into the genome.

[0075] “Naturally-occurring” or “wild-type” refers to the form found in nature. For example, a naturally occurring or wild-type polypeptide or polynucleotide sequence is a sequence present in an organism that can be isolated from a source in nature and which has not been intentionally modified by human manipulation. A wild-type organism refers to an organism that has not been intentionally modified by human manipulation.

[0076] “Codon optimized” refers to changes in the codons of the polynucleotide encoding a protein to those preferentially used in a particular organism such that the encoded protein is efficiently expressed in the organism of interest. Although the genetic code is degenerate in that most amino acids are represented by several codons, called “synonyms” or “synonymous” codons, it is well known that codon usage by particular organisms is nonrandom and biased towards particular codon triplets. This codon usage bias may be higher in reference to a given gene, genes of common function or ancestral origin, highly expressed proteins versus low copy number proteins, and the aggregate protein coding regions of an organism’s genome. In some embodiments, the polynucleotides encoding enzymes may be codon optimized for optimal production from the host organism selected for expression.

[0077] “Preferred, optimal, high codon usage bias codons” refers interchangeably to codons that are used at higher frequency in the protein coding regions than other codons that code for the same amino acid. The preferred codons may be determined in relation to codon usage in a single gene, a set of genes of common function or origin, highly expressed genes, the codon frequency in the aggregate protein coding regions of the whole organism, codon frequency in the aggregate protein coding regions of related organisms, or combinations thereof. Codons whose frequency increases with the level of gene expression are typically optimal codons for expression. A variety of methods are known for determining the codon frequency (e.g., codon usage, relative synonymous codon usage) and codon preference in specific organisms, including multivariate analysis, for example, using cluster analysis or correspondence analysis, and the effective number of codons used in a gene (See GCG Codon Preference, Genetics Computer Group Wisconsin Package; CodonW, John Peden, University of Nottingham; McInerney, J. O, 1998, Bioinformatics 14:372-73; Stenico et al., 1994, Nucleic Acids Res. 22:2437-46; Wright, F., 1990, Gene 87:23-29). Codon usage tables are available for a growing list of organisms (see for example, Wada et al., 1992, Nucleic Acids Res. 20:2111-2118; Nakamura et al., 2000, Nucl. Acids Res. 28:292; Duret, et al., supra; Henaut and Danchin, “*Escherichia coli* and *Salmonella*,” 1996, Neidhardt, et al. Eds., ASM Press, Washington D.C., p. 2047-2066), all of which are incorporated herein by reference. The data source for obtaining codon usage may rely on any available nucleotide sequence capable

of coding for a protein. These data sets include nucleic acid sequences actually known to encode expressed proteins (e.g., complete protein coding sequences-CDS), expressed sequence tags (ESTs), or predicted coding regions of genomic sequences (see for example, Mount, D., Bioinformatics: Sequence and Genome Analysis, Chapter 8, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y., 2001; Uberbacher, E. C., 1996, Methods Enzymol. 266:259-281; Tiwari et al., 1997, Comput. Appl. Biosci. 13:263-270), all of which are incorporated herein by reference.

[0078] “Expression” as used herein may include any step involved in the production of the FAR polypeptides including but not limited to transcription and translation.

[0079] “Control sequence” is defined herein to include all components, which are necessary or advantageous for the expression of a polypeptide of the present disclosure. Each control sequence may be native or foreign to the nucleic acid sequence encoding the polypeptide. Such control sequences include, but are not limited to, a leader, polyadenylation sequence, propeptide sequence, promoter, signal peptide sequence, and transcription terminator. At a minimum, the control sequences include a promoter, and transcriptional and translational stop signals. The control sequences may be provided with linkers for the purpose of introducing specific restriction sites facilitating ligation of the control sequences with the coding region of the nucleic acid sequence encoding a polypeptide.

[0080] “Operably linked” and “operably associated” are defined herein as a configuration in which a control sequence is appropriately placed at a position relative to the coding sequence of the DNA sequence such that the control sequence directs the expression of a polynucleotide and/or polypeptide.

[0081] “Promoter sequence” is a nucleic acid sequence that is recognized by a host cell for expression of the coding region. The control sequence may comprise an appropriate promoter sequence. The promoter sequence contains transcriptional control sequences, which mediate the expression of the polypeptide. The promoter may be any nucleic acid sequence which shows transcriptional activity in the host cell of choice including mutant, truncated, and hybrid promoters, and may be obtained from genes encoding extracellular or intracellular polypeptides either endogenous or heterologous to the host cell.

[0082] As used herein the term “transformed” or “transformation” used in reference to a cell means a cell has a non-native nucleic acid sequence integrated into its genome or as a plasmid that is maintained through multiple generations.

[0083] It should be noted that the indefinite articles “a” and “an” and the definite article “the” are used in the present application to mean one or more unless the context clearly dictates otherwise. Further, the term “or” is used in the present application to mean the disjunctive “or” and the conjunctive “and”.

#### FAR Enzymes Useful in the Methods of the Disclosure:

[0084] In one aspect the present disclosure provides FAR enzymes derived from or obtained from gammaproteobacteria (such as marine gammaproteobacteria) and/or functional fragments thereof having FAR activity. In some embodiments, the FAR enzyme is isolated directly from the gammaproteobacteria in which it occurs naturally. In some embodiments, the FAR enzyme and/or a function fragment is isolated from a heterologous host microorganism engineered to express the FAR enzyme or functional fragment. The FAR

enzyme and/or functional fragment can be derived or obtained from virtually any genus of marine gammaproteobacteria. In certain embodiments, the FAR enzyme for use in the methods disclosed herein is endogenous to a marine bacterium, i.e., a bacterium mined from a marine environment. In various embodiments, the FAR enzyme for use in the methods disclosed herein is endogenous to an organism other than a marine bacterium.

[0085] In certain embodiments, the FAR enzyme and/or functional fragment can be derived or obtained from a gammaproteobacterium of the order Alteromonadales. In some embodiments, the FAR enzyme and/or functional fragment can be derived from or obtained from the Alteromonadales family Alteromonadaceae. In certain embodiments, the FAR enzyme and/or functional fragment can be derived from or obtained from an Alteromonadaceae genus such as but not limited to the Alteromonadaceae genus *Marinobacter*. In certain specific embodiments, the FAR enzyme and/or functional fragment can be derived from the *Marinobacter* species *algicola*. In a particular embodiment, the FAR enzyme and/or functional fragment can be derived from or obtained from the *M. algicola* species strain DG893. In some specific embodiments, the FAR enzyme for use in the methods disclosed herein is from the marine bacterium *Marinobacter algicola* (e.g., *M. algicola*) DG893 (SEQ ID NO: 2) ("FAR\_Maa").

[0086] In some embodiments, the FAR enzyme and/or functional fragment is derived or obtained from a species of *Marinobacter* including, but not limited to, a species selected from *M. algicola*, *M. alkaliphilus*, *M. aquaeolei*, *M. arcticus*, *M. bryozorum*, *M. daeponensis*, *M. excellens*, *M. flavigaris*, *M. guadonensis*, *M. hydrocarbonoclasticus*, *M. koreensis*, *M. lipolyticus*, *M. litoralis*, *M. lutaoensis*, *M. maritimus*, *M. sediminum*, *M. squalenivirans* and *M. vinifirmus* and equivalent and synonymous species thereof.

[0087] In one specific embodiment, the FAR enzyme is derived or obtained from *M. algicola* strain DG893 and has an amino acid sequence that is at least 30% identical, at least 50% identical, at least 60% identical, at least 70% identical, at least 75%, at least 80% identical, at least 85% identical, at least 90% identical, at least 93% identical at least 95% identical, at least 97% identical and/or at least 98% identical to SEQ ID NO: 2 or a functional fragment thereof. In another specific embodiment, the isolated FAR enzyme has an amino acid sequence that is identical to SEQ ID NO: 2.

[0088] In one specific embodiment, the FAR enzyme is derived or obtained from *Marinobacter aquaeolei* and has an amino acid sequence that is at least 30% identical, at least 50% identical, at least 60% identical, at least 70% identical, at least 75%, at least 80% identical, at least 85% identical, at least 90% identical, at least 93% identical, at least 95% identical, at least 97% identical and/or at least 98% identical to SEQ ID NO: 14 or a functional fragment thereof. In another specific embodiment, the isolated FAR enzyme has an amino acid sequence that is identical to SEQ ID NO: 14.

[0089] In various embodiments, the isolated FAR enzyme and/or functional fragment is obtained or derived from a marine bacterium selected from the group of *Meptunibacter caesariensis* species strain MED92, *Reinekea* sp. strain MED297, *Marinomonas* sp. strain MED121, unnamed gammaproteobacterium strain HTCC2207 and *Marinobacter* sp. strain ELB17 and equivalents and synonymous species thereof.

[0090] In various embodiments, the FAR enzyme and/or functional fragment can be derived or obtained from a gam-

maproteobacterium of the order Oceanospirillales. In some embodiments, the FAR enzyme and/or functional fragment can be derived from or obtained from the Oceanospirillales family Oceanospirillaceae. In certain embodiments, the FAR enzyme and/or functional fragment can be derived from or obtained from a Oceanospirillaceae genus, such as but not limited to *Oceanobacter*. In a particular embodiment, the FAR enzyme and/or functional fragment can be derived from or obtained from the *Oceanobacter* species strain RED65 and has an amino acid sequence that is at least 30% identical, at least 50% identical, at least 60% identical, at least 70% identical, at least 75% identical, at least 80% identical, at least 85% identical, at least 90% identical, at least 93% identical, at least 95% identical, at least 97% identical and/or at least 98% identical to SEQ ID NO: 6 or a functional fragment thereof. In another specific embodiment, the FAR enzyme for use in the methods disclosed herein comprises or consists of a sequence having 100% identity to the sequence of SED ID NO: 6 ("FAR\_Ocs"). In other specific embodiments, the isolated FAR enzyme or functional fragment is obtained or derived from *Oceanobacter kriegii*. In still other specific embodiments, the isolated FAR enzyme or functional fragment is obtained or derived from *Oceanobacter* strain WH099.

[0091] In various embodiments, the FAR enzyme is from a marine bacterium and is selected from the group consisting of FAR\_Hch (*Hahella chejuensis* KCTC 2396 GenBank YP\_436183.1); FAR\_Mac (from marine *actinobacterium* strain PHSC20C1), FAR\_JVC (JCVI\_ORF\_1096697648832, GenBank Accession No. EDD40059.1; from a marine metagenome), FAR\_Fer (JCVI\_SCAF\_1101670217388; from a marine bacterium found at a depth of 12 m in an upwelling in the area of Fernandina Island, the Galapagos Islands, Ecuador), FAR\_Key (JCVI\_SCAF\_1097205236585, from a marine bacterium found at a depth of 1.7 m off the coast of Key West Fla.), and FAR\_Gal (JCVI\_SCAF\_1101670289386, at a depth of 0.1 m at Isabella Island, Galapagos Islands, Ecuador). Approximate sequence identity to *M. algicola* DG893 (FAR\_Maa) and *Oceanobacter* sp. RED65 (FAR\_Ocs) is given in the table below.

FAR Gene	% Sequence Identity to FAR_Maa (SEQ ID NO: 2/4)	% Sequence Identity to FAR_Ocs (SEQ ID NO: 6/8)
FAR_Maa	100	46
FAR_Mac	32	31
FAR_Fer	61	36
FAR_Gal	25	25
FAR_JVC	34	30
FAR_Key	32	30
FAR_Maq	78	45
FAR_Hch	54	47

[0092] In one particular embodiment, the FAR enzyme is isolated or derived from the marine bacterium FAR\_Gal. In other embodiments, the FAR enzyme or functional fragment is isolated or derived from an organism selected from the group consisting of *Vitis vinifera* (GenBank Accession No. CAO22305.1 or CAO67776.1), *Desulfatibacillum alkenivorans* (GenBank Accession No. NZ\_ABII01000018.1), *Stigmatella aurantiaca* (NZ\_AAMD01000005.1) and *Phytophthora ramorum* (GenBank Accession No.: AAQX01001105.1).

[0093] In certain embodiments, a FAR enzyme or functional fragment thereof that is especially suitable for the production of fatty alcohols is identified by the presence of one or more domains, which are found in proteins with FAR activity. In various embodiments, the one or more domains is identified by multiple sequence alignments using hidden Markov models (“HMMs”) to search large collections of protein families, for example, the Pfam collection available at <http://pfam.sanger.ac.uk/>. See R. D. Finn et al. (2008) *Nucl. Acids Res.* Database Issue 36:D281-D288.

[0094] In certain embodiments, the one or more protein domains by which FAR enzymes are identified belongs to a family of NAD binding domains found in the male sterility proteins of *arabidopsis* and *drosophila*, as well as in the fatty acyl reductase enzyme from the jojoba plant (JJFAR). See Aarts M G et al. (1997) *Plant J.* 12:615-623. This family of binding domains is designated “NAD\_binding\_4” (PF07993; see <http://pfam.sanger.ac.uk/family?acc=PF07993>). In various embodiments, the NAD\_binding\_4 domain is found near the N-terminus of the putative FAR enzyme. In various embodiments, the one or more protein domains by which enzymes with FAR activity are identified belongs to a family of domains known as a “sterile” domain (PF03015; see <http://pfam.sanger.ac.uk/family?acc=PF03015>), which are also found in the male sterility proteins of *Arabidopsis* species and a number of other organisms. See Aarts M G et al. (1997) *Plant J.* 12:615-623. In particular embodiments, the sterile domain is found near the C-terminus of the putative FAR enzyme. In certain specific embodiments, a FAR enzyme for use in the methods described herein is identified by the presence of at least one NAD\_binding\_4 domain near the N-terminus and the presence of at least one sterile domain near the C-terminus.

[0095] In certain embodiments, the NAD\_binding\_4 domain of the putative FAR enzyme has an amino acid sequence that is at least 10%, such as at least 15%, such as at least 20%, such as at least 25%, such as at least 30%, such as at least 35%, such as at least 40%, such as at least 45%, such as at least 50%, such as at least 60%, such as at least 70%, such as at least 80%, such as at least 85%, such as at least 90% or more identical to the amino acid sequence of a known NAD\_binding\_4 domain. See, e.g., Aarts M G et al. (1997) *Plant J.* 12:615-623. In various embodiments, the sterile domain of the putative FAR enzyme has an amino acid sequence that is at least 10%, such as at least 15%, such as at least 20%, such as at least 25%, such as at least 30%, such as at least 35%, such as at least 40%, such as at least 45%, such as at least 50% or more identical to the amino acid sequence of a known sterile domain. See id.

[0096] In some embodiments, the NAD\_binding\_4 domain of the putative FAR enzyme has an amino acid sequence that is at least 10%, such as at least 15%, such as at least 20%, such as at least 25%, such as at least 30%, such as at least 35%, such as at least 40%, such as at least 45%, such as at least 50%, such as at least 60%, such as at least 70%, such as at least 80%, such as at least 85%, such as at least 90% or more identical to the amino acid sequence of one or more example polypeptides that form the definition of the NAD\_binding\_4 Pfam domain (PF07993). In certain embodiments, the sterile domain of the putative FAR enzyme has an amino acid sequence that is at least 10%, such as at least 15%, such as at least 20%, such as at least 25%, such as at least 30%, such as at least 35%, such as at least 40%, such as at least 45%, such as at least 50% or more identical to the amino acid sequence of a known sterile domain.

acid sequence of one or more example polypeptides that form the definition of the sterile Pfam domain (PF03015). In various embodiments, the NAD\_binding\_4 domain or the sterile domain of the putative FAR enzyme is identified by an E-value of  $1 \times 10^{-4}$  or less, such as an E-value of  $1 \times 10^{-5}$ , such as an E-value of  $1 \times 10^{-10}$ , such as an E-value of  $1 \times 10^{-15}$ , such as an E-value of  $1 \times 10^{-20}$ , such as an E-value of  $1 \times 10^{-25}$ , such as an E-value of  $1 \times 10^{-30}$  or lower. As used herein, the term E-value (expectation value) is the number of hits that would be expected to have a score equal or better than a particular hit by chance alone. Accordingly, the E-value is a criterion by which the significance of a database search hit can be evaluated. See, e.g., <http://pfam.sanger.ac.uk/help>; <http://www.csb.yale.edu/userguides/seq/hmmer/docs/node5.html>.

[0097] The FAR enzymes described herein have not previously been recognized as FAR enzymes because of the low homology of the FAR coding sequences to the sequences coding for proteins with known FAR activity, such as the FAR enzymes from *S. chinensis* ((FAR Sim); GenBank Accession no. AAD38039.1; gil5020215|gb|AAD38039.1|AF149917\_1 acylCoA reductase [*Simmondsia chinensis*]-Plant Physiol. 2000 March; 122(3):635-44. Purification of a jojoba embryo fatty acyl-coenzyme A reductase and expression of its cDNA in high erucic acid rapeseed; Metz J G, Pollard M R, Anderson L, Hayes T R, Lassner M W. PMID: 10712526), *B. mori* ((FAR Bom); GenBank Accession no. BAC79425.1; gil33146307|dbj|BAC79425.1|fatty-acyl reductase [*Bombyx mori*]; Proc Natl Acad Sci USA 2003 Aug. 5; 100(16):9156-61. Epub 2003 Jul. 18. Pheromone gland-specific fatty-acyl reductase of the silkworm, *Bombyx mori*. Moto K, Yoshiga T, Yamamoto M, Takahashi S, Okano K, Ando T, Nakata T, Matsumoto S. PMID: 12871998), *Arabidopsis thaliana* (GenBank Accession no. DQ446732.1 or NM\_115529.1; gil91806527|gb|DQ446732.1|Arabidopsis thaliana clone pENTR221-At3g44560; gil18410556|ref|NM\_115529.1|Arabidopsis thaliana male sterility protein, putative (AT3G56700); Plant Physiol. 2009 May 15; 166(8):787-96. Epub 2008 Dec. 4. Functional expression of five *Arabidopsis* fatty acyl-CoA reductase genes in *Escherichia coli*. Doan T T, Carlsson A S, Hamberg M, Büllow L, Stymne S, Olsson P. PMID: 19062129) or *Ostrinia scapulalis* (GenBank Accession no. EU817405.1; gil210063138|gb|EU817405.1|Ostrinia scapulalis FAR-like protein XIII; Insect Biochem. Mol. Biol. 2009 February; 39(2):90-5. Epub 2008 Oct. 26 Pheromone-gland-specific fatty-acyl reductase in the adzuki bean borer, *Ostrinia scapulalis* (*Lepidoptera: Crambidae*) Antony B, Fujii T, Moto K, Matsumoto S, Fukuzawa M, Nakano R, Tatsuki S, Ishikawa Y.).

[0098] In certain embodiments, the FAR enzyme for use in the disclosed methods of producing fatty alcohols is endogenous to a marine bacterium and has an amino acid sequence having less than about 35% sequence identity to the amino acid sequence of a known FAR enzyme, such as those described above. In various embodiments, the FAR enzyme for use in the methods disclosed herein is endogenous to a marine bacterium and has an amino acid sequence having less than about 33% sequence identity, such as less than 31% sequence identity, such as less than about 29% sequence identity, such as less than about 27% sequence identity, such as less than about 25% sequence identity, such as less than about 23% sequence identity, even such as less than about

22% sequence identity to the amino acid sequence of a polypeptide previously recognized as possessing FAR enzyme activity.

[0099] In various embodiments, the FAR enzyme for use in the disclosed methods of producing fatty alcohols is endogenous to an organism selected from the group consisting *Vitis vinifera*, *Desulfatibacillum alkenivorans*, *Stigmatella aurantiaca*, and *Phytophthora ramorum* and has an amino acid sequence having less than about 65% sequence identity to the amino acid sequence of a known FAR enzyme, such as those described above. In various embodiments, the FAR enzyme for use in the methods disclosed herein is endogenous to a marine bacterium and has an amino acid sequence having less than about 60% sequence identity, such as less than 55% sequence identity, such as less than about 50% sequence identity, such as less than about 45% sequence identity, such as less than about 40% sequence identity, such as less than about 35% sequence identity, even such as less than about 34% sequence identity, such as less than about 33% sequence identity, such as less than about 32% sequence identity, such as less than about 31% sequence identity, such as less than about 30% sequence identity to the amino acid sequence of a polypeptide previously recognized as possessing FAR enzyme activity.

[0100] In certain embodiments, the FAR enzyme endogenous to *Marinobacter algicola* DG893 for use in the methods disclosed herein is the full-length polypeptide having the amino acid set forth in SEQ ID NO: 2. In various embodiments, the full-length FAR enzyme has an amino acid sequence that is at least about 75% identical, such as at least about 80% identical, such as at least about 85% identical, such as at least about 90% identical, such as at least about 91%, such as at least about 92%, such as at least about 93%, such as at least about 94%, such as at least about 95%, such as at least about 96%, such as at least about 97%, such as at least about 98%, such as at least about 99% identical to the sequence set forth in SEQ ID NO: 2. In certain preferred embodiments, where the amino acid sequence is less than 100% identical to that shown in SEQ ID NO: 2, all amino acid substitutions are conservative.

[0101] In various embodiments, the FAR enzyme endogenous to *M. algicola* DG893 for use in the methods disclosed herein is a functional fragment of the full-length polypeptide, the amino acid sequence of which is set forth in SEQ ID NO: 2. In certain embodiments, the amino acid sequence of the functional fragment is at least about 70%, such as at least about 75%, such as at least about 80%, such as at least about 90%, such as at least about 91%, such as at least about 92%, such as at least about 93%, such as at least about 94%, such as at least about 95%, such as at least about 96%, such as at least about 97%, such as at least about 98%, such as at least about 99% identical to the amino acid sequence of the corresponding region of SEQ ID NO: 2.

[0102] In certain embodiments, the FAR enzyme endogenous to *Marinobacter aquaeolei* for use in the methods disclosed herein is the full-length polypeptide having the amino acid set forth in SEQ ID NO: 14. In various embodiments, the full-length FAR enzyme has an amino acid sequence that is at least about 75% identical, such as at least about 80% identical, such as at least about 85% identical, such as at least about 90% identical, such as at least about 91%, such as at least about 92%, such as at least about 93%, such as at least about 94%, such as at least about 95%, such as at least about 96%,

such as at least about 97%, such as at least about 98%, such as at least about 99% identical to the sequence set forth in SEQ ID NO: 14. In certain preferred embodiments, where the amino acid sequence is less than 100% identical to that shown in SEQ ID NO: 14, all amino acid substitutions are conservative.

[0103] In various embodiments, the FAR enzyme endogenous to *M. aquaeolei* for use in the methods disclosed herein is a functional fragment of the full-length polypeptide, the amino acid sequence of which is set forth in SEQ ID NO: 14. In certain embodiments, the amino acid sequence of the functional fragment is at least about 70%, such as at least about 75%, such as at least about 80%, such as at least about 90%, such as at least about 91%, such as at least about 92%, such as at least about 93%, such as at least about 94%, such as at least about 95%, such as at least about 96%, such as at least about 97%, such as at least about 98%, such as at least about 99% identical to the amino acid sequence of the corresponding region of SEQ ID NO: 14.

[0104] In certain embodiments, the FAR enzyme endogenous to *Oceanobacter* sp. RED65 for use in the methods disclosed herein is the full-length polypeptide having the amino acid set forth in SEQ ID NO: 6. In various embodiments, the FAR enzyme has an amino acid sequence that is at least about 75% identical, such as at least 80% identical, such as at least about 85% identical, such as at least about 90% identical, such as at least about 91%, such as at least about 92%, such as at least about 93%, such as at least about 94%, such as at least about 95%, such as at least about 96%, such as at least about 97%, such as at least about 98%, such as at least about 99% identical to the sequence set forth in SEQ ID NO: 6. In certain preferred embodiments, where the sequence identity is less than 100%, all amino acid substitutions are conservative.

[0105] In various embodiments, the FAR enzyme endogenous to *Oceanobacter* sp. RED65 for use in the methods disclosed herein is a functional fragment of the full-length polypeptide, the amino acid sequence of which is set forth in SEQ ID NO: 6. In certain embodiments, the functional fragment has an amino acid sequence that is at least about 70%, such as at least about 75%, such as at least about 80%, such as at least about 90%, such as at least about 91%, such as at least about 92%, such as at least about 93%, such as at least about 94%, such as at least about 95%, such as at least about 96%, such as at least about 97%, such as at least about 98%, such as at least about 99% identical to the amino acid sequence of the corresponding region of SEQ ID NO: 6.

#### Nucleic Acids for Expression of FAR Enzymes:

[0106] In another aspect, the present disclosure provides polynucleotides encoding a FAR enzyme, or functional fragment thereof, as described above. The polynucleotide can be a DNA or RNA, and can be single-stranded or double-stranded. The polynucleotide can be isolated from a naturally occurring microorganism, or prepared wholly or partially via synthetic means. In some embodiments, the polynucleotide is a construct designed to provide for expression of the encoded FAR enzyme or functional fragment in a host microorganism. The FAR enzyme or the specific or functional fragment encoded by the polynucleotide construct may be endogenous or heterologous to the host microorganism.

[0107] In various aspects of the disclosure, the availability of a polypeptide sequence of a specific FAR enzyme provides a description of all polynucleotides capable of encoding the

polypeptide of known sequence because of the known correspondence of particular codons and the amino acids they encode. In certain embodiments, the degeneracy of the genetic code is used to produce a large number of polynucleotides that encode the gamma proteobacterial FAR polypeptides described herein. Thus, for example, the codons AGA, AGG, CGA, CGC, CGG, and CGU all encode the amino acid arginine (R). Thus, at every position in the nucleic acids of the invention where an arginine is specified by a codon, the codon can be altered to any of the corresponding codons described above without altering the encoded polypeptide. In some embodiments, the polynucleotides that encode the FAR enzymes described herein are codon optimized for expression in a recombinant host cells. In particular embodiments, the polynucleotides that encode the FAR enzymes described herein are codon optimized for expression in bacteria, yeast or filamentous fungi. In various embodiments, the polynucleotides that encode the FAR enzymes described herein are codon optimized for expression in oleaginous yeast. In certain specific embodiments, the polynucleotides that encode the FAR enzymes are codon optimized for expression in *E. coli*, *S. cerevisiae* or *Y. lipolytica*.

[0108] In certain embodiments, the present disclosure provides an isolated nucleic acid encoding a FAR enzyme from a marine bacterium. In some embodiments, the nucleic acid encodes a FAR enzyme having an amino acid sequence that is at least about 75% identical, at least about 80% identical, such as at least about 85% identical, such as at least about 90% identical, such as at least about 91%, such as at least about 92%, such as at least about 93%, such as at least about 94%, such as at least about 95%, such as at least about 96%, such as at least about 97%, such as at least about 98%, such as at least about 99% identical to the sequence set forth in SEQ ID NO: 2. In some embodiments, the nucleic acid encodes a FAR enzyme having an amino acid sequence that is at least about 75% identical, at least about 80% identical, such as at least about 85% identical, such as at least about 90% identical, such as at least about 91%, such as at least about 92%, such as at least about 93%, such as at least about 94%, such as at least about 95%, such as at least about 96%, such as at least about 97%, such as at least about 98%, such as at least about 99% identical to the sequence set forth in SEQ ID NO: 14. In other embodiments, the nucleic acid encodes a FAR enzyme having an amino acid sequence that is at least about 75% identical, such as at least 80% identical, such as at least about 85% identical, such as at least about 90% identical, such as at least about 91%, such as at least about 92%, such as at least about 93%, such as at least about 94%, such as at least about 95%, such as at least about 96%, such as at least about 97%, such as at least about 98%, such as at least about 99% identical to the sequence set forth in SEQ ID NO: 6.

[0109] In various embodiments, the present disclosure provides DNA constructs, vectors and polynucleotides encoding FAR enzymes for expression in heterologous recombinant host cells. In certain embodiments, the DNA constructs, vectors and polynucleotides comprise a sequence that encodes a FAR enzyme endogenous to a marine bacterium. In some embodiments, the DNA constructs, vectors and polynucleotides comprise a sequence that encodes a FAR enzyme endogenous to a *Marinobacter* species. In various embodiments, the DNA constructs, vectors and polynucleotides comprise a sequence that encodes a FAR enzyme endogenous to an *Oceanobacter* species. In some particular embodiments, the DNA constructs, vectors and polynucleotides comprise a

sequence that encodes a FAR enzyme endogenous to *M. algicola* DG893 or *M. aquaeolei*. In other particular embodiments, the DNA constructs, vectors and polynucleotides comprise a sequence that encodes a FAR enzyme endogenous to *Oceanobacter* sp. RED65. In some particular embodiments, the polynucleotide is a codon optimized polynucleotide, such as a polynucleotide having at least 90% (at least 93%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99% and even 100%) sequence identity to SEQ ID NOs: 1, 3, 5, 7 or 13.

[0110] In certain embodiments, the DNA constructs, vectors and polynucleotides comprise a sequence that encodes the full length FAR polypeptide encompassed by the invention. In some embodiments, the DNA construct, vector or polynucleotide comprises a sequence that encodes a full length FAR endogenous to a *Marinobacter* strain. In some embodiments the strain is *M. algicola* DG893 and in other embodiments the strain is a *M. aquaeolei*. In other embodiments, the DNA constructs, vectors and polynucleotides comprise a sequence that encodes a functional fragment of the FAR polypeptide of *M. algicola* DG893 or a *M. aquaeolei*. In various embodiments, the DNA constructs, vectors and polynucleotides comprise a sequence that encodes the full length FAR polypeptide endogenous to *Oceanobacter* sp. RED65. In some embodiments, the DNA constructs, vectors and polynucleotides comprise a sequence that encodes a functional fragment of the FAR polypeptide of *Oceanobacter* sp. RED65.

[0111] In certain embodiments, the DNA constructs, vectors and polynucleotides comprise a sequence that encodes a full-length FAR polypeptide endogenous to *M. algicola* DG893 having an amino acid sequence that is at least about 75% identical, such as at least 80% identical, such as at least about 85% identical, such as at least about 90% identical, such as at least about 91%, such as at least about 92%, such as at least about 93%, such as at least about 94%, such as at least about 95%, such as at least about 96%, such as at least about 97%, such as at least about 98%, such as at least about 99% identical to the amino acid sequence set forth in SEQ ID NO: 2.

[0112] In some embodiments, the DNA constructs, vectors and polynucleotides comprise a sequence that encodes a functional fragment of the FAR enzyme having an amino acid sequence that is at least about 80% identical, such as at least about 85% identical, such as at least about 90% identical, such as at least about 91%, such as at least about 92%, such as at least about 93%, such as at least about 94%, such as at least about 95%, such as at least about 96%, such as at least about 97%, such as at least about 98%, such as at least about 99% identical to the amino acid sequence of the corresponding region of SEQ ID NO: 2.

[0113] In certain embodiments, the DNA constructs, vectors and polynucleotides comprise a sequence that encodes a full-length FAR polypeptide endogenous to *M. aquaeolei* having an amino acid sequence that is at least about 75% identical, such as at least 80% identical, such as at least about 85% identical, such as at least about 90% identical, such as at least about 91%, such as at least about 92%, such as at least about 93%, such as at least about 94%, such as at least about 95%, such as at least about 96%, such as at least about 97%, such as at least about 98%, such as at least about 99% identical to the amino acid sequence set forth in SEQ ID NO: 14.

[0114] In some embodiments, the DNA constructs, vectors and polynucleotides comprise a sequence that encodes a

functional fragment of the FAR enzyme having an amino acid sequence that is at least about 80% identical, such as at least about 85% identical, such as at least about 90% identical, such as at least about 91%, such as at least about 92%, such as at least about 93%, such as at least about 94%, such as at least about 95%, such as at least about 96%, such as at least about 97%, such as at least about 98%, such as at least about 99% identical to the amino acid sequence of the corresponding region of SEQ ID NO: 14.

[0115] In various embodiments, the DNA constructs, vectors and polynucleotides comprise a sequence that encodes a full-length FAR polypeptide endogenous to *Oceanobacter* sp. RED65 having an amino acid sequence that is at least about 80% identical, such as at least about 85% identical, such as at least about 90% identical, such as at least about 91%, such as at least about 92%, such as at least about 93%, such as at least about 94%, such as at least about 95%, such as at least about 96%, such as at least about 97%, such as at least about 98%, such as at least about 99% identical to the amino acid sequence set forth in SEQ ID NO: 6.

[0116] In some embodiments, the DNA constructs, vectors and polynucleotides comprise a sequence that encodes a functional fragment of the FAR enzyme that has an amino acid sequence that is at least about 80% identical, such as at least about 85% identical, such as at least about 90% identical, such as at least about 91%, such as at least about 92%, such as at least about 93%, such as at least about 94%, such as at least about 95%, such as at least about 96%, such as at least about 97%, such as at least about 98%, such as at least about 99% identical to the amino acid sequence of the corresponding region of SEQ ID NO: 6.

[0117] In still other embodiments, the present disclosure provides DNA constructs, vectors and polynucleotides encoding FAR enzymes endogenous to a marine bacterium, wherein the FAR enzyme is selected from the group consisting of FAR\_Mac (from marine actinobacterium species PHSC20C1), FAR\_JVC (JCVI\_ORF\_1096697648832, GenBank Accession No. EDD40059.1; from a marine metagenome), FAR\_Fer (JCVI\_SCAF\_1101670217388; from a marine bacterium found at a depth of 12 m in an upwelling in the area of Fernandina Island, the Galapagos Islands, Ecuador), FAR\_Key (JCVI\_SCAF\_1097205236585, from a marine bacterium found at a depth of 1.7 m off the coast of Key West Fla.), and FAR\_Gal (JCVI\_SCAF\_1101670289386, at a depth of 0.1 m at Isabella Island, Galapagos Islands, Ecuador).

[0118] In still other embodiments, the present disclosure provides DNA constructs, vectors and polynucleotides encoding FAR enzymes endogenous to a marine bacterium selected from the group consisting of *M. alkaliphilus*, *M. aquaeolei*, *M. arcticus*, *M. bryozoorum*, *M. daeponensis*, *M. excellens*, *M. flavimaris*, *M. guadonensis*, *M. hydrocarbonoclasticus*, *M. koreensis*, *M. lipolyticus*, *M. litoralis*, *M. lutaoensis*, *M. maritimus*, *M. sediminum*, *M. squalenivirans*, *M. vinifirmus*, *Meptunibacter caesariensis* species strain MED92, *Reinekea* sp. strain MED297, *Marinomonas* sp. strain MED121, unnamed gammaproteobacterium strain HTCC2207, *Marinobacter* sp. strain ELB17, *Oceanobacter kriegii*, and *Oceanobacter* strain WH099.

[0119] In yet other embodiments, the present disclosure provides DNA constructs, vectors and polynucleotides encoding a FAR enzyme endogenous to an organism selected

from the group consisting of *Vitis vinifera*, *Desulfatibacillum alkenivorans*, *Stigmatella aurantiaca* and *Phytophthora ramorum*.

[0120] In some embodiments, the polynucleotide of the invention encodes a FAR enzyme encompassed by the invention and comprises a nucleic acid comprising at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 93%, at least 95%, at least 96%, at least 97%, and at least 99% sequence identity with SEQ ID NOs: 1, 3, 5, 7, or 13 and hybridizes with SEQ ID NOs: 1, 3, 5, 7, or 13. Nucleic acids “hybridize” when they associate, typically in solution. Nucleic acids hybridize due to a variety of well-characterized physico-chemical forces, such as hydrogen bonding, solvent exclusion, base stacking and the like. As used herein, the term “stringent hybridization wash conditions” in the context of nucleic acid hybridization experiments, such as Southern and Northern hybridizations, are sequence dependent, and are different under different environmental parameters. An extensive guide to the hybridization of nucleic acids is found in Tijssen, 1993, “Laboratory Techniques in Biochemistry and Molecular Biology-Hybridization with Nucleic Acid Probes,” Part I, Chapter 2 (Elsevier, New York), which is incorporated herein by reference. For polynucleotides of at least 100 nucleotides in length, low to very high stringency conditions are defined as follows: prehybridization and hybridization at 42° C. in 5×SSPE, 0.3% SDS, 200 µg/ml sheared and denatured salmon sperm DNA, and either 25% formamide for low stringencies, 35% formamide for medium and medium-high stringencies, or 50% formamide for high and very high stringencies, following standard Southern blotting procedures. For polynucleotides of at least 100 nucleotides in length, the carrier material is finally washed three times each for 15 minutes using 2×SSC, 0.2% SDS at least at 50° C. (low stringency), at least at 55° C. (medium stringency), at least at 60° C. (medium-high stringency), at least at 65° C. (high stringency), and at least at 70° C. (very high stringency).

[0121] The polynucleotides encoding FAR enzymes for expression in heterologous recombinant host cells as described herein are operably linked to a promoter, and optionally, to other control sequences. Suitable promoters include constitutive promoters, regulated promoters, and inducible promoters. Appropriate promoter sequences can be obtained from genes encoding extracellular or intracellular polypeptides which are either endogenous or heterologous to the host cell. Methods for the isolation, identification and manipulation of promoters of varying strengths are available in or readily adapted from the art. See e.g., Nevoigt et al. (2006) *Appl. Environ. Microbiol.* 72:5266-5273, the disclosure of which is herein incorporated by reference in its entirety.

[0122] In certain embodiments, the DNA constructs, vectors and polynucleotides are suitable for expression of a heterologous FAR enzyme in bacteria. For bacterial host cells, suitable promoters for directing transcription of the nucleic acid constructs of the present disclosure, include the promoters obtained from the *E. coli* lac operon, *Streptomyces coelicolor* agarase gene (dagA), *Bacillus subtilis* levansucrase gene (sacB), *Bacillus licheniformis* alpha-amylase gene (amyL), *Bacillus stearothermophilus* maltogenic amylase gene (amyM), *Bacillus amyloliquefaciens* alpha-amylase gene (amyQ), *Bacillus licheniformis* penicillinase gene (penP), *Bacillus subtilis* xylA and xylB genes, *Bacillus megaterium* promoters, and prokaryotic beta-lactamase gene

(Villa-Kamaroff et al., *Proc. Natl. Acad. Sci. USA* 75: 3727-3731 (1978)), as well as the tac promoter (DeBoer et al., *Proc. Natl. Acad. Sci. USA* 80: 21-25 (1993)). Additional promoters include trp promoter, phage lambda PL, T7 promoter, promoters found at PromEC (<http://margalit.huji.ac.il/promec/index.html>) and the like. Promoters suitable for use in the present disclosure are described in "Useful proteins from recombinant bacteria" in *Scientific American* 242:74-94 (1980); and in Sambrook et al (2001) *Molecular Cloning: A Laboratory Manual*, 3<sup>rd</sup> ed., Cold Spring Harbor Laboratory Press, New York.

[0123] In various embodiments, the DNA constructs, vectors and polynucleotides are suitable for expression of a heterologous FAR enzyme in yeast. In certain embodiments, the DNA constructs, vectors and polynucleotides are suitable for expression of a heterologous FAR enzyme in oleaginous yeast. In various embodiments, the DNA constructs, vectors and polynucleotides are suitable for expression of a heterologous FAR enzyme in the oleaginous yeast *Y. lipolytica*. In certain embodiments, the DNA constructs, vectors and polynucleotides are suitable for expression of a heterologous FAR enzyme in *S. cerevisiae*. For yeast host cells, suitable promoters for directing transcription of the nucleic acid constructs of the present disclosure are known to the skilled artisan and include, but are not limited to, an enolase (ENO-1\_gene) promoter, a galactokinase (GAL1) promoter, an alcohol dehydrogenase/glyceraldehyde-3-phosphate dehydrogenase (ADH2/GAP) promoter, a translation elongation factor EF-1 alpha (TEF1) promoter as well as those described by Romanos et al. (1992) *Yeast* 8:423-488. In other embodiments, promoters include the TEF1 promoter and an RPS7 promoter.

[0124] For filamentous fungal host cells, suitable promoters for directing the transcription of the nucleic acid constructs of the present disclosure include promoters obtained from the genes for *Aspergillus oryzae* TAKA amylase, *Rhizomucor miehei* aspartic proteinase, *Aspergillus niger* neutral alpha-amylase, *Aspergillus niger* acid stable alpha-amylase, *Aspergillus niger* or *Aspergillus awamori* glucoamylase (glaA), *Rhizomucor miehei* lipase, *Aspergillus oryzae* alkaline protease, *Aspergillus oryzae* triose phosphate isomerase, *Aspergillus nidulans acetamidase*, and *Fusarium oxysporum* trypsin-like protease (WO 96/00787), as well as the NA2-tpi promoter (a hybrid of the promoters from the genes for *Aspergillus niger* neutral alpha-amylase and *Aspergillus oryzae* triose phosphate isomerase), and mutant, truncated, and hybrid promoters thereof. Examples of suitable promoters useful for directing the transcription of the nucleotide constructs of the present invention in a filamentous fungal host cell are promoters such as cbh1, cbh2, egl1, egl2, pepA, hfb1, hfb2, xyn1, amy, and glaA (Nunberg et al., *Mol. Cell. Biol.*, 4:2306-2315 (1984), Boel et al., *EMBO J.* 3:1581-1585 (1984) and EPA 137280).

[0125] In various embodiments, an expression vector optionally contains a ribosome binding site (RBS) for translation initiation, and a transcription terminator, such as PinII. The vector also optionally includes appropriate sequences for amplifying expression, e.g., an enhancer.

[0126] In various embodiments, the polynucleotides useful for expressing heterologous FAR enzymes in recombinant host cells are operably linked to other control sequences, including but not limited to, a transcription terminator sequence, a signal sequence that when translated directs the expressed polypeptide into the secretory pathway of the

recombinant host cell, and a polyadenylation sequence (eukaryotes). The choice of appropriate control sequences for use in the polynucleotide constructs of the present disclosure is within the skill in the art and in various embodiments is dependent on the recombinant host cell used and the desired method of recovering the fatty alcohol compositions produced.

[0127] The instant disclosure further relates to recombinant expression vectors for use in the methods described herein. A recombinant expression vector can be any vector, e.g., a plasmid or a virus, which can be manipulated by recombinant DNA techniques to facilitate expression of a heterologous FAR enzyme in a recombinant host cell. In certain embodiments, the expression vector is integrated into the chromosome of the recombinant host cell and comprises one or more heterologous genes operably linked to one or more control sequences useful for production of a FAR enzyme. In other embodiments, the expression vector is an extra chromosomal replicative DNA molecule, e.g., a linear or closed circular plasmid, that is found either in low copy number (e.g., from about 1 to about 10 copies per genome equivalent) or in high copy number (e.g., more than about 10 copies per genome equivalent). In various embodiments, the expression vector includes a selectable marker, such as a gene that confers antibiotic resistance (e.g., ampicillin, kanamycin, chloramphenicol or tetracycline resistance) to the recombinant host organism that comprises the vector.

[0128] Expression vectors which, in certain embodiments, are useful for expressing FAR enzymes as disclosed herein are commercially available, e.g., from Sigma-Aldrich Chemicals, St. Louis Mo. and Stratagene, LaJolla Calif. In some embodiments, examples of suitable expression vectors are plasmids which are derived from pBR322 (Gibco BRL), pUC (Gibco BRL), pREP4, pCEP4 (Invitrogen) or pPoly (Lathe et al., 1987, *Gene* 57:193-201).

[0129] In certain embodiments, the present disclosure provides a plasmid for expression of heterologous genes in *E. coli*. Expression vector pCK110900, which comprises a P15A origin of replication (P15A ori), lac a CAP binding site, a lac promoter, a T7 ribosomal binding site (T7g10 RBS) and a chloramphenicol resistance gene (camR). This expression vector is depicted in FIG. 3 of U.S. Patent Publication No. 2006/0195947, which is incorporated herein by reference in its entirety.

[0130] In particular embodiments, the present disclosure provides an autonomous replicating plasmid for expression of heterologous genes in *Yarrowia*, and particularly in *Y. lipolytica*. This plasmid vector (pCEN411; FIG. 1) was engineered with cassettes for expression of a gene encoding resistance to hygromycin (HygB<sup>R</sup>) and for expression of a FAR gene optimized for *Y. lipolytica*. In these embodiments, expression of each gene is independently regulated by a strong, constitutive promoter isolated from *Y. lipolytica*: pTEF1 for FAR expression and pRPS7 for HygB<sup>R</sup> expression. When this plasmid was transformed into *Y. lipolytica*, it conferred resistance to hygromycin and directed expression of FAR. This plasmid can be further modified for expression of heterologous genes useful for fatty alcohol production in yeast, inter alia, *Y. lipolytica*.

[0131] In some embodiments, expression vectors as described herein are adapted for over expression of genes encoding enzymes other than FAR that are directly involved in fatty acid biosynthesis. In particular embodiments, the over expressed gene encodes a protein selected from a fatty acid

synthase (FAS), an acyl-ACP thioesterase (TE), a fatty acyl-CoA synthase (FACS) and an acetyl-CoA carboxylase (ACC). In some embodiments, the expression vector encoding the FAR enzyme and the expression vector encoding a second enzyme (e.g., an FAS, TE, FACS or ACC) are separate nucleic acids. In other embodiments, the heterologous FAR enzyme and the second enzyme are encoded on the same expression vector, and expression of each enzyme is independently regulated by a different promoter.

[0132] Methods, reagents and tools for transforming host cells described herein, such as bacteria, yeast (including oleaginous yeast) and filamentous fungi are known in the art. General methods, reagents and tools for transforming, e.g., bacteria can be found, for example, in Sambrook et al (2001) *Molecular Cloning: A Laboratory Manual*, 3<sup>rd</sup> ed., Cold Spring Harbor Laboratory Press, New York. Methods, reagents and tools for transforming yeast are described in "Guide to Yeast Genetics and Molecular Biology," C. Guthrie and G. Fink, Eds., *Methods in Enzymology* 350 (Academic Press, San Diego, 2002). Methods, reagents and tools for transforming *Y. lipolytica* are found in "Yarrowia lipolytica," C. Madzak, J. M. Nicaud and C. Gaillardin in "Production of Recombinant Proteins. Novel Microbial and Eucaryotic Expression Systems," G. Gellissen, Ed. 2005. In some embodiments, introduction of the DNA construct or vector of the present invention into a host cell can be effected by calcium phosphate transfection, DEAE-Dextran mediated transfection, electroporation, or other common techniques (See Davis et al., 1986, *Basic Methods in Molecular Biology*, which is incorporated herein by reference). The present invention also provides a method for producing a recombinant host cell, wherein the method comprises: (a) providing a nucleic acid construct of the present invention, wherein the nucleic acid construct comprises a polynucleotide encoding a FAR polypeptide as described herein; and (b) transforming a host cell with the nucleic acid construct to produce a recombinant cell. In particular embodiments, the host cell will be a bacterial cell and in other particular embodiments the host cell will be a yeast cell.

#### Recombinant Microorganisms for Production of Fatty Alcohols:

[0133] In yet another aspect, the present disclosure provides recombinant microorganisms engineered to express a heterologous FAR enzyme, or functional fragment thereof, as described herein. In certain particular aspects, the present disclosure provides recombinant microorganisms engineered to express a heterologous FAR enzyme from a marine gammaproteobacterium, or functional fragment thereof, as described herein for the production of fatty alcohols from fatty acyl-ACP substrates. In other embodiments, the present disclosure provides recombinant microorganisms engineered to express a heterologous FAR enzyme from an organism selected from the group consisting of *Vitis vinifera*, *Desulfatibacillum alkenivorans*, *Stigmatella aurantiaca* and *Phytophthora ramorum*. In certain embodiments, the FAR enzyme can be expressed in the host cell in which it is found naturally. In various embodiments, the FAR enzyme can be expressed in a heterologous host cell. In particular, when expressed in heterologous host cells described herein, the FAR enzymes are capable of generating high yields of total and secreted fatty alcohols from fatty acyl-ACP substrates.

[0134] In various embodiments, the heterologous host cells useful for expressing the FAR enzymes described herein are

selected from bacteria, yeast or filamentous fungi. In certain embodiments, the yeast is an oleaginous yeast.

[0135] In certain embodiments, microorganisms useful as recombinant host cells are wild-type microorganisms.

[0136] In various embodiments, microorganisms useful as recombinant host cells are genetically modified. As used herein, "genetically modified" microorganisms include microorganisms having one or more endogenous genes removed, microorganisms having one or more endogenous genes with reduced expression compared to the parent or wild-type microorganism, or microorganisms having one or more genes over expressed compared to the parent or wild-type microorganism. In certain embodiments, the one or more genes that are over expressed are endogenous to the microorganism. In some embodiments, the one or more genes that are over expressed are heterologous to the microorganism.

[0137] In certain embodiments, the genetically modified microorganism comprises an inactivated or silenced endogenous gene that codes for a protein involved in the biosynthesis of fatty acyl-CoA substrates. In particular embodiments, the inactive or silenced gene encodes a fatty acyl-ACP thioesterase or a fatty acyl-CoA synthetase (FACS).

[0138] In some embodiments, the host cell is a prokaryotic cell. Suitable prokaryotic cells include gram positive, gram negative and gram-variable bacterial cells. In certain embodiments, host cells include, but are not limited to, species of a genus selected from the group consisting of *Agrobacterium*, *Alicyclobacillus*, *Anabaena*, *Anacystis*, *Acinetobacter*, *Acidothermus*, *Arthrobacter*, *Azobacter*, *Bacillus*, *Bifidobacterium*, *Brevibacterium*, *Butyrivibrio*, *Buchnera*, *Campestris*, *Campilobacter*, *Clostridium*, *Corynebacterium*, *Chromatium*, *Coprococcus*, *Cyanobacteria*, *Escherichia*, *Enterococcus*, *Enterobacter*, *Erwinia*, *Fusobacterium*, *Faecalibacterium*, *Francisella*, *Flavobacterium*, *Geobacillus*, *Haemophilus*, *Helicobacter*, *Klebsiella*, *Lactobacillus*, *Lactococcus*, *Ilyobacter*, *Micrococcus*, *Microbacterium*, *Mesorhizobium*, *Methylobacterium*, *Methylobacterium*, *Mycobacterium*, *Neisseria*, *Pantoea*, *Pseudomonas*, *Prochlorococcus*, *Rhodobacter*, *Rhodopseudomonas*, *Rhodopseudomonas*, *Roseburia*, *Rhodospirillum*, *Rhodococcus*, *Scenedesmus*, *Streptomyces*, *Streptococcus*, *Synechococcus*, *Synechocystis*, *Saccharomonospora*, *Staphylococcus*, *Serratia*, *Salmonella*, *Shigella*, *Thermoanaerobacterium*, *Tropheryma*, *Tularensis*, *Tremecula*, *Thalassiosira*, *Thermosynechococcus*, *Thermococcus*, *Ureaplasma*, *Xanthomonas*, *Xylella*, *Yersinia* and *Zymomonas*. In particular embodiments, the host cell is a species of a genus selected from the group consisting of *Agrobacterium*, *Arthrobacter*, *Clostridium*, *Corynebacterium*, *Escherichia*, *Erwinia*, *Geobacillus*, *Klebsiella*, *Lactobacillus*, *Mycobacterium*, *Pantoea*, *Rhodococcus*, *Streptomyces* and *Zymomonas*.

[0139] In certain embodiments, the recombinant host cell is an industrial bacterial strain. Numerous bacterial industrial strains are known and suitable for use in the methods disclosed herein. In some embodiments, the bacterial host cell is a species of the genus *Bacillus*, e.g., *B. thuringiensis*, *B. anthracis*, *B. megaterium*, *B. subtilis*, *B. lenthilis*, *B. circulans*, *B. pumilus*, *B. laetus*, *B. coagulans*, *B. brevis*, *B. firmus*, *B. alkaophilus*, *B. licheniformis*, *B. clausii*, *B. stearothermophilus*, *B. halodurans* and *B. amyloliquefaciens*. In particular embodiments, the host cell is a species of the genus *Bacillus* and is selected from the group consisting of *B. subtilis*, *B. pumilus*, *B. licheniformis*, *B. clausii*, *B. stearothermophilus*, *B. megaterium* and *B. amyloliquefaciens*.

[0140] In some embodiments the bacterial host cell is a species of the genus *Erwinia*, e.g., *E. uredovora*, *E. carotovora*, *E. ananas*, *E. herbicola*, *E. punctata* or *E. terreus*.

[0141] In other embodiments the bacterial host cell is a species of the genus *Pantoea*, e.g., *P. citrea* or *P. agglomerans*.

[0142] In still other embodiments, the bacterial host cell is a species of the genus *Streptomyces*, e.g., *S. ambofaciens*, *S. achromogenes*, *S. avermitilis*, *S. coelicolor*, *S. aureofaciens*, *S. aureus*, *S. fungicidicus*, *S. griseus* or *S. lividans*.

[0143] In further embodiments, the bacterial host cell is a species of the genus *Zymomonas*, e.g., *Z. mobilis* or *Z. lipolytica*.

[0144] In further embodiments, the bacterial host cell is a species of the genus *Rhodococcus*, e.g. *R. opacus*.

[0145] In particular embodiments, the bacterial host cell is a species of the genus *Escherichia*, e.g., *E. coli*. In certain embodiments, the *E. coli* is a wild-type bacterium. In various embodiments, the wild-type *E. coli* bacterial strain useful in the processes described herein is selected from, but not limited to, strain W3110, strain MG1655 and strain BW25113. In other embodiments, the *E. coli* is genetically modified. Examples of genetically modified *E. coli* useful as recombinant host cells include, but are not limited to, genetically modified *E. coli* found in the Keio Collection, available from the National BioResource Project at NBRP *E. coli*, Microbial Genetics Laboratory, National Institute of Genetics 1111 Yata, Mishima, Shizuoka, 411-8540 Japan (www at shigen.nig.ac.jp/ecoli/strain/top/top.jsp).

[0146] In particular embodiments, the genetically modified *E. coli* comprises an inactivated or silenced endogenous fadD gene, which codes for an acyl-CoA synthetase protein. In other embodiments the genetically modified *E. coli* comprises an inactivated or silenced endogenous fadK gene, which codes for an endogenous short-chain acyl-CoA synthetase. In still other embodiments, the genetically modified *E. coli* comprises an inactivated or silenced endogenous fadD gene and an inactivated or silenced endogenous fadK gene. In other embodiments, the genetically modified *E. coli* comprises an endogenous fadD gene that has reduced expression compared to the parent or wild-type strain. In various embodiments, the genetically modified *E. coli* comprises an endogenous fadK gene that has reduced expression compared to the parent or wild-type strain.

[0147] In certain embodiments, the recombinant host cell is a yeast. In various embodiments, the yeast host cell is a species of a genus selected from the group consisting of *Candida*, *Hansenula*, *Saccharomyces*, *Schizosaccharomyces*, *Pichia*, *Kluyveromyces*, and *Yarrowia*. In particular embodiments, the yeast host cell is a species of a genus selected from the group consisting of *Saccharomyces*, *Candida*, *Pichia* and *Yarrowia*.

[0148] In various embodiments, the yeast host cell is selected from the group consisting of *Hansenula polymorpha*, *Saccharomyces cerevisiae*, *Saccharomyces carlsbergenensis*, *Saccharomyces diastaticus*, *Saccharomyces norbensis*, *Saccharomyces kluyveri*, *Schizosaccharomyces pombe*, *Pichia pastoris*, *Pichia finlandica*, *Pichia trehalophila*, *Pichia fernieri*, *Pichia kodamae*, *Pichia membranefaciens*, *Pichia opuntiae*, *Pichia thermotolerans*, *Pichia salicifolia*, *Pichia quercuum*, *Pichia piperi*, *Pichia stipitis*, *Pichia methanolica*, *Pichia angusta*, *Kluyveromyces lactis*, *Candida albicans*, *Candida krusei*, *Candida ethanolic* and *Yarrowia lipolytica* and synonyms or taxonomic equivalents thereof.

[0149] In certain embodiments, the yeast host cell is a wild-type cell. In various embodiments, the wild-type yeast cell strain is selected from, but not limited to, strain BY4741, strain FL100a, strain INVSC1, strain NRRL Y-390, strain NRRL Y-1438, strain NRRL YB-1952, strain NRRL Y-5997, strain NRRL Y-7567, strain NRRL Y-1532, strain NRRL YB-4149 and strain NRRL Y-567. In other embodiments, the yeast host cell is genetically modified. Examples of genetically modified yeast useful as recombinant host cells include, but are not limited to, genetically modified yeast found in the Open Biosystems collection found at <http://www.openbiosystems.com/GeneExpression/Yeast/YKO/>. See Winzeler et al. (1999) *Science* 285:901-906.

[0150] In other embodiments, the recombinant host cell is an oleaginous yeast. Oleaginous yeasts are organisms that accumulate lipids such as tri-acyl glycerols. Examples of oleaginous yeast include, but are not limited to, organisms selected from the group consisting of *Yarrowia lipolytica*, *Yarrowia paralipolytica*, *Candida revkaufi*, *Candida pulcherrima*, *Candida tropicalis*, *Candida utilis*, *Candida curvata D*, *Candida curvata R*, *Candida diddensiae*, *Candida boldinii*, *Rhodotorula glutinous*, *Rhodotorula graminis*, *Rhodotorula mucilaginosa*, *Rhodotorula minuta*, *Rhodotorula bacarum*, *Rhodosporidium toruloides*, *Cryptococcus (terricolus) albidus* var. *albidus*, *Cryptococcus laurentii*, *Trichosporon pullans*, *Trichosporon cutaneum*, *Trichosporon cutancum*, *Trichosporon pullulans*, *Lipomyces starkeyii*, *Lipomyces lipoferus*, *Lipomyces tetrasporus*, *Endomycopsis vernalis*, *Hansenula ciferri*, *Hansenula saturnus*, and *Trigonopsis variables*. In particular embodiments, the oleaginous yeast is *Y. lipolytica*. In certain embodiments, *Yarrowia lipolytica* strains include, but are not limited to, DSMZ 1345, DSMZ 3286, DSMZ 8218, DSMZ 70561, DSMZ 70562, DSMZ 21175, and also strains available from the Agricultural Research Service (NRRL) such as but not limited to NRRL YB-421, NRRL YB-423, NRRL YB-423-12 and NRRL YB-423-3.

[0151] In certain embodiments, the oleaginous yeast is a wild-type organism. In other embodiments, the oleaginous yeast is genetically modified.

[0152] In yet other embodiments, the recombinant host cell is a filamentous fungus. In certain embodiments, the filamentous fungal host cell is a species of a genus selected from the group consisting of *Achlya*, *Acremonium*, *Aspergillus*, *Aureobasidium*, *Bjerkandera*, *Ceriporiopsis*, *Cephalosporium*, *Chrysosporium*, *Cochliobolus*, *Corynascus*, *Cryphonectria*, *Cryptococcus*, *Coprinus*, *Coriolus*, *Diplodia*, *Endothia*, *Fusarium*, *Gibberella*, *Gliocladium*, *Humicola*, *Hypocreahyphomycetes*, *Myceliophthora*, *Mucor*, *Neurospora*, *Penicillium*, *Podospora*, *Phlebia*, *Piromyces*, *Pyricularia*, *Rhizomucor*, *Rhizopus*, *Schizophyllum*, *Scutellidium*, *Sporotrichum*, *Talaromyces*, *Thermoascus*, *Thielavia*, *Trametes*, *Tolypocladium*, *Trichoderma*, *Verticillium*, *Volvariella*, and teleomorphs, synonyms or taxonomic equivalents thereof.

[0153] In some embodiments, the filamentous fungal host cell is an *Aspergillus* species, a *Chrysosporium* species, a *Corynascus* species, a *Fusarium* species, a *Humicola* species, a *Myceliophthora* species, a *Neurospora* species, a *Penicillium* species, a *Tolypocladium* species, a *Trametes* species, or *Trichoderma* species. In other embodiments, the *Trichoderma* species is selected from *T. longibrachiatum*, *T. viride*, *Hypocreahyphomycetes*, *reesei*; the *Aspergillus* species is selected from *A. awamori*, *A. fumigatus*, *A. japonicus*, *A. nidulans*, *A. niger*, *A. aculeatus*, *A. foetidus*, *A. oryzae*, *A. sojae*, and *A.*

*kawachi*; the *Chrysosporium* species is *C. lucknowense*; the *Fusarium* species is selected from *F. graminum*, *F. oxysporum* and *F. venenatum*; the *Myceliophthora* species is *M. thermophilia*; the *Neurospora* species is *N. crassa*; the *Hemicoloma* species is selected from *H. insolens*, *H. grisea*, and *H. lanuginosa*; the *Penicillium* species is selected from *P. purpurogenum*, *P. chrysogenum*, and *P. verruculosum*; the *Thielavia* species is *T. terrestris*; and the *Trametes* species is selected from *T. villosa* and *T. versicolor*.

[0154] In some embodiments, the filamentous fungal host is a wild-type organism. In other embodiments, the filamentous fungal host is genetically modified.


[0155] In certain particular embodiments, recombinant host cells for use in the methods described herein are derived from strains of *Escherichia coli*, *Bacillus*, *Saccharomyces*, *Streptomyces* and *Yarrowia*.

[0156] In various embodiments, recombinant host cells which are useful in the practice of the present disclosure include prokaryotic and eukaryotic cells which are readily accessible from a number of culture collections, e.g., the American Type Culture Collection (ATCC), Deutsche Sammlung von Mikroorganismen und Zellkulturen GmbH (DSMZ) (German Collection of Microorganisms and Cell Culture), Centraalbureau Voor Schimmelcultures (CBS), and Agricultural Research Service Patent Culture Collection, Northern Regional Research Center (NRRL).

[0157] In various embodiments, the recombinant microorganism over expresses a gene that encodes a FAR enzyme. In some embodiments, the FAR enzyme is endogenous to the microorganism. In other embodiments, the FAR enzyme is heterologous to the microorganism.

[0158] In certain embodiments, the recombinant microorganism over expresses a gene that encodes one or more proteins other than a FAR enzyme. In various embodiments, the one or more over expressed protein increases the rate at which the recombinant cell produces the acyl-thioester FAR substrate, e.g., the compound of formula (I) shown above. In some embodiments, the one or more over expressed genes encodes a protein directly involved in fatty acid biosynthesis. In particular embodiments, the one or more over expressed genes encode a protein selected from a fatty acid synthase (FAS), an acyl-ACP thioesterase (TE), a fatty acyl-CoA synthase (FACS) and an acetyl-CoA carboxylase (ACC). In some embodiments, the over expressed gene is endogenous to the microorganism. In other embodiments, the over expressed gene is heterologous to the microorganism.


[0159] “Fatty acid synthase (FAS)” refers to an enzyme or enzyme complex that catalyzes the conversion of acetyl-CoA and malonyl-CoA units to fatty acyl-ACP as set forth in Scheme II, and reference is also made to FIG. 13:




wherein R has the same meaning as set forth above in the definitions. In certain embodiments, the FAS is composed of

more than one distinct enzymatic activity. In various embodiments, the distinct enzymatic activities reside in separate polypeptides. In some embodiments, the separate polypeptides form one or more protein complexes.


[0160] The term “acyl-ACP thioesterase (TE)” refers to an enzyme that catalyzes the cleavage of acyl-ACP to form a fatty acid, as shown in Scheme III and reference is also made to FIG. 13, wherein R has the same meaning as set forth above:



[0161] “FACS” refers to an enzyme that catalyzes the formation of the covalent complex between the acyl portion of the fatty acid and Coenzyme A as shown below in Scheme IV, wherein R has the same meaning as set forth above:



[0162] The term “acetyl-CoA carboxylase (ACC)” refers to an enzyme that catalyzes the conversion of acetyl-CoA to malonyl-CoA as shown below in Scheme V:



Synthesis of Fatty Alcohols by an Acyl-CoA Independent Pathway:

[0163] FIG. 13 illustrates the biosynthetic pathways for fatty alcohol formation. FAR enzymes can act directly on fatty acyl-CoA substrates to form fatty alcohols. However, as described in more detail herein, the present inventors have discovered that the amount of fatty alcohols produced in certain microorganisms comprising a gene encoding a FAR encompassed by the invention and lacking a functional endogenous fatty acyl-CoA synthetase (FACS) is comparable to the amount of fatty alcohol produced by the microorganism that has a functional endogenous FACS gene. Referring to FIG. 13, a recombinant microorganism that lacks a functional FACS gene does not catalyze the formation of fatty acyl-CoA substrates from fatty acids. Accordingly, the present inventors have posited that FAR enzymes (such as marine gammapro-

teobacterium FAR enzymes) as described herein catalyze the biosynthesis of fatty alcohols from fatty acyl-ACP substrates via an acyl-CoA independent pathway in lieu of, or in addition to, using fatty acyl-CoA substrates.

[0164] In certain embodiments, the present disclosure relates to the use of a FAR enzyme as described herein, e.g., a FAR enzyme from a marine bacterium, for producing a fatty alcohol composition. The discovery that FAR enzymes described herein utilized fatty acyl-ACP substrates instead of or in addition to, fatty acyl-CoA substrates provided for more efficient methods for producing fatty alcohol. In hosts, such as *E. coli* which convert fatty acyl-ACP to fatty acyl-CoA via free fatty acids, the formation of fatty acyl-CoA complexes requires ATP. Thus, in the acyl-CoA independent pathway because production of fatty alcohols by FAR enzymes (e.g., marine bacterial FAR enzymes) from fatty acyl-ACP as described herein bypasses this step, less energy is used to form the fatty alcohol compositions. The present inventors' discovery of the mechanism of action of FARs as encompassed by the invention further allows for the production of fatty alcohols in recombinant cells that do not produce fatty acyl-CoA substrates or that produce low levels of the acyl-CoA substrates compared to wild-type or parent microorganisms. Accordingly fatty alcohols can be produced by a recombinant microorganism that lacks a functional endogenous enzyme involved in the biosynthesis of fatty acyl-CoA substrates. Thus in some aspects, fatty alcohols can be produced by a recombinant microorganism that express a gene encoding a FAR enzyme described herein and that lacks a gene encoding a fatty acyl-CoA synthetase (FACS) and/or a gene encoding a fatty acyl-ACP thioesterase (TE).

[0165] Thus, in some embodiments, the present disclosure provides for a mechanism of action for the production of fatty alcohols by an acyl-CoA independent pathway, wherein certain FAR enzymes encompassed by the invention may act on fatty acyl-ACP substrates in lieu of, or in addition to, acting on fatty acyl-CoA substrates, to produce fatty alcohols. In some embodiments of this aspect, the FAR enzymes from as encompassed by the invention herein including but not limited to FAR enzymes from *Marinobacter* species, such as *Marinobacter* species *algicola* (DG893) and *Marinobacter* *aquaeolei*, and *Oceanobacter* species strain RED65 or functional fragments thereof can produce levels of fatty alcohols in the absence of, or in the presence of low levels of fatty acyl-CoA substrates, which were thought to be the main substrates of FAR enzymes that are comparable to the levels of fatty alcohols produced in the presence of normal levels of fatty acyl-CoA substrates. In further embodiments of this aspect, the invention relates to a recombinant host cell comprising a nucleic acid encoding a FAR enzyme according to the invention and an inactivated or silenced endogenous fatty acyl-ACP thioesterase gene, an inactivated or silenced endogenous fatty acyl-CoA synthetase gene or both. In some embodiments, the recombinant microorganisms provided herein above do not produce detectable levels of fatty acyl-CoA substrates and in other embodiments the recombinant microorganisms produce reduced levels of fatty acyl-CoA substrates as compared to wild-type cells.

[0166] In one preferred embodiment, fatty alcohols are produced by an acyl-CoA independent pathway comprising culturing a recombinant bacterial cells under suitable culture conditions, wherein the recombinant bacterial cell comprises a gene encoding a heterologous FAR polypeptide encompassed by the invention and wherein the recombinant bacte-

rial cell comprises one or more endogenous polynucleotides that have been inactivated or silenced selected from a fadD gene, which codes for an acyl-CoA synthetase protein, a fadK gene, which codes for an endogenous short-chain acyl-CoA synthetase or both; allowing expression of the gene encoding the heterologous FAR enzyme and producing fatty alcohols by the conversion of fatty acyl-ACP substrates.

[0167] In certain embodiments, the recombinant host cell is an *E. coli* cell. In some embodiments of this aspect, the FAR enzymes are derived from *Marinobacter algicola* (DG893), *Marinobacter aquaeolei*, and *Oceanobacter* species strain RED65 or functional fragments thereof. In certain embodiments, the FAR enzyme for use in the methods disclosed herein has an amino acid that has an amino acid sequence that is at least about 75% identical, at least about 80% identical, such as at least about 85% identical, such as at least about 90% identical, such as at least about 91%, such as at least about 92%, such as at least about 93%, such as at least about 94%, such as at least about 95%, such as at least about 96%, such as at least about 97%, such as at least about 98%, such as at least about 99% identical to the sequence set forth in SEQ ID NO: 2, SEQ ID NO: 6, SEQ ID NO: 14, or functional fragments thereof.

#### Methods of Producing Fatty Alcohols and Compositions of Fatty Alcohols:

[0168] In various aspects of the present disclosure, fatty alcohols can be produced in cell-free systems. In certain embodiments, a FAR enzyme as described herein is provided with a substrate and all necessary cofactors to carry out the production of a fatty alcohol from a fatty acyl-thioesterase substrate. In certain aspects, fatty alcohols can be produced by recombinant microorganisms that express a heterologous FAR enzyme as described herein. In particular aspects, fatty alcohols can be produced by recombinant microorganisms that express a heterologous FAR enzyme and one or more proteins other than a FAR enzyme. In some embodiments, the fatty alcohols produced by the recombinant microorganisms are secreted into the nutrient medium. In other embodiments, the fatty alcohols produced by the recombinant microorganisms are extracted from the microorganisms.

[0169] In certain embodiments, the present disclosure relates to the use of a FAR enzyme as described herein, e.g., a FAR enzyme from a marine gamma proteobacterium, for producing a fatty alcohol composition in a cell-free system. In some embodiments, the FAR enzyme is provided with a substrate, e.g., a fatty acyl-thioester of the formula (I) and NAD(P)H under suitable conditions of temperature, pH, and ionic strength and time sufficient for the production of a fatty alcohol composition. In some embodiments, the FAR enzyme is provided with a composition of a fatty acid, Coenzyme A, ATP, NAD(P)H, and a fatty acyl-CoA synthase under suitable conditions of temperature, pH and ionic strength and time sufficient for production of the fatty acyl-CoA substrate of the FAR enzyme by the fatty acyl-CoA synthase and for production of a fatty alcohol composition from the fatty acyl-CoA substrate.

[0170] In certain embodiments of the present disclosure, a fatty alcohol composition is produced by a recombinant host cell comprising a heterologous gene encoding a FAR enzyme as described above, which is cultured in an aqueous nutrient medium comprising an assimilable source of carbon under conditions suitable for production of a fatty alcohol composition. In some embodiments, the fatty alcohol is produced by

a recombinant host cell comprising a heterologous gene encoding a FAR enzyme as described above and the recombinant host cell comprises an inactivated or silenced gene selected from an acyl-ACP thioesterase and a fatty acyl-CoA synthetase.

[0171] In various embodiments, the recombinant host cell is a bacterium. In particular embodiments, the bacterium is *E. coli*. In other embodiments, the recombinant host cell is a yeast. In particular embodiments, the yeast is *S. cerevisiae*. In certain embodiments, the yeast is an oleaginous yeast. In certain particular embodiments, the oleaginous yeast is *Y. lipolytica*.

[0172] In various embodiments, a fatty alcohol composition is produced by a recombinant host cell comprising a heterologous gene encoding a FAR enzyme from a marine bacterium and which over expresses a gene encoding one or more proteins that increases the rate at which the recombinant host cell produces the acyl-thioester FAR substrate, e.g., the compound of formula (I) above. In some embodiments, the one or more over expressed genes encode a protein directly involved in fatty acid biosynthesis. In particular embodiments, the one or more over expressed genes encode a protein selected from a fatty acid synthase (FAS), an acyl-ACP thioesterase (TE), a fatty acyl-CoA synthase (FACS) and an acetyl-CoA carboxylase (ACC).

[0173] "Culturing" or "cultivation" refers to growing a population of microbial cells under suitable conditions in a liquid or solid medium. In particular embodiments, culturing refers to the fermentative bioconversion of a substrate to an end-product. Culturing media are well known and individual components of such culture media are available from commercial sources, e.g., under the Difco™ and BBL™ trademarks. In one non-limiting example, the aqueous nutrient medium is a "rich medium" comprising complex sources of nitrogen, salts, and carbon, such as YP medium, comprising 10 g/L of peptone and 10 g/L yeast extract of such a medium.

[0174] In other non-limiting embodiments, the aqueous nutrient medium comprises a mixture of Yeast Nitrogen Base (Difco) supplemented with an appropriate mixture of amino acids, e.g. SC medium. In particular aspects of this embodiment, the amino acid mixture lacks one or more amino acids, thereby imposing selective pressure for maintenance of an expression vector within the recombinant host cell.

[0175] Fermentation of the recombinant host cell comprising a heterologous FAR gene for production of fatty alcohols is carried out under suitable conditions and for a time sufficient for production of fatty alcohols. Conditions for the culture and production of cells, including bacterial and yeast cells, are readily available. Cell culture media in general are set forth in Atlas and Parks (eds.) *The Handbook of Microbiological Media* (1993) CRC Press, Boca Raton, Fla., which is incorporated herein by reference. Additional information for cell culture is found in available commercial literature such as the *Life Science Research Cell Culture Catalogue* (1998) from Sigma-Aldrich, Inc (St Louis, Mo.) ("Sigma-LSRCCC") and, for example, *The Plant Culture Catalogue* and supplement (1997) also from Sigma-Aldrich, Inc (St Louis, Mo.) ("Sigma-PCCS"), all of which are incorporated herein by reference.

[0176] In some embodiments, cells expressing the heterologous FAR genes as described herein are grown under batch or continuous fermentations conditions. Classical batch fermentation is a closed system, wherein the compositions of the medium is set at the beginning of the fermentation and is

not subject to artificial alternations during the fermentation. A variation of the batch system is a fed-batch fermentation which also finds use in the present invention. In this variation, the substrate is added in increments as the fermentation progresses. Fed-batch systems are useful when catabolite repression is likely to inhibit the metabolism of the cells and where it is desirable to have limited amounts of substrate in the medium. Batch and fed-batch fermentations are common and well known in the art. Continuous fermentation is an open system where a defined fermentation medium is added continuously to a bioreactor and an equal amount of conditioned medium is removed simultaneously for processing. Continuous fermentation generally maintains the cultures at a constant high density where cells are primarily in log phase growth. Continuous fermentation systems strive to maintain steady state growth conditions. Methods for modulating nutrients and growth factors for continuous fermentation processes as well as techniques for maximizing the rate of product formation are well known in the art of industrial microbiology.

[0177] In some embodiments, fermentations are carried out a temperature within the range of from about 10° C. to about 60° C., from about 15° C. to about 50° C., from about 20° C. to about 45° C., and from about 25° C. to about 40° C. In a particular aspect, the fermentation is carried out at a temperature of from about 28° C. and also from about 30° C. In other embodiments, the fermentation is carried out for a period of time within the range of from about 8 hours to 240 hours, from about 8 hours to about 168 hours, from about 16 hours to about 144 hours, from about 16 hours to about 120 hours, or from about 24 hours to about 72 hours. It will be understood that, in certain embodiments where thermostable host cells are used, fermentations may be carried out at higher temperatures. In other embodiments, the fermentation will be carried out at a pH in the range of 4 to 8, in the range of 4.5 to 7.5, in the range of 5 to 7, and also in the range of 5.5 to 6.5.

[0178] Carbon sources useful in the aqueous fermentation medium or broth of the disclosed process in which the recombinant microorganisms are grown are those assimilable by the recombinant host strain. Assimilable carbon sources are available in many forms and include renewable carbon sources and the cellulosic and starch feedstock substrates obtained there from. Such examples include for example monosaccharides, disaccharides, oligosaccharides, saturated and unsaturated fatty acids, succinate, acetate and mixtures thereof. Further carbon sources include, without limitation, glucose, galactose, sucrose, xylose, fructose, glycerol, arabinose, mannose, raffinose, lactose, maltose, and mixtures thereof. In some embodiments, the term "fermentable sugars" is used interchangeably with the term "assimilable carbon source". In one aspect, fermentation is carried out with a mixture of glucose and galactose as the assimilable carbon source. In another aspect, fermentation is carried out with glucose alone to accumulate biomass, after which the glucose is substantially removed and replaced with an inducer, e.g., galactose for induction of expression of one or more heterologous genes involved in fatty alcohol production. In still another aspect, fermentation is carried out with an assimilable carbon source that does not mediate glucose repression, e.g., raffinose, to accumulate biomass, after which the inducer, e.g., galactose, is added to induce expression of one or more heterologous genes involved in fatty alcohol production. In some preferred embodiments, the assimilable carbon source is from cellulosic and starch feedstock derived from but not

limited to, wood, wood pulp, paper pulp, grain, corn stover, corn fiber, rice, paper and pulp processing waste, woody or herbaceous plants, fruit or vegetable pulp, distillers grain, grasses, rice hulls, wheat straw, cotton, hemp, flax, sisal, corn cobs, sugar cane bagasse, switch grass and mixtures thereof.

[0179] In various embodiments, the present disclosure relates to methods of producing a fatty alcohol composition comprising culturing a recombinant host cell comprising a gene expressing a heterologous FAR enzyme in an aqueous nutrient medium comprising an assimilable source of carbon under conditions in which a fatty alcohol composition of produced, and isolating the fatty alcohols.

[0180] In certain embodiments of the disclosed processes, at least 10%, such as at least 20%, such as at least 25%, such as at least 30%, such as at least 40%, such as at least 50%, such as at least 60%, such as at least 70%, such as at least 80%, such as at least 90% of the fatty alcohols produced by the methods described herein are secreted into the culture medium. In particular embodiments, when the recombinant microorganism is *E. coli*, *Yarrowia* (e.g., *Y. lipolytica*) or *Saccharomyces* (e.g., *S. cerevisiae*) at least 25% and also at least 50% of the fatty alcohols produced by the methods encompassed by the invention will be secreted into the culture medium.

[0181] In various embodiments, fatty alcohols produced by the methods of the invention are further recovered or isolated. Recovery or isolating the produced fatty alcohols refers to substantially separating the fatty alcohols from other components of the culture medium or fermentation process. Recovery or isolation may be by solvent extraction of the aqueous nutrient medium with a suitable water immiscible solvent. Extraction may occur simultaneously with fatty alcohol production and extraction may be continuous. Phase separation followed by solvent removal provides the fatty alcohol which may then be further purified and fractionated using methods and equipment known in the art. In other aspects of the disclosure, the secreted fatty alcohols coalesce to form a water immiscible phase that can be directly separated from the aqueous nutrient medium either during the fermentation or after its completion.

[0182] In some embodiments, less than 60%, less than about 50%, such as less than about 40%, such as less than about 30%, such as less than about 20%, such as less than about 10% of the fatty alcohols produced by the methods described herein are secreted from the recombinant host cells. In some embodiments, at least about 30%, such as at least about 40%, such as at least about 45%, such as at least about 50%, such as at least about 60%, such as at least about 65%, such as at least about 70%, such as at least about 75%, such as at least about 80%, such as at least about 85%, such as at least about 90% of the fatty alcohols produced by the methods described herein are secreted from the recombinant host cells. In certain embodiments, fatty alcohols are isolated by separating the cells from the aqueous nutrient medium, for example by centrifugation, resuspension and extraction of the fatty alcohols from the recombinant host cells using an organic solvent or solvent mixture. Suitable protocols for recovering fatty alcohols from recombinant host cells and/or culture medium are known to the skilled artisan. In some embodiments, fatty alcohols may be recovered by first lysing the cells to release the fatty alcohols and then extracting the fatty alcohol from the lysate using conventional means. Reference is also made to Yeast Protocols Handbook, (2009)

Clontech Laboratories, Inc. A Takara Bio Company, Mt. View Calif. 94043 and PNAS 2003 Vol. 100, 16:9156-9161.

[0183] In some embodiments, the present disclosure relates to fatty alcohol compositions produced by recombinant host cells comprising a gene encoding any one of the FAR enzymes as encompassed by the invention and described hereinabove. In some particular embodiments, the FAR enzyme is derived from a *Marinobacter* or *Oceanobacter* species of gammaproteobacterium. In some embodiments, the FAR enzyme is derived from a *Marinobacter* species selected from, but not limited to, *M. algicola* DG893, *M. alkaliphilis*, *M. aquaeoli*, *M. arcticus*, *M. bryozoorum*, *M. daepoensis*, *M. excellens*, *M. flavimaris*, *M. guadonensis*, *M. hydrocarbonoclasticus*, *M. koreensis*, *M. lipolyticus*, *M. litoralis*, *M. lutoensis*, *M. maritimus*, *M. sediminum*, *M. squalenivirans* and *M. vinifirmus*. In various embodiments, the FAR enzyme is obtained from a marine bacterium selected from the group consisting of *Meptuniibacter caesariensis* species strain MED92, *Reinekea* sp. strain MED297, *Marinomonas* sp. strain MED121, unnamed gammaproteobacterium strain HTCC2207 and *Marinobacter* sp. strain ELB17. In some specific embodiments, the FAR enzyme is derived from or endogenous to an *Oceanobacter* species selected from, but not limited to, *Oceanobacter kriegii* and *Oceanobacter* strain WH099.

[0184] In various embodiments, the compositions produced by the methods described herein comprise both saturated and unsaturated fatty alcohols. In certain embodiments, the unsaturated fatty alcohols are monounsaturated fatty alcohols. In some embodiments, the fatty alcohol compositions comprise both saturated and unsaturated fatty alcohols, and the amount of unsaturated fatty alcohols is less than about 30%, such as less than about 20%, such as less than about 10%, such as less than about 5%, such as less than about 1% of the fatty alcohols present in the composition. In other embodiments, the fatty alcohol compositions comprise both saturated and unsaturated fatty alcohols, and the amount of saturated fatty alcohols is less than about 30%, such as less than about 20%, such as less than about 10%, such as less than about 5%, such as less than about 1% of the fatty alcohols present in the composition.

[0185] In some typical embodiments, the fatty alcohol compositions produced by the methods described herein comprise one or more alcohols selected from the group consisting of 1-octanol (C8:0), 1-decanol (C10:0), 1-dodecanol (C12:0), 1-tetradecanol (C14:0), 1-hexadecanol (C16:0), 1-octadecanol (C18:0), 1-icosanol (C20:0), 1-docosanol, 1-tetracosanol, cis Δ<sup>9</sup>-1-hexadecanol, and cis Δ<sup>11</sup>-1-octadecanol.

[0186] In typical embodiments, C8 to C20 fatty alcohols comprise at least about 80%, such as at least about 85%, such as at least about 90%, such as at least about 92%, such as at least about 95%, such as at least about 97%, such as at least about 99% by weight of the total isolated fatty alcohols. In certain embodiments, C10 to C18 fatty alcohols comprise about 80%, such as at least about 85%, such as at least about 90%, such as at least about 92%, such as at least about 95%, such as at least about 97%, such as at least about 99% by weight of the total isolated fatty alcohols. In certain embodiments, C14 to C18 fatty alcohols comprise about 80%, such as at least about 85%, such as at least about 90%, such as at least about 92%, such as at least about 95%, such as at least about 97%, such as at least about 99% by weight of the total isolated fatty alcohols. It is understood that a reference to a

"Cx fatty alcohol" includes both saturated and unsaturated fatty alcohols having "x" carbon atoms.

[0187] In various embodiments, C16 to C18 fatty alcohols comprise at least about 80%, such as at least about 85%, such as at least about 90%, such as at least about 91%, such as at least about 92%, such as at least about 93%, such as at least about 94%, such as at least about 95%, such as at least about 96%, such as at least about 97%, such as at least about 98%, such as at least about 99% by weight of the total isolated fatty alcohols. In certain embodiments, the C16 to C18 fatty alcohols are saturated. In other embodiments the C16 to C18 fatty alcohols are a mixture of saturated and unsaturated fatty alcohols.

[0188] In certain embodiments, the amount of fatty alcohols produced by the methods described herein comprise saturated and/or unsaturated C8 to C24 alcohols produced by a recombinant host cell comprising a heterologous gene encoding a FAR as described herein in a range of about 10 mg/L to about 50 g/L of aqueous nutrient medium, such as in a range of about 10 mg/L to about 5 g/L, such as in a range of about 10 mg/L to about 2 g/L of medium by routine modification of culturing conditions. In particular embodiments, the amount of fatty alcohols produced by the methods described herein is at least about 0.5 g/L, such as at least about 1 g/L, such as at least about 1.5 g/L, such as at least about 2.0 g/L, such as at least about 2.5 g/L, such as at least about 3 g/L, such as at least about 3.5 g/L, such as at least about 4 g/L, such as at least about 4.5 g/L, such as at least about 5 g/L, such as at least about 10 g/L of medium. In various embodiments, the amount of fatty alcohols produced by the methods described herein is at least about 20 g/L, such as at least about 30 g/L, such as at least about 40 g/L, such as at least about 50 g/L of medium. In particular embodiments, the recombinant microorganism comprising a gene encoding FAR enzyme as encompassed by the invention is *E. coli*, *Yarrowia lipolytica* or *Saccharomyces cerevisiae* and the amount of fatty alcohol produced is at least 1.0 g/L, at least 5.0 g/L, at least 10 g/L, at least 15 g/L, at least 20 g/L, at least 25 g/L and at least 30 g/L of medium.

[0189] In some embodiments, the amount of fatty alcohols produced by the methods described herein is in the range of about 100 mg/g to about 5 g/g of dry cell weight. In other embodiments, the amount of fatty alcohols produced by the methods described herein is in the range of about 1 g/g to about 4 g/g of dry cell weight, such as in the range of about 2 g/g to about 3 g/g of dry cell weight by routine modification of culturing conditions.

[0190] In certain embodiments, the amount of fatty alcohols produced by the methods described herein is in the range of about 10% to about 20% of dry cell weight, such as in the range of about 20% to about 30% of dry cell weight, such as in the range of about 30% to about 40% of dry cell weight, such as in the range of about 40% to about 50% of dry cell weight, such as in the range of about 50% to about 60% of dry cell weight, such as in the range of about 60% to about 70% of dry cell weight, such as in the range of about 70% to about 80% of dry cell weight by routine modification of culturing conditions. Fatty alcohol compositions produced by the methods described herein may also include fatty acid derived components, such as but not limited to esters (e.g. acetate and waxes) and fatty acids.

#### Methods of Producing Alkane and/or Alkene Compositions:

[0191] In various embodiments, the fatty alcohol compositions produced by the methods described herein are subjected

to reduction to produce alkanes and/or alkenes having the same carbon chain length as the fatty alcohol starting materials. Without being bound by any particular theory, the hydroxyl group of an alcohol is a poor leaving group, and therefore, in principle a chemical moiety that binds to the oxygen atom of the hydroxyl group to make it a better leaving group can be used to reduce the fatty alcohols described herein.

[0192] Any method known in the art can be used to reduce the fatty alcohols present in the compositions described herein. In some embodiments, reduction of fatty alcohols can be carried out chemically, for example, by a Barton deoxygenation (or Barton-McCombie deoxygenation), a two-step reaction in which the alcohol is first converted to a methyl xanthate or thioimidazoyl carbamate, and the xanthate or thioimidazoyl carbamate is reduced with a tin hydride or trialkylsilane reagent under radical conditions to produce the alkane and/or alkene. See J. J. Li, C. Limberakis, D. A. Pflum *Modern Organic Synthesis in the Laboratory* (Oxford University Press, 2007) at pp. 81-83.

[0193] In some embodiments, reduction of fatty alcohols to the corresponding alkanes and/or alkenes can be accomplished using a microorganism that has a biosynthetic pathway for reducing fatty alcohols. In certain embodiments, the microorganism is a bacterium. In specific embodiments, the bacterium is *Vibrio furnissii* strain M1. In some embodiments, the fatty alcohol compositions produced by the methods described herein are contacted with the appropriate microorganism for reduction to alkanes and/or alkenes. In other embodiments, the fatty alcohol compositions produced by the methods described herein are contacted with membrane fractions from the appropriate microorganism so that the reduction is carried out in a cell free system. See e.g., Park (2005) *J. Bacteriol.* 187(4): 1426-1429.

[0194] In certain embodiments, alkanes and/or alkenes produced by the reduction of fatty alcohols described herein are isolated from the reaction mixture and unreacted fatty alcohol starting materials to produce a composition that comprises substantially all alkanes and/or alkenes. In some embodiments, the alkanes and/or alkenes produced by the reduction of fatty alcohols described herein and the unreacted fatty alcohol starting materials are isolated from the reaction mixture to produce a composition comprising alkanes and/or alkenes and fatty alcohols.

[0195] In certain embodiments, the resulting compositions comprise at least about 60% alkanes and/or alkenes, such as at least about 70% alkanes and/or alkenes, such as at least about 80% alkanes and/or alkenes, such as at least about 85% alkanes and/or alkenes, such as at least about 90% alkanes and/or alkenes, such as at least about 92% alkanes and/or alkenes, such as at least about 95% alkanes and/or alkenes, such as at least about 96% alkanes and/or alkenes, such as at least about 97% alkanes and/or alkenes, such as at least about 98% alkanes and/or alkenes, such as at least about 99% alkanes and/or alkenes by weight of the composition after reduction.

[0196] In other embodiments, the resulting compositions comprise at least about 10% alkanes and/or alkenes, such as at least about 20% alkanes and/or alkenes, such as at least about 30% alkanes and/or alkenes, such as at least about 40% alkanes and/or alkenes, such as at least about 50% alkanes and/or alkenes by weight of the composition after reduction.

[0197] In some typical embodiments, the compositions produced by the methods described herein comprise one or

more alkanes selected from the group consisting of octane, decane, dodecane, tetradecane, hexadecane, octadecane, icosane, docosane, and tetracosane. In other typical embodiments, the compositions produced by the methods described herein comprise one or more alkenes selected from the group consisting of octene, decene, dodecene, tetradecene, hexadecene, octadecene, icosene, docosene, and tetracosene.

[0198] In typical embodiments, C<sub>8</sub> to C<sub>20</sub> alkanes and/or alkenes comprise at least about 80%, such as at least about 85%, such as at least about 90%, such as at least about 92%, such as at least about 95%, such as at least about 97%, such as at least about 99% by weight of the total alkanes and/or alkenes in the composition. In certain embodiments, C10 to C18 alkanes and/or alkenes comprise about 80%, such as at least about 85%, such as at least about 90%, such as at least about 92%, such as at least about 95%, such as at least about 97%, such as at least about 99% by weight of the total alkanes and/or alkenes in the composition.

[0199] In certain embodiments, alkanes and/or alkenes having particular carbon chain lengths can be isolated from longer and/or shorter alkanes and/or alkenes, for example by HPLC. In certain embodiments, alkane and/or alkene compositions that are suitable, e.g., for use in jet fuels, comprise C10 to C14 alkanes and/or alkenes. In other embodiments, alkane and/or alkene compositions that are suitable, e.g., for use in diesel fuels comprise alkanes and/or alkenes that have 16 or more carbons (e.g., C16 or longer-chain alkanes and/or alkenes).

#### Fuel Compositions:

[0200] In certain embodiments, the fatty alcohol compositions described herein and compounds derived there from can be used as components of fuel compositions. In certain embodiments, the fatty alcohol compositions produced by the methods described above can be used directly in fuel compositions. In various embodiments, the fatty alcohols can be reacted with a carboxylic acid to produce acid esters. In particular embodiments, the acid esters are used as components of biodiesel fuel compositions. In other embodiments, the fatty alcohols are reacted with a reducing agent to produce alkanes and/or alkenes. In some embodiments, alkanes and/or alkenes derived from the fatty alcohol compositions are used as components of jet fuel compositions. In other embodiments, alkanes and/or alkenes derived from fatty alcohol compositions are used as components of rocket fuel. In still other embodiments, alkanes and/or alkenes derived from the fatty alcohol compositions are used as components in petro-diesel-like fuel compositions.

[0201] In some embodiments, the fuel compositions comprise an alkane and/or alkene derived from the fatty alcohol compositions described herein. In certain embodiments, the alkanes and/or alkenes have from 6 to 16 carbons and the fuel composition is a kerosene-like fuel composition. In various embodiments, the kerosene-like fuel compositions are included in jet fuel compositions. In particular embodiments, the kerosene-like fuel compositions are included in various grades of jet fuel, including but not limited to, grades Avtur, Jet A, Jet A-1, Jet B, JP-4, JP-5, JP-7 and JP-8. In other embodiments, the kerosene-like fuel compositions are included in fuel compositions for heating. In still other embodiments, the kerosene-like fuel compositions derived from the fatty alcohol compositions described above are

burned with liquid oxygen to provide rocket fuel. In particular embodiments, the kerosene-like fuel compositions are used in RP-1 rocket fuel.

[0202] In some embodiments, the alkanes and/or alkenes derived from the fatty alcohol compositions described herein are used in fuel compositions that are similar to petro-diesel fuel compositions, e.g., that contain saturated and aromatic hydrocarbons. In certain embodiments, the fuel compositions comprise only alkanes and/or alkenes derived from the fatty alcohol compositions described herein. In other embodiments, the fuel compositions comprise alkanes and/or alkenes derived from the fatty alcohol compositions described herein mixed with other components, such as petro-diesel fuel.

[0203] In certain embodiments, the fatty alcohols are further processed with a carboxylic acid to form acid esters. Esterification reactions of fatty alcohols are well-known in the art. In certain embodiments, the transesterification reaction is carried out in the presence of a strong catalyst, e.g., a strong alkaline such as sodium hydroxide. In other embodiments, the reaction is carried out enzymatically using an enzyme that catalyzes the conversion of fatty alcohols to acid esters, such as lipoprotein lipase. See, e.g., Tsujita et al. (1999) *J. Biochem.* 126(6):1074-1079. In various embodiments, the acid esters are used as biodiesel fuel without being mixed with other components. In certain embodiments, the fatty acid esters are mixed with other components, such as petro-diesel fuel.

[0204] In certain embodiments, fatty alcohols, or acid esters or alkanes and/or alkenes derived there from, are combined with other fuels or fuel additives to produce compositions having desired properties for their intended use. Exemplary fuels and fuel additives for particular applications are well-known in the art. Exemplary fuels which can be combined with the compositions described herein include, but are not limited to, traditional fuels such as ethanol and petroleum-based fuels. Exemplary fuel additives which can be combined with the compositions described herein include, but are not limited to, cloud point lowering additives, surfactants, anti-oxidants, metal deactivators, corrosion inhibitors, anti-icing additives, anti-wear additives, deposit-modifying additives and octane enhancers.

#### EXAMPLES

[0205] Various features and embodiments of the disclosure are provided in the following representative examples of the disclosure are provided in the following representative examples, which are intended to be illustrative and not limiting.

##### Example 1

###### Gene Acquisition

[0206] Wild-type *B. mori*, *M. algicola* DG893, and *Oceanobacter* sp. RED65 fatty acyl reductase (FAR) genes were designed for expression in *E. coli*, *S. cerevisiae*, and *Y. lipolytica* based on the reported amino acid sequences (*B. mori* FAR: Moto et al. (2003) *Proc. Nat'l Acad. Sci.* 100: 9156-61; Reiser (1997) GenBank accession number BAC79425.1; *M. algicola* DG893 FAR: GenBank genome accession number NZ\_ABCP01000001.1; and *Oceanobacter* sp RED65 FAR: GenBank genome accession number NZ\_AAQH01000001.1). *M. aquaeolei* FAR gene having GenBank accession number YP\_959486 was codon optimized for expression in *E. coli*. Codon optimization was performed using an algorithm as described in Example 1 of

WO2008/042876, incorporated herein by reference in its entirety. Input codon usage information for the algorithm was either a hybrid of *E. coli*, *S. cerevisiae*, and *Y. lipolytica* usages, or the codon usage for the individual organisms. The genes were synthesized by Genscript (Piscataway, N.J.) with flanking restriction sites for cloning into *E. coli* vector pCK 110900 described in US Patent Publication No. 2006/0195947. Nucleotide sequences for SfiI restriction sites were added to the 5' end and the 3' end of the gene as well as to the t7g10 RBS in front of the ATG start codon.

(SEQ ID NO: 15)  
5' ACAATCTGGATCCGCCAGCCTGGCCATAAGGAGATACAT  
and

(SEQ ID NO: 16)  
3' TAATGAGGCCAAACTGGCCGTCGACACCAGTATG.

The genes were provided in the vector pUC57 by Genscript (Piscataway, N.J.) and the sequences verified by DNA sequencing. The sequences of the codon optimized genes for recombinant host cells for *M. algicola* DG893 FAR, *M. aquaeolei*, *Oceanobacter* sp. RED65 FAR, and *B. mori* FAR are set forth respectively in SEQ ID NOs: 1 and 3 for FAR\_Maa, SEQ ID NO: 13 for FAR\_Maq, SEQ ID NOs: 5 and 7 for FAR\_Osc and SEQ ID NO: 11 for *B. mori* and the corresponding polypeptide sequences are set forth respectively in SEQ ID NOs: 2 and 4 for FAR\_Maa, SEQ ID NO: 14 for FAR\_Maq, SEQ ID NOs: 6 and 8 for FAR\_Osc and SEQ ID NO: 12 for *B. mori*.

### Example 2

Expression and Activity of FARs in *E. coli*

[0207] a. Construction of Vectors to Express FARs in *E. coli*  
[0208] The FAR genes were cloned into the vector pCK110900 (shown in FIG. 3 of US Patent Publication No. 2006/0195947) under the control of a lac promoter using the Sfi I restriction sites. The expression vector also contained the P15a origin of replication and the chloramphenicol resistance gene. The resulting plasmids were introduced into *E. coli* BW25113 ( $\Delta$ fadE) (Baba et al., Molecular Systems Biology, 2006 doi:10.1038/msb4100050, Article Number: 2006.0008) using routine transformation methods.

b. In Vivo Activity of FARs in Recombinant *E. coli* Using Shake Flasks

**[0209]** Recombinant *E. coli* host strains comprising a plasmid containing a heterologous gene encoding either the *B. mori*, *M. algicola* DG893, *M. aquaeolei* or *Oceanobacter* sp. RED65 fatty acyl-CoA reductase, were grown in Luria Bertani Broth (LB) medium supplemented with 1% glucose and 30 µg/mL chloramphenicol (CAM), for approximately 16-18

hours (overnight) at 30° C., 200 rpm. A 5% inoculum was used to initiate fresh 50 mL 2xYT Broth culture supplemented with 30 µg/mL CAM and 0-0.4% glucose. The culture was incubated in a shaker for 2.5 hours at 30° C. and 200 rpm to an OD<sub>600</sub> of about 0.6 to about 0.8, at which point expression of the heterologous FAR was induced with isopropyl-β-D-thiogalactoside (IPTG) (1 mM final concentration). Incubation was continued for about 16 hours (overnight) under the same conditions. Cells were collected by centrifugation for 10 minutes at 6000 rpm in F15B-8×50C rotor. The cell pellets were resuspended in 0.5 mL of 6.7% Na<sub>2</sub>SO<sub>4</sub> and then extracted with 1 mL of isopropanol:methyl t-butyl ether (4/6 ratio) for 2 hrs. The extract was centrifuged and analyzed either directly by GC-FID or GC-MS or derivatized with BSTFA before analysis. For derivatization, a 400 µL sample was taken off the top organic layer, evaporated under a nitrogen stream and the residue was derivatized with 100 µL N,O-Bis(trimethylsilyl)trifluoroacetamide (BSTFA) at 37° C. for 1 hour, and then diluted with 100 µL of heptanes before analysis by GC-FID or GC-MS. 0.5 mL of the culture medium (after removal of cells by centrifugation) was also extracted with 1 mL methyl t-butyl ether for 1 hr. The organic phase was either analyzed directly by GC-FID or GC-MS or derivatized with BSTFA as described above before analysis. In addition, 0.5 mL of the cell culture (before removal of cells by centrifugation) was directly extracted with 1 mL of isopropanol:hexane (4:6 ratio) for 2 hrs. The organic phase was either analyzed directly by GC-FID or GC-MS or derivatized with BSTFA as described above before analysis.

**[0210]** A 1  $\mu$ L sample was analyzed by GC-FID with the split ratio 1:10 using the following conditions: GC-6890N from Agilent Technologies equipped with FID detector and HP-5 column (length 30 m, I.D. 0.32 mm, film 0.25  $\mu$ m). GC method: start at 100°C., increase the temperature with a rate of 25°C./min to 246°C. and hold for 1.96 min. Total run time, 7.8 min. Under the above GC conditions the approximate retention times (min) of produced fatty alcohols and acids are as follows: 5.08, C14:0-OH; 5.40; C14:0-OOOH; 5.74, C16:1-OH, 5.93; C16:0-OH, 6.11; C16:0-OOMe (internal standard); 6.16, C16:1-OOH, 6.29; C16:0-OOH, 6.80; C18:1-OH, 6.90; C18:0-OH, 7.3; C18:0- and C18:1-OOH. Under the conditions tested, expression of the *B. mori*, *M. algicola* DG893 and *Oceanobacter* sp. RED65, FARs in *E. coli* BW25113  $\Delta$ fadE resulted production of fatty alcohols (see Table 2). Identification of individual fatty alcohol was done by comparison to commercial standards (Sigma Chemical Company, 6050 Spruce St. Louis, Mo. 63103). In addition the total production of fatty alcohols by FAR\_Maq was similar to that obtained from FAR\_Maa (*M. algicola* DG893) and produced fatty alcohols include C14:0 (2.5%), C16:1 (10%), C16:0 (34%), C18:1 (53%) and C18:0 (0.5%).

TABLE 2

TABLE 2-continued

Fatty alcohol profile exhibited by recombinant <i>E. coli</i> BW25113 ΔfadE host cells over expressing heterologous FAR genes							Total <sup>b</sup>	Secreted <sup>b</sup>
FAR Enzyme	C12:0	C14:0	C16:1	C16:0	C18:1	C18:0	(mg/L)	(mg/L)
<i>M. algincola</i> DG893	ND	8	30	30	32	ND	820	720

<sup>a</sup>The relative amounts of each fatty alcohol component are expressed as a % of the total fatty alcohols detected via GC-FID. Fatty alcohols include: C12:0 (1-dodecanol), no C12:1 (1-dodecenol) was detected, C14:0 (1-tetradecanol), no C14:1 (1-tetradecenol) was detected, C16:1 (cis A<sup>9</sup>-1-hexadecenol), C16:0 (1-hexahexanol), C18:1 (cis A<sup>11</sup>-1-octadecenol), 18:0 (1-octadecanol). ND: not detected.

<sup>b</sup>Enzyme productivity was estimated using both internal and external standards.

c. In Vivo Activity of FARs in Recombinant *E. coli* Using Fermentors

[0211] In an aerated, agitated stirred tank fermentor, 1.0 L of growth medium containing 11.28 g/L 5×M9 minimal salts (BD, Franklin Lakes, N.J.); 2 g/L yeast extract (BD, Franklin Lakes, N.J.); 3 g/L glucose (Sigma Chemical Co., St. Louis, Mo.); 2 mM MgSO<sub>4</sub> (Sigma Chemical Co., St. Louis, Mo.); 0.1 mM CaCl<sub>2</sub> (Sigma Chemical Co., St. Louis, Mo.); and 30 µg/ml chloroamphenicol (Sigma Chemical Co., St. Louis, Mo.) was brought to a temperature of 30° C. The fermentor was inoculated with a late exponential culture of *E. coli* BW25113 ΔfadE equipped with plasmid containing *M. algincola* DG893 (as described in Example 2b) to a starting optical density at 600 nm (OD<sub>600</sub>) of 0.8 to 1.0. The inoculum was grown in a shake flask containing TB (BD, Franklin Lakes, N.J.); 0.4% glycerol (Sigma Chemical Co., St. Louis, Mo.), and 30 µg/ml chloroamphenicol (Sigma Chemical Co., St. Louis, Mo.) at 30° C. and 250 rpm until OD<sub>600</sub> reached around 5.0. The fermentor was agitated at 400-1000 rpm and air supplied at 1.0 L/min to maintain a minimum dissolved oxygen level of 30% of saturation. The pH of the culture was controlled at 7.0 initially by addition of mixture of 14% v/v ammonium hydroxide and 10% w/v sodium hydroxide. The base for pH control was switched to 20% w/v sodium hydroxide when ammonium concentration in the broth reached 1.5-2.0 g/L. After the culture reached an OD<sub>600</sub> of 5.0, the expression of *M. algincola* DG893 FAR was induced by addition of isopropyl-β-D-thiogalactoside (IPTG) (Sigma Chemical Corp., St. Louis, Mo.) to a final concentration of 1 mM. Growth of the culture and production of fatty alcohol were maintained by addition of a feed solution containing 500 g/L glucose. Addition of glucose (6 g/L per pulse) was triggered whenever pH reached 7.2. Culture was then grown for another 135-140 hours at 30° C. 3 g/L yeast extract shots were added at 55 and 95 hours. Samples were taken at different time points for analysis. Extraction and quantification of fatty alcohols were performed as described in Example 2b. Under the conditions tested, total production and secretion of fatty alcohols were estimated to be ~12-15 and 9-13 g/L, respectively.

### Example 3

#### Expression and Activity of FAR Enzymes in Various *E. coli* Strains

[0212] a. Construction of Vectors to Express FARs in Various *E. coli* Strains

[0213] The FAR genes were cloned into the vector PCK110900-I (shown in FIG. 3 of U.S. Patent Publication

No. 2006/0195947) under the control of a lac promoter as described above. The resulting plasmids were introduced into *E. coli* BW25113 parent and knock-out strains (ΔfadE and ΔfadD, ΔfadK) derived from BW25113 (Baba et al., Molecular Systems Biology (2006) doi:10.1038/msb4100050, Article No. 2006.0008; Morgan-Kiss et al., J. Biol. Chem. (2004), 279: 37324-37333 and Campbell, et al., Molecular Microbiology (2003), 47(3): 793-805) by routine transformation methods.

b. In Vivo Activity of FARs in Various Recombinant *E. coli* Strains Using Shake Flasks

[0214] Recombinant *E. coli* BW25113 strains (parent, ΔfadE, ΔfadD, and ΔfadK knock-outs) comprising a plasmid containing a heterologous gene encoding either *M. algincola* DG893 or *Oceanobacter* sp. RED65 fatty acyl reductase (FAR), were grown on Luria Bertani (LB) medium supplements with 1% glucose and 30 µg/ml chloramphenicol (CAM) for approximately 16-18 hours (overnight) at 30° C., 200 rpm. A 5% inoculum was used to initiate fresh 50 mL LB culture supplemented with 30 µg/mL CAM. The culture was incubated for 2.5 hrs at 30° C. and 200 rpm to an OD<sub>600</sub> of 0.6-0.8, at which point expression of the heterologous FAR gene was induced with isopropyl-β-D-thiogalactoside (IPTG) (1 mM final concentration). Incubation was continued for about 24 hours under the same conditions. Extraction and quantitation of fatty alcohols were performed as described in Example 2b except the use of following GC method: start at 80° C. for 3 mins, increase the temperature at a rate of 50° C./min to 200° C., then increase the temperature at a rate of 10° C./min to 270° C. and finally, increase at 20° C./min to 300° C. and hold for 5 min (total run time 18.8 min). Under the above GC conditions the approximate retention times (mins) of produced fatty alcohols and acids were as follows: 6.90, C12:0-OH; 8.04, C14:0-OH; 8.42, d27-C14:0-OOH (internal standard); 8.53, C14:0-OOH; 9.11, C16:1-OH; 9.23, C16:0-OH; 9.68, C16:1-OOH; 9.80; C16:0-OOH; 10; 46, C18:1-OH; 10.57, C18:0-OH; 11.10, C18:1-OOH, and 11.25, C18:0-OOH. Identification of individual fatty alcohol was done by comparison to commercial standards (Sigma Chemical Company, St Louis, Mo. 63103). Under the conditions tested, expression of the *M. algincola* DG893 and *Oceanobacter* sp. RED65 FAR in *E. coli* BW25113 and the corresponding ΔfadD, ΔfadE and ΔfadK strains resulted in production of fatty alcohols (Tables 3a and 3b). Typical shake flask titer for *E. coli* BW25113 expressing *M. algincola* DG893 and *Oceanobacter* sp. RED65 FARs was 100-200

mg/L. Since both FARs are able to produce significant amounts of fatty alcohols in the absence of the fadD and fadK acyl-CoA synthetases in *E. coli*, these FARs are able to reduce non acyl-CoA substrates.

TABLE 3A

Fatty alcohol profile exhibited by recombinant <i>E. coli</i> host cells expressing a heterologous <i>M. algicola</i> FAR gene		
Recombinant <i>E. coli</i> cells	Deleted protein	Relative production of fatty alcohols <sup>a</sup>
BW25113	—	1.0
BW25113 ΔfadE	Acyl-CoA dehydrogenase	1.0
BW25113 ΔfadD	Acyl-CoA synthetase	0.78
BW25113 ΔfadKlydID	Short chain acyl-CoA synthetase, anaerobic	0.64 <sup>b</sup>

<sup>a</sup>Fatty alcohols include: 15-17% C14:0 (1-tetradecanol), approximately 38% C16:1 (cis Δ<sup>9</sup>-1-hexadecanol), 23-24% C16:0 (1-hexacecanol), 21-22% C18:1 (cis Δ<sup>11</sup>-1-octadecanol), 0-1% C18:0 (1-octadecanol).

<sup>b</sup>Data obtained from evaluation of strains in microtiter plates.

TABLE 3B

Fatty alcohol profile exhibited by recombinant <i>E. coli</i> host cells expressing a heterologous <i>Oceanobacter</i> sp. RED65 FAR		
Recombinant <i>E. coli</i> cells	Deleted protein	Relative production of fatty alcohols <sup>a</sup>
BW25113	—	1.0
BW25113 ΔfadE	Acyl-CoA dehydrogenase	0.78
BW25113 ΔfadD	Acyl-CoA synthetase	0.60

<sup>a</sup>Fatty alcohols include: 0-2% C14:0 (1-tetradecanol), 2-10% C16:1 (cis Δ<sup>9</sup>-1-hexadecanol), 32-36% C16:0 (1-hexacecanol), 56-62% C18:1 (cis Δ<sup>11</sup>-1-octadecanol), 0-1% C18:0 (1-octadecanol).

### c. Extraction and Quantitation of acyl-CoAs from *E. coli* Strains

[0215] Strains were grown as described above in Example 3b. 10 OD<sub>600</sub> units of cells were resuspended in 0.6 ml of ice-cold freshly made extraction buffer (100 mM KH<sub>2</sub>PO<sub>4</sub>, pH 4.9:isopropanol, 1:1), containing 200 ng/mL C17:0-CoA (heptadecanoyl-CoA) as an internal standard, and 0.6 ml of acetonitrile. The cell suspension was vortexed for 5 minutes and centrifuged at 4° C. and 14,000 rpm for 20 minutes. The supernatant was collected and evaporated under a nitrogen stream. The dry residue was resuspended in 200 μl of isopropanol/1 mM acetic acid (4:1) and centrifuged as described above. Supernatant was transferred into a glass vial, and 10 μl was injected into an LC-MS system (AB Sciex 5500 QTRAP) equipped with a Zorbax Extend C18, 4.6 mm×50 mm, 18 μm analytical column. HPLC separation was conducted using a binary gradient with 10% acetonitrile/90% water containing 0.1% NH<sub>4</sub>OH (A) and acetonitrile containing 0.1% NH<sub>4</sub>OH (B) at flow rate 0.5 mL/min. The chromatographic conditions were as follows: 0-1 min 5% solvent B, 1-7 min linear gradient to 95% solvent B, 7-9 min 95% solvent B, The post equilibrium time was 2.5 min. Total run time was 11.5 minutes. LC-MS/MS was performed in a negative mode using parameters described in Table 4. All fatty acyl-CoA standards were obtained from Sigma (St. Louis, Mo.).

TABLE 4

MS parameters for determination of fatty acyl-CoAs					
Fatty acyl-CoA	Ms/MS transition	Declustering potential	Entrance potential	Collision energy	Collision cell exit potential
C14:1	486.7/79	-90	-10	-20	-11
C14:0	487.7/79	-90	-10	-106	-11
C16:1	500.7/79	-75	-10	-100	-9
C16:0	501.7/134	-70	-10	-34	-10
C17:0	508.7/79	-70	-10	-100	-10
C18:1	514.6/79	-70	-10	-20	-10
C18:0	515.6/134	-70	-10	-34	-10

[0216] Relative fatty-acyl-CoA concentrations of different *E. coli* strains are shown in Table 5. Results indicate that disruption of ΔfadD significantly reduces the intracellular concentration of acyl-CoAs. In this example the level of acyl-CoA is reduced by ~8 fold in *E. coli* BW25113 ΔfadD containing *M. algicola* DG893 FAR compared to the strain without the deletion. However, fatty alcohol production, as reported above, is not substantially affected. Therefore, *M. algicola* DG893 FAR appears to be capable of reducing non-CoA substrates, most likely acyl-ACPs, to fatty alcohols.

TABLE 5

Fatty acyl-CoA profile exhibited by recombinant <i>E. coli</i> cells with and without expressing a heterologous <i>M. algicola</i> DG893 FAR			
Recombinant <i>E. coli</i> cells	Deleted protein	FAR enzyme	Total amount of acyl-CoA (ng/OD <sub>600</sub> ) <sup>a</sup>
BW25113	—	—	4.7
BW25113	—	<i>M. algicola</i> DG893	1.5
BW25113 ΔfadE	Acyl-CoA dehydrogenase	—	4.3
BW25113 ΔfadD	Acyl-CoA synthetase	—	Not detectable/trace

<sup>a</sup>Fatty acyl-CoA production was estimated using both internal and external standards. Fatty acyl-CoAs detected by LC-MS/MS include: 11-16% C14:0 (tetradecanoyl-CoA), 2-11% C14:1 (tetradecenoyl-CoA), 28-31% C16:1 (cis Δ<sup>9</sup>-1-hexadecenoyl-CoA), 14-26% C16:0 (hexadecanoyl-CoA), 14-20% C18:1 (cis Δ<sup>11</sup>-1-octadecenoyl-CoA), 9-15% C18:0 (octadecanoyl-CoA).

### Example 4

#### Expression and Activity of FARs in *S. cerevisiae*

[0217] a. Construction of Vectors to Express FARs in *S. cerevisiae*

[0218] The FAR genes were PCR amplified and cloned downstream of the TEF1 promoter with BamHII and Sall sites into pCEN318 to create pCEN319 (*B. mori* FAR), pCEN328 (*M. algicola* DG896 FAR), and pCEN333 (*Oceanobacter* sp. RED65 FAR). pCEN318 was constructed by replacing the KanMX gene of p427-TEF (Sunrise Science Products, San Diego, Calif.) with the hygromycin resistance gene. All FAR plasmids were transformed into *S. cerevisiae* FL100 by routine transformation methods.

b. In Vivo Activity of FARs in Recombinant *S. cerevisiae* Using Shake Flasks

[0219] The recombinant *S. cerevisiae* strains comprising a plasmid containing a heterologous gene encoding either the *B. mori*, *M. algicola* DG893 or *Oceanobacter* sp. RED65 FAR were inoculated into 5 ml of YPD containing 200 μg/mL hygromycin, grown at 30° C. for 48 hours (OD ~8-10).

Approximately 2.5 ml were subcultured into 50 ml of YPD media (20× dilution) containing 200 µg/mL hygromycin and grown in a shaker at 30° C. and 250 rpm for 96 hours. Cell cultures were centrifuged at ~3000-4000 rpm (F15B-8×50C rotor) for 10 minutes, the supernatant was separated from cells. The pellets were washed with 20 ml of 50 mM Tris-HCl pH 7.5. Extraction and analysis of fatty alcohols were performed as described in Example 2b. Under the conditions tested, expression of *B. mori*, *M. algicola* DG893 and *Oceanobacter* sp. RED65 FAR genes in *S. cerevisiae* FL100 resulted in the production of fatty alcohols as shown in Table 6.

TABLE 6

Fatty Alcohol profile exhibited by recombinant <i>S. cerevisiae</i> FL100 host cells over-expressing the heterologous FAR enzyme genes						
FAR	Fatty alcohol composition <sup>a</sup>				Total (mg/L)	Secreted (mg/L)
	C14:0	C16:0	C16:1	C18:0		
<i>B. mori</i>	8	88	1	3	67	20
<i>Oceanobacter</i> sp.	trace	48	trace	52	157	78
RED65						
<i>M. algicola</i> DG893	trace	52	trace	48	300	150

<sup>a</sup>The relative amounts of each fatty alcohol component are expressed as a % of the total fatty alcohols detected using via GC-FID or GC-MS. Endogenous fatty alcohols include: C14:0 (1-tetradecanol), C16:0 (1-hexadecanol), and 18:0 (1-octadecanol). No unsaturated fatty alcohols were detected.

<sup>b</sup>Enzyme productivity was estimated using internal and external standards.

### Example 5

#### Expression and Activity of FAR Enzymes in *Yarrowia lipolytica*

[0220] a. Construction of Vectors to Express FARs in *Yarrowia lipolytica*

[0221] An autonomous replicating plasmid for expression of genes in *Y. lipolytica* was engineered with two antibiotic selection marker cassettes for resistance to hygromycin and phleomycin (HygB(R) or Ble(R), respectively) (named plasmid pCEN354). Expression of each cassette was independently regulated by a strong, constitutive promoter isolated from *Y. lipolytica*: pTEF1 for Ble(R) expression and pRPS7 for HygB(R) expression. Plasmid pCEN354 was used to assemble *Y. lipolytica* expression plasmids. Using “restriction free cloning” methodology, the FAR gene was inserted into pCEN354 to provided plasmid pCEN411 (FIG. 1). In pCEN411, heterologous gene expression is under control of the constitutive TEF1 promoter. The HygB<sup>R</sup> gene allows for selection in media containing hygromycin. Ars18 is an autonomous replicating sequence isolated from *Y. lipolytica* genomic DNA. The resulting plasmid was transformed into *Y. lipolytica* 1345 obtained from the German Resource Centre for Biological Material (DSMZ) using routine transformation methods [(Madzak, C. et al. (2003) *Yarrowia lipolytica*. In G. Gellissen (Ed.), *Production of Recombinant Proteins Novel Microbial and Eukaryotic Expression Systems* (p 163-189)].

b. In Vivo Activity of Heterologous *M. algicola* FAR in Recombinant *Y. lipolytica* Using Shake Flasks

[0222] The recombinant *Y. lipolytica* strain comprising plasmid containing a heterologous gene encoding *M. algicola* DG893 FAR, was inoculated in 200 mL YPD media containing 8% glucose and 500 µg/mL hygromycin. The cultures

were grown at 30° C. to an OD600 of 4-7. Cells were then harvested by centrifugation and washed with 20 ml of 50 mM Tris-HCl pH 7.5. Extraction and identification of intra- and extra-cellular fatty alcohols were performed as described in Example 2b. Under the conditions tested 14 mg/L of 1-hexadecanol was detected. Secreted fatty alcohols were not detected. Using 8% glycerol as carbon source, the total production and secretion of fatty alcohols was estimated to be ~0.8 and 0.2 g/L, respectively. Fatty alcohols include: 30% C16:0 (1-hexadecanol), 56% 18:0 (1-octadecanol) and 12% C18:1 (cis Δ<sup>9</sup>-1-octadecenol).

#### c. In Vivo Activity of Integrated *M. algicola* FAR in *Y. lipolytica* Genome Using 96-Well Microtiter Plates

[0223] An improved *Y. lipolytica* strain (YL1415) for production of fatty alcohols was identified from a library of UV-mutagenized strains defective in growth on media with hexadecane as the sole carbon source. When transformed with pCEN411, YL1415 strain showed 7-10-fold increase in fatty alcohol titer and a significant reduction in the rate of degradation of exogenous 1-hexadecanol in YPD media containing 8% glucose and 500 µg/mL hygromycin compared to that detected from *Y. lipolytica* 1345. Expression of *M. algicola* DG893 FAR was improved in the YL1415 strain by integrating the *M. algicola* DG893 FAR gene at random locations in the genome by non-homologous recombination. An integration cassette consisting of *M. algicola* FAR under control of the TEF promoter and the HygBR gene under control of the RPS promoter was amplified from pCEN411 using primers 411-F1 (AGAGACCGGGTTGGCGG) (SEQ ID NO: 17) and 411-R1 (CATTGCCATTCTGTAACGCTG) (SEQ ID NO: 18).

[0224] Transformation of YL1415 with this integration cassette yielded a library of hygromycin-resistant strains with FAR integrated at various locations in the *Y. lipolytica* genome [(Madzak, C. et al. (2003) *Yarrowia lipolytica*. In G. Gellissen (Ed.), *Production of Recombinant Proteins Novel Microbial and Eukaryotic Expression Systems* (p 163-189)]. This collection of strains with integrated *M. algicola* DG893 FAR in the YL1415 UV mutant strain was grown in 96-well Axygen 96-well plates containing 250 µL YPD supplemented with the total of 2% glucose and 500 µg/mL hygromycin. Plates were incubated in a Kuhner shaker for approximately 40-48 hours at 30° C., 200 rpm and 85% relative humidity. The cell cultures were diluted by transferring 50 µL of overnight grown cultures into the Axygen 96-well plates containing 250 µL YPD supplemented with the total of 2% glucose and 500 µg/mL hygromycin. The plates were incubated for approximately 24-28 hours in a Kuhner shaker under the similar conditions. 20 µL of the cell cultures were then transferred into deep 96-well plates containing 380 µL YPD supplemented with the total of 8% glucose and 500 µg/mL hygromycin. The plates were incubated for approximately 22-26 hours under similar conditions. Cells were collected by centrifugation for 10 minutes at 3500 rpm. Cell pellets were then resuspended in 400 µL of nitrogen limitation media (1.7 g/L yeast nitrogen base, 1.4 g/L (NH<sub>4</sub>)<sub>2</sub>SO<sub>4</sub>, 30 g/L glucose) containing 500 µg/mL Hygromycin and incubated for 22-26 hours in a Kuhner shaker at 30° C., 200 rpm and 85% relative humidity. The cell cultures were then extracted with 1 mL of isopropanol:hexane (4:6 ratio) for 2 hrs. The extracts were centrifuged and the upper organic phase was transferred into polypropylene 96-well plates. Samples were then analyzed using GC-FID method as described in Example 2b. Under the conditions tested, *M. algicola* FAR integrant strains with up

to ~5× improvement in total fatty alcohol production (~0.5-1 g/L) compared to plasmid-based expression were identified. Fatty alcohols include: 70-80% C16:0 (1-hexahecanol), 10-15% 18:0 (1-octadecanol) and 10-15% C18:1 (cis Δ<sup>9</sup>-1-octadecenol).

[0225] All publications, patents, patent applications and other documents cited in this application are hereby incorporated by reference in their entireties for all purposes to the

same extent as if each individual publication, patent, patent application or other document were individually indicated to be incorporated by reference for all purposes.

[0226] While various specific embodiments have been illustrated and described, it will be appreciated that various changes can be made without departing from the spirit and scope of the invention(s).

## SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 18

<210> SEQ ID NO 1  
<211> LENGTH: 1539  
<212> TYPE: DNA  
<213> ORGANISM: Artificial  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic  
<220> FEATURE:  
<221> NAME/KEY: CDS  
<222> LOCATION: (1)...(1539)

<400> SEQUENCE: 1

atg gct act caa caa caa cag aac ggt gca tct gca tcc ggc gtc ttg	48
Met Ala Thr Gln Gln Gln Asn Gly Ala Ser Ala Ser Gly Val Leu	
1                   5                   10                   15	
gaa caa ctt cgt gga aag cac gtt ctt atc aca ggt act acc gga ttt	96
Glu Gln Leu Arg Gly Lys His Val Leu Ile Thr Gly Thr Thr Gly Phe	
20                   25                   30	
ttg ggc aaa gtt gtt ctg gaa aag ttg att cgt act gtt ccg qat att	144
Leu Gly Lys Val Val Leu Glu Lys Leu Ile Arg Thr Val Pro Asp Ile	
35                   40                   45	
gga ggt att cat ctg ctg att cgt ggc aat aaa cgt cat cca gcc gct	192
Gly Gly Ile His Leu Leu Ile Arg Gly Asn Lys Arg His Pro Ala Ala	
50                   55                   60	
cgt gaa cgt ttc ctg aac gaa att gcg tcc tcc tcc gtc ttc gaa cgt	240
Arg Glu Arg Phe Leu Asn Glu Ile Ala Ser Ser Val Phe Glu Arg	
65                   70                   75                   80	
ttg cgt cac gat gat aat gaa gcc ttc gag acc ttc ttg gaa gaa cgt	288
Leu Arg His Asp Asp Asn Glu Ala Phe Glu Thr Phe Leu Glu Glu Arg	
85                   90                   95	
gtt cac tgt att acc ggt gag gtt act gaa tcc cgt ttt ggt ttg aca	336
Val His Cys Ile Thr Gly Glu Val Thr Glu Ser Arg Phe Gly Leu Thr	
100                105                110	
cct gaa cgt ttt cgt gct ttg gcc ggt cag gtt gac gct ttt att aac	384
Pro Glu Arg Phe Arg Ala Leu Ala Gly Gln Val Asp Ala Phe Ile Asn	
115                120                125	
agc gct gca agc gtg aac ttt cgt gag gaa ttg gat aaa gcc ctg aaa	432
Ser Ala Ala Ser Val Asn Phe Arg Glu Glu Leu Asp Lys Ala Leu Lys	
130                135                140	
atc aac acc ttg tgt ctt gaa aat gtt gct gtt gca gaa ttg aac	480
Ile Asn Thr Leu Cys Leu Glu Asn Val Ala Ala Leu Ala Glu Leu Asn	
145                150                155                160	
tcc gct atg gtc att cag gtt tcc act tgt tac gtt aac ggt aaa	528
Ser Ala Met Ala Val Ile Gln Val Ser Thr Cys Tyr Val Asn Gly Lys	
165                170                175	
aac tcc ggt caa att acc gaa tcc gtc att aaa cct gct ggc gaa tcc	576
Asn Ser Gly Gln Ile Thr Glu Ser Val Ile Lys Pro Ala Gly Glu Ser	
180                185                190	

---

-continued

---

att ccc cgt tcc act gac ggt tac tac gag atc gaa gaa ttg gtc cat	624
Ile Pro Arg Ser Thr Asp Gly Tyr Tyr Glu Ile Glu Glu Leu Val His	
195 200 205	
ctg ttg caa gac aag att tcc gat gtt aaa gct cgt tac tcc ggc aaa	672
Leu Leu Gln Asp Lys Ile Ser Asp Val Lys Ala Arg Tyr Ser Gly Lys	
210 215 220	
gtt ctg gag aaa aaa ttg gtt gat ttg ggt att cgt gag gcc aat aat	720
Val Leu Glu Lys Leu Val Asp Leu Gly Ile Arg Glu Ala Asn Asn	
225 230 235 240	
tac gga tgg tcc gac acc tac aca ttc acc aaa tgg ttg ggt gaa caa	768
Tyr Gly Trp Ser Asp Thr Tyr Thr Phe Thr Lys Trp Leu Gly Glu Gln	
245 250 255	
ctg ctg atg aag gcc ttg tct ggt cgt tct ttg act att gtg cgt ccc	816
Leu Leu Met Lys Ala Leu Ser Gly Arg Ser Leu Thr Ile Val Arg Pro	
260 265 270	
tct att att gag tcc gct ttg gaa gaa cct tcc cct ggt tgg atc gaa	864
Ser Ile Ile Glu Ser Ala Leu Glu Pro Ser Pro Gly Trp Ile Glu	
275 280 285	
ggc gtt aaa gtt gcc gat gcc att atc ttg gct tat gcc cgt gaa aaa	912
Gly Val Lys Val Ala Asp Ala Ile Ile Leu Ala Tyr Ala Arg Glu Lys	
290 295 300	
gtt agc ctg ttc cct gga aaa cgt tcc ggc att att gat gtt att cct	960
Val Ser Leu Phe Pro Gly Lys Arg Ser Gly Ile Ile Asp Val Ile Pro	
305 310 315 320	
gtc gat ttg gtt gcg aac tcc atc atc ttg tct ctg gct gag gcg ttg	1008
Val Asp Leu Val Ala Asn Ser Ile Ile Leu Ser Leu Ala Glu Ala Leu	
325 330 335	
tct ggt tct ggt caa cgt cgt att tat caa tgt tgc agc ggt ggt tct	1056
Ser Gly Ser Gly Gln Arg Arg Ile Tyr Gln Cys Cys Ser Gly Gly Ser	
340 345 350	
aat cca atc tcc ctg ggt aag ttc att gat tat ttg atg gcc gag gct	1104
Asn Pro Ile Ser Leu Gly Lys Phe Ile Asp Tyr Leu Met Ala Glu Ala	
355 360 365	
aag acc aac tat gct gcc tac gat caa ctg ttt tat cgt cgt cct act	1152
Lys Thr Asn Tyr Ala Ala Tyr Asp Gln Leu Phe Tyr Arg Arg Pro Thr	
370 375 380	
aaa cct ttc gtc gcc gtc aac cgt aaa ttg ttt gac gtt gtt gtt ggt	1200
Lys Pro Phe Val Ala Val Asn Arg Lys Leu Phe Asp Val Val Gly	
385 390 395 400	
ggt atg cgt gtt cct ctt tct att gcc ggt aaa gct atg cgt ttg gct	1248
Gly Met Arg Val Pro Leu Ser Ile Ala Gly Lys Ala Met Arg Leu Ala	
405 410 415	
ggt caa aat cgt gag ttg aaa gtt ctt aag aac ctt gat acg acc cgt	1296
Gly Gln Asn Arg Glu Leu Lys Val Leu Lys Asn Leu Asp Thr Thr Arg	
420 425 430	
tcc ctt gca acc att ttt ggc ttc tat act gct ccc gac tat atc ttc	1344
Ser Leu Ala Thr Ile Phe Gly Phe Tyr Thr Ala Pro Asp Tyr Ile Phe	
435 440 445	
cgt aac gat agc ttg atg gcc ctg gct tct cgt atg ggt gaa ttg gat	1392
Arg Asn Asp Ser Leu Met Ala Leu Ala Ser Arg Met Gly Glu Leu Asp	
450 455 460	
cgt gtt ctt ttc cca gtt gat gct cgt caa att gat ttg cag ttg tac	1440
Arg Val Leu Phe Pro Val Asp Ala Arg Gln Ile Asp Trp Gln Leu Tyr	
465 470 475 480	
ttg tgg aaa att cat ttg ggt ggt ctg aac cgt tac gct ttg aag gaa	1488
Leu Cys Lys Ile His Leu Gly Gly Leu Asn Arg Tyr Ala Leu Lys Glu	
485 490 495	

---

-continued

---

cgt aaa ctg tat tct ttg cgt gct gat act cgt aaa aaa gct gcc	1536
Arg Lys Leu Tyr Ser Leu Arg Ala Ala Asp Thr Arg Lys Lys Ala Ala	
500	505
510	
 taa	 1539
 <b>&lt;210&gt; SEQ ID NO 2</b>	
<211> LENGTH: 512	
<212> TYPE: PRT	
<213> ORGANISM: Artificial	
<220> FEATURE:	
<223> OTHER INFORMATION: Synthetic Construct	
 <400> SEQUENCE: 2	
Met Ala Thr Gln Gln Gln Asn Gly Ala Ser Ala Ser Gly Val Leu	
1	5
	10
	15
Glu Gln Leu Arg Gly Lys His Val Leu Ile Thr Gly Thr Thr Gly Phe	
20	25
	30
Leu Gly Lys Val Val Leu Glu Lys Leu Ile Arg Thr Val Pro Asp Ile	
35	40
	45
Gly Gly Ile His Leu Leu Ile Arg Gly Asn Lys Arg His Pro Ala Ala	
50	55
	60
Arg Glu Arg Phe Leu Asn Glu Ile Ala Ser Ser Ser Val Phe Glu Arg	
65	70
	75
	80
Leu Arg His Asp Asp Asn Glu Ala Phe Glu Thr Phe Leu Glu Glu Arg	
85	90
	95
Val His Cys Ile Thr Gly Glu Val Thr Glu Ser Arg Phe Gly Leu Thr	
100	105
	110
Pro Glu Arg Phe Arg Ala Leu Ala Gly Gln Val Asp Ala Phe Ile Asn	
115	120
	125
Ser Ala Ala Ser Val Asn Phe Arg Glu Glu Leu Asp Lys Ala Leu Lys	
130	135
	140
Ile Asn Thr Leu Cys Leu Glu Asn Val Ala Ala Leu Ala Glu Leu Asn	
145	150
	155
	160
Ser Ala Met Ala Val Ile Gln Val Ser Thr Cys Tyr Val Asn Gly Lys	
165	170
	175
Asn Ser Gly Gln Ile Thr Glu Ser Val Ile Lys Pro Ala Gly Glu Ser	
180	185
	190
Ile Pro Arg Ser Thr Asp Gly Tyr Tyr Glu Ile Glu Glu Leu Val His	
195	200
	205
Leu Leu Gln Asp Lys Ile Ser Asp Val Lys Ala Arg Tyr Ser Gly Lys	
210	215
	220
Val Leu Glu Lys Lys Leu Val Asp Leu Gly Ile Arg Glu Ala Asn Asn	
225	230
	235
	240
Tyr Gly Trp Ser Asp Thr Tyr Thr Phe Thr Lys Trp Leu Gly Glu Gln	
245	250
	255
Leu Leu Met Lys Ala Leu Ser Gly Arg Ser Leu Thr Ile Val Arg Pro	
260	265
	270
Ser Ile Ile Glu Ser Ala Leu Glu Glu Pro Ser Pro Gly Trp Ile Glu	
275	280
	285
Gly Val Lys Val Ala Asp Ala Ile Ile Leu Ala Tyr Ala Arg Glu Lys	
290	295
	300
Val Ser Leu Phe Pro Gly Lys Arg Ser Gly Ile Ile Asp Val Ile Pro	
305	310
	315
	320

---

-continued

---

Val Asp Leu Val Ala Asn Ser Ile Ile Leu Ser Leu Ala Glu Ala Leu  
325 330 335

Ser Gly Ser Gly Gln Arg Arg Ile Tyr Gln Cys Cys Ser Gly Gly Ser  
340 345 350

Asn Pro Ile Ser Leu Gly Lys Phe Ile Asp Tyr Leu Met Ala Glu Ala  
355 360 365

Lys Thr Asn Tyr Ala Ala Tyr Asp Gln Leu Phe Tyr Arg Arg Pro Thr  
370 375 380

Lys Pro Phe Val Ala Val Asn Arg Lys Leu Phe Asp Val Val Val Gly  
385 390 395 400

Gly Met Arg Val Pro Leu Ser Ile Ala Gly Lys Ala Met Arg Leu Ala  
405 410 415

Gly Gln Asn Arg Glu Leu Lys Val Leu Lys Asn Leu Asp Thr Thr Arg  
420 425 430

Ser Leu Ala Thr Ile Phe Gly Phe Tyr Thr Ala Pro Asp Tyr Ile Phe  
435 440 445

Arg Asn Asp Ser Leu Met Ala Leu Ala Ser Arg Met Gly Glu Leu Asp  
450 455 460

Arg Val Leu Phe Pro Val Asp Ala Arg Gln Ile Asp Trp Gln Leu Tyr  
465 470 475 480

Leu Cys Lys Ile His Leu Gly Gly Leu Asn Arg Tyr Ala Leu Lys Glu  
485 490 495

Arg Lys Leu Tyr Ser Leu Arg Ala Ala Asp Thr Arg Lys Lys Ala Ala  
500 505 510

<210> SEQ ID NO 3  
<211> LENGTH: 1536  
<212> TYPE: DNA  
<213> ORGANISM: Artificial  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic  
<220> FEATURE:  
<221> NAME/KEY: CDS  
<222> LOCATION: (1)..(1536)

<400> SEQUENCE: 3

atg gcc acc cag cag cag aac ggt gca tcc gct tcg ggc gtt ctg	48
Met Ala Thr Gln Gln Gln Asn Gly Ala Ser Ala Ser Gly Val Leu	
1 5 10 15	
gag cag ctt aga ggc aag cat gtc ttg att acc ggt act aca gga ttt	96
Glu Gln Leu Arg Gly Lys His Val Leu Ile Thr Gly Thr Thr Gly Phe	
20 25 30	
ctg gga aag gtg gtt ctg gag aag ctg atc cga acc gtg cct gac atc	144
Leu Gly Lys Val Val Leu Glu Lys Leu Ile Arg Thr Val Pro Asp Ile	
35 40 45	
ggc ggt att cat ctg ctg att aga ggc aac aag aga cat cct gct gcc	192
Gly Gly Ile His Leu Leu Ile Arg Gly Asn Lys Arg His Pro Ala Ala	
50 55 60	
aga gaa aga ttc ttg aac gaa atc gcc tct tcc tct gtc gag cggtt	240
Arg Glu Arg Phe Leu Asn Glu Ile Ala Ser Ser Val Phe Glu Arg	
65 70 75 80	
ctt aga cat gac gac aac gaa gcc ttt gag act ttc ctg gag gag cgt	288
Leu Arg His Asp Asp Asn Glu Ala Phe Glu Thr Phe Leu Glu Glu Arg	
85 90 95	
gtg cac tgc atc acc gga gaa gtc acc gag tcg aga ttt ggc ctt act	336
Val His Cys Ile Thr Gly Glu Val Thr Glu Ser Arg Phe Gly Leu Thr	
100 105 110	

---

-continued

---

cct gag cgg ttc cga gcc ctt gct ggc caa gtg gat gcc ttc atc aat Pro Glu Arg Phe Arg Ala Leu Ala Gly Gln Val Asp Ala Phe Ile Asn 115 120 125	384
tcc gcc gcc tct gtt aac ttc aga gag gag ctg gac aag gca ctc aag Ser Ala Ala Ser Val Asn Phe Arg Glu Glu Leu Asp Lys Ala Leu Lys 130 135 140	432
atc aac acc ctg tgt ctg gag aac gtg gct gct ctg gcc gaa ctt aac Ile Asn Thr Leu Cys Leu Glu Asn Val Ala Ala Leu Ala Glu Leu Asn 145 150 155 160	480
tcc gct atg gca gtg atc caa gtt tcc acc tgt tac gtg aac ggc aag Ser Ala Met Ala Val Ile Gln Val Ser Thr Cys Tyr Val Asn Gly Lys 165 170 175	528
aac tct gga cag atc acc gag tcc gtt atc aag ccc gct ggc gaa tcc Asn Ser Gly Gln Ile Thr Glu Ser Val Ile Lys Pro Ala Gly Glu Ser 180 185 190	576
atc ccc aga tcc aca gat ggc tac tac gag atc gag gag ctg gtc cac Ile Pro Arg Ser Thr Asp Gly Tyr Tyr Glu Ile Glu Glu Leu Val His 195 200 205	624
ctt ctg caa gac aag atc tcc gac gtg aag gct cga tac tct ggc aag Leu Leu Gln Asp Lys Ile Ser Asp Val Lys Ala Arg Tyr Ser Gly Lys 210 215 220	672
gtg ttg gag aag aag ctg gtg gac ctg ggc atc cga gag gcg aac aac Val Leu Glu Lys Leu Val Asp Leu Gly Ile Arg Glu Ala Asn Asn 225 230 235 240	720
tac ggc tgg tct gac acc tac acc ttc acc aaa tgg ctc gga gag cag Tyr Gly Trp Ser Asp Thr Tyr Thr Phe Thr Lys Trp Leu Gly Glu Gln 245 250 255	768
ctt ctg atg aaa gct ctg tcc gga aga tcc ctg act atc gtg cgg oct Leu Leu Met Lys Ala Leu Ser Gly Arg Ser Leu Thr Ile Val Arg Pro 260 265 270	816
tcc atc atc gag tcg gct ctt gaa gag cct tct cca ggt tgg atc gag Ser Ile Ile Glu Ser Ala Leu Glu Pro Ser Pro Gly Trp Ile Glu 275 280 285	864
ggc gtg aag gtt gct gac gcc atc atc ctt gcg tac gcc aga gag aag Gly Val Lys Val Ala Asp Ala Ile Ile Leu Ala Tyr Ala Arg Glu Lys 290 295 300	912
gtt tcg ttg ttc ccc ggc aag cga tct ggc atc atc gac gtt atc ccc Val Ser Leu Phe Pro Gly Lys Arg Ser Gly Ile Ile Asp Val Ile Pro 305 310 315 320	960
gtg gat ctg gtg gcc aac tct atc att ctc tct ctt gct gaa gcc ctt Val Asp Leu Val Ala Asn Ser Ile Ile Leu Ser Leu Ala Glu Ala Leu 325 330 335	1008
tct gga tct ggc cag cgt aga atc tac caa tgt tgt tct ggc ggt tct Ser Gly Ser Gly Gln Arg Arg Ile Tyr Gln Cys Ser Gly Gly Ser 340 345 350	1056
aac ccg att tct ctg ggc aag ttc atc gac tac ctt atg gcc gaa gcc Asn Pro Ile Ser Leu Gly Lys Phe Ile Asp Tyr Leu Met Ala Glu Ala 355 360 365	1104
aag acc aac tat gct gcc tac gac cag ctc ttc tac cga cga ccc acc Lys Thr Asn Tyr Ala Ala Tyr Asp Gln Leu Phe Tyr Arg Arg Pro Thr 370 375 380	1152
aag ccc ttc gtc gct gtg aac cga aag ctg ttc gat gtt gtc gtg gga Lys Pro Phe Val Ala Val Asn Arg Lys Leu Phe Asp Val Val Gly 385 390 395 400	1200
gga atg cga gtg cct tcc att gct ggc aag gcc atg aga ttg gcg Gly Met Arg Val Pro Leu Ser Ile Ala Gly Lys Ala Met Arg Leu Ala 405 410 415	1248

---

-continued

---

ggc cag aat cga gaa ttg aag gtt ctc aag aac ctt gac act act cga	1296
Gly Gln Asn Arg Glu Leu Lys Val Leu Lys Asn Leu Asp Thr Thr Arg	
420 425 430	
tcg ctc gct act atc ttt gga ttc tac act gct cct gac tac atc ttc	1344
Ser Leu Ala Thr Ile Phe Gly Phe Tyr Thr Ala Pro Asp Tyr Ile Phe	
435 440 445	
cgg aat gag tct ctg atg gct ctt gtc cga atg gga gaa ctc gat	1392
Arg Asn Asp Ser Leu Met Ala Leu Ala Ser Arg Met Gly Glu Leu Asp	
450 455 460	
cgt gtg ctg ttc cct gtt gac gct cga cag atc gac tgg cag ctc tac	1440
Arg Val Leu Phe Pro Val Asp Ala Arg Gln Ile Asp Trp Gln Leu Tyr	
465 470 475 480	
ttg tgt aag atc cac ctg ggc ggc ctg aac cga tat gct ctg aaa gaa	1488
Leu Cys Lys Ile His Leu Gly Gly Leu Asn Arg Tyr Ala Leu Lys Glu	
485 490 495	
cga aag ctg tac agc ctt aga gcc gct gat acc cga aag aag gct gct	1536
Arg Lys Leu Tyr Ser Leu Arg Ala Ala Asp Thr Arg Lys Lys Ala Ala	
500 505 510	

&lt;210&gt; SEQ ID NO 4

&lt;211&gt; LENGTH: 512

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Artificial

&lt;220&gt; FEATURE:

&lt;223&gt; OTHER INFORMATION: Synthetic Construct

&lt;400&gt; SEQUENCE: 4

```

Met Ala Thr Gln Gln Gln Asn Gly Ala Ser Ala Ser Gly Val Leu
1           5           10          15

Glu Gln Leu Arg Gly Lys His Val Leu Ile Thr Gly Thr Thr Gly Phe
20          25           30

Leu Gly Lys Val Val Leu Glu Lys Leu Ile Arg Thr Val Pro Asp Ile
35          40           45

Gly Gly Ile His Leu Leu Ile Arg Gly Asn Lys Arg His Pro Ala Ala
50          55           60

Arg Glu Arg Phe Leu Asn Glu Ile Ala Ser Ser Ser Val Phe Glu Arg
65          70           75           80

Leu Arg His Asp Asp Asn Glu Ala Phe Glu Thr Phe Leu Glu Glu Arg
85          90           95

Val His Cys Ile Thr Gly Glu Val Thr Glu Ser Arg Phe Gly Leu Thr
100         105          110

Pro Glu Arg Phe Arg Ala Leu Ala Gly Gln Val Asp Ala Phe Ile Asn
115         120          125

Ser Ala Ala Ser Val Asn Phe Arg Glu Glu Leu Asp Lys Ala Leu Lys
130         135          140

Ile Asn Thr Leu Cys Leu Glu Asn Val Ala Ala Leu Ala Glu Leu Asn
145         150          155          160

Ser Ala Met Ala Val Ile Gln Val Ser Thr Cys Tyr Val Asn Gly Lys
165         170          175

Asn Ser Gly Gln Ile Thr Glu Ser Val Ile Lys Pro Ala Gly Glu Ser
180         185          190

Ile Pro Arg Ser Thr Asp Gly Tyr Tyr Glu Ile Glu Glu Leu Val His
195         200

Leu Leu Gln Asp Lys Ile Ser Asp Val Lys Ala Arg Tyr Ser Gly Lys
210         215          220

```

---

-continued

Val	Leu	Glu	Lys	Lys	Leu	Val	Asp	Leu	Gly	Ile	Arg	Glu	Ala	Asn	Asn
225					230			235							240
Tyr	Gly	Trp	Ser	Asp	Thr	Tyr	Thr	Phe	Thr	Lys	Trp	Leu	Gly	Glu	Gln
					245			250							255
Leu	Leu	Met	Lys	Ala	Leu	Ser	Gly	Arg	Ser	Leu	Thr	Ile	Val	Arg	Pro
					260			265							270
Ser	Ile	Ile	Glu	Ser	Ala	Leu	Glu	Glu	Pro	Ser	Pro	Gly	Trp	Ile	Glu
					275			280							285
Gly	Val	Lys	Val	Ala	Asp	Ala	Ile	Ile	Leu	Ala	Tyr	Ala	Arg	Glu	Lys
					290			295							300
Val	Ser	Leu	Phe	Pro	Gly	Lys	Arg	Ser	Gly	Ile	Ile	Asp	Val	Ile	Pro
					305			310							320
Val	Asp	Leu	Val	Ala	Asn	Ser	Ile	Ile	Leu	Ser	Leu	Ala	Glu	Ala	Leu
					325			330							335
Ser	Gly	Ser	Gly	Gln	Arg	Arg	Ile	Tyr	Gln	Cys	Cys	Ser	Gly	Gly	Ser
					340			345							350
Asn	Pro	Ile	Ser	Leu	Gly	Lys	Phe	Ile	Asp	Tyr	Leu	Met	Ala	Glu	Ala
					355			360							365
Lys	Thr	Asn	Tyr	Ala	Ala	Tyr	Asp	Gln	Leu	Phe	Tyr	Arg	Arg	Pro	Thr
					370			375							380
Lys	Pro	Phe	Val	Ala	Val	Asn	Arg	Lys	Leu	Phe	Asp	Val	Val	Val	Gly
					385			390							400
Gly	Met	Arg	Val	Pro	Leu	Ser	Ile	Ala	Gly	Lys	Ala	Met	Arg	Leu	Ala
					405			410							415
Gly	Gln	Asn	Arg	Glu	Leu	Lys	Val	Leu	Lys	Asn	Leu	Asp	Thr	Thr	Arg
					420			425							430
Ser	Leu	Ala	Thr	Ile	Phe	Gly	Tyr	Thr	Ala	Pro	Asp	Tyr	Ile	Phe	
					435			440							445
Arg	Asn	Asp	Ser	Leu	Met	Ala	Leu	Ala	Ser	Arg	Met	Gly	Glu	Leu	Asp
					450			455							460
Arg	Val	Leu	Phe	Pro	Val	Asp	Ala	Arg	Gln	Ile	Asp	Trp	Gln	Leu	Tyr
					465			470							480
Leu	Cys	Lys	Ile	His	Leu	Gly	Gly	Leu	Asn	Arg	Tyr	Ala	Leu	Lys	Glu
					485			490							495
Arg	Lys	Leu	Tyr	Ser	Leu	Arg	Ala	Ala	Asp	Thr	Arg	Lys	Lys	Ala	Ala
					500			505							510

<210> SEQ ID NO 5  
<211> LENGTH: 1545  
<212> TYPE: DNA  
<213> ORGANISM: Artificial  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic  
<220> FEATURE:  
<221> NAME/KEY: CDS  
<222> LOCATION: (1)..(1545)

<400> SEQUENCE: 5

atg	tct	cag	tac	tcc	ggc	ttc	tcc	gtt	tct	caa	tct	ttg	aag	ggc	aaa
Met	Ser	Gln	Tyr	Ser	Ala	Phe	Ser	Val	Ser	Gln	Ser	Leu	Lys	Gly	Lys
1						5			10						15
cat	atc	ttc	ttg	aca	ggt	gtt	acg	ggt	ttc	ttg	ggc	aag	gcg	att	ctg
His	Ile	Phe	Leu	Thr	Gly	Val	Thr	Gly	Phe	Leu	Gly	Lys	Ala	Ile	Leu
						20			25						30

-continued

---

gag aaa ctg ttg tac tcc gtt cca caa ttg gct cag att cat atc ctg Glu Lys Leu Leu Tyr Ser Val Pro Gln Leu Ala Gln Ile His Ile Leu 35 40 45	144
gtc cgt ggt ggt aaa gtt agc gct aag aag cgt ttc caa cat gac atc Val Arg Gly Gly Lys Val Ser Ala Lys Lys Arg Phe Gln His Asp Ile 50 55 60	192
ttg ggt tct tcc atc ttt gag cgt ctt aag gaa caa cat ggc gaa cat Leu Gly Ser Ser Ile Phe Glu Arg Leu Lys Glu Gln His Gly Glu His 65 70 75 80	240
ttc gaa gaa tgg gtt caa agc aag atc aac ctt gtc gag ggc gaa ctt Phe Glu Glu Trp Val Gln Ser Lys Ile Asn Leu Val Glu Gly Glu Leu 85 90 95	288
act caa cct atg ttc gat ttg cct tct gct gag ttc gct ggc ttg gct Thr Gln Pro Met Phe Asp Leu Pro Ser Ala Glu Phe Ala Gly Leu Ala 100 105 110	336
aac caa ttg gat ctg atc atc aat agc gct gct tct gtc aac ttt cgt Asn Gln Leu Asp Leu Ile Ile Asn Ser Ala Ala Ser Val Asn Phe Arg 115 120 125	384
gag aac ttg gag aag gct ctg aac atc aat acc ctg tgt ctg aac aac Glu Asn Leu Glu Lys Ala Leu Asn Ile Asn Thr Leu Cys Leu Asn Asn 130 135 140	432
atc att gcg ctg gct cag tac aac gtc gct gct caa act cct gtt atg Ile Ile Ala Leu Ala Gln Tyr Asn Val Ala Gln Thr Pro Val Met 145 150 155 160	480
caa atc tcc acc tgc tat gtc aac ggc ttc aac aaa ggc caa atc aac Gln Ile Ser Thr Cys Tyr Val Asn Gly Phe Asn Lys Gly Gln Ile Asn 165 170 175	528
gag gaa gtt gtt ggt cct gct tct ggt att cct cag ttg tcc caa Glu Glu Val Val Gly Pro Ala Ser Gly Leu Ile Pro Gln Leu Ser Gln 180 185 190	576
gac tgc tac gac att gat tcc gtc ttc aag cgt gtt cat tcc cag att Asp Cys Tyr Asp Ile Asp Ser Val Phe Lys Arg Val His Ser Gln Ile 195 200 205	624
gaa caa gtc aag aaa cgt aag acc gac att gaa cag caa gaa caa gcc Glu Gln Val Lys Lys Arg Lys Thr Asp Ile Glu Gln Gln Glu Gln Ala 210 215 220	672
ttg atc aaa ctg ggt atc aag acc tct caa cat ttc ggc tgg aat gat Leu Ile Lys Leu Gly Ile Lys Thr Ser Gln His Phe Gly Trp Asn Asp 225 230 235 240	720
acc tac acc ttc acc aag tgg ctg ggt gaa caa ttg ttg att cag aag Thr Tyr Thr Phe Thr Lys Trp Leu Gly Glu Gln Leu Leu Ile Gln Lys 245 250 255	768
ttg ggc aag caa tct ttg aca att ctt cgt ccg tcc atc atc gaa tct Leu Gly Lys Gln Ser Leu Thr Ile Leu Arg Pro Ser Ile Ile Glu Ser 260 265 270	816
gct gtt cgt gaa cct gct cct ggc tgg gtt gaa ggc gtg aaa gtt gcg Ala Val Arg Glu Pro Ala Pro Gly Trp Val Glu Gly Val Lys Val Ala 275 280 285	864
gat gcc ttg atc tat gct tac gct aaa ggt cgt gtt agc atc ttt ccc Asp Ala Leu Ile Tyr Ala Tyr Ala Lys Gly Arg Val Ser Ile Phe Pro 290 295 300	912
ggt cgt gat gaa ggt atc ttg gat gtc atc cct gtt gat ttg gtt gct Gly Arg Asp Glu Gly Ile Leu Asp Val Ile Pro Val Asp Leu Val Ala 305 310 315 320	960
aat gcc gcc gca ttg tcc gct gct cag ctt atg gaa tct aac cag caa Asn Ala Ala Ala Leu Ser Ala Ala Gln Leu Met Glu Ser Asn Gln Gln 325 330 335	1008

---

-continued

---

act ggc tat cgt atc tat caa tgc tgt tct ggt tcc cgt aat ccg atc Thr Gly Tyr Arg Ile Tyr Gln Cys Cys Ser Gly Ser Arg Asn Pro Ile 340 345 350	1056
aag ctg aaa gag ttc att cgt cac att cag aat gtt gct caa gcc cgt Lys Leu Lys Glu Phe Ile Arg His Ile Gln Asn Val Ala Gln Ala Arg 355 360 365	1104
tac caa gaa tgg cca aag ttg ttt gct gac aaa cct cag gaa gcc ttc Tyr Gln Glu Trp Pro Lys Leu Phe Ala Asp Lys Pro Gln Glu Ala Phe 370 375 380	1152
aag acc gtg tct cct aaa cgt ttc aag ctg tac atg tct ggt ttc act Lys Thr Val Ser Pro Lys Arg Phe Lys Leu Tyr Met Ser Gly Phe Thr 385 390 395 400	1200
gct atc aca tgg gct aag acc atc att ggt cgt gtc ttt gga tct aat Ala Ile Thr Trp Ala Lys Thr Ile Ile Gly Arg Val Phe Gly Ser Asn 405 410 415	1248
gct gcc tcc caa cac atg ttg aag gct aag acc aca gca tct ttg gcg Ala Ala Ser Gln His Met Leu Lys Ala Lys Thr Thr Ala Ser Leu Ala 420 425 430	1296
aac atc ttt ggt ttc tac act gct cct aac tac cgt ttc tcc tct cag Asn Ile Phe Gly Phe Tyr Thr Ala Pro Asn Tyr Arg Phe Ser Ser Gln 435 440 445	1344
aag ttg gaa cag ttg gtc aaa cag ttc gat act acc gaa caa cgt ctt Lys Leu Glu Gln Leu Val Lys Gln Phe Asp Thr Thr Glu Gln Arg Leu 450 455 460	1392
tac gac att cgt gct gac cat ttc gat ttg aag tac tac ttg cag gaa Tyr Asp Ile Arg Ala Asp His Phe Asp Trp Lys Tyr Tyr Leu Gln Glu 465 470 475 480	1440
gtt cac atg gat ggt ctt cac aag tat gcg ttg gcc gat cgt caa gaa Val His Met Asp Gly Leu His Lys Tyr Ala Leu Ala Asp Arg Gln Glu 485 490 495	1488
ttg aag ccc aaa cat gtg aag aag cgt aaa cgt gaa acg att cgt caa Leu Lys Pro Lys His Val Lys Lys Arg Lys Arg Glu Thr Ile Arg Gln 500 505 510	1536
gct gct taa Ala Ala	1545

```

<210> SEQ ID NO 6
<211> LENGTH: 514
<212> TYPE: PRT
<213> ORGANISM: Artificial
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 6

Met Ser Gln Tyr Ser Ala Phe Ser Val Ser Gln Ser Leu Lys Lys
1 5 10 15

His Ile Phe Leu Thr Gly Val Thr Gly Phe Leu Gly Lys Ala Ile Leu
20 25 30

Glu Lys Leu Leu Tyr Ser Val Pro Gln Leu Ala Gln Ile His Ile Leu
35 40 45

Val Arg Gly Gly Lys Val Ser Ala Lys Lys Arg Phe Gln His Asp Ile
50 55 60

Leu Gly Ser Ser Ile Phe Glu Arg Leu Lys Glu Gln His Gly Glu His
65 70 75 80

Phe Glu Glu Trp Val Gln Ser Lys Ile Asn Leu Val Glu Gly Glu Leu
85 90 95

Thr Gln Pro Met Phe Asp Leu Pro Ser Ala Glu Phe Ala Gly Leu Ala

```

---

-continued

---

100	105	110
Asn Gln Leu Asp Leu Ile Ile Asn Ser Ala Ala Ser Val Asn Phe Arg		
115	120	125
Glu Asn Leu Glu Lys Ala Leu Asn Ile Asn Thr Leu Cys Leu Asn Asn		
130	135	140
Ile Ile Ala Leu Ala Gln Tyr Asn Val Ala Ala Gln Thr Pro Val Met		
145	150	155
Gln Ile Ser Thr Cys Tyr Val Asn Gly Phe Asn Lys Gly Gln Ile Asn		
165	170	175
Glu Glu Val Val Gly Pro Ala Ser Gly Leu Ile Pro Gln Leu Ser Gln		
180	185	190
Asp Cys Tyr Asp Ile Asp Ser Val Phe Lys Arg Val His Ser Gln Ile		
195	200	205
Glu Gln Val Lys Lys Arg Lys Thr Asp Ile Glu Gln Gln Glu Gln Ala		
210	215	220
Leu Ile Lys Leu Gly Ile Lys Thr Ser Gln His Phe Gly Trp Asn Asp		
225	230	235
Thr Tyr Thr Phe Thr Lys Trp Leu Gly Glu Gln Leu Ile Gln Lys		
245	250	255
Leu Gly Lys Gln Ser Leu Thr Ile Leu Arg Pro Ser Ile Ile Glu Ser		
260	265	270
Ala Val Arg Glu Pro Ala Pro Gly Trp Val Glu Gly Val Lys Val Ala		
275	280	285
Asp Ala Leu Ile Tyr Ala Tyr Ala Lys Gly Arg Val Ser Ile Phe Pro		
290	295	300
Gly Arg Asp Glu Gly Ile Leu Asp Val Ile Pro Val Asp Leu Val Ala		
305	310	315
Asn Ala Ala Ala Leu Ser Ala Ala Gln Leu Met Glu Ser Asn Gln Gln		
325	330	335
Thr Gly Tyr Arg Ile Tyr Gln Cys Cys Ser Gly Ser Arg Asn Pro Ile		
340	345	350
Lys Leu Lys Glu Phe Ile Arg His Ile Gln Asn Val Ala Gln Ala Arg		
355	360	365
Tyr Gln Glu Trp Pro Lys Leu Phe Ala Asp Lys Pro Gln Glu Ala Phe		
370	375	380
Lys Thr Val Ser Pro Lys Arg Phe Lys Leu Tyr Met Ser Gly Phe Thr		
385	390	395
Ala Ile Thr Trp Ala Lys Thr Ile Ile Gly Arg Val Phe Gly Ser Asn		
405	410	415
Ala Ala Ser Gln His Met Leu Lys Ala Lys Thr Thr Ala Ser Leu Ala		
420	425	430
Asn Ile Phe Gly Phe Tyr Thr Ala Pro Asn Tyr Arg Phe Ser Ser Gln		
435	440	445
Lys Leu Glu Gln Leu Val Lys Gln Phe Asp Thr Thr Glu Gln Arg Leu		
450	455	460
Tyr Asp Ile Arg Ala Asp His Phe Asp Trp Lys Tyr Tyr Leu Gln Glu		
465	470	475
Val His Met Asp Gly Leu His Lys Tyr Ala Leu Ala Asp Arg Gln Glu		
485	490	495
Leu Lys Pro Lys His Val Lys Lys Arg Lys Arg Glu Thr Ile Arg Gln		
500	505	510

---

-continued

---

Ala Ala

```

<210> SEQ ID NO 7
<211> LENGTH: 1542
<212> TYPE: DNA
<213> ORGANISM: Artificial
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic
<220> FEATURE:
<221> NAME/KEY: CDS
<222> LOCATION: (1)..(1542)

<400> SEQUENCE: 7

atg tcc cag tac tcg gct ttc tct gtt tct caa tct ctg aag ggc aag      48
Met Ser Gln Tyr Ser Ala Phe Ser Val Ser Gln Ser Leu Lys Gly Lys
1           5          10          15

cac atc ttt ctc act ggt gtc act ggc ttt ctc gga aag gca att ctg      96
His Ile Phe Leu Thr Gly Val Thr Gly Phe Leu Gly Lys Ala Ile Leu
20          25          30

gag aag ctc ttg tac tcg gtt ccc cag ctg gca cag atc cat atc ctt      144
Glu Lys Leu Leu Tyr Ser Val Pro Gln Leu Ala Gln Ile His Ile Leu
35          40          45

gtg aga ggc ggc aaa gtt tct gcc aag aag cgg ttt cag cac gac atc      192
Val Arg Gly Gly Lys Val Ser Ala Lys Lys Arg Phe Gln His Asp Ile
50          55          60

ctg ggc tct agc atc ttc gag aga ctt aag gag caa cac gga gag cac      240
Leu Gly Ser Ser Ile Phe Glu Arg Leu Lys Glu Gln His Gly Glu His
65          70          75          80

ttt gag gaa tgg gtt cag tcc aag atc aac ctt gtg gag gga gaa ctg      288
Phe Glu Glu Trp Val Gln Ser Lys Ile Asn Leu Val Glu Gly Glu Leu
85          90          95

act caa cca atg ttt gac ctc cct tct gct ggt gga ctt gct gtc aat      336
Thr Gln Pro Met Phe Asp Leu Pro Ser Ala Glu Phe Ala Gly Leu Ala
100         105         110

aac cag ctg gac ctc atc aac tct gcc gct tcg gtt aac ttt cga      384
Asn Gln Leu Asp Leu Ile Ile Asn Ser Ala Ala Ser Val Asn Phe Arg
115         120         125

gag aac ctg gag aag gct ctg aac atc aac acc ctg tgc ctg aac aac      432
Glu Asn Leu Glu Lys Ala Leu Asn Ile Asn Thr Leu Cys Leu Asn Asn
130         135         140

atc atc gct ctg gct cag tac aat gtc gct gcc cag act cct gtg atg      480
Ile Ile Ala Leu Ala Gln Tyr Asn Val Ala Ala Gln Thr Pro Val Met
145         150         155         160

cag att tcc acc tgc tac gtg aac ggc ttc aac aag ggc cag atc aac      528
Gln Ile Ser Thr Cys Tyr Val Asn Gly Phe Asn Lys Gly Gln Ile Asn
165         170         175

gaa gaa gtt gtg gga cct gct tct gga ctg atc cct cag ctg tct caa      576
Glu Glu Val Val Gly Pro Ala Ser Gly Leu Ile Pro Gln Leu Ser Gln
180         185         190

gat tgc tac gac atc gac tcc gtc ttc aag aag aga gtg cat tcc cag att      624
Asp Cys Tyr Asp Ile Asp Ser Val Phe Lys Arg Val His Ser Gln Ile
195         200         205

gag cag gtg aag aag aag aac gac att gag cag cag gaa caa ggc      672
Glu Gln Val Lys Lys Arg Lys Thr Asp Ile Glu Gln Gln Glu Gln Ala
210         215         220

ctt atc aag ctc ggt atc aag act tcc cag cac ttt ggc tgg aat gac      720
Leu Ile Lys Leu Gly Ile Lys Thr Ser Gln His Phe Gly Trp Asn Asp
225         230         235         240

```

---

-continued

---

act tac acc ttc acc aaa tgg ctc ggc gaa cag ctg ttg att cag aag Thr Tyr Thr Phe Thr Lys Trp Leu Gly Glu Gln Leu Leu Ile Gln Lys 245 250 255	768
ctc ggc aag caa tct ctc acc att ctt cga cct tcg atc att gag tct Leu Gly Lys Gln Ser Leu Thr Ile Leu Arg Pro Ser Ile Ile Glu Ser 260 265 270	816
gct gtg aga gag cct gcg ccc gga tgg gtc gaa gga gtg aaa gtc gct Ala Val Arg Glu Pro Ala Pro Gly Trp Val Glu Gly Val Lys Val Ala 275 280 285	864
gac gcc ctt atc tac gct tat gct aag gga aga gtc tcg atc ttt cct Asp Ala Leu Ile Tyr Ala Tyr Ala Lys Gly Arg Val Ser Ile Phe Pro 290 295 300	912
gga aga gac gag ggc atc ctt gat gtg att ccc gtt gac ctt gtt gcc Gly Arg Asp Glu Gly Ile Leu Asp Val Ile Pro Val Asp Leu Val Ala 305 310 315 320	960
aat gct gct gct ctt tct gct gca caa ctc atg gag tcc aac caa cag Asn Ala Ala Ala Leu Ser Ala Ala Gln Leu Met Glu Ser Asn Gln Gln 325 330 335	1008
act ggc tac cga atc tac caa tgc tgc tct gga tct cga aac ccc atc Thr Gly Tyr Arg Ile Tyr Gln Cys Cys Ser Gly Ser Arg Asn Pro Ile 340 345 350	1056
aag ctg aag gag ttc atc cga cac atc cag aac gtg gct cag gcc cga Lys Leu Lys Glu Phe Ile Arg His Ile Gln Asn Val Ala Gln Ala Arg 355 360 365	1104
tac cag gaa tgg cct aag ctg ttt gcc gat aag cct caa gag gcc ttc Tyr Gln Glu Trp Pro Lys Leu Phe Ala Asp Lys Pro Gln Glu Ala Phe 370 375 380	1152
aag act gtc tct ccc aag aga ttc aag ctg tac atg tcc ggc ttc act Lys Thr Val Ser Pro Lys Arg Phe Lys Leu Tyr Met Ser Gly Phe Thr 385 390 395 400	1200
gcc atc acc tgg gct aag acc atc att ggc cga gtg ttc ggc tct aac Ala Ile Thr Trp Ala Lys Thr Ile Ile Gly Arg Val Phe Gly Ser Asn 405 410 415	1248
gca gcc tcc cag cac atg ctg aag gca aag acc aca gct tcc ctg gcc Ala Ala Ser Gln His Met Leu Lys Ala Lys Thr Thr Ala Ser Leu Ala 420 425 430	1296
aac atc ttc ggc ttc tac act gca ccc aac tac cgg ttc tct tcc cag Asn Ile Phe Gly Phe Tyr Thr Ala Pro Asn Tyr Arg Phe Ser Ser Gln 435 440 445	1344
aag ttg gaa cag ctg gtg aag cag ttc gac act acc gag cag cga ctg Lys Leu Glu Gln Leu Val Lys Gln Phe Asp Thr Thr Glu Gln Arg Leu 450 455 460	1392
tac gac att cga gct gac cac ttc gac tgg aag tac tac ctg caa gag Tyr Asp Ile Arg Ala Asp His Phe Asp Trp Lys Tyr Tyr Leu Gln Glu 465 470 475 480	1440
gtt cac atg gat ggc ctt cac aag tac gct ttg gcc gat cgt cag gag Val His Met Asp Gly Leu His Lys Tyr Ala Leu Ala Asp Arg Gln Glu 485 490 495	1488
ctg aaa ccc aag cac gtg aag aag aga aag cgg gag act atc cga caa Leu Lys Pro Lys His Val Lys Lys Arg Lys Arg Glu Thr Ile Arg Gln 500 505 510	1536
gct gcg Ala Ala	1542

&lt;210&gt; SEQ ID NO 8

&lt;211&gt; LENGTH: 514

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Artificial

---

-continued

&lt;220&gt; FEATURE:

&lt;223&gt; OTHER INFORMATION: Synthetic Construct

&lt;400&gt; SEQUENCE: 8

```

Met Ser Gln Tyr Ser Ala Phe Ser Val Ser Gln Ser Leu Lys Gly Lys
1           5          10          15

His Ile Phe Leu Thr Gly Val Thr Gly Phe Leu Gly Lys Ala Ile Leu
20          25          30

Glu Lys Leu Leu Tyr Ser Val Pro Gln Leu Ala Gln Ile His Ile Leu
35          40          45

Val Arg Gly Gly Lys Val Ser Ala Lys Lys Arg Phe Gln His Asp Ile
50          55          60

Leu Gly Ser Ser Ile Phe Glu Arg Leu Lys Glu Gln His Gly Glu His
65          70          75          80

Phe Glu Glu Trp Val Gln Ser Lys Ile Asn Leu Val Glu Gly Glu Leu
85          90          95

Thr Gln Pro Met Phe Asp Leu Pro Ser Ala Glu Phe Ala Gly Leu Ala
100         105         110

Asn Gln Leu Asp Leu Ile Ile Asn Ser Ala Ala Ser Val Asn Phe Arg
115         120         125

Glu Asn Leu Glu Lys Ala Leu Asn Ile Asn Thr Leu Cys Leu Asn Asn
130         135         140

Ile Ile Ala Leu Ala Gln Tyr Asn Val Ala Ala Gln Thr Pro Val Met
145         150         155         160

Gln Ile Ser Thr Cys Tyr Val Asn Gly Phe Asn Lys Gly Gln Ile Asn
165         170         175

Glu Glu Val Val Gly Pro Ala Ser Gly Leu Ile Pro Gln Leu Ser Gln
180         185         190

Asp Cys Tyr Asp Ile Asp Ser Val Phe Lys Arg Val His Ser Gln Ile
195         200         205

Glu Gln Val Lys Lys Arg Lys Thr Asp Ile Glu Gln Gln Glu Gln Ala
210         215         220

Leu Ile Lys Leu Gly Ile Lys Thr Ser Gln His Phe Gly Trp Asn Asp
225         230         235         240

Thr Tyr Thr Phe Thr Lys Trp Leu Gly Glu Gln Leu Leu Ile Gln Lys
245         250         255

Leu Gly Lys Gln Ser Leu Thr Ile Leu Arg Pro Ser Ile Ile Glu Ser
260         265         270

Ala Val Arg Glu Pro Ala Pro Gly Trp Val Glu Gly Val Lys Val Ala
275         280         285

Asp Ala Leu Ile Tyr Ala Tyr Ala Lys Gly Arg Val Ser Ile Phe Pro
290         295         300

Gly Arg Asp Glu Gly Ile Leu Asp Val Ile Pro Val Asp Leu Val Ala
305         310         315         320

Asn Ala Ala Ala Leu Ser Ala Ala Gln Leu Met Glu Ser Asn Gln Gln
325         330         335

Thr Gly Tyr Arg Ile Tyr Gln Cys Cys Ser Gly Ser Arg Asn Pro Ile
340         345         350

Lys Leu Lys Glu Phe Ile Arg His Ile Gln Asn Val Ala Gln Ala Arg
355         360         365

Tyr Gln Glu Trp Pro Lys Leu Phe Ala Asp Lys Pro Gln Glu Ala Phe
370         375         380

```

-continued

Lys	Thr	Val	Ser	Pro	Lys	Arg	Phe	Lys	Leu	Tyr	Met	Ser	Gly	Phe	Thr
385					390				395					400	
Ala	Ile	Thr	Trp	Ala	Lys	Thr	Ile	Ile	Gly	Arg	Val	Phe	Gly	Ser	Asn
					405			410						415	
Ala	Ala	Ser	Gln	His	Met	Leu	Lys	Ala	Lys	Thr	Thr	Ala	Ser	Leu	Ala
					420			425						430	
Asn	Ile	Phe	Gly	Phe	Tyr	Thr	Ala	Pro	Asn	Tyr	Arg	Phe	Ser	Ser	Gln
					435			440						445	
Lys	Leu	Glu	Gln	Leu	Val	Lys	Gln	Phe	Asp	Thr	Thr	Glu	Gln	Arg	Leu
					450			455						460	
Tyr	Asp	Ile	Arg	Ala	Asp	His	Phe	Asp	Trp	Lys	Tyr	Tyr	Leu	Gln	Glu
					465			470						480	
Val	His	Met	Asp	Gly	Leu	His	Lys	Tyr	Ala	Leu	Ala	Asp	Arg	Gln	Glu
					485			490						495	
Leu	Lys	Pro	Lys	His	Val	Lys	Lys	Arg	Lys	Arg	Glu	Thr	Ile	Arg	Gln
					500			505						510	

```

<210> SEQ ID NO 9
<211> LENGTH: 1383
<212> TYPE: DNA
<213> ORGANISM: Artificial
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic
<220> FEATURE:
<221> NAME/KEY: CDS
<222> LOCATION: (1)..(1383)

<400> SEQUENCE: 9

atg tcc cac aat ggt act ctt gat gag cat tat cag aca gtt cgt gaa 48
Met Ser His Asn Gly Thr Leu Asp Glu His Tyr Gln Thr Val Arg Glu
1 5 10 15

ttt tat gac ggt aag agc gtt ttt att act ggc gcc act ggc ttt ctt 96
Phe Tyr Asp Gly Lys Ser Val Phe Ile Thr Gly Ala Thr Gly Phe Leu
20 25 30

gga aaa gcc tac gtt gag aaa ctg gca tat tct tgt ccc gga att gtc 144
Gly Lys Ala Tyr Val Glu Lys Leu Ala Tyr Ser Cys Pro Gly Ile Val
35 40 45

tct att tat att ttg att cgt gat aaa aag ggc tcc aac acg gag gag 192
Ser Ile Tyr Ile Leu Ile Arg Asp Lys Lys Gly Ser Asn Thr Glu Glu
50 55 60

cgt atg cgt aaa tat ttg gac caa ccc att ttc tct cgt att aaa tat 240
Arg Met Arg Lys Tyr Leu Asp Gln Pro Ile Phe Ser Arg Ile Lys Tyr
65 70 75 80

gag cat cca gag tac ttc aaa aag att atc ccc att tct ggc gat att 288
Glu His Pro Glu Tyr Phe Lys Lys Ile Ile Pro Ile Ser Gly Asp Ile
85 90 95

acc gcc cct aaa ctt ggt ctt tgc gac gag gag cgt aac atc ctg att 336
Thr Ala Pro Lys Leu Gly Leu Cys Asp Glu Glu Arg Asn Ile Leu Ile
100 105 110

aat gaa gtg tcc atc gtt atc cac tct gct gcc agc gtt aaa ctg aac 384
Asn Glu Val Ser Ile Val Ile His Ser Ala Ala Ser Val Lys Leu Asn
115 120 125

gat cat ctt aaa ttc acc ttg aac acc aac gtt ggt ggt act atg aaa 432
Asp His Leu Lys Phe Thr Leu Asn Thr Asn Val Gly Gly Thr Met Lys
130 135 140

```

-continued

---

gtc ttg gag ctt gtt aag gag atg aaa aat ttg gcg atg ttt gtg tac Val Leu Glu Leu Val Lys Glu Met Lys Asn Leu Ala Met Phe Val Tyr 145 150 155 160	480
gtt tct acg gct tat tct aac acc agc caa cgt att ttg gaa gaa aaa Val Ser Thr Ala Tyr Ser Asn Thr Ser Gln Arg Ile Leu Glu Glu Lys 165 170 175	528
ttg tac cct cag tct ctg aat ttg aac gaa att cag aaa ttc gct gaa Leu Tyr Pro Gln Ser Leu Asn Leu Asn Glu Ile Gln Lys Phe Ala Glu 180 185 190	576
gag cac tac att ctg ggt aag gat aac gac gaa atg att aaa ttc att Glu His Tyr Ile Leu Gly Lys Asp Asn Asp Glu Met Ile Lys Phe Ile 195 200 205	624
gga aac cat cct aac act tac gcc tac acg aaa gcc ctg gct gag aat Gly Asn His Pro Asn Thr Tyr Ala Tyr Thr Lys Ala Leu Ala Glu Asn 210 215 220	672
ctt gtc gct gaa gaa cat gga gaa att cct act att att cgt cct Leu Val Ala Glu Glu His Gly Glu Ile Pro Thr Ile Ile Arg Pro 225 230 235 240	720
tct atc atc aca gcc tcc gag gaa ccc gtt cgt gga ttt gtt gat Ser Ile Ile Thr Ala Ser Ala Glu Glu Pro Val Arg Gly Phe Val Asp 245 250 255	768
tct tgg agc gga gcc acg gct atg gct gca ttt gca ctt aaa ggc tgg Ser Trp Ser Gly Ala Thr Ala Met Ala Ala Phe Ala Leu Lys Gly Trp 260 265 270	816
aac aac atc atg tat tcc acc ggt gaa gag aac att gac ttg atc ccc Asn Asn Ile Met Tyr Ser Thr Gly Glu Glu Asn Ile Asp Leu Ile Pro 275 280 285	864
ctg gat tac gtt gtc aac ttg aca ctg gtg gct att gct aaa tat aag Leu Asp Tyr Val Val Asn Leu Thr Leu Val Ala Ile Ala Lys Tyr Lys 290 295 300	912
cca acg aaa gag gtt acg gtg tac cat gtt aca acg acg gac ttg aac Pro Thr Lys Glu Val Thr Val Tyr His Val Thr Ser Asp Leu Asn 305 310 315 320	960
ccg att agc atc cgt cgt att ttc atc aaa ctg agc gag ttc gcc tcc Pro Ile Ser Ile Arg Arg Ile Phe Ile Lys Leu Ser Glu Phe Ala Ser 325 330 335	1008
aaa aac cca act agc aac gct gcc cca ttc gct gcc act aca ttg ctg Lys Asn Pro Thr Ser Asn Ala Ala Pro Phe Ala Ala Thr Thr Leu Leu 340 345 350	1056
acc aaa cag aaa ccg ctt att aaa ctg gtg aca ttt ctt atg cag acc Thr Lys Gln Lys Pro Leu Ile Lys Leu Val Thr Phe Leu Met Gln Thr 355 360 365	1104
aca cca gct ttt ttg gca gac ttg tgg atg aaa acg cag cgt aaa gag Thr Pro Ala Phe Leu Ala Asp Leu Trp Met Lys Thr Gln Arg Lys Glu 370 375 380	1152
gcc aag ttc gtg aaa cag cac aac ttg gtc gtt cgt tct cgt gat cag Ala Lys Phe Val Lys Gln His Asn Leu Val Val Arg Ser Arg Asp Gln 385 390 395 400	1200
ctg gag ttc ttt aca tct cag agc tgg ctt ttg cgt tgt gag cgt gca Leu Glu Phe Thr Ser Gln Ser Trp Leu Leu Arg Cys Glu Arg Ala 405 410 415	1248
cgt gtc ctg tcc gcg gcc ttg tcc gat tcc gac cgt gct gtc ttc cgt Arg Val Leu Ser Ala Ala Leu Ser Asp Ser Asp Arg Ala Val Phe Arg 420 425 430	1296
tgc gat cct tcc acg att gat tgg gat cag tat ctt cct atc tac ttc Cys Asp Pro Ser Thr Ile Asp Trp Asp Gln Tyr Leu Pro Ile Tyr Phe 435 440 445	1344

-continued

gaa ggt att aat aaa cac ctg ttc aaa aat aag ttg tag  
 Glu Gly Ile Asn Lys His Leu Phe Lys Asn Lys Leu  
 450 455 460

<210> SEQ ID NO 10  
 <211> LENGTH: 460  
 <212> TYPE: PRT  
 <213> ORGANISM: Artificial  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 10

Met Ser His Asn Gly Thr Leu Asp Glu His Tyr Gln Thr Val Arg Glu  
 1 5 10 15

Phe Tyr Asp Gly Lys Ser Val Phe Ile Thr Gly Ala Thr Gly Phe Leu  
 20 25 30

Gly Lys Ala Tyr Val Glu Lys Leu Ala Tyr Ser Cys Pro Gly Ile Val  
 35 40 45

Ser Ile Tyr Ile Leu Ile Arg Asp Lys Lys Gly Ser Asn Thr Glu Glu  
 50 55 60

Arg Met Arg Lys Tyr Leu Asp Gln Pro Ile Phe Ser Arg Ile Lys Tyr  
 65 70 75 80

Glu His Pro Glu Tyr Phe Lys Lys Ile Ile Pro Ile Ser Gly Asp Ile  
 85 90 95

Thr Ala Pro Lys Leu Gly Leu Cys Asp Glu Glu Arg Asn Ile Leu Ile  
 100 105 110

Asn Glu Val Ser Ile Val Ile His Ser Ala Ala Ser Val Lys Leu Asn  
 115 120 125

Asp His Leu Lys Phe Thr Leu Asn Thr Asn Val Gly Gly Thr Met Lys  
 130 135 140

Val Leu Glu Leu Val Lys Glu Met Lys Asn Leu Ala Met Phe Val Tyr  
 145 150 155 160

Val Ser Thr Ala Tyr Ser Asn Thr Ser Gln Arg Ile Leu Glu Glu Lys  
 165 170 175

Leu Tyr Pro Gln Ser Leu Asn Leu Asn Glu Ile Gln Lys Phe Ala Glu  
 180 185 190

Glu His Tyr Ile Leu Gly Lys Asp Asn Asp Glu Met Ile Lys Phe Ile  
 195 200 205

Gly Asn His Pro Asn Thr Tyr Ala Tyr Thr Lys Ala Leu Ala Glu Asn  
 210 215 220

Leu Val Ala Glu Glu His Gly Glu Ile Pro Thr Ile Ile Ile Arg Pro  
 225 230 235 240

Ser Ile Ile Thr Ala Ser Ala Glu Glu Pro Val Arg Gly Phe Val Asp  
 245 250 255

Ser Trp Ser Gly Ala Thr Ala Met Ala Ala Phe Ala Leu Lys Gly Trp  
 260 265 270

Asn Asn Ile Met Tyr Ser Thr Gly Glu Glu Asn Ile Asp Leu Ile Pro  
 275 280 285

Leu Asp Tyr Val Val Asn Leu Thr Leu Val Ala Ile Ala Lys Tyr Lys  
 290 295 300

Pro Thr Lys Glu Val Thr Val Tyr His Val Thr Thr Ser Asp Leu Asn  
 305 310 315 320

Pro Ile Ser Ile Arg Arg Ile Phe Ile Lys Leu Ser Glu Phe Ala Ser  
 325 330 335

-continued

Lys	Asn	Pro	Thr	Ser	Asn	Ala	Ala	Pro	Phe	Ala	Ala	Thr	Thr	Leu	Leu
						340			345					350	
Thr	Lys	Gln	Lys	Pro	Leu	Ile	Lys	Leu	Val	Thr	Phe	Leu	Met	Gln	Thr
						355			360				365		
Thr	Pro	Ala	Phe	Leu	Ala	Asp	Leu	Trp	Met	Lys	Thr	Gln	Arg	Lys	Glut
						370			375			380			
Ala	Lys	Phe	Val	Lys	Gln	His	Asn	Leu	Val	Val	Arg	Ser	Arg	Asp	Gln
						385			390			395			400
Leu	Glu	Phe	Phe	Thr	Ser	Gln	Ser	Trp	Leu	Leu	Arg	Cys	Glu	Arg	Ala
						405			410				415		
Arg	Val	Leu	Ser	Ala	Ala	Leu	Ser	Asp	Ser	Asp	Arg	Ala	Val	Phe	Arg
						420			425				430		
Cys	Asp	Pro	Ser	Thr	Ile	Asp	Trp	Asp	Gln	Tyr	Leu	Pro	Ile	Tyr	Phe
						435			440				445		
Glu	Gly	Ile	Asn	Lys	His	Leu	Phe	Lys	Asn	Lys	Leu				
						450			455			460			

```
<210> SEQ ID NO 11
<211> LENGTH: 1380
<212> TYPE: DNA
<213> ORGANISM: Artificial
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic
<220> FEATURE:
<221> NAME/KEY: CDS
<222> LOCATION: (1)..(1380)
```

---

-continued

---

145	150	155	160	
gtg tcc acc gct tac tcc aac acc tct cag aga atc ctg gag gag aag Val Ser Thr Ala Tyr Ser Asn Thr Ser Gln Arg Ile Leu Glu Glu Lys 165	170	175		528
ctg tac cct cag tcc ctg aac ctg aac gag atc cag aag ttc gcc gag Leu Tyr Pro Gln Ser Leu Asn Leu Asn Glu Ile Gln Lys Phe Ala Glu 180	185	190		576
gaa cac tac atc ctg ggc aag gac aac gac gag atg atc aag ttc atc Glu His Tyr Ile Leu Gly Lys Asp Asn Asp Glu Met Ile Lys Phe Ile 195	200	205		624
ggc aac cac ccc aac acc tac gct tac acc aaa gcc ctg gct gag aac Gly Asn His Pro Asn Thr Tyr Ala Tyr Thr Lys Ala Leu Ala Glu Asn 210	215	220		672
ctt gtt gcc gaa gag cat gga gag atc ccc acc atc atc atc cga ccc Leu Val Ala Glu Glu His Gly Glu Ile Pro Thr Ile Ile Ile Arg Pro 225	230	235	240	720
tct atc att act gca tcc gcg gag gag cct gtg cga ggc ttc gtc gat Ser Ile Ile Thr Ala Ser Ala Glu Glu Pro Val Arg Gly Phe Val Asp 245	250	255		768
tct tgg tct ggc gct act gct atg gct gct ttc gct ctc aag gga tgg Ser Trp Ser Gly Ala Thr Ala Met Ala Ala Phe Ala Leu Lys Gly Trp 260	265	270		816
aac aac atc atg tat tcc acc ggc gaa gag aac att gac ctg atc cct Asn Asn Ile Met Tyr Ser Thr Gly Glu Glu Asn Ile Asp Leu Ile Pro 275	280	285		864
ctc gac tac gtg gtg aac ctt acc ctg gtg gcc atc gct aag tac aag Leu Asp Tyr Val Val Asn Leu Thr Leu Val Ala Ile Ala Lys Tyr Lys 290	295	300		912
cct acc aaa gag gtg acc gtg tac cac gtc acg act tcg gat ctg aat Pro Thr Lys Glu Val Thr Val Tyr His Val Thr Thr Ser Asp Leu Asn 305	310	315	320	960
ccc atc tcc atc cgg cga atc ttc atc aag ctg tct gag ttt gct tct Pro Ile Ser Ile Arg Arg Ile Phe Ile Lys Leu Ser Glu Phe Ala Ser 325	330	335		1008
aag aac ccg act tct aat gct gct ctt gtc gct gcc act act ctg ctt Lys Asn Pro Thr Ser Asn Ala Ala Pro Phe Ala Ala Thr Thr Leu Leu 340	345	350		1056
acc aag cag aaa ccc ctg atc aag ctg gtt acg ttt ctg atg caa acc Thr Lys Gln Lys Pro Leu Ile Lys Leu Val Thr Phe Leu Met Gln Thr 355	360	365		1104
act cct gcc ttc ctc gct gac ctg tgg atg aag acc cag cga aag gag Thr Pro Ala Phe Leu Ala Asp Leu Trp Met Lys Thr Gln Arg Lys Glu 370	375	380		1152
gcc aag ttc gtc aaa cag cat aac ctc gtc gtt aga tcg cga gat cag Ala Lys Phe Val Lys Gln His Asn Leu Val Val Arg Ser Arg Asp Gln 385	390	395	400	1200
ttg gag ttc ttt acc tcc cag tcc tgg ctg ctt cgt tgt gaa aga gcc Leu Glu Phe Phe Thr Ser Gln Ser Trp Leu Leu Arg Cys Glu Arg Ala 405	410	415		1248
aga gtg ctg tct gct gct ctt agc gac tct gat cgt gcc gtg ttt aga Arg Val Leu Ser Ala Ala Leu Ser Asp Ser Asp Arg Ala Val Phe Arg 420	425	430		1296
tgt gac ccc tcg aca atc gac tgg gat cag tac ctg ccc atc tac ttc Cys Asp Pro Ser Thr Ile Asp Trp Asp Gln Tyr Leu Pro Ile Tyr Phe 435	440	445		1344
gag ggc atc aac aag cac ctg ttc aag aac aag ctc Glu Gly Ile Asn Lys His Leu Phe Lys Asn Lys Leu				1380

---

-continued

---

450

455

460

```

<210> SEQ ID NO 12
<211> LENGTH: 460
<212> TYPE: PRT
<213> ORGANISM: Artificial
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 12

Met Ser His Asn Gly Thr Leu Asp Glu His Tyr Gln Thr Val Arg Glu
1           5          10          15

Phe Tyr Asp Gly Lys Ser Val Phe Ile Thr Gly Ala Thr Gly Phe Leu
20          25          30

Gly Lys Ala Tyr Val Glu Lys Leu Ala Tyr Ser Cys Pro Gly Ile Val
35          40          45

Ser Ile Tyr Ile Leu Ile Arg Asp Lys Lys Gly Ser Asn Thr Glu Glu
50          55          60

Arg Met Arg Lys Tyr Leu Asp Gln Pro Ile Phe Ser Arg Ile Lys Tyr
65          70          75          80

Glu His Pro Glu Tyr Phe Lys Lys Ile Ile Pro Ile Ser Gly Asp Ile
85          90          95

Thr Ala Pro Lys Leu Gly Leu Cys Asp Glu Glu Arg Asn Ile Leu Ile
100         105         110

Asn Glu Val Ser Ile Val Ile His Ser Ala Ala Ser Val Lys Leu Asn
115         120         125

Asp His Leu Lys Phe Thr Leu Asn Thr Asn Val Gly Gly Thr Met Lys
130         135         140

Val Leu Glu Leu Val Lys Glu Met Lys Asn Leu Ala Met Phe Val Tyr
145         150         155         160

Val Ser Thr Ala Tyr Ser Asn Thr Ser Gln Arg Ile Leu Glu Glu Lys
165         170         175

Leu Tyr Pro Gln Ser Leu Asn Leu Asn Glu Ile Gln Lys Phe Ala Glu
180         185         190

Glu His Tyr Ile Leu Gly Lys Asp Asn Asp Glu Met Ile Lys Phe Ile
195         200         205

Gly Asn His Pro Asn Thr Tyr Ala Tyr Thr Lys Ala Leu Ala Glu Asn
210         215         220

Leu Val Ala Glu Glu His Gly Glu Ile Pro Thr Ile Ile Ile Arg Pro
225         230         235         240

Ser Ile Ile Thr Ala Ser Ala Glu Glu Pro Val Arg Gly Phe Val Asp
245         250         255

Ser Trp Ser Gly Ala Thr Ala Met Ala Ala Phe Ala Leu Lys Gly Trp
260         265         270

Asn Asn Ile Met Tyr Ser Thr Gly Glu Glu Asn Ile Asp Leu Ile Pro
275         280         285

Leu Asp Tyr Val Val Asn Leu Thr Leu Val Ala Ile Ala Lys Tyr Lys
290         295         300

Pro Thr Lys Glu Val Thr Val Tyr His Val Thr Thr Ser Asp Leu Asn
305         310         315         320

Pro Ile Ser Ile Arg Arg Ile Phe Ile Lys Leu Ser Glu Phe Ala Ser
325         330         335

Lys Asn Pro Thr Ser Asn Ala Ala Pro Phe Ala Ala Thr Thr Leu Leu

```

---

-continued

---

340	345	350	
Thr Lys Gln Lys Pro Leu Ile Lys Leu Val Thr Phe Leu Met Gln Thr			
355	360	365	
Thr Pro Ala Phe Leu Ala Asp Leu Trp Met Lys Thr Gln Arg Lys Glu			
370	375	380	
Ala Lys Phe Val Lys Gln His Asn Leu Val Val Arg Ser Arg Asp Gln			
385	390	395	400
Leu Glu Phe Phe Thr Ser Gln Ser Trp Leu Leu Arg Cys Glu Arg Ala			
405	410	415	
Arg Val Leu Ser Ala Ala Leu Ser Asp Ser Asp Arg Ala Val Phe Arg			
420	425	430	
Cys Asp Pro Ser Thr Ile Asp Trp Asp Gln Tyr Leu Pro Ile Tyr Phe			
435	440	445	
Glu Gly Ile Asn Lys His Leu Phe Lys Asn Lys Leu			
450	455	460	

<210> SEQ ID NO 13  
<211> LENGTH: 1542  
<212> TYPE: DNA  
<213> ORGANISM: Artificial  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic  
<220> FEATURE:  
<221> NAME/KEY: CDS  
<222> LOCATION: (1)...(1542)

<400> SEQUENCE: 13

atg gct atc cag cag gtt cat cac gcc gac aca tcc tcc tct aaa gtc	48		
Met Ala Ile Gln Gln Val His His Ala Asp Thr Ser Ser Ser Lys Val			
1	5	10	15
ctg ggt caa ctt cgt ggt aaa cgt gtc ttg att acc ggc act act gga	96		
Leu Gly Gln Leu Arg Gly Lys Arg Val Leu Ile Thr Gly Thr Thr Gly			
20	25	30	
ttc ttg ggt aaa gtc gtc ttg gaa cgt ttg att cgt gcc gtt cct gac	144		
Phe Leu Gly Lys Val Val Leu Arg Leu Ile Arg Ala Val Pro Asp			
35	40	45	
atc ggt gct atc tac ctg ctg att cgt ggt aac aag cgt cac ccg gat	192		
Ile Gly Ala Ile Tyr Leu Leu Ile Arg Gly Asn Lys Arg His Pro Asp			
50	55	60	
gct cgt tct cgt ttg gag gag att gct acc tcc tct gtc ttt gat	240		
Ala Arg Ser Arg Phe Leu Glu Glu Ile Ala Thr Ser Ser Val Phe Asp			
65	70	75	80
cgt ttg cgt gaa gct gat tcc gaa ggt ttc gat gct ttc ctg gaa gaa	288		
Arg Leu Arg Glu Ala Asp Ser Glu Gly Phe Asp Ala Phe Leu Glu Glu			
85	90	95	
cgt att cac tgt gtt act ggt gaa gtt act gaa gct ggt ttc ggt att	336		
Arg Ile His Cys Val Thr Gly Glu Val Thr Glu Ala Gly Phe Gly Ile			
100	105	110	
ggt caa gag gac tat cgt aag ttg gcc acc gaa ttg gac gca gtc atc	384		
Gly Gln Glu Asp Tyr Arg Lys Leu Ala Thr Glu Leu Asp Ala Val Ile			
115	120	125	
aat tct gct gcc gtc aac ttc cgt gag gag ttg gat aag gct ctg	432		
Asn Ser Ala Ala Ser Val Asn Phe Arg Glu Glu Leu Asp Lys Ala Leu			
130	135	140	
gcc atc aac act ctg tgt ttg cgt aac atc gct ggt atg gtg gat ctt	480		
Ala Ile Asn Thr Leu Cys Leu Arg Asn Ile Ala Gly Met Val Asp Leu			
145	150	155	160

-continued

---

aac cct aag ctg gcc gtt ctt caa gtc tct acg tgt tac gtc aac ggt Asn Pro Lys Leu Ala Val Leu Gln Val Ser Thr Cys Tyr Val Asn Gly 165 170 175	528
atg aac tct ggt caa gtt act gaa tcc gtc atc aaa cca gct ggt gaa Met Asn Ser Gly Gln Val Thr Glu Ser Val Ile Lys Pro Ala Gly Glu 180 185 190	576
gct gtt cct cgt tct cct gat gga ttc tac gag atc gag gaa ttg gtt Ala Val Pro Arg Ser Pro Asp Gly Phe Tyr Glu Ile Glu Glu Leu Val 195 200 205	624
cgt ctg ctg caa gac aag att gaa gac gtt caa gca cgt tac tct ggt Arg Leu Leu Gln Asp Lys Ile Glu Asp Val Gln Ala Arg Tyr Ser Gly 210 215 220	672
aag gtg ttg gag cgt aag ttg gtt gat ttg ggt att cgt gag gct aat Lys Val Leu Glu Arg Lys Leu Val Asp Leu Gly Ile Arg Glu Ala Asn 225 230 235 240	720
cgt tac ggt tgg tct gat aca tac acc ttc acg aaa tgg ttg ggt gaa Arg Tyr Gly Trp Ser Asp Thr Tyr Thr Phe Thr Lys Trp Leu Gly Glu 245 250 255	768
caa ctt ctg atg aaa gcc ttg aat ggt cgt acc ttg act att ctg cgt Gln Leu Leu Met Lys Ala Leu Asn Gly Arg Thr Leu Thr Ile Leu Arg 260 265 270	816
cct agc atc att gaa tct gct ttg gaa gaa cca gca cct ggt tgg att Pro Ser Ile Ile Glu Ser Ala Leu Glu Glu Pro Ala Pro Gly Trp Ile 275 280 285	864
gaa ggc gtg aaa gtt gca gat gcg atc atc ttg gct tat gct cgt gag Glu Gly Val Lys Val Ala Asp Ala Ile Ile Leu Ala Tyr Ala Arg Glu 290 295 300	912
aag gtt act ttg ttt ccg ggt aaa cgt tct ggt atc att gat gtg att Lys Val Thr Leu Phe Pro Gly Lys Arg Ser Gly Ile Ile Asp Val Ile 305 310 315 320	960
cct gtt gac ttg gtt gcc aat tcc atc atc ttg tct ttg gct gag gct Pro Val Asp Leu Val Ala Asn Ser Ile Ile Leu Ser Leu Ala Glu Ala 325 330 335	1008
ctg ggc gaa cct ggt cgt cgt atc tac caa tgt tgt tct ggt ggt Leu Gly Glu Pro Gly Arg Arg Arg Ile Tyr Gln Cys Cys Ser Gly Gly 340 345 350	1056
ggt aat cct atc tcc ctg ggc gag ttc att gat cac ctg atg gct gaa Gly Asn Pro Ile Ser Leu Gly Glu Phe Ile Asp His Leu Met Ala Glu 355 360 365	1104
tcc aaa gcc aac tat gcc gca tac gat cat ctg ttc tac cgt caa occ Ser Lys Ala Asn Tyr Ala Ala Tyr Asp His Leu Phe Tyr Arg Gln Pro 370 375 380	1152
tcc aag cct ttc ctt gct gtc aac cgt gct ttg ttc gac ttg gtt atc Ser Lys Pro Phe Leu Ala Val Asn Arg Ala Leu Phe Asp Leu Val Ile 385 390 395 400	1200
tct ggt gtc cgt ctg cct ttg tct ttg acc gac cgt gtc ttg aag ctg Ser Gly Val Arg Leu Pro Leu Ser Leu Thr Asp Arg Val Leu Lys Leu 405 410 415	1248
ctg ggc aac tcc cgt gac ctg aag atg ctg cgt aac ctg gat act acg Leu Gly Asn Ser Arg Asp Leu Lys Met Leu Arg Asn Leu Asp Thr Thr 420 425 430	1296
caa tcc ctg gct act atc ttt ggc ttc tac aca gcc ccc gac tac atc Gln Ser Leu Ala Thr Ile Phe Gly Phe Tyr Thr Ala Pro Asp Tyr Ile 435 440 445	1344
ttc cgt aat gac gag ttg atg gcc ctg gct aac cgt atg ggc gag gtt Phe Arg Asn Asp Glu Leu Met Ala Leu Asn Arg Met Gly Glu Val 450 455 460	1392

---

-continued

---

gat aag ggt ttg ttc ccc gtt gat gct cgt ctg att gat tgg gaa ttg	1440
Asp Lys Gly Leu Phe Pro Val Asp Ala Arg Leu Ile Asp Trp Glu Leu	
465                   470                   475                   480	

tac ctg cgt aag att cac ctg gct ggt ttg aac cgt tac gcc ttg aag	1488
Tyr Leu Arg Lys Ile His Leu Ala Gly Leu Asn Arg Tyr Ala Leu Lys	
485                   490                   495	

gag cgt aag gtt tac tct ttg aag aca gcc cgt cag cgt aag aag gca	1536
Glu Arg Lys Val Tyr Ser Leu Lys Thr Ala Arg Gln Arg Lys Lys Ala	
500                   505                   510	

gct taa	1542
Ala	

<210> SEQ ID NO 14

<211> LENGTH: 513

<212> TYPE: PRT

<213> ORGANISM: Artificial

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 14

Met Ala Ile Gln Gln Val His His Ala Asp Thr Ser Ser Ser Lys Val	
1                   5                   10                   15	

Leu Gly Gln Leu Arg Gly Lys Arg Val Leu Ile Thr Gly Thr Thr Gly	
20                   25                   30	

Phe Leu Gly Lys Val Val Leu Glu Arg Leu Ile Arg Ala Val Pro Asp	
35                   40                   45	

Ile Gly Ala Ile Tyr Leu Leu Ile Arg Gly Asn Lys Arg His Pro Asp	
50                   55                   60	

Ala Arg Ser Arg Phe Leu Glu Glu Ile Ala Thr Ser Ser Val Phe Asp	
65                   70                   75                   80	

Arg Leu Arg Glu Ala Asp Ser Glu Gly Phe Asp Ala Phe Leu Glu Glu	
85                   90                   95	

Arg Ile His Cys Val Thr Gly Glu Val Thr Glu Ala Gly Phe Gly Ile	
100                   105                   110	

Gly Gln Glu Asp Tyr Arg Lys Leu Ala Thr Glu Leu Asp Ala Val Ile	
115                   120                   125	

Asn Ser Ala Ala Ser Val Asn Phe Arg Glu Glu Leu Asp Lys Ala Leu	
130                   135                   140	

Ala Ile Asn Thr Leu Cys Leu Arg Asn Ile Ala Gly Met Val Asp Leu	
145                   150                   155                   160	

Asn Pro Lys Leu Ala Val Leu Gln Val Ser Thr Cys Tyr Val Asn Gly	
165                   170                   175	

Met Asn Ser Gly Gln Val Thr Glu Ser Val Ile Lys Pro Ala Gly Glu	
180                   185                   190	

Ala Val Pro Arg Ser Pro Asp Gly Phe Tyr Glu Ile Glu Glu Leu Val	
195                   200                   205	

Arg Leu Leu Gln Asp Lys Ile Glu Asp Val Gln Ala Arg Tyr Ser Gly	
210                   215                   220	

Lys Val Leu Glu Arg Lys Leu Val Asp Leu Gly Ile Arg Glu Ala Asn	
225                   230                   235                   240	

Arg Tyr Gly Trp Ser Asp Thr Tyr Thr Phe Thr Lys Trp Leu Gly Glu	
245                   250                   255	

Gln Leu Leu Met Lys Ala Leu Asn Gly Arg Thr Leu Thr Ile Leu Arg	
260                   265                   270	

---

-continued

```

Pro Ser Ile Ile Glu Ser Ala Leu Glu Glu Pro Ala Pro Gly Trp Ile
275          280          285

Glu Gly Val Lys Val Ala Asp Ala Ile Ile Leu Ala Tyr Ala Arg Glu
290          295          300

Lys Val Thr Leu Phe Pro Gly Lys Arg Ser Gly Ile Ile Asp Val Ile
305          310          315          320

Pro Val Asp Leu Val Ala Asn Ser Ile Ile Leu Ser Leu Ala Glu Ala
325          330          335

Leu Gly Glu Pro Gly Arg Arg Arg Ile Tyr Gln Cys Cys Ser Gly Gly
340          345          350

Gly Asn Pro Ile Ser Leu Gly Glu Phe Ile Asp His Leu Met Ala Glu
355          360          365

Ser Lys Ala Asn Tyr Ala Ala Tyr Asp His Leu Phe Tyr Arg Gln Pro
370          375          380

Ser Lys Pro Phe Leu Ala Val Asn Arg Ala Leu Phe Asp Leu Val Ile
385          390          395          400

Ser Gly Val Arg Leu Pro Leu Ser Leu Thr Asp Arg Val Leu Lys Leu
405          410          415

Leu Gly Asn Ser Arg Asp Leu Lys Met Leu Arg Asn Leu Asp Thr Thr
420          425          430

Gln Ser Leu Ala Thr Ile Phe Gly Phe Tyr Thr Ala Pro Asp Tyr Ile
435          440          445

Phe Arg Asn Asp Glu Leu Met Ala Leu Ala Asn Arg Met Gly Glu Val
450          455          460

Asp Lys Gly Leu Phe Pro Val Asp Ala Arg Leu Ile Asp Trp Glu Leu
465          470          475          480

Tyr Leu Arg Lys Ile His Leu Ala Gly Leu Asn Arg Tyr Ala Leu Lys
485          490          495

Glu Arg Lys Val Tyr Ser Leu Lys Thr Ala Arg Gln Arg Lys Lys Ala
500          505          510

Ala

```

```

<210> SEQ ID NO 15
<211> LENGTH: 42
<212> TYPE: DNA
<213> ORGANISM: Artificial
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

```

```
<400> SEQUENCE: 15
```

```
acaatctgga tccggccagc ctggccataa ggagatatac at           42
```

```

<210> SEQ ID NO 16
<211> LENGTH: 34
<212> TYPE: DNA
<213> ORGANISM: Artificial
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

```

```
<400> SEQUENCE: 16
```

```
taatgaggcc aaaactggccg tcgacaccag tatg           34
```

```

<210> SEQ ID NO 17
<211> LENGTH: 17
<212> TYPE: DNA
<213> ORGANISM: Artificial

```

---

-continued

---

<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic primer

<400> SEQUENCE: 17

agagaccggg ttggcgg

17

<210> SEQ ID NO 18  
<211> LENGTH: 21  
<212> TYPE: DNA  
<213> ORGANISM: Artificial  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic primer

<400> SEQUENCE: 18

catttgcatt tcgttaacgct g

21

---

What is claimed is:

1. A method of producing a fatty alcohol composition comprising:
  - a) culturing a recombinant microorganism in a suitable culture medium, wherein the recombinant microorganism comprises a gene encoding a heterologous fatty acyl reductase (FAR) enzyme having at least 30% sequence identity to SEQ ID NO: 2, SEQ ID NO: 6, SEQ ID NO: 14 or a functional fragment thereof, and
  - b) allowing expression of said gene, wherein said expression results in the production of a composition of fatty alcohols.
2. The method according to claim 1, wherein the recombinant microorganism is selected from the group of a bacterium, a filamentous fungus, a yeast, and an algae.
3. The method according to claim 2, wherein the recombinant microorganism is a bacterium.
4. The method according to claim 3, wherein the recombinant bacterium is an *E. coli* strain.
5. The method according to claim 2, wherein the recombinant microorganism is a yeast.
6. The method according to claim 5, wherein the yeast is a strain of *Saccharomyces*, *Candida*, *Pichia* or *Yarrowia*.
7. The method according to claim 1, wherein the FAR enzyme has at least 75% sequence identity to SEQ ID NO: 2, SEQ ID NO: 6, SEQ ID NO: 14 or functional fragments thereof.
8. The method according to claim 1, wherein the FAR enzyme has at least 85% sequence identity to SEQ ID NO: 2, SEQ ID NO: 6, SEQ ID NO: 14 or functional fragments thereof.
9. The method according to claim 1, further comprising isolating the composition of fatty alcohols.
10. A composition comprising the isolated fatty alcohols produced according to the method of claim 1.
11. The method according to claim 1, wherein the produced fatty alcohols include C12-C18 fatty alcohols.
12. The method according to claim 1, wherein the produced fatty alcohols include a mixture of at least two of C14:0, C16:0, C18:0, C16:1 and C18:1 fatty alcohols.
13. The method according to claim 1, wherein at least 25% of the produced fatty alcohols are secreted.
14. The method according to claim 13, wherein at least 50% of the produced fatty alcohols are secreted.

**15.** The method according to claim 1, wherein the recombinant microorganism produces at least 0.5 g/L of fatty alcohol.

**16.** The method according to claim 15, wherein the recombinant microorganism produces at least 5 g/L of fatty alcohols.

**17.** The method according to claim 16, wherein the recombinant microorganism produces at least 15 g/L of fatty alcohol.

**18.** The method according to claim 1, wherein the gene encoding the FAR comprises a codon optimized nucleic acid sequence.

**19.** The method according to claim 18, wherein the nucleic acid sequence is a sequence having at least 70% sequence identity to SEQ ID NOS: 1, 3, 5, 7, or 13.

**20.** The method according to claim 1, wherein the nucleic acid sequence is integrated into the chromosome of the recombinant microorganism.

**21.** The method according to claim 1, wherein the nucleic acid is present on a vector.

**22.** The method according to claim 1, wherein the culturing conditions comprise a temperature in the range of 20° C. to 45° C.; a pH in the range of 5 to 7, a time period in the range of 16 hours to 144 hours; and an assimilable carbon source comprising fermentable sugars obtained from a cellulosic feedstock.

**23.** The method according to claim 1, wherein the fatty alcohol is produced by an acyl-CoA independent pathway.

**24.** The method according to claim 1 further comprising reducing the fatty alcohol composition to yield an alkane composition.

**25.** A fuel composition comprising the alkanes produced according to the method of claim 24.

**26.** The method according to claim 1 further comprising esterification of the fatty alcohol composition to produce fatty esters.

**27.** A fuel composition comprising the fatty esters produced according to the method of claim 26.

**28.** An isolated nucleic acid sequence encoding a fatty acyl-CoA reductase (FAR) enzyme comprising an amino acid sequence having at least 30% sequence identity to SEQ ID NO: 2, SEQ ID NO: 6, SEQ ID NO: 14 or functional fragments thereof.

**29.** The isolated nucleic acid of claim **28**, wherein the FAR enzyme has at least 80% sequence identity to SEQ ID NO: 2, SEQ ID NO: 6, SEQ ID NO: 14 or a functional fragment thereof.

**30.** The isolated nucleic acid of claim **29**, wherein the FAR enzyme has at least 90% sequence identity to SEQ ID NO: 2, SEQ ID NO: 6, SEQ ID NO: 14 or a functional fragment thereof.

**31.** The isolated nucleic acid of claim **29**, wherein the polynucleotide is codon optimized for expression in *E. coli*.

**32.** The isolated nucleic acid of claim **29**, wherein the polynucleotide is codon optimized for expression in yeast.

**33.** The isolated nucleic acid of claim **32**, wherein the polynucleotide is codon optimized for expression in *Yarrowia* or *Saccharomyces*.

**34.** A vector comprising the nucleic acid of claim **28**.

**35.** A host cell transformed with the vector of claim **34**.

**36.** The host cell of claim **35**, wherein said host cell is *E. coli*, *Yarrowia*, or *S. cerevisiae*.

**37.** An isolated fatty acyl reductase (FAR) enzyme comprising an amino acid sequence that is at least 30% identical to SEQ ID NO: 2, SEQ ID NO: 6, SEQ ID NO: 14 or a functional fragment thereof.

**38.** The isolated FAR enzyme of claim **37**, wherein the FAR enzyme comprises an amino acid sequence that is at least 75% identical to SEQ ID NO: 2, SEQ ID NO: 6, SEQ ID NO: 14 or a functional fragment thereof.

**39.** The isolated FAR enzyme of claim **38**, wherein the FAR enzyme comprises an amino acid sequence that is at least 90% identical to SEQ ID NO: 2, SEQ ID NO: 6, SEQ ID NO: 14 or a functional fragment thereof.

**40.** A method of producing a fatty alcohol composition comprising culturing a recombinant *E. coli* in a nutrient medium under suitable conditions, wherein said recombinant *E. coli* comprises a nucleic acid sequence encoding a heterologous fatty acyl-CoA reductase (FAR) enzyme comprising an amino acid sequence having at least 75% sequence identity to SEQ ID NO: 2, SEQ ID NO: 6, SEQ ID NO: 14 or functional fragments thereof; allowing expression of said FAR enzyme; and producing a fatty alcohol composition.

**41.** The method according to claim **40**, wherein the recombinant *E. coli* further comprises an inactivated or silenced endogenous fatty acyl-ACP thioesterase gene, an inactivated or silenced endogenous fatty acyl-CoA synthetase gene or both.

\* \* \* \* \*