

Table 1: Variable means by Task (df_{pp}) with ANOVA results

Variable	Task 0	Task 1	Task 2	Task 3	Task 4	Welch ANOVA p	Kruskal-Wallis p
Cumulative Score Diff	5.49	5.43	4.00	2.87	3.22	0.000	0.000
End Score Diff	1.19	1.30	1.60	1.02	1.17	0.387	0.587
Nonhammer Win	0.23	0.25	0.14	0.23	0.20	0.593	0.674
Hammer 2+	0.44	0.44	0.50	0.42	0.47	0.910	0.906
Handle	0.50	0.47	0.38	0.54	0.52	0.473	0.474
Points	3.11	3.64	3.88	2.96	2.63	0.000	0.000
Score Hammer	1.50	1.59	1.76	1.36	1.47	0.430	0.622
Score Nonhammer	0.31	0.28	0.17	0.33	0.30	0.437	0.800

Table 2: Variable means by HDBSCAN Cluster (df_{pp2}) with ANOVA results

Variable	Cluster 0	Cluster 1	Cluster 3	Welch ANOVA p	Kruskal-Wallis p
Cumulative Score Diff	3.51	6.00	4.84	0.015	0.029
End Score Diff	1.00	1.07	1.24	0.774	0.910
Nonhammer Win	0.26	0.23	0.21	0.893	0.886
Hammer 2+	0.40	0.40	0.40	0.999	0.999
Handle	0.06	0.96	0.60	0.000	0.000
Points	3.46	3.28	3.67	0.012	0.003
Score Hammer	1.40	1.43	1.52	0.875	0.958
Score Nonhammer	0.40	0.36	0.28	0.682	0.904