
a
1. Cross-lineage shared
2. Likely shared
PBMC
Myeloid
Lymphoid
3. Lineage-specific
T
4. T-cell-specific
Mono
DC
NK
B
CD4+
CD8+
5. Single cell-type
b eGenes
Cross-lineage shared (n = 10,594)
Likely shared (n = 4,681)
Lineage-specific (n = 1,827)
T-cell specific (n = 868)
Single cell-type (n = 2,859)
51%
30%
12%
22%
6%
9%
6%
4%
14%
50%
13%
4%
10%
Distinct variants
consistent, heterogeneous, opposite effects
c caPeaks
Cross-lineage shared (n = 68,438)
Likely shared (n = 93,181)
Lineage-specific (n = 15,449)
T-cell specific (n = 4,729)
Single cell-type (n = 28,787)
32%
18%
9%
44%
14%
8%
7%
2%
14%
5%
43%
13%
2%
14%
Distinct
consistent, heterogeneous, opposite effects
d
IKZF1
CD4+ TEM
CD8+ TEM
Normalized expression
rs6976036 (ALT)
e
FOXO1
CD14+ Mono
CD4+ Naive
rs17061503 (ALT)
f
NFATC1
B
Mono
rs68734 (ALT)
g
NFATC2
B
CD4+ T
−log10(P)
51.4
51.5
51.6
NFATC2
ATP9A
Chromosome 20 (Mb)
h
ZBTB1
CD14+ Mono
CD4+ TCM
Normalized expression
rs9323451 (ALT)
i
ZNF804A
B
Mono
rs10497655 (ALT)
j
ZNF7
CD14+ Mono
CD4+ Naive
rs1209879 (ALT)
k
ZNF430
ZNF66
ZNF100
ZNF43
ZNF85
ZNF431
ZNF626
ZNF429
ZNF708
ZNF493
ZNF257
ZNF208
B
NK
CD4+ T
CD8+ T
DC
Mono
rs55778393 (ALT)
−0.4
−0.3
−0.2
−0.1
0
l
Mono
i
ii
Norm. CA
56.327
56.328
56.329
Chromosome 5 (Mb)
m
B
n
CD4+ T
o
i. chr5:56327396−56328237
ii. chr5:56328850−56329349
CD4+ T
CD8+ T
B
NK
DC
Mono
−1.0
−0.5
0.0
0.5
1.0
Effect size (rs158488)