

Oceanographic features delineate growth zonation in Northeast Pacific sablefish

Kapur, M.¹, Haltuch, M.², Connors, B.³, Rogers, L.⁴, Berger, A.², Koontz, E.⁵, Cope, J.², Echave, K.⁶, Fenske, K.⁶, Hanselman, D.⁶, Punt, A.E.¹

¹ University of Washington, School of Aquatic and Fisheries Sciences. 1122 NE Boat St, Seattle WA 98105

² Fisheries Resource and Monitoring Division, Northwest Fisheries Science Center, National Marine Fisheries Service National Oceanic and Atmospheric Administration. 2725 Montlake Blvd. E. Seattle, WA 98112

³ Institute of Ocean Sciences, Fisheries and Oceans Canada. 9860 W. Saanich Rd. Sidney, B.C. Canada V8L 5T5

⁴ Fisheries and Oceans Canada, School of Resource and Environmental Management, Simon Fraser University. 8888 University Drive, Burnaby, British Columbia, Canada, V5A 1S6

⁵ Center for Quantitative Science, Ocean Teaching Building, Suite 300, Box 357941, Seattle, WA 98195.

⁶ Auke Bay Laboratories, Alaska Fisheries Science Center, National Marine Fisheries Service National Oceanic and Atmospheric Administration. 17101 Pt. Lena Loop Rd. Juneau, AK 99801

Corresponding author: kapurm@uw.edu

Keywords: growth, von Bertalanffy, ecosystem-based fisheries management, sablefish, spatiotemporal

Abstract

Renewed interest in the estimation of spatial and temporal variation in fish traits, such as body size, is a result of computing advances and the development of spatially-explicit management frameworks. However, many attempts to quantify spatial structure or the distribution of traits utilize *a priori* approaches, which involve pre-designated geographic regions and thus cannot detect unanticipated spatial patterns. We developed a new, **model-based** method that uses the first derivative of the spatial smoothing term of a generalized additive model to identify spatial zones of variation in fish length-at-age. We use simulation testing to evaluate the method across a variety of synthetic, stratified age and length datasets, and then apply it to survey data for Northeast Pacific sablefish (*Anoplopoma fimbria*). Simulation testing illustrates the robustness of the method across a variety of scenarios related to spatially or temporally stratified length-at-age data, including strict boundaries, overlapping zones and changes at the extreme of the range. Results indicate that length-at-age for Northeast Pacific sablefish increases with latitude, which is consistent with previous work from the western United States. Model-detected spatial breakpoints corresponded to major oceanographic features, including the northern end of the Southern California Bight and the bifurcation of the North Pacific Current. This method has the potential to improve detection of large-scale patterns in fish growth, and aid in the development of spatiotemporally structured population dynamics models to inform ecosystem-based fisheries management.