Simplicty and uniqueness of trace of group C^* -algebras Bachelor thesis defense

Malthe Munk Karbo

Advisor: Mikael Rørdam

February 3, 2017

Table of Contents

1 Introduction, history and motivation

Some notation and preliminaries

Malthe Munk Karbo February 3, 2017 2 /

Table of Contents

1 Introduction, history and motivation

2 Some notation and preliminaries

Malthe Munk Karbo February 3, 2017 4 /

The main subject of this thesis is a theory describing the ideal structure of something called *crossed products* associated to a certain type of dynamical systems.

The main subject of this thesis is a theory describing the ideal structure of something called *crossed products* associated to a certain type of dynamical systems.

A natural question is: What is the all the fuzz about?

The main subject of this thesis is a theory describing the ideal structure of something called *crossed products* associated to a certain type of dynamical systems.

A natural question is: What is the all the fuzz about? The crossed product construction associates to a C^* -dynamical system (A, G, α) a C^* -algebra $A \rtimes_{\alpha} G$.

The main subject of this thesis is a theory describing the ideal structure of something called *crossed products* associated to a certain type of dynamical systems.

A natural question is: What is the all the fuzz about? The crossed product construction associates to a C^* -dynamical system (A, G, α) a C^* -algebra $A \rtimes_{\alpha} G$. This construction is important:

The main subject of this thesis is a theory describing the ideal structure of something called *crossed products* associated to a certain type of dynamical systems.

A natural question is: What is the all the fuzz about? The crossed product construction associates to a C^* -dynamical system (A, G, α) a C^* -algebra $A \rtimes_{\alpha} G$. This construction is important:

• It provides many important and interesting examples of new C^* -algebras.

The main subject of this thesis is a theory describing the ideal structure of something called *crossed products* associated to a certain type of dynamical systems.

A natural question is: What is the all the fuzz about? The crossed product construction associates to a C^* -dynamical system (A, G, α) a C^* -algebra $A \rtimes_{\alpha} G$. This construction is important:

- It provides many important and interesting examples of new C^* -algebras.
- It encodes quite a bit of information about both G and A and the action $G \stackrel{\alpha}{\sim} A$.

The main subject of this thesis is a theory describing the ideal structure of something called *crossed products* associated to a certain type of dynamical systems.

A natural question is: What is the all the fuzz about? The crossed product construction associates to a C^* -dynamical system (A, G, α) a C^* -algebra $A \rtimes_{\alpha} G$. This construction is important:

- It provides many important and interesting examples of new C^* -algebras.
- It encodes quite a bit of information about both G and A and the action $G \stackrel{\alpha}{\sim} A$.
- The construction served as a way of answering many questions in the history of operator algebras.

The main subject of this thesis is a theory describing the ideal structure of something called *crossed products* associated to a certain type of dynamical systems.

A natural question is: What is the all the fuzz about? The crossed product construction associates to a C^* -dynamical system (A, G, α) a C^* -algebra $A \rtimes_{\alpha} G$. This construction is important:

- It provides many important and interesting examples of new C^* -algebras.
- It encodes quite a bit of information about both G and A and the action $G \stackrel{\alpha}{\sim} A$.
- The construction served as a way of answering many questions in the history of operator algebras.
- Ties together with classic theory of dynamics.

The main subject of this thesis is a theory describing the ideal structure of something called *crossed products* associated to a certain type of dynamical systems.

A natural question is: What is the all the fuzz about? The crossed product construction associates to a C^* -dynamical system (A, G, α) a C^* -algebra $A \rtimes_{\alpha} G$. This construction is important:

- It provides many important and interesting examples of new C^* -algebras.
- It encodes quite a bit of information about both G and A and the action $G \stackrel{\alpha}{\sim} A$.
- The construction served as a way of answering many questions in the history of operator algebras.
- Ties together with classic theory of dynamics.
- Ties together with many areas in C^* -theory, including the group C^* -algebra construction.

The main subject of this thesis is a theory describing the ideal structure of something called *crossed products* associated to a certain type of dynamical systems.

A natural question is: What is the all the fuzz about? The crossed product construction associates to a C^* -dynamical system (A, G, α) a C^* -algebra $A \rtimes_{\alpha} G$. This construction is important:

- It provides many important and interesting examples of new C^* -algebras.
- It encodes quite a bit of information about both G and A and the action $G \stackrel{\alpha}{\sim} A$.
- The construction served as a way of answering many questions in the history of operator algebras.
- Ties together with classic theory of dynamics.
- Ties together with many areas in C^* -theory, including the group C^* -algebra construction.

Okay, but why do we care about the ideal structure of this construction?

The main subject of this thesis is a theory describing the ideal structure of something called *crossed products* associated to a certain type of dynamical systems.

A natural question is: What is the all the fuzz about? The crossed product construction associates to a C^* -dynamical system (A, G, α) a C^* -algebra $A \rtimes_{\alpha} G$. This construction is important:

- It provides many important and interesting examples of new C^* -algebras.
- It encodes quite a bit of information about both G and A and the action $G \stackrel{\alpha}{\sim} A$.
- The construction served as a way of answering many questions in the history of operator algebras.
- Ties together with classic theory of dynamics.
- Ties together with many areas in C^* -theory, including the group C^* -algebra construction.

Okay, but why do we care about the ideal structure of this construction? Many answers, most notably we want to do classification.

The main subject of this thesis is a theory describing the ideal structure of something called *crossed products* associated to a certain type of dynamical systems.

A natural question is: What is the all the fuzz about? The crossed product construction associates to a C^* -dynamical system (A, G, α) a C^* -algebra $A \rtimes_{\alpha} G$. This construction is important:

- It provides many important and interesting examples of new C^* -algebras.
- It encodes quite a bit of information about both G and A and the action $G \stackrel{\alpha}{\sim} A$.
- The construction served as a way of answering many questions in the history of operator algebras.
- Ties together with classic theory of dynamics.
- Ties together with many areas in C^* -theory, including the group C^* -algebra construction.

Okay, but why do we care about the ideal structure of this construction? Many answers, most notably we want to do <u>classification</u>.

And the most important question:

The main subject of this thesis is a theory describing the ideal structure of something called *crossed products* associated to a certain type of dynamical systems.

A natural question is: What is the all the fuzz about? The crossed product construction associates to a C^* -dynamical system (A, G, α) a C^* -algebra $A \rtimes_{\alpha} G$. This construction is important:

- It provides many important and interesting examples of new C^* -algebras.
- It encodes quite a bit of information about both G and A and the action $G \stackrel{\alpha}{\curvearrowright} A$.
- The construction served as a way of answering many questions in the history of operator algebras.
- Ties together with classic theory of dynamics.
- Ties together with many areas in C^* -theory, including the group C^* -algebra construction.

Okay, but why do we care about the ideal structure of this construction? Many answers, most notably we want to do <u>classification</u>.

And the most important question: Real world application?

The main subject of this thesis is a theory describing the ideal structure of something called *crossed products* associated to a certain type of dynamical systems.

A natural question is: What is the all the fuzz about? The crossed product construction associates to a C^* -dynamical system (A, G, α) a C^* -algebra $A \rtimes_{\alpha} G$. This construction is important:

- It provides many important and interesting examples of new C^* -algebras.
- It encodes quite a bit of information about both G and A and the action $G \stackrel{\alpha}{\curvearrowright} A$.
- The construction served as a way of answering many questions in the history of operator algebras.
- Ties together with classic theory of dynamics.
- Ties together with many areas in C^* -theory, including the group C^* -algebra construction.

Okay, but why do we care about the ideal structure of this construction? Many answers, most notably we want to do <u>classification</u>.

And the most important question: Real world application? (money?)

The main subject of this thesis is a theory describing the ideal structure of something called *crossed products* associated to a certain type of dynamical systems.

A natural question is: What is the all the fuzz about? The crossed product construction associates to a C^* -dynamical system (A, G, α) a C^* -algebra $A \rtimes_{\alpha} G$. This construction is important:

- It provides many important and interesting examples of new C^* -algebras.
- It encodes quite a bit of information about both G and A and the action $G \stackrel{\alpha}{\curvearrowright} A$.
- The construction served as a way of answering many questions in the history of operator algebras.
- Ties together with classic theory of dynamics.
- Ties together with many areas in C^* -theory, including the group C^* -algebra construction.

Okay, but why do we care about the ideal structure of this construction? Many answers, most notably we want to do classification.

And the most important question: Real world application? (money?) Probably not

Malthe Munk Karbo

February 3, 2017

To a C^* -dynamical system (A, G, α) , our goal is to answer the following question:

To a C^* -dynamical system (A, G, α) , our goal is to answer the following question:

Question 1: What properties P describes the ideal structure of $A \rtimes_{\alpha} G$?

To a C^* -dynamical system (A, G, α) , our goal is to answer the following question:

Question 1: What properties P describes the ideal structure of $A \rtimes_{\alpha} G$?

During this talk, the hope is that we will provide some answers to this question.

To a C^* -dynamical system (A, G, α) , our goal is to answer the following question:

Question 1: What properties P describes the ideal structure of $A \rtimes_{\alpha} G$?

During this talk, the hope is that we will provide some answers to this question.

Answering question 1 in the above form is <u>hard</u>.

Instead, we break it into special cases

To a C^* -dynamical system (A, G, α) , our goal is to answer the following question:

Question 1: What properties P describes the ideal structure of $A \rtimes_{\alpha} G$?

During this talk, the hope is that we will provide some answers to this question.

Answering question 1 in the above form is $\underline{\text{hard}}$.

Instead, we break it into special cases Some cases discussed today will be:

To a C^* -dynamical system (A, G, α) , our goal is to answer the following question:

Question 1: What properties P describes the ideal structure of $A \rtimes_{\alpha} G$?

During this talk, the hope is that we will provide some answers to this question. Answering question 1 in the above form is hard.

Instead, we break it into special cases Some cases discussed today will be:

• Which properties P answers Question 1 when A is abelian?

To a C^* -dynamical system (A, G, α) , our goal is to answer the following question:

Question 1: What properties P describes the ideal structure of $A \rtimes_{\alpha} G$?

During this talk, the hope is that we will provide some answers to this question.

Answering question 1 in the above form is <u>hard</u>.

Instead, we break it into special cases Some cases discussed today will be:

- Which properties P answers Question 1 when A is abelian?
- Which properties P answers Question 1 when G is abelian?

To a C^* -dynamical system (A, G, α) , our goal is to answer the following question:

Question 1: What properties P describes the ideal structure of $A \rtimes_{\alpha} G$?

During this talk, the hope is that we will provide some answers to this question. Answering question 1 in the above form is hard.

Instead, we break it into special cases Some cases discussed today will be:

- Which properties P answers Question 1 when A is abelian?
- Which properties P answers Question 1 when G is abelian?
- Which properties P answers Question 1 when $A = \mathbb{C}$?

To a C^* -dynamical system (A, G, α) , our goal is to answer the following question:

Question 1: What properties P describes the ideal structure of $A \rtimes_{\alpha} G$?

During this talk, the hope is that we will provide some answers to this question.

Answering question 1 in the above form is $\underline{\text{hard}}$.

Instead, we break it into special cases Some cases discussed today will be:

- Which properties P answers Question 1 when A is abelian?
- Which properties P answers Question 1 when G is abelian?
- Which properties P answers Question 1 when $A = \mathbb{C}$?
- Which properties *P* answers Question 1 when *G* is discrete?

To a C^* -dynamical system (A, G, α) , our goal is to answer the following question:

Question 1: What properties P describes the ideal structure of $A \rtimes_{\alpha} G$?

During this talk, the hope is that we will provide some answers to this question.

Answering question 1 in the above form is hard.

Instead, we break it into special cases Some cases discussed today will be:

- Which properties P answers Question 1 when A is abelian?
- Which properties P answers Question 1 when G is abelian?
- Which properties P answers Question 1 when $A = \mathbb{C}$?
- Which properties P answers Question 1 when G is discrete?

And of course, we have overlaps between the above cases.

Table of Contents

1 Introduction, history and motivation

Some notation and preliminaries

Disclaimer

Malthe Munk Karbo February 3, 2017 7 / 13

Disclaimer

The theory discussed today is quite heavy and dull for the uninitiated.

Malthe Munk Karbo February 3, 2017 7 /

Disclaimer

The theory discussed today is quite heavy and dull for the uninitiated. Sorry.

Malthe Munk Karbo February 3, 2017 7 /

A short primer on C^* -dynamical systems.

A short primer on C^* -dynamical systems.

Lots of prerequisite theory needed to understand todays talk.

Lots of prerequisite theory needed to understand todays talk. Not possible to cover all.

Lots of prerequisite theory needed to understand todays talk. Not possible to cover all.

Definitin

An abstract C^* -algebra A is a Banach algebra A equipped with an involution such that

$$||a||^2 = ||a^*a||,$$

for all $a \in A$.

Lots of prerequisite theory needed to understand todays talk. Not possible to cover all.

Definitin

An abstract C^* -algebra A is a Banach algebra A equipped with an involution such that

$$||a||^2 = ||a^*a||,$$

for all $a \in A$.

Notable examples include

Lots of prerequisite theory needed to understand todays talk. Not possible to cover all.

Definitin

An abstract C^* -algebra A is a Banach algebra A equipped with an involution such that

$$||a||^2 = ||a^*a||,$$

for all $a \in A$.

Notable examples include

• The algebra $C_0(X)$ of continuous functions vanishing at infinity on a locally compact Hausdorff space X.

Lots of prerequisite theory needed to understand todays talk. Not possible to cover all.

Definitin

An abstract C^* -algebra A is a Banach algebra A equipped with an involution such that

$$||a||^2 = ||a^*a||,$$

for all $a \in A$.

Notable examples include

- The algebra $C_0(X)$ of continuous functions vanishing at infinity on a locally compact Hausdorff space X.
- The algebra $\mathbb{B}(\mathcal{H})$ of bounded linear operators on a Hilbert space \mathcal{H} .

Lots of prerequisite theory needed to understand todays talk. Not possible to cover all.

Definitin

An abstract C^* -algebra A is a Banach algebra A equipped with an involution such that

$$||a||^2 = ||a^*a||,$$

for all $a \in A$.

Notable examples include

- The algebra $C_0(X)$ of continuous functions vanishing at infinity on a locally compact Hausdorff space X.
- ullet The algebra $\mathbb{B}(\mathcal{H})$ of bounded linear operators on a Hilbert space $\mathcal{H}.$
- The group C^* -algebras $C^*_r(G)$ and $C^*(G)$ of a locally compact Hausdorff group G.

Lots of prerequisite theory needed to understand todays talk. Not possible to cover all.

Definitin

An abstract C^* -algebra A is a Banach algebra A equipped with an involution such that

$$||a||^2 = ||a^*a||,$$

for all $a \in A$.

Notable examples include

- The algebra $C_0(X)$ of continuous functions vanishing at infinity on a locally compact Hausdorff space X.
- ullet The algebra $\mathbb{B}(\mathcal{H})$ of bounded linear operators on a Hilbert space \mathcal{H} .
- The group C^* -algebras $C^*_r(G)$ and $C^*(G)$ of a locally compact Hausdorff group G.

All of these C^* -algebras play a crucial role in the theory of crossed products.

Definitin

An abstract C^* -algebra A is an involutive Banach algebra A satisfying

$$||a||^2 = ||a^*a||,$$

for all $a \in A$.

Definitin

An abstract C^* -algebra A is an involutive Banach algebra A satisfying

$$||a||^2 = ||a^*a||,$$

for all $a \in A$.

Definition

A representation of a C^* -algebra A is a pair (π, \mathcal{H}) consisting of a bounded *-homomorphism $\pi \colon A \to \mathbb{B}(\mathcal{H})$.

Definitin

An abstract C^* -algebra A is an involutive Banach algebra A satisfying

$$||a||^2 = ||a^*a||,$$

for all $a \in A$.

Definition

A representation of a C^* -algebra A is a pair (π, \mathcal{H}) consisting of a bounded *-homomorphism $\pi \colon A \to \mathbb{B}(\mathcal{H})$.

The Gelfand-Neimark-Segal result

The Gelfand-Neimark-Segal result is an important result stating that all C^* -algebras A are faithfully represented on a Hilbert space.

Malthe Munk Karbo February 3, 2017

Definition

A topological group G is topological space equipped with a group structure such that the maps $G \times G \ni (s,t) \mapsto st$ and $G \ni t \mapsto t^{-1} \in G$ are continuous for all $s,t \in G$.

Definition

A topological group G is topological space equipped with a group structure such that the maps $G \times G \ni (s,t) \mapsto st$ and $G \ni t \mapsto t^{-1} \in G$ are continuous for all $s,t \in G$. By G we will **always** denote a topological group G with a locally compact Hausdorff topology.

Definition

A topological group G is topological space equipped with a group structure such that the maps $G \times G \ni (s,t) \mapsto st$ and $G \ni t \mapsto t^{-1} \in G$ are continuous for all $s,t \in G$. By G we will **always** denote a topological group G with a locally compact Hausdorff topology.

Important cases include \mathbb{Z} , \mathbb{R} and S^1 .

Definition

A topological group G is topological space equipped with a group structure such that the maps $G \times G \ni (s,t) \mapsto st$ and $G \ni t \mapsto t^{-1} \in G$ are continuous for all $s,t \in G$. By G we will **always** denote a topological group G with a locally compact Hausdorff topology.

Important cases include \mathbb{Z} , \mathbb{R} and S^1 . Topological group are important and arises naturally in many fields of mathematics.

Malthe Munk Karbo February 3, 2017 10 / 13

Definition

A topological group G is topological space equipped with a group structure such that the maps $G \times G \ni (s,t) \mapsto st$ and $G \ni t \mapsto t^{-1} \in G$ are continuous for all $s,t \in G$. By G we will **always** denote a topological group G with a locally compact Hausdorff topology.

Important cases include \mathbb{Z} , \mathbb{R} and S^1 . Topological group are important and arises naturally in many fields of mathematics. We will need results from the theory of abstract harmonic analysis, representation theory and Fourier analysis

Definition

A topological group G is topological space equipped with a group structure such that the maps $G \times G \ni (s,t) \mapsto st$ and $G \ni t \mapsto t^{-1} \in G$ are continuous for all $s,t \in G$. By G we will **always** denote a topological group G with a locally compact Hausdorff topology.

Important cases include \mathbb{Z} , \mathbb{R} and S^1 . Topological group are important and arises naturally in many fields of mathematics. We will need results from the theory of **abstract** harmonic analysis, representation theory and Fourier analysis

Definition

A unitary representation of a group G is a pair (u,\mathcal{H}) consisting of a group homomorphism $u\colon G\to \mathcal{U}(\mathcal{H})$ such that for all $\xi\in\mathcal{H}$ the map $t\mapsto u_t\xi$ is continuous (= u is strongly continuous).

Malthe Munk Karbo February 3, 2017

Definition

A topological group G is topological space equipped with a group structure such that the maps $G \times G \ni (s,t) \mapsto st$ and $G \ni t \mapsto t^{-1} \in G$ are continuous for all $s,t \in G$. By G we will **always** denote a topological group G with a locally compact Hausdorff topology.

Definition

A unitary representation of a group G is a pair (u,\mathcal{H}) consisting of a group homomorphism $u\colon G\to \mathcal{U}(\mathcal{H})$ such that for all $\xi\in\mathcal{H}$ the map $t\mapsto u_t\xi$ is continuous (= u is strongly continuous).

Definition

A topological group G is topological space equipped with a group structure such that the maps $G \times G \ni (s,t) \mapsto st$ and $G \ni t \mapsto t^{-1} \in G$ are continuous for all $s,t \in G$. By G we will **always** denote a topological group G with a locally compact Hausdorff topology.

Definition

A unitary representation of a group G is a pair (u,\mathcal{H}) consisting of a group homomorphism $u\colon G\to \mathcal{U}(\mathcal{H})$ such that for all $\xi\in\mathcal{H}$ the map $t\mapsto u_t\xi$ is continuous (= u is strongly continuous).

An important example is the *left-regular representation* of *G*:

Definition

A topological group G is topological space equipped with a group structure such that the maps $G \times G \ni (s,t) \mapsto st$ and $G \ni t \mapsto t^{-1} \in G$ are continuous for all $s,t \in G$. By G we will **always** denote a topological group G with a locally compact Hausdorff topology.

Definition

A unitary representation of a group G is a pair (u,\mathcal{H}) consisting of a group homomorphism $u\colon G\to \mathcal{U}(\mathcal{H})$ such that for all $\xi\in\mathcal{H}$ the map $t\mapsto u_t\xi$ is continuous (= u is strongly continuous).

An important example is the *left-regular representation* of *G*:

Definition

The left-regular representation of G is the tuple $(\lambda, L^2(G))$, where

$$\lambda_t \xi(s) = \xi(t^{-1}s),$$

for $\xi \in L^2(G)$ and $s, t \in G$.

Malthe Munk Karbo February 3, 2017 12 / 13

Definition

A C^* -dynamical system is a triple (A,G,α) consisting of

Definition

A C^* -dynamical system is a triple (A,G,α) consisting of

• A C^* -algebra A,

Definition

A C^* -dynamical system is a triple (A,G,α) consisting of

- A C*-algebra A,
- A topological group G and

Definition

A C^* -dynamical system is a triple (A, G, α) consisting of

- A C*-algebra A,
- A topological group G and
- a strongly continuous group homomorphism $\alpha \colon G \to \operatorname{Aut}(A)$ $(\forall a \in A \colon G \ni t \mapsto \alpha_t(a) \in A \text{ is continuous}).$

Definition

A C^* -dynamical system is a triple (A, G, α) consisting of

- A C*-algebra A,
- A topological group G and
- a strongly continuous group homomorphism $\alpha \colon G \to \operatorname{Aut}(A)$ $(\forall a \in A \colon G \ni t \mapsto \alpha_t(a) \in A \text{ is continuous}).$

Definition

A C^* -dynamical system is a triple (A, G, α) consisting of

- A C*-algebra A,
- A topological group G and
- a strongly continuous group homomorphism $\alpha \colon G \to \operatorname{Aut}(A)$ ($\forall a \in A \colon G \ni t \mapsto \alpha_t(a) \in A$ is continuous).

Definition

A covariant representation of (A,G,α) is a triple (π,u,\mathcal{H}) consisting of representations of A and G as bounded operators on a common Hilbert space \mathcal{H} satisfying the relation

$$u_t\pi(a)u_t^*=\pi(\alpha_t(a))$$
 for all $a\in A$ and $t\in G$.

Malthe Munk Karbo February 3, 2017

Definition

A C^* -dynamical system is a triple (A, G, α) consisting of

- A C*-algebra A,
- A topological group G and
- a strongly continuous group homomorphism $\alpha \colon G \to \operatorname{Aut}(A)$ $(\forall a \in A \colon G \ni t \mapsto \alpha_t(a) \in A \text{ is continuous}).$

Definition

A covariant representation of (A,G,α) is a triple (π,u,\mathcal{H}) consisting of representations of A and G as bounded operators on a common Hilbert space \mathcal{H} satisfying the relation

$$u_t\pi(a)u_t^*=\pi(\alpha_t(a))$$
 for all $a\in A$ and $t\in G$.

When G is discrete and A is unital, it is easy to associate to such a triple a **nice** new C^* -algebra:

When G is discrete and A is unital, it is easy to associate to such a triple a **nice** new C^* -algebra:

When G is discrete and A is unital, it is easy to associate to such a triple a **nice** new C^* -algebra:

Discrete crossed product cookbook recipe

• Given (A, G, α) with G discrete and A unital, consider the *-algebra

$$C_c(G,A) = \left\{ \sum_{t \in F} a_t \delta_t \mid F \subseteq G \text{ finite} \right\}.$$

Malthe Munk Karbo February 3, 2017

When G is discrete and A is unital, it is easy to associate to such a triple a **nice** new C^* -algebra:

Discrete crossed product cookbook recipe

1 Given (A, G, α) with G discrete and A unital, consider the *-algebra

$$C_c(G,A) = \left\{ \sum_{t \in F} a_t \delta_t \mid F \subseteq G \text{ finite} \right\}.$$

② Pick a covariant representation (π, u, \mathcal{H}) of (A, G, α) (they exist!)

Malthe Munk Karbo February 3, 2017

When G is discrete and A is unital, it is easy to associate to such a triple a **nice** new C^* -algebra:

Discrete crossed product cookbook recipe

• Given (A, G, α) with G discrete and A unital, consider the *-algebra

$$C_c(G,A) = \left\{ \sum_{t \in F} a_t \delta_t \mid F \subseteq G \text{ finite} \right\}.$$

- ② Pick a covariant representation (π, u, \mathcal{H}) of (A, G, α) (they exist!)
- **9** Define a representation $\pi \times u$ of $C_c(G, A)$ by

$$\pi
times u\left(\sum \mathsf{a}_t\delta_t
ight):=\sum \pi(\mathsf{a}_t)u_t$$

Malthe Munk Karbo February 3, 2017

When G is discrete and A is unital, it is easy to associate to such a triple a **nice** new C^* -algebra:

Discrete crossed product cookbook recipe

• Given (A, G, α) with G discrete and A unital, consider the *-algebra

$$C_c(G,A) = \left\{ \sum_{t \in F} a_t \delta_t \mid F \subseteq G \text{ finite} \right\}.$$

- 2 Pick a covariant representation (π, u, \mathcal{H}) of (A, G, α) (they exist!)
- Define a representation $\pi \times u$ of $C_c(G,A)$ by

$$\pi
times u\left(\sum a_t\delta_t
ight):=\sum \pi(a_t)u_t$$

• Take the closure of $\pi \rtimes u(C_c(G,A))$

When G is discrete and A is unital, it is easy to associate to such a triple a **nice** new C^* -algebra:

Discrete crossed product cookbook recipe

• Given (A, G, α) with G discrete and A unital, consider the *-algebra

$$C_c(G,A) = \left\{ \sum_{t \in F} a_t \delta_t \mid F \subseteq G \text{ finite} \right\}.$$

- ② Pick a covariant representation (π, u, \mathcal{H}) of (A, G, α) (they exist!)
- **3** Define a representation $\pi \rtimes u$ of $C_c(G, A)$ by

$$\pi
times u\left(\sum a_t\delta_t\right):=\sum \pi(a_t)u_t$$

• Take the closure of $\pi \times u(C_c(G, A))$

This is the building stone for constructing crossed products.

When G is discrete and A is unital, it is easy to associate to such a triple a **nice** new C^* -algebra:

Discrete crossed product cookbook recipe

• Given (A, G, α) with G discrete and A unital, consider the *-algebra

$$C_c(G,A) = \left\{ \sum_{t \in F} a_t \delta_t \mid F \subseteq G \text{ finite} \right\}.$$

- ② Pick a covariant representation (π, u, \mathcal{H}) of (A, G, α) (they exist!)
- **3** Define a representation $\pi \rtimes u$ of $C_c(G, A)$ by

$$\pi
times u\left(\sum a_t\delta_t\right):=\sum \pi(a_t)u_t$$

• Take the closure of $\pi \rtimes u(C_c(G,A))$

This is the building stone for constructing crossed products. Fails when G is non-discrete.

Malthe Munk Karbo February 3, 2017

Table of Contents

Introduction, history and motivation

Some notation and preliminaries