

# Simplicity and uniqueness of trace of group $C^*$ -algebras

Bachelor thesis defense

Malthe Munk Karbo

Advisor: Mikael Rørdam

February 3, 2017

- 1 Introduction, history and motivation
- 2 An introduction to various topics
  - Group  $C^*$ -algebras
  - Amenability
  - Group actions and the Reduced crossed product
  - Dixmier property
- 3 Boundary actions
- 4  $C^*$ -simplicity using boundary actions

- 1 Introduction, history and motivation
- 2 An introduction to various topics
  - Group  $C^*$ -algebras
  - Amenability
  - Group actions and the Reduced crossed product
  - Dixmier property
- 3 Boundary actions
- 4  $C^*$ -simplicity using boundary actions

# Motivation and background

- In 1975, Robert T. Powers proved that  $\mathbb{F}_2$  was  $C^*$ -simple and that this implied the unique trace property of  $\mathbb{F}_2$ .

## Motivation and background

- In 1975, Robert T. Powers proved that  $\mathbb{F}_2$  was  $C^*$ -simple and that this implied the unique trace property of  $\mathbb{F}_2$ .
- The proof Powers gave was generalised to a larger class of groups - Powers groups.

## Motivation and background

- In 1975, Robert T. Powers proved that  $\mathbb{F}_2$  was  $C^*$ -simple and that this implied the unique trace property of  $\mathbb{F}_2$ .
- The proof Powers gave was generalised to a larger class of groups - Powers groups.

We want to study a group  $G$  by studying group  $C^*$ -algebras, for an example by studying  $C_r^*(G)$ .

# Motivation and background

- In 1975, Robert T. Powers proved that  $\mathbb{F}_2$  was  $C^*$ -simple and that this implied the unique trace property of  $\mathbb{F}_2$ .
- The proof Powers gave was generalised to a larger class of groups - Powers groups.

We want to study a group  $G$  by studying group  $C^*$ -algebras, for an example by studying  $C_r^*(G)$ .

Was open question if  $C_r^*(G)$  simple  $\iff C_r^*(G)$  unique trace  $\iff$  trivial amenable radical of  $G$ .



# Motivation and background

- In 1975, Robert T. Powers proved that  $\mathbb{F}_2$  was  $C^*$ -simple and that this implied the unique trace property of  $\mathbb{F}_2$ .
- The proof Powers gave was generalised to a larger class of groups - Powers groups.

We want to study a group  $G$  by studying group  $C^*$ -algebras, for an example by studying  $C_r^*(G)$ .

Was open question if  $C_r^*(G)$  simple  $\iff C_r^*(G)$  unique trace  $\iff$  trivial amenable radical of  $G$ .

Today we see for discrete countable group  $G$  :

$$C_r^*(G) \text{ simple} \implies C_r^*(G) \text{ unique trace} \iff \text{trivial amenable radical of } G.$$

- 1 Introduction, history and motivation
- 2 An introduction to various topics
  - Group  $C^*$ -algebras
  - Amenability
  - Group actions and the Reduced crossed product
  - Dixmier property
- 3 Boundary actions
- 4  $C^*$ -simplicity using boundary actions

## A three minute crash-course on group $C^*$ -algebras

We are interested in describing discrete groups  $G$  using theory of  $C^*$ -algebras. How do we do this?

## A three minute crash-course on group $C^*$ -algebras

We are interested in describing discrete groups  $G$  using theory of  $C^*$ -algebras. How do we do this?

### Unitary representations

A *unitary representation* of a discrete group  $G$  is a pair  $(u, \mathcal{H})$ , where  $\mathcal{H}$  is a Hilbert space and  $u: G \rightarrow \mathcal{U}(\mathcal{H})$  is a group homomorphism into the group of unitary operators on  $\mathcal{H}$ . We use the following notation for unitary representations:

$$u(g) := u_g, \quad g \in G.$$

## A three minute crash-course on group $C^*$ -algebras

We are interested in describing discrete groups  $G$  using theory of  $C^*$ -algebras. How do we do this?

### Unitary representations

A *unitary representation* of a discrete group  $G$  is a pair  $(u, \mathcal{H})$ , where  $\mathcal{H}$  is a Hilbert space and  $u: G \rightarrow \mathcal{U}(\mathcal{H})$  is a group homomorphism into the group of unitary operators on  $\mathcal{H}$ . We use the following notation for unitary representations:

$$u(g) := u_g, \quad g \in G.$$

### Group $C^*$ -algebra

Given a unitary representation  $(u, \mathcal{H})$  of a discrete group  $G$ , we define the *group  $C^*$ -algebra* associated to  $u$  by  $C_u^*(G) := C^*(\{u_g : g \in G\}) \subseteq B(\mathcal{H})$ , the  $C^*$ -subalgebra of  $B(\mathcal{H})$  generated by the unitaries  $\{u_g\}_{g \in G}$ .

# A three minute crash-course on group $C^*$ -algebras

## An important example

## An important example

The pair  $(\lambda, \ell^2(G))$ , where  $\lambda: G \rightarrow \mathcal{U}(\ell^2(G))$  is defined by

$$(\lambda_g \xi)(s) = \xi(g^{-1}s), \quad s, g \in G, \quad \xi \in \ell^2(G),$$

is called the *left-regular representation* of  $G$ ,

## A three minute crash-course on group $C^*$ -algebras

### An important example

The pair  $(\lambda, \ell^2(G))$ , where  $\lambda: G \rightarrow \mathcal{U}(\ell^2(G))$  is defined by

$$(\lambda_g \xi)(s) = \xi(g^{-1}s), \quad s, g \in G, \quad \xi \in \ell^2(G),$$

is called the *left-regular representation* of  $G$ , and the associated group  $C^*$ -algebra,

$$C_r^*(G) := C_\lambda^*(G) \subseteq B(\ell^2(G)),$$

called the *reduced group  $C^*$ -algebra* of  $G$ .



## A three minute crash-course on group $C^*$ -algebras

The set  $\{\delta_g\}_{g \in G} \subseteq \ell^2(G)$ , where for  $s, g \in G$  we define

$$\delta_s(g) = \begin{cases} 1 & \text{if } s = g \\ 0 & \text{else} \end{cases},$$

is an orthonormal basis for  $\ell^2(G)$ .

## A three minute crash-course on group $C^*$ -algebras

The set  $\{\delta_g\}_{g \in G} \subseteq \ell^2(G)$ , where for  $s, g \in G$  we define

$$\delta_s(g) = \begin{cases} 1 & \text{if } s = g \\ 0 & \text{else} \end{cases},$$

is an orthonormal basis for  $\ell^2(G)$ .

### Lemma

For discrete group  $G$ , the map  $\tau_0: C_r^*(G) \rightarrow \mathbb{C}$  defined by  $\tau_0(x) = \langle x\delta_e, \delta_e \rangle$  is a faithful tracial state on  $C_r^*(G)$ .

## A three minute crash-course on group $C^*$ -algebras

The set  $\{\delta_g\}_{g \in G} \subseteq \ell^2(G)$ , where for  $s, g \in G$  we define

$$\delta_s(g) = \begin{cases} 1 & \text{if } s = g \\ 0 & \text{else} \end{cases},$$

is an orthonormal basis for  $\ell^2(G)$ .

### Lemma

For discrete group  $G$ , the map  $\tau_0: C_r^*(G) \rightarrow \mathbb{C}$  defined by  $\tau_0(x) = \langle x\delta_e, \delta_e \rangle$  is a faithful tracial state on  $C_r^*(G)$ .

### Definition

For a discrete group  $G$ , we say that  $G$  is  $C^*$ -*simple* if the reduced group  $C^*$ -algebra,  $C_r^*(G)$ , is simple. We say that  $G$  has the *unique trace property* if  $\tau_0$ , defined as above, is the only tracial state on  $C_r^*(G)$ .

Let  $G$  be a countable discrete group.

Let  $G$  be a countable discrete group.

## Definition

A group  $G$  is *amenable* if there exists a finitely additive left-invariant probability measure  $\mu$  on  $G$ .

# Amenability

Let  $G$  be a countable discrete group.

## Definition

A group  $G$  is *amenable* if there exists a finitely additive left-invariant probability measure  $\mu$  on  $G$ .

Amenability is stable under usual group operations such as passing to subgroups and quotients.

Let  $G$  be a countable discrete group.

## Definition

A group  $G$  is *amenable* if there exists a finitely additive left-invariant probability measure  $\mu$  on  $G$ .

Amenability is stable under usual group operations such as passing to subgroups and quotients.

## Proposition

Any group  $G$  has a largest amenable normal subgroup,  $\text{AR}_G$ , called the *amenable radical* of  $G$ .

# Amenability

Let  $G$  be a countable discrete group.

## Definition

A group  $G$  is *amenable* if there exists a finitely additive left-invariant probability measure  $\mu$  on  $G$ .

Amenability is stable under usual group operations such as passing to subgroups and quotients.

## Proposition

Any group  $G$  has a largest amenable normal subgroup,  $\text{AR}_G$ , called the *amenable radical* of  $G$ .

Amenability is a strong property to satisfy, and has many consequences.



## Group actions and the Reduced crossed product

Let  $G$  be a countable discrete group and  $\mathcal{A}$  a unital  $C^*$ -algebra.

# Group actions and the Reduced crossed product

Let  $G$  be a countable discrete group and  $\mathcal{A}$  a unital  $C^*$ -algebra.

## Definition

An *action* of  $G$  on  $\mathcal{A}$  is a group homomorphism  $\alpha: G \rightarrow \text{Aut}(\mathcal{A})$  from  $G$  to the group of  $*$ -automorphism on  $\mathcal{A}$ . Abbreviated  $G \overset{\alpha}{\curvearrowright} \mathcal{A}$ .

# Group actions and the Reduced crossed product

Let  $G$  be a countable discrete group and  $\mathcal{A}$  a unital  $C^*$ -algebra.

## Definition

An *action* of  $G$  on  $\mathcal{A}$  is a group homomorphism  $\alpha: G \rightarrow \text{Aut}(\mathcal{A})$  from  $G$  to the group of  $*$ -automorphism on  $\mathcal{A}$ . Abbreviated  $G \curvearrowright^\alpha \mathcal{A}$ .

Whenever  $G \curvearrowright^\alpha \mathcal{A} \curvearrowright$  we make a  $C^*$ -algebra  $\mathcal{A} \rtimes_{\alpha,r} G$  which contains  $\mathcal{A}$  and  $C_r^*(G)$  in a nice way. This  $C^*$ -algebra is called the reduced crossed product.

## Dixmier property

Let  $\mathcal{A}$  be a unital  $C^*$ -algebra.

## Dixmier property

Let  $\mathcal{A}$  be a unital  $C^*$ -algebra. For  $a \in \mathcal{A}$  define  $D(a) := \text{conv} \{ uau^* : u \in \mathcal{U}(\mathcal{A}) \}$ .

# Dixmier property

Let  $\mathcal{A}$  be a unital  $C^*$ -algebra. For  $a \in \mathcal{A}$  define  $D(a) := \text{conv} \{ uau^* : u \in \mathcal{U}(\mathcal{A}) \}$ .

## Definition

We say that  $\mathcal{A}$  has the *Dixmier property* if

$$\overline{D(a)} \cap \mathbb{C}1_{\mathcal{A}} \neq \emptyset \quad \forall a \in \mathcal{A}.$$

## Dixmier property

Let  $\mathcal{A}$  be a unital  $C^*$ -algebra. For  $a \in \mathcal{A}$  define  $D(a) := \text{conv} \{uau^* : u \in \mathcal{U}(\mathcal{A})\}$ .

### Definition

We say that  $\mathcal{A}$  has the *Dixmier property* if

$$\overline{D(a)} \cap \mathbb{C}1_{\mathcal{A}} \neq \emptyset \quad \forall a \in \mathcal{A}.$$

### Lemma

Let  $\mathcal{A}$  be a unital  $C^*$ -algebra with a tracial state  $\tau$ . If  $\mathcal{A}$  has the Dixmier property, then  $\tau$  is unique.

## Dixmier property

Let  $\mathcal{A}$  be a unital  $C^*$ -algebra. For  $a \in \mathcal{A}$  define  $D(a) := \text{conv} \{uau^* : u \in \mathcal{U}(\mathcal{A})\}$ .

### Definition

We say that  $\mathcal{A}$  has the *Dixmier property* if

$$\overline{D(a)} \cap \mathbb{C}1_{\mathcal{A}} \neq \emptyset \quad \forall a \in \mathcal{A}.$$

### Lemma

Let  $\mathcal{A}$  be a unital  $C^*$ -algebra with a tracial state  $\tau$ . If  $\mathcal{A}$  has the Dixmier property, then  $\tau$  is unique.

### Proposition

Let  $\mathcal{A}$  be a unital  $C^*$ -algebra with a faithful tracial state  $\tau$ . If  $\mathcal{A}$  has the Dixmier property, then  $\mathcal{A}$  is simple.



- 1 Introduction, history and motivation
- 2 An introduction to various topics
  - Group  $C^*$ -algebras
  - Amenability
  - Group actions and the Reduced crossed product
  - Dixmier property
- 3 Boundary actions
- 4  $C^*$ -simplicity using boundary actions

## Boundary actions: $G$ -spaces

In the following, let  $G$  be a second countable locally compact group and  $X$  a compact Hausdorff space.

## Boundary actions: $G$ -spaces

In the following, let  $G$  be a second countable locally compact group and  $X$  a compact Hausdorff space.

### Definition

A  $G$ -action of  $G$  on  $X$  is a continuous group homomorphism  $\alpha: G \rightarrow \text{Homeo}(X)$ . We write  $G \curvearrowright X$  to mean that  $G$  acts on  $X$ , and as before, we write  $\alpha(g)(x) := g.x$  for  $x \in X$  and  $g \in G$ .

## Boundary actions: $G$ -spaces

In the following, let  $G$  be a second countable locally compact group and  $X$  a compact Hausdorff space.

### Definition

A  $G$ -action of  $G$  on  $X$  is a continuous group homomorphism  $\alpha: G \rightarrow \text{Homeo}(X)$ . We write  $G \curvearrowright X$  to mean that  $G$  acts on  $X$ , and as before, we write  $\alpha(g)(x) := g.x$  for  $x \in X$  and  $g \in G$ . A space equipped with a  $G$ -action is called a  $G$ -space.

## Boundary actions: $G$ -spaces

In the following, let  $G$  be a second countable locally compact group and  $X$  a compact Hausdorff space.

### Definition

A  $G$ -action of  $G$  on  $X$  is a continuous group homomorphism  $\alpha: G \rightarrow \text{Homeo}(X)$ . We write  $G \curvearrowright X$  to mean that  $G$  acts on  $X$ , and as before, we write  $\alpha(g)(x) := g.x$  for  $x \in X$  and  $g \in G$ . A space equipped with a  $G$ -action is called a  $G$ -space.

$\text{Prob}(X) = \mathcal{S}(C(X))$  with weak\* topology.

## Boundary actions: $G$ -spaces

In the following, let  $G$  be a second countable locally compact group and  $X$  a compact Hausdorff space.

### Definition

A  $G$ -action of  $G$  on  $X$  is a continuous group homomorphism  $\alpha: G \rightarrow \text{Homeo}(X)$ . We write  $G \curvearrowright X$  to mean that  $G$  acts on  $X$ , and as before, we write  $\alpha(g)(x) := g.x$  for  $x \in X$  and  $g \in G$ . A space equipped with a  $G$ -action is called a  $G$ -space.

$\text{Prob}(X) = \mathcal{S}(C(X))$  with weak\* topology.

④  $G \curvearrowright X \rightsquigarrow G \curvearrowright C(X)$  by equipping  $G$  with discrete topology and defining

$$g.f(x) := f(g^{-1}.x), \quad g \in G, \quad x \in X \text{ and } f \in C(X).$$

## Boundary actions: $G$ -spaces

In the following, let  $G$  be a second countable locally compact group and  $X$  a compact Hausdorff space.

### Definition

A  $G$ -action of  $G$  on  $X$  is a continuous group homomorphism  $\alpha: G \rightarrow \text{Homeo}(X)$ . We write  $G \curvearrowright X$  to mean that  $G$  acts on  $X$ , and as before, we write  $\alpha(g)(x) := g.x$  for  $x \in X$  and  $g \in G$ . A space equipped with a  $G$ -action is called a  $G$ -space.

$\text{Prob}(X) = \mathcal{S}(C(X))$  with weak\* topology.

- ①  $G \curvearrowright X \rightsquigarrow G \curvearrowright C(X)$  by equipping  $G$  with discrete topology and defining

$$g.f(x) := f(g^{-1}.x), \quad g \in G, \quad x \in X \text{ and } f \in C(X).$$

- ②  $G \curvearrowright X \rightsquigarrow G \curvearrowright \text{Prob}(X)$  by equipping  $G$  with discrete topology and defining

$$g.\mu(E) := \mu(g^{-1}.E), \quad \mu \in \text{Prob}(X), \quad g \in G \text{ and } E \subseteq X \text{ Borel},$$

### Definition

A  $G$ -space  $X$  is said to be minimal if every  $G$ -orbit is dense in  $X$ .



### Definition

A  $G$ -space  $X$  is said to be minimal if every  $G$ -orbit is dense in  $X$ .

### Definition

A subset  $Y \subseteq X$  of a  $G$ -space  $X$  is *minimal* if it is non-empty, closed and  $G$ -invariant and a minimal element in the set of non-empty, closed and  $G$ -invariant subsets of  $X$ .

### Definition

A  $G$ -space  $X$  is said to be minimal if every  $G$ -orbit is dense in  $X$ .

### Definition

A subset  $Y \subseteq X$  of a  $G$ -space  $X$  is *minimal* if it is non-empty, closed and  $G$ -invariant and a minimal element in the set of non-empty, closed and  $G$ -invariant subsets of  $X$ .

An application of Zorn's lemma yields that every compact  $G$ -space  $X$  has a minimal subspace. Moreover a subset  $Y$  is minimal if and only if the action  $G \curvearrowright Y$  is minimal

## Definition

For a  $G$ -space  $X$ , the action of  $G$  is *strongly proximal* if for each  $\mu \in \text{Prob}(X)$  we have

$$\overline{G \cdot \mu}^{w^*} \cap \{\delta_x : x \in X\} \neq \emptyset.$$

## Definition

For a  $G$ -space  $X$ , the action of  $G$  is *strongly proximal* if for each  $\mu \in \text{Prob}(X)$  we have

$$\overline{G \cdot \mu}^{w*} \cap \{\delta_x : x \in X\} \neq \emptyset.$$

## Definition

If  $X$  is a compact  $G$ -space, we say that the action  $G \curvearrowright X$  is a *boundary action* if it is minimal and strongly proximal. A compact  $G$ -space  $X$  for which the action is a boundary action is called a  *$G$ -boundary*.

There are quite a few different equivalent conditions for a compact  $G$ -space  $X$  to be a  $G$ -boundary.

There are quite a few different equivalent conditions for a compact  $G$ -space  $X$  to be a  $G$ -boundary.

### Proposition

For a compact  $G$ -space  $X$ , TFAE:

- ❶  $X$  is a  $G$ -boundary,
- ❷ For every  $\nu \in \text{Prob}(X)$  we have  $\{\delta_x : x \in X\} \subseteq \overline{G \cdot \nu}^{w^*}$  and
- ❸  $\text{Prob}(X)$  admits no non-trivial closed convex subsets which are  $G$ -invariant under the induced action  $G \curvearrowright \text{Prob}(X)$ .

There are quite a few different equivalent conditions for a compact  $G$ -space  $X$  to be a  $G$ -boundary.

### Proposition

For a compact  $G$ -space  $X$ , TFAE:

- ❶  $X$  is a  $G$ -boundary,
- ❷ For every  $\nu \in \text{Prob}(X)$  we have  $\{\delta_x : x \in X\} \subseteq \overline{G \cdot \nu}^{w^*}$  and
- ❸  $\text{Prob}(X)$  admits no non-trivial closed convex subsets which are  $G$ -invariant under the induced action  $G \curvearrowright \text{Prob}(X)$ .

In laymans terms, a  $G$ -boundary is a space such that every  $G$ -orbit in  $\text{Prob}(X)$  becomes a black hole for all the point masses when going to weak\* closure.

## Properties of different actions

It turns out that some of the properties we've seen behave nicely and have certain properties:



It turns out that some of the properties we've seen behave nicely and have certain properties:

- The  $G$ -boundary property is preserved under quotients,

It turns out that some of the properties we've seen behave nicely and have certain properties:

- The  $G$ -boundary property is preserved under quotients,
- Strongly proximality is preserved under taking products,

It turns out that some of the properties we've seen behave nicely and have certain properties:

- The  $G$ -boundary property is preserved under quotients,
- Strongly proximality is preserved under taking products,
- If  $G \curvearrowright Y$  is minimal and  $X$  is a  $G$ -boundary, then any continuous  $G$ -map  $\varphi: Y \rightarrow X$  will be unique and surjective.

We now proceed to construct a largest  $G$ -boundary, in the sense that it has the universal property that every other  $G$ -boundary is a quotient of it.

We now proceed to construct a largest  $G$ -boundary, in the sense that it has the universal property that every other  $G$ -boundary is a quotient of it.

### Definition

A compact  $G$ -boundary  $Z$  is a *universal boundary* for  $G$  if every other  $G$ -boundary is a quotient of it.

We now proceed to construct a largest  $G$ -boundary, in the sense that it has the universal property that every other  $G$ -boundary is a quotient of it.

## Definition

A compact  $G$ -boundary  $Z$  is a *universal boundary* for  $G$  if every other  $G$ -boundary is a quotient of it.

**remark:** Will be unique up to isomorphism.

We now proceed to construct a largest  $G$ -boundary, in the sense that it has the universal property that every other  $G$ -boundary is a quotient of it.

## Definition

A compact  $G$ -boundary  $Z$  is a *universal boundary* for  $G$  if every other  $G$ -boundary is a quotient of it.

**remark:** Will be unique up to isomorphism.

## Proposition

There exists a universal boundary of  $G$ , called the Furstenburg boundary of  $G$ , denoted by  $\partial_F G$ .

We now proceed to construct a largest  $G$ -boundary, in the sense that it has the universal property that every other  $G$ -boundary is a quotient of it.

## Definition

A compact  $G$ -boundary  $Z$  is a *universal boundary* for  $G$  if every other  $G$ -boundary is a quotient of it.

**remark:** Will be unique up to isomorphism.

## Proposition

There exists a universal boundary of  $G$ , called the Furstenburg boundary of  $G$ , denoted by  $\partial_F G$ .

The proof is roughly:



We now proceed to construct a largest  $G$ -boundary, in the sense that it has the universal property that every other  $G$ -boundary is a quotient of it.

## Definition

A compact  $G$ -boundary  $Z$  is a *universal boundary* for  $G$  if every other  $G$ -boundary is a quotient of it.

**remark:** Will be unique up to isomorphism.

## Proposition

There exists a universal boundary of  $G$ , called the Furstenberg boundary of  $G$ , denoted by  $\partial_F G$ .

The proof is roughly:

- 1 There is a bound on the cardinality of every compact  $G$ -boundary.

We now proceed to construct a largest  $G$ -boundary, in the sense that it has the universal property that every other  $G$ -boundary is a quotient of it.

## Definition

A compact  $G$ -boundary  $Z$  is a *universal boundary* for  $G$  if every other  $G$ -boundary is a quotient of it.

**remark:** Will be unique up to isomorphism.

## Proposition

There exists a universal boundary of  $G$ , called the Furstenberg boundary of  $G$ , denoted by  $\partial_F G$ .

The proof is roughly:

- 1 There is a bound on the cardinality of every compact  $G$ -boundary.
- 2 Index them up to isomorphism by their cardinality and form their product. It will be strongly proximal and compact.

We now proceed to construct a largest  $G$ -boundary, in the sense that it has the universal property that every other  $G$ -boundary is a quotient of it.

## Definition

A compact  $G$ -boundary  $Z$  is a *universal boundary* for  $G$  if every other  $G$ -boundary is a quotient of it.

**remark:** Will be unique up to isomorphism.

## Proposition

There exists a universal boundary of  $G$ , called the Furstenberg boundary of  $G$ , denoted by  $\partial_F G$ .

The proof is roughly:

- 1 There is a bound on the cardinality of every compact  $G$ -boundary.
- 2 Index them up to isomorphism by their cardinality and form their product. It will be strongly proximal and compact.
- 3 Use Zorn's lemma to pick a minimal subset.

## Amenable normal subgroups and boundary actions

It turns out that amenability plays a role in how groups act on boundary spaces.

It turns out that amenability plays a role in how groups act on boundary spaces.

## Proposition

A closed amenable normal subgroup  $N \trianglelefteq G$  of a second countable locally compact group  $G$  acts trivially on every compact  $G$ -boundary  $X$ .

# Amenable normal subgroups and boundary actions

It turns out that amenability plays a role in how groups act on boundary spaces.

## Proposition

A closed amenable normal subgroup  $N \trianglelefteq G$  of a second countable locally compact group  $G$  acts trivially on every compact  $G$ -boundary  $X$ .

# Amenable normal subgroups and boundary actions

It turns out that amenability plays a role in how groups act on boundary spaces.

## Proposition

~~A closed amenable normal subgroup  $N \trianglelefteq G$  of a second countable locally compact group  $G$~~  An amenable normal subgroup  $N \trianglelefteq G$  of a countable discrete group  $G$  acts trivially on every compact  $G$ -boundary  $X$ .

# Amenable normal subgroups and boundary actions

It turns out that amenability plays a role in how groups act on boundary spaces.

## Proposition

~~A closed amenable normal subgroup  $N \trianglelefteq G$  of a second countable locally compact group  $G$~~  An amenable normal subgroup  $N \trianglelefteq G$  of a countable discrete group  $G$  acts trivially on every compact  $G$ -boundary  $X$ .

And more generally we have:



# Amenable normal subgroups and boundary actions

It turns out that amenability plays a role in how groups act on boundary spaces.

## Proposition

~~A closed amenable normal subgroup  $N \trianglelefteq G$  of a second countable locally compact group  $G$~~  An amenable normal subgroup  $N \trianglelefteq G$  of a countable discrete group  $G$  acts trivially on every compact  $G$ -boundary  $X$ .

And more generally we have:

## Proposition (Furman)

For any countable discrete group  $G$  with  $t \in G$ , it holds that  $t \in \text{AR}_G$  if and only if  $t \in \ker(G \curvearrowright X)$  for all compact  $G$ -boundaries  $X$ .

# Table of Contents

- 1 Introduction, history and motivation
- 2 An introduction to various topics
  - Group  $C^*$ -algebras
  - Amenability
  - Group actions and the Reduced crossed product
  - Dixmier property
- 3 Boundary actions
- 4  $C^*$ -simplicity using boundary actions

## Some remarks and notation

From now  $G$  will be a discrete and countable group and  $X$  a compact Hausdorff space.

## Some remarks and notation

From now  $G$  will be a discrete and countable group and  $X$  a compact Hausdorff space. Before continuing onward, we recall:

## Some remarks and notation

From now  $G$  will be a discrete and countable group and  $X$  a compact Hausdorff space. Before continuing onward, we recall:

$$\textcircled{1} \quad G \curvearrowright X \rightsquigarrow G \curvearrowright C(X)$$

## Some remarks and notation

From now  $G$  will be a discrete and countable group and  $X$  a compact Hausdorff space. Before continuing onward, we recall:

- ❶  $G \curvearrowright X \rightsquigarrow G \curvearrowright C(X)$
- ❷  $G \curvearrowright X \rightsquigarrow G \curvearrowright \text{Prob}(X)$ .

## Some remarks and notation

From now  $G$  will be a discrete and countable group and  $X$  a compact Hausdorff space. Before continuing onward, we recall:

- ❶  $G \curvearrowright X \rightsquigarrow G \curvearrowright C(X)$
- ❷  $G \curvearrowright X \rightsquigarrow G \curvearrowright \text{Prob}(X)$ .
- ❸ If  $G \overset{\alpha}{\curvearrowright} C(X)$  we form  $C(X) \rtimes_{\alpha, r} G$ .

## Some remarks and notation

From now  $G$  will be a discrete and countable group and  $X$  a compact Hausdorff space. Before continuing onward, we recall:

- ❶  $G \curvearrowright X \rightsquigarrow G \curvearrowright C(X)$
- ❷  $G \curvearrowright X \rightsquigarrow G \curvearrowright \text{Prob}(X)$ .
- ❸ If  $G \overset{\alpha}{\curvearrowright} C(X)$  we form  $C(X) \rtimes_{\alpha, r} G$ .
- ❹  $G \curvearrowright \mathcal{S}(C_r^*(G))$  by  $t.\varphi(a) = \varphi(\lambda_t a \lambda_t^*)$ ,  $\varphi \in \mathcal{S}(C_r^*(G))$ ,  $a \in C_r^*(G)$  and  $t \in G$



## Some remarks and notation

From now  $G$  will be a discrete and countable group and  $X$  a compact Hausdorff space. Before continuing onward, we recall:

- ❶  $G \curvearrowright X \rightsquigarrow G \curvearrowright C(X)$
- ❷  $G \curvearrowright X \rightsquigarrow G \curvearrowright \text{Prob}(X)$ .
- ❸ If  $G \overset{\alpha}{\curvearrowright} C(X)$  we form  $C(X) \rtimes_{\alpha, r} G$ .
- ❹  $G \curvearrowright \mathcal{S}(C_r^*(G))$  by  $t.\varphi(a) = \varphi(\lambda_t a \lambda_t^*)$ ,  $\varphi \in \mathcal{S}(C_r^*(G))$ ,  $a \in C_r^*(G)$  and  $t \in G$
- ❺  $G \curvearrowright X \rightsquigarrow G \curvearrowright \mathcal{S}(C(X) \rtimes_r G)$ : for  $t \in G$  and any state  $\varphi$  define  $t.\varphi$  by

$$t.\varphi(a) = \varphi(\lambda_t a \lambda_t^*), \quad a \in C(X) \rtimes_r G.$$

We now start to show some of results mentioned in the beginning.

We now start to show some of results mentioned in the beginning.

### Theorem

Let  $G$  be a group and  $t \in G$ . Then  $\tau(\lambda_t) = 0$  for all tracial states  $\tau$  on  $C_r^*(G)$  if and only if  $t \notin \text{AR}_G$ .

We now start to show some of results mentioned in the beginning.

### Theorem

Let  $G$  be a group and  $t \in G$ . Then  $\tau(\lambda_t) = 0$  for all tracial states  $\tau$  on  $C_r^*(G)$  if and only if  $t \notin \text{AR}_G$ .

### Corollary

A group  $G$  has the unique trace property if and only if it has trivial amenable radical.

We now start to show some of results mentioned in the beginning.

### Theorem

Let  $G$  be a group and  $t \in G$ . Then  $\tau(\lambda_t) = 0$  for all tracial states  $\tau$  on  $C_r^*(G)$  if and only if  $t \notin \text{AR}_G$ .

### Corollary

A group  $G$  has the unique trace property if and only if it has trivial amenable radical.

## $C^*$ -simplicity using boundary actions

We now start to show some of results mentioned in the beginning.

### Theorem

Let  $G$  be a group and  $t \in G$ . Then  $\tau(\lambda_t) = 0$  for all tracial states  $\tau$  on  $C_r^*(G)$  if and only if  $t \notin \text{AR}_G$ .

### Corollary

A group  $G$  has the unique trace property if and only if it has trivial amenable radical.

The proof of this relies on the fact that given a normal amenable subgroup  $N \trianglelefteq G$ , there is a  $*$ -homomorphism  $\pi: C_r^*(G) \rightarrow C_r^*(G/N)$  such that

$$\pi(\lambda_G(t)) = \lambda_{G/N}([t]), \quad \text{for all } t \in G.$$

## $C^*$ -simplicity using boundary actions

We now start to show some of results mentioned in the beginning.

### Theorem

Let  $G$  be a group and  $t \in G$ . Then  $\tau(\lambda_t) = 0$  for all tracial states  $\tau$  on  $C_r^*(G)$  if and only if  $t \notin \text{AR}_G$ .

### Corollary

A group  $G$  has the unique trace property if and only if it has trivial amenable radical.

The proof of this relies on the fact that given a normal amenable subgroup  $N \trianglelefteq G$ , there is a  $*$ -homomorphism  $\pi: C_r^*(G) \rightarrow C_r^*(G/N)$  such that

$$\pi(\lambda_G(t)) = \lambda_{G/N}([t]), \quad \text{for all } t \in G.$$

Using this, one creates a trace  $\tau$  on  $C_r^*(G)$  which is 0 on  $\lambda_t$  for all  $t \notin \text{AR}_G$ .

We now start to show some of results mentioned in the beginning.

### Theorem

Let  $G$  be a group and  $t \in G$ . Then  $\tau(\lambda_t) = 0$  for all tracial states  $\tau$  on  $C_r^*(G)$  if and only if  $t \notin \text{AR}_G$ .

### Corollary

A group  $G$  has the unique trace property if and only if it has trivial amenable radical.



## $C^*$ -simplicity using boundary actions

We now start to show some of results mentioned in the beginning.

### Theorem

Let  $G$  be a group and  $t \in G$ . Then  $\tau(\lambda_t) = 0$  for all tracial states  $\tau$  on  $C_r^*(G)$  if and only if  $t \notin \text{AR}_G$ .

### Corollary

A group  $G$  has the unique trace property if and only if it has trivial amenable radical.

### Theorem

Let  $G$  be a group and  $t \in G$ . Then  $t \notin \text{AR}_G$  if and only if

$$0 \in \overline{\text{conv}} \{ \lambda_{sts^{-1}} : s \in G \}.$$

Proposition [Matthew Kennedy, Emmanuel Breuillard, Merhdad Kalatar and Narutaka Ozawa, 2014]

A discrete group  $G$  is  $C^*$ -simple if and only if there is a compact  $G$ -boundary  $X$  such that the action is free.

Proposition [Matthew Kennedy, Emmanuel Breuillard, Merhdad Kalatar and Narutaka Ozawa, 2014]

A discrete group  $G$  is  $C^*$ -simple if and only if there is a compact  $G$ -boundary  $X$  such that the action is free.

What is this? Why is this interesting?

## $C^*$ -simplicity and uniqueness of trace using boundary actions

Using this, we may prove the following:

Using this, we may prove the following:

## Theorem

For a group  $G$  TFAE:

Using this, we may prove the following:

## Theorem

For a group  $G$  TFAE:

- 1  $C_r^*(G)$  is simple,

# $C^*$ -simplicity and uniqueness of trace using boundary actions

Using this, we may prove the following:

## Theorem

For a group  $G$  TFAE:

- ❶  $C_r^*(G)$  is simple,
- ❷  $\tau_0 \in \overline{\{s.\varphi : s \in G\}}^{w^*}$  for all states  $\varphi$  on  $C_r^*(G)$ ,
- ❸  $\tau_0 \in \overline{\text{conv}}^{w^*} \{s.\varphi : s \in G\}$  for all states  $\varphi$  on  $C_r^*(G)$ ,
- ❹  $\omega(1)\tau_0 \in \overline{\text{conv}} \{s.\omega : s \in G\}$  for all  $\omega \in (C_r^*(G))^*$ ,
- ❺ For all  $t_1, t_2, \dots, t_m \in G \setminus \{e\}$ , we have

$$0 \in \overline{\text{conv}}^{w^*} \{ \lambda_s (\lambda_{t_1} + \lambda_{t_2} + \dots + \lambda_{t_m}) \lambda_s^* : s \in G \},$$

- ❻ For all  $t_1, t_2, \dots, t_m \in G \setminus \{e\}$  and  $\varepsilon > 0$ , there exists  $s_1, s_2, \dots, s_n \in G$  such that

$$\left\| \sum_{k=1}^n \frac{1}{n} \lambda_{s_k t_j s_k^{-1}} \right\| < \varepsilon,$$

for all  $1 \leq j \leq m$ .

To summarize everything, we have



To summarize everything, we have

## Theorem

Let  $G$  be a discrete group and  $t \in G$ . TFAE:

- ①  $t \notin \text{AR}_G$ ;
- ② there is a boundary action  $G \curvearrowright X$  such that  $t$  acts non-trivially on  $X$ ;
- ③  $\lambda_t \in \ker(\tau)$  for all tracial states  $\tau$  on  $C_r^*(G)$ ;
- ④  $0 \in \overline{\text{conv}} \{ \lambda_{sts^{-1}} : s \in G \}$ .

To summarize everything, we have

## Theorem

Let  $G$  be a discrete group and  $t \in G$ . TFAE:

- ❶  $t \notin \text{AR}_G$ ;
- ❷ there is a boundary action  $G \curvearrowright X$  such that  $t$  acts non-trivially on  $X$ ;
- ❸  $\lambda_t \in \ker(\tau)$  for all tracial states  $\tau$  on  $C_r^*(G)$ ;
- ❹  $0 \in \overline{\text{conv}} \{ \lambda_{sts^{-1}} : s \in G \}$ .

## Theorem

Let  $G$  be a discrete group. TFAE:

- ❶  $C_r^*(G)$  has unique tracial state;
- ❷  $G$  admits a faithful boundary action;
- ❸  $\text{AR}_G = \{e\}$ ;
- ❹ for all  $t \in G \setminus \{e\}$  and all  $\varepsilon > 0$ , there is  $s_1, s_2, \dots, s_n \in G$  such that

$$\left\| \sum_{j=1}^n \frac{1}{n} \lambda_{s_k t s_k^{-1}} \right\| < \varepsilon.$$

To summarize everything, we have

To summarize everything, we have

## Theorem

Let  $G$  be a discrete group. TFAE:

- ❶  $C_r^*(G)$  is simple;
- ❷  $G$  admits a free boundary action;
- ❸  $\tau_0 \in \overline{\{s.\varphi : g \in G\}}^{w^*}$ , for all states  $\varphi$  on  $C_r^*(G)$ ;
- ❹  $\tau_0 \in \overline{\text{conv}}^{w^*} \{s.\varphi : s \in G\}$ , for all states  $\varphi$  on  $C_r^*(G)$ ;
- ❺ for all  $t_1, t_2, \dots, t_m \in G \setminus \{e\}$  and all  $\varepsilon > 0$ , there exist  $s_1, s_2, \dots, s_n \in G$  such that

$$\left\| \frac{1}{n} \sum_{j=1}^n \lambda_{s_j t_k s_j^{-1}} \right\| < \varepsilon,$$

for all  $k = 1, 2, \dots, m$ ;

- ❻  $C_r^*(G)$  has the Dixmier property.

From these we have

From these we have

### Theorem

$\text{AR}_G = \{e_G\} \iff G \text{ has unique trace property} \iff$  for all  $t \in G \setminus \{e\}$  and all  $\varepsilon > 0$ , there is  $s_1, s_2, \dots, s_n \in G$  such that

$$\left\| \sum_{j=1}^n \frac{1}{n} \lambda_{s_k t s_k^{-1}} \right\| < \varepsilon.$$

From these we have

### Theorem

$\text{AR}_G = \{e_G\} \iff G$  has unique trace property  $\iff$  for all  $t \in G \setminus \{e\}$  and all  $\varepsilon > 0$ , there is  $s_1, s_2, \dots, s_n \in G$  such that

$$\left\| \sum_{j=1}^n \frac{1}{n} \lambda_{s_k t s_k^{-1}} \right\| < \varepsilon.$$

### Theorem

$G$  is  $C^*$ -simple  $\iff$  for all  $t_1, t_2, \dots, t_m \in G \setminus \{e\}$  and all  $\varepsilon > 0$ , there exist  $s_1, s_2, \dots, s_n \in G$  such that

$$\left\| \frac{1}{n} \sum_{j=1}^n \lambda_{s_j t_k s_j^{-1}} \right\| < \varepsilon,$$

for all  $k = 1, 2, \dots, m$

From these we have

### Theorem

$\text{AR}_G = \{e_G\} \iff G$  has unique trace property  $\iff$  for all  $t \in G \setminus \{e\}$  and all  $\varepsilon > 0$ , there is  $s_1, s_2, \dots, s_n \in G$  such that

$$\left\| \sum_{j=1}^n \frac{1}{n} \lambda_{s_k t s_k^{-1}} \right\| < \varepsilon.$$

### Theorem

$G$  is  $C^*$ -simple  $\iff$  for all  $t_1, t_2, \dots, t_m \in G \setminus \{e\}$  and all  $\varepsilon > 0$ , there exist  $s_1, s_2, \dots, s_n \in G$  such that

$$\left\| \frac{1}{n} \sum_{j=1}^n \lambda_{s_j t_k s_j^{-1}} \right\| < \varepsilon,$$

for all  $k = 1, 2, \dots, m$

Thus

$C_r^*(G)$  simple  $\implies C_r^*(G)$  unique trace  $\iff$  trivial amenable radical of  $G$ .



The end.  
thanks for listening