FEAR

PHOBIA

ANXIETY

DEPRESSION

CHRONIC PAIN

UNDERSTANDING PATIENT'S PSYCHOLOGY

ANXIETY IS AN EMOTION WITH NEGATIVE IMPACTS THAT MAY HINDER THE ABILITY TO MAKE DECISIONS AND PERFORM DAILY TASKS.

ANXIETY IS THE ANTICIPATION OF FUTURE THREAT

WHAT IS FEAR ??

HOW THE FEAR IS DIFFERENT FROM ANXIETY ?

PANIC ATTACKS

- A panic attack is a sudden, intense episode of fear or discomfort that peaks within minutes, often accompanied by physical and cognitive symptoms. Panic attacks can occur unexpectedly (un-cued) or be triggered by specific situations (cued), (APA,2022).
- Panic attacks can occur in the context of any situation of anxiety or anxiety disorder as well as other mental disorders. (Depression, PTSD, OCD etc.) and some medical conditions (e.g., cardiac, respiratory, vestibular, gastrointestinal).
- A panic attack is characterized by at least four (or more) of the following symptoms that develop abruptly and peak within minutes: (DSM, 2022)

SYMPTOMS

Palpitations, pounding heart, or accelerated heart rate.	Sweating.	Trembling or shaking.
Sensations of shortness of breath or smothering.	Feelings of choking.	Chest pain or discomfort.
Nausea or abdominal distress.	Feeling dizzy, unsteady, light-headed, or faint.	Chilis or heat sensations.
Derealization (feelings of unreality) or depersonalization (being detached from oneself).	Fear of losing control or “going crazy.”	Fear of dying.

MAJOR DEPRESSIVE DISORDER

How that influence in health care ?

MDD



- Major Depressive Disorder (MDD) is a serious mental health condition characterized by persistent sadness, loss of interest in activities, and a range of cognitive and physical symptoms that impair daily functioning. It is one of the leading causes of disability worldwide (WHO, 2023).
- According to the DSM-5-TR (APA, 2022), MDD is diagnosed when five (or more) symptoms are present for at least two weeks, causing significant distress or impairment. At least one of the symptoms must be (1) depressed mood or (2) loss of interest or pleasure (anhedonia).
- Prevalence: About 5% of the global population suffers from MDD, with a lifetime prevalence of 16-20% (Kessler et al., 2003; WHO, 2023).

SYMPTOMS

A. Five (or more) of the following symptoms have been present during the same 2-week period and represent a change from previous functioning: at least one of the symptoms is either (1) depressed mood or (2) loss of interest or pleasure.

Note: Do not include symptoms that are clearly attributable to another medical condition.

1. Depressed mood most of the day, nearly every day, as indicated by either subjective report (e.g., feels sad, empty, hopeless) or observation made by others (e.g., appears tearful). (Note: In children and adolescents, can be irritable mood.)

SYMPTOMS

2. Markedly diminished interest or pleasure in all, or almost all, activities most of the day, nearly every day (as indicated by either subjective account or observation).

3. Significant weight loss when not dieting or weight gain (e.g., a change of more than 5% of body weight in a month), or decrease or increase in appetite nearly every day.

(Note: In children, consider failure to make expected weight gain).

4. Insomnia or hypersomnia nearly every day.

5. Psychomotor agitation or retardation nearly every day (observable by others, not merely subjective feelings of restlessness or being slowed down).

OBSESSIVE COMPULSIVE DISORDER

- How that influence in health care ?

OCD

- OCD is characterized by the presence of obsessions and/or compulsions.
- *Obsessions* are recurrent and persistent thoughts, urges, or images that are experienced as intrusive and unwanted.
- *Compulsions* are repetitive behaviors or mental acts that an individual feels driven to perform in response to an obsession or according to rules that must be applied rigidly.DSM-5-TR (APA, 2022).

TYPES OF OBSESSION

- Contamination
- symmetry, incompleteness, and ordering
- harm (to self and others), shame, and guilt (this is sometimes categorized under taboo thoughts)
- taboo thoughts around violence, religion, and sexual behaviors

TYPES OF OBSESSION



SYMPTOMS

A. Presence of obsessions, compulsions, or both:
Obsessions are defined by (1) and (2):

- 1. Recurrent and persistent thoughts urges, or images that are experienced, at some time during the disturbance, as intrusive and unwanted, and that in most individuals cause marked anxiety or distress.

- 2. The individual attempts to ignore or suppress such thoughts, urges, or images, or to neutralize them with some other thought or action (i.e., by performing a compulsion).

B Compulsions are defined by (1) and (2):

- 1. Repetitive behaviors (e.g., hand washing, ordering, checking) or mental acts (e.g., praying, counting, repeating words silently) that the individual feels driven to perform in response to an obsession or according to rules that must be applied rigidly.

- 2. The behaviors or mental acts are aimed at preventing or reducing anxiety or distress, or preventing some dreaded event or situation; however, these behaviors or mental acts are not connected in a realistic way with what they are designed to neutralize or prevent, or are

SYMPTOMS

Note: Young children may not be able to articulate the aims of these behaviors or mental acts.

B. The obsessions or compulsions are time-consuming (e.g., take more than 1 hour per day) or cause clinically significant distress or impairment in social, occupational, or other important areas of functioning.

Compulsions are defined by (1) and (2):

- 1. Repetitive behaviors (e.g., hand washing, ordering, checking) or mental acts (e.g., praying, counting, repeating words silently) that the individual feels driven to perform in response to an obsession or according to rules that must be applied rigidly.

- 2. The behaviors or mental acts are aimed at preventing or reducing anxiety or distress, or preventing some dreaded event or situation; however, these behaviors or mental acts are not connected in a realistic way with what they are designed to neutralize or prevent, or are clearly excessive.

CHRONIC PAIN & IT'S PSYCHOLOGICAL EFFECTS

- How that influence in health care ?

CHRONIC PAIN

- Pain that persists or recurs for more than 3 months
- Pain is an unpleasant sensory and emotional experience associated with, or resembling that associated with, actual or potential tissue damage. (WHO, 2022).
- Chronic pain, one of the most common reasons adults seek medical care, has been linked to activity limitations, dependence on opioids, anxiety and depression, and reduced quality of life
- If your chronic pain makes it difficult to contribute at work or home, you might experience low self-worth or even shame



CHRONIC PAIN

- Chronic pain and mental health disorders often occur together.
- People living with chronic pain are at heightened risk for mental health problems, including depression, anxiety, and substance use disorders.
- Pain can also be a common symptom among people with an anxiety disorder, particularly generalized anxiety disorder, according to the Anxiety and Depression Association of America (ADAA).
- Anxiety, depression, and other mood disorders commonly occur at the same time as chronic pain from conditions like fibromyalgia, back problems, migraines and arthritis.
- According to DSM-5-TR Mental wellbeing always effects the physical wellbeing and vice versa

THANK YOU