Bcqs
facial nerve trigeminal nerve hypothalamus

- 1) a patient has facial deviation along with loss of sensation on anterior 2/3rd of tongue the lesion lies in
- A) facial canal
- B) internal aunditory meatus
- C) facial nerve nucleus
- D) stylomastoid foramen

2) Parasympathetic nerves are

- A) 3,7,9,10
- B) 4,5,7,8
- C) 9,10,11,12
- D) 2,4,6,8



- 3) from which foramen facial nerve enters temporal lobe
- A) internal acoustic meatus
- B) styloid foramen
- C) foramen ovale
- D)foramen spinosum

- 4) regarding intracranial part of facial nerve what is true
- A) give rise to greater petrosal nerve
- B) nerve to submandibular gland
- C) nerve to tensor tympani
- D) doesn’t pass through posterior cranial fossa



- 5) all are parasympathetic except
- A) 3
- B) 5
- C) 7
- D) 10

- 6) which among the following nuclei is not related to facial nerve
- A) main dorsal nucleus
- B) nucleus of tractus solitarius
- C) nucleus abiguis
- D) sensory nucleus

- 7)a 65 yr old patient presented with hemiplegia and facial paralysis on examination lower part of face is more severly affected with loss of voluntary movements of facial muscles what could be the site of lesion
- A) facial canal

B) petrous temporal

c) Supranuclear

d) Cerebropontine angle

- 8) facial nerve supplies
- A) stapedius muscle
- B) muscle of facial expressions except orbicularis occuli
- C) sensation from face
- D) none of the above

- 9) upper motor neuron lesion of facial nerve
- A) forehead spared
- B) deviation of angle of mouth towards side of lesion
- C) one eye spared
- D) none of the above

10) trigeminal nerve

- A) parasympathetic
- B) supplies temporalis muscle
- C) it has 4 main division
- D) all are incorrect

- 11) patient with injury to left 8th cervial segment of spinal cord will not show which of the following sign
- A) decrease sense of position
- B) vibration below lesion on same side
- C) extensor planter on left side
- D) power of muscle below the lesion on same side
- E) decrease sense of pain and temperature below the lesion on same side

12)Which of the following structures travel through the substance of the parotid gland?

A) The maxillary artery

b) The maxillary artery and retromandibular vein

c) The maxillary artery, retromandibular vein and facial artery

d) The maxillary artery, retromandibular vein, facial artery and buccal branch of the mandibular nerve

13) a patient with damage to their cervical sympathetic chain will show the following signs and symptoms.

a) Pupillary constriction

b) Pupillary constriction and vasodilation of facial vessels

c) Pupillary constriction, vasodilation of facial vessels and increased facial sweating

d) Pupillary constriction, vasodilation of facial vessels, increased facial sweating and decreased lacrimation

14) Lesion causing compression of the facial nerve at the stylomastoid foramen will cause ipsilateral

a) paralysis of the facial muscles.

B) paralysis of the facial muscles and loss of taste.

C) paralysis of the facial muscles, loss of taste and lacrimation.

- D) paralysis of the facial muscles, loss of taste, lacrimation and decreased salivation

15) Which of these branches of the trigeminal nerve contain somatic motor processes?

A) The supraorbital nerve

b) The infraorbital nerve

c) The mental nerve

d) None of the above

16) On which surfaces of the teeth is dental plaque most likely to accumulate in the mouth of a patient with poor oral hygiene?

A) The palatal surfaces of the upper molars and lingual surfaces of the lower incisors.

B) The buccal surfaces of the upper molars and lingual surfaces of the lower incisors.

C) The palatal surfaces of the upper molars and labial surfaces of the lower incisors.

- D) The buccal surfaces of the upper molars and labial surfaces of the lower incisors

17) The parotid duct runs

a) deep to masseter and enters the mouth opposite the upper second molar.

B) superficial to masseter and enters the mouth opposite the upper second molar.

C) deep to masseter and enters the mouth opposite the upper second premolar.

D) superficial to masseter and enters the mouth opposite the upper second premolar

18) What connects hypothalamus to the pituitary?

A) Infundibulam

b) Corpus callosum

c) Anterior

- d) Cerebral cortex

19) Hypothalamus is a part of _____

a) Midbrain

b) Forebrain

c) Hind brain

- d) Future brain

- 20) Which of the following events could be a result of damage to hypothalamus portal system?
  a) Decreased secretion of ADH
  b) Decreased secretion oxytocin
  c) Decreased secretion TSH
  d) Decreased secretion PTH



- 21) the following statement concern the hyp
- A) it lies below the thalamus in the tectum c
- B) it is not related to limbic system
- C) the nuclei of hypothalamus are divided b imaginary plane formed by the columns of f mamilothalamic tract into medial and lateral
- D) the suprachaismatic nucleus does not re nerve fibers from the retina
- E)the lateral boundary of hypothalamus is f external capsule

- 22)the following statemts concern the hypothalamo hypophyseal tract

- A) oxytocin inhibts the contraction of smooth muscle of uterus

- B) nerve cells of supra optic and para ventricular nuclei produce hormones vasopressin and oxytocin

- C) hormones travel in lymph vessels with protein carries neurophysins

- D) vasopressin stimulates the proximal convulated tubules of kidney causing increased absorbtion of water from urine

- E) hormones are abosrbed into blood stream in cappilaries of the anterior lobe of hypophysis

- 23) explain what is meant by the terms hypothalamic hpophyseal tract and hypophyseal portal system

- 24) enlist the main functions of hormones
- Supra optic
- Para ventricular
- Preoptic and anterior nucleus
- Lateral hypothalamic nuclei
- Supra chiasmatic