DR. ISHRAT -UL -EBAD KHAN INSTITUTE OF ORAL HEALTH SCIENCES
DOW UNIVERSITY OF HEALTH SCIENCES
DEPARMENT OF ANATOMY (3rd BCQ- CLASS TEST)

Name: M. Ahmed Malick
Roll no: 90

Max Marks: 26
Min Marks: 13

Date: 21-09-2024

Each BCQ carries one (01) mark. There will be no negative marking.

Q1) Which of the following structures does not pass through foramen magnum?
a) Anterior spinal artery.
b) Posterior spinal arteries.
c) Spinal accessory nerves.
d) Vagus nerves.
e) Vertebral arteries.

Q2) Maxillary nerve exits the cranial cavity through which opening?
a) Foramen lacerum.
b) Foramen rotundum.
c) Foramen spinosum.
d) Inferior orbital fissure.
e) Internal acoustic meatus.

Q3) Carotid canal is present in the:
a) Basilar part of occipital bone.
b) Greater wing of sphenoid bone.
c) Petrous part of temporal bone.
d) Pterygoid fossa of sphenoid bone.
e) Squamous part of temporal bone.

Q4) A person is brought to casualty with bleeding and discharge of CSF from the nose, followed by a banging injury to his forehead. His C.T scan reveals a fracture in his anterior cranial fossa most likely involving which of the following bones?
a) Ethmoid.
b) Frontal.
c) Lacrimal.
d) Sphenoid.
e) Temporal.

Q5) Regarding mandible, mark the statement which is false.
a) Buccinator arises from oblique line on lateral surface of body of mandible.
b) Coronoid process is usually larger than head in an infant's mandible.
c) Geniohyoid arises from lower genial tubercle/mental spine.
d) Sphenomandibular ligament is attached to lingula.
e) Temporalis is inserted into the pterygoid fovea.

Q6) Emissary veins, mainly responsible for transmitting any infection from scalp to intracranial venous sinuses, are present in which layer of the scalp?

a) Galea aponeurotica.
b) Loose areolar tissue.
c) Pericranium.
d) Skin.
e) Superficial fascia.

Q7) The collection of blood deep to which layer of the scalp causes cephalhaematoma?
a) Connective tissue/superficial fascia.
b) Epicranial aponeurosis.
c) Loose areolar tissue.
d) Pericranium.
e) Skin.

Q8) Regarding buccinator muscle, choose the odd statement out:
a) It is a sphincter of oral fissure.
b) It is a whistling muscle.
c) It is pierced by the parotid duct.
d) It is supplied by the mandibular nerve.
e) It originates from maxilla, mandible and pterygomandibular raphe.

Q9) Which muscle of face protrudes and everts the lower lip making an expression of "doubt"?
a) Depressor anguli oris.
b) Depressor labii inferioris.
c) Mentalis.
d) Zygomaticus major.
e) Zygomaticus minor.

Q10) Which of the following muscles of face dilates the lacrimal sac?
a) Corrugator supercilia.
b) Levator palpebrae superioris.
c) Nasalis.
d) Orbicularis oculi.
e) Procerus.

Q11) Branches of cervical plexus give cutaneous (sensory) supply to which areas of face?
a) Anterior part of temple.
b) Lower lip and chin.
c) Side of head.
d) Skin over the angle of mandible and parotid gland.
e) Upper 2/3rd of lateral surface of auricle.

Q12) Regarding venous drainage of face, mark the statement which is true?
a) Common facial vein is formed by the union of anterior division of retromandibular vein & facial vein.
b) Facial vein drains into external jugular vein.
c) Facial vein has no direct connection with pterygoid venous plexus.
d) Pterygoid venous plexus drains into superficial temporal vein.
e) Retromandibular vein is formed by the union of supraorbital & supratrochlear veins.

Q13) Regarding nasal cavity, mark the statement which is false:
a) Bony part of the nasal septum is formed by the sphenoid bone.
b) It has a communication with the pterygopalatine fossa.
c) Its floor is formed by the maxilla and palatine bone.
d) Its lateral walls are featured by the conchae.
e) Its upper 1/3rd is lined by the olfactory mucosa.

Q14) Which of the following is considered as the chief artery of epistaxis?
a) Anterior ethmoidal.
b) Greater palatine.
c) Sphenoplalatine.
d) Superior labial.
e) Transverse facial.

Q15) The superior meatus receives the opening of the:
a) Anterior ethmoidal sinus.
b) Maxillary sinus.
c) Middle ethmoidal sinus.
d) Nasolacrimal duct.
e) Posterior ethmoidal sinus.

Q16) The pterygopalatine fossa:
a) Communicates anteriorly with the orbit through superior orbital fissure.
b) Contains infraorbital and posterior superior alveolar branches of maxillary nerve.
c) Communicates laterally with the infratemporal fossa via pterygomaxillary fissure.
d) Is occluded inferiorly by body of sphenoid bone.
e) Contains second part of maxillary artery.

Q17) Preganglionic parasympathetic fibers are carried to pterygopalatine ganglion via which branch of facial nerve?
a) Chorda tympani branch.
b) Greater petrosal branch.
c) Nerve to stapedius.
d) Posterior auricular branch.
e) Stylohyoid branch.

Q18) The floor of the bony orbit is formed by which of the following set of bones?
a) Frontal and lesser wing of sphenoid.
b) Frontal, lacrimal and ethmoid.
c) Maxilla and zygomatic.
d) Maxilla, zygomatic, and palatine.
e) Zygomatic and greater wing of sphenoid.

Q19) Which of the following is not a component of the pupillary (light) reflex pathway?
a) Edinger Westphal nucleus.
b) Medial geniculate body.
c) Optic tract.
d) Pretectal nucleus.
e) Ganglion cells in the retina.

Q20) Eye examination of a patient indicates that he has a medially directed/convergent squint. This could be due to damage to the:
a) Abducent nerve.
b) Oculomotor nerve.
c) Ophthalmic division of trigeminal nerve.
d) Optic nerve.
e) Trochlear nerve.

Q21) In primary position, what are the actions of the superior rectus muscle on the eyeball?
a) Abduction and adduction.
b) Depression, abduction, and intortion.
c) Depression, adduction and extortion.
d) Elevation, abduction, and extortion.
e) Elevation, adduction, and intortion.

Q22) Regarding layers/structures of the eye ball, mark the statement which is false:
a) The anterior chamber contains the vitreous body.
b) The blind spot of the retina is devoid of rods and cones.
c) The middle coat contains the choroid, ciliary body and iris.

d) The junction of sclera and cornea is called the limbus.
e) The outer most fibrous coat consists of the sclera and cornea.

Q23) The aqueous humor of the eyeball is normally drained by the:
a) Canal of Schlemm.
b) Central vein of retina.
c) Hyaloid canal.
d) Nasolacrimal duct.
e) Scleral venous plexus.

Q24) Choose the correct pair from the following sets of pharyngeal arches/pouches and thei
a) 1st pharyngeal arch----------stylohyoid muscle.
b) 1st pharyngeal pouch----------tonsillar fossa.
c) 2nd pharyngeal arch----------muscles of mastication.
d) 3rd pharyngeal pouch----------thymus.
e) 4th pharyngeal arch----------stylopharyngeus muscle.

Q25) Which of the following do not arise from frontonasal prominence?
a) Alae of nose.
b) Bridge of nose.
c) Forehead.
d) Lateral nasal prominences.
e) Medial nasal prominences.

Q26) Regarding development of face, choose the odd step out from the following:
a) Appearance of nasal placodes on the frontonasal prominence.
b) Appearance of nasal pits in the placodes.
c) Formation of cheeks by the maxillary prominences.
d) Fusion of the two medial nasal prominences to form philtrum of upper lip.
e) Fusion of the medial and lateral nasal prominences to form the nasolacrimal duct.

----------X----------X----------