Regarding First cervical vertebrae which statement is incorrect

- It is ring shaped
- It has long anterior arch
- Does not have a body and spine
- It forms ellipsoid joint with occipital bone

Lingula gives attachment to

- Stylomandibular ligament
- Lateral pterygoid
- Sphenomandibular ligament
- Mylohoid muscle

Frontal belly of galea aponeurotica is supplied by

- Trigeminal Nerve
- Zygomaticofacial Nerve
- Facial nerve
- Oculomotor nerve

Compression of brain in scalp injury is prevented by

- Connective tissue layer
- Aponeurosis
- Loose areolar tissue
- Pericranium

Which muscle helps in lowering the mandible

- Depressor anguli oris
- Platysma
- Depressor labii inferioris
- Depressor septi

A person suffers a nerve lesion after which his voluntary facial movements are compromised while emotional expressions remain normal, he is suffering from

- Frey's syndrome
- Carotid sinus syndrome
- Bell's palsy
- Medial Medullary syndrome

Digastric is supplied by

- V C. N

- C1 +XII C. N
- IX C. N+ V C. N
- VII C. N+ V C. N

Find the Odd one out in terms of drainage

- Superior thyroid vein
- Common facial vein
- Middle thyroid vein
- Both B and C

Parathormone is secreted by

- Oxyphilic cells
- Chief cells
- Parafollicular cells
- C cells

Lateral Part Of eye drains into

- Mastoid lymph nodes
- Submandibular lymph nodes
- Retro Pharyngeal lymph nodes
- Parotid lymph nodes

Main artery of tonsils

- Ascending palatine artery
- Lesser palatine artery
- Tonsillar artery
- Ascending Pharyngeal artery

Terminal branch of maxillary artery

- Middle meningeal artery
- Temporal artery
- Inferior alveolar artery
- Sphenopalalatine artery

Trunk of trigeminal nerve supplies all except

- Buccal pads
- Meninges
- Medial pterygoid muscle
- Tensor tympani

All are parts of middle ear except

- Stapedius
- Tympanic plexus
- Anvil
- Cristae

Muscle for the abduction of vocal cord

- Aryepiglottic
- Post cricothyroid
- Thyroepiglottic
- Post cricoarytenoid

What is false about lacrimal gland

- It is a j shaped gland
- It is located in lateral part of superior palbabrae
- It's postganglionic fibres are carried by maxillary nerve
- Ciliary ganglion is where preganglionic fibres relay

Which ganglion is known as ganglion of hay fever

- Pterygopalatine
- Otic
- Ciliary
- Submandibular

Maxillary artery is divided by

- Medial pterygoid
- Lateral pterygoid
- Mylohoid
- Buccinator

Transverse facial artery is a branch of

- Angular artery
- Superficial temporal artery
- Greater palatine artery
- Maxillary artery

A bicyclist was hit by a car, after falling down his head hit the foot path, there is no sign of external bleeding but his reports show extradural hemorrhage.which artery must be responsible for the bleed

- Anterior cerebral artery

- Middle cerebral artery
- Temporal artery
- Middle meningeal artery

Hoarseness of voice is a result of lesion of which nerve

- Vagus nerve
- Recurrent laryngeal nerve
- Superior laryngeal nerve
- External laryngeal nerve

Hypoglossal nerve of right side suffers a lesion, outcome will be

- Deviation of the tongue on the opposite side
- Deviation of the tongue on the same side
- Deviation of uvula on the opposite side
- Deviation of uvula on the same side

Secretomotor root of the parotid gland

- Facial nerve
- Maxillary nerve
- Auriculotemporal nerve
- Glossopharyngeal nerve

Bifurcation of common carotid artery takes place at the level of

- Thyroid cartilage
- Cricoid cartilage
- C9
- T4

A patient comes to the clinic and he suffers from medial deviation of eye, which muscle of this patient is paralyzed

- Superior oblique
- Medial rectus
- Lateral rectus
- Both a and b

In children Lumbar puncture is done at the level of

- C3 and C4
- L3 and l4
- L5 and L6
- T12 and L1

Increase amount of amniotic fluid in the last stages of pregnancy is due to

- Renal agenesis
- Cardiac failure
- Esophageal atresia
- Pulmonary disease

About Stylohyoid muscle all are true except

- Is innervated by facial nerve
- Pull hyoid bone upwards
- Draw larynx downwards
- Lies on the upper border of posterior belly of diagastric

Lymph from tongue drains finally into

- Jugulodiagastric lymph nodes
- Jugulomohyoid lymph nodes
- Tonsillar lymph nodes
- Submandibular lymph nodes

Life saving muscle of Tongue helps in

- Protruding the tongue
- Deviation of tongue
- Falling back of tongue
- Narrowing of tongue

Floor of posterior triangle is formed by

- Pretracheal fascia
- Investing layer
- Buccopharyngeal fascia
- Prevertebral fascia

To avoid excessive bleeding thyroidectomy should be done along with

- False capsule
- Isthmus
- True capsule
- Superior thyroid artery

Metopic suture fuses at the age of

- 6 years
- 3 years
- 9 years
- 7 years

Sinus which is formed after birth

- Maxillary sinus
- Anterior ethmoidal sinus
- Frontal Sinus
- Middle ethmoidal sinus

Dangerous area of face includes

- Lower eyelid
- Upper lip region
- Cheek Area
- Both b and c

About vocal Fold, which statement is false

- It is vascular
- It produces sound
- It is below vestibular fold
- Vocal process of arytenoid cartilage is a part
-

Plane perpendicular to coronal plane is

- Sagittal plane
- Transverse plane
- Oblique plane
- Cardinal plane

Complete section Of optic chiasma results in

- Nasal hemianopia
- Bitemporal hemianopia
- Total blindness in one eye
- Total blindness in both the eyes

Efferent gag reflex is lost due to lesiok of

- Vagus nerve
- Hypoglossal nerve
- Abducent nerve
- Glossopharyngeal nerve

Nerve which originates from the psot surface of midbrain is

- Abducent nerve
- Trochlear nerve
- Oculomotor nerve
- Vestibulocochlear nerve

Hassel Corpuscle are characterstic feature of

- Spleen
- Thymus
- Tonsils
- Lymph nodes

In thymus

- Pals is present
- Germinal Centre do not appear normally
- It shrinks with age
- Lymphocytes are present in paracortex

- Palatine tonsils have
- Pleats
- Crypts
- Invaginations
- Both A and B

a. Subscapular triangle
b. Sub occipital triangle
c. Carotid triangle
d. Submental triangle
e. Supraclavicular triangle
f. Occipital triangle
g. Submandibular triangle
h. Mascular triangle

1.After a Gun shot injury a person is unable to rotate his head on the opposite side and bend his head towards the shoulder of the same side. The location of the gun shot would be

2.Common carotid artery is present in

3.Post and anterior belly of diagstric form it's bonudaries

4.Tip of tongue drains into lymph nodes of

5.Thymus is present in

a. Simple squamous epithelium
b. Transitional epithelium
c. Stratified cuboidal epithelium
d. Stratified squamous Keratinized epithelium
e. Simple columnar epithelium
f. Pseudostratified columnar epithelium
g. Neuroepithelium
h. Stratified squamous non Keratinized epithelium

1.Cornea

2.Hard palate

3.Tracheal Epithelium

4.Endothelium