## Questions

### Section One (MCQs A- Type)

### Skull and Cervical Vertebrae

1. In the anterior view of the viscerocranium, which one of the following bones is single?
   a. Nasal.
   b. Frontal.
   c. Lacrimal.
   d. Maxillary.
   e. Zygomatic.
2. What is the strongest bone among the bones of the face?
   a. Nasal.
   b. Frontal.
   c. Maxillary.
   d. Zygomatic.
   e. Mandibular.
3. The superior orbital margin is formed by the ____ bone:
   a. Frontal.
   b. Maxillary.
   c. Zygomatic.
   d. Ethmoidal.
   e. Sphenoidal.
4. Which part of the frontal bone contributes to the floor of the anterior cranial fossa?
   a. Nasal.
   b. Squamous.
   c. Orbital plate.
   d. Maxillary process.
   e. Zygomatic process.
5. Which part of the frontal bone articulates at the Pterion?
   a. Nasal.
   b. Squamous.
   c. Orbital plate.
   d. Maxillary process.
   e. Zygomatic process.
6. The metopic suture of the frontal bone normally disappears at about:
   a. 2 months.
   b. 4 months.
   c. 5 years.
   d. 8 years.
   e. 15 years.
7. In the anterior cranial fossa, the cribriform plate is part of the ___ bone:
   a. Frontal.
   b. Maxillary.
   c. Zygomatic.
   d. Ethmoidal.
   e. Sphenoidal.
8. In the walls of the orbit, the ethmoid bone contributes to the formation of the:
   a. Medial orbital wall.
   b. Inferior orbital wall.
   c. Lateral orbital margin.
   d. Superior orbital margin.
   e. Floor of the orbital cavity.
9. The ___ artery exits the middle cranial fossa through the optic canal:
   a. Buccal.
   b. Pterygoid.
   c. Masseteric.
   d. Ophthalmic.
   e. Deep temporal.
10. The Posterolateral fontanelle is located at the:
    a. Inion.
    b. Pterion.
    c. Bregma.
    d. Lambda.
    e. Asterion.
11. The presence of the fontanelles is due to the delayed ossification of the four angels of the ___ bone:
    a. Frontal.
    b. Parietal.
    c. Occipital.
    d. Sphenoid.
    e. Temporal.
12. The nerve that provides motor supply to the tongue emerges from the cranium through a canal in the ___ bone:
    a. Nasal.
    b. Maxilla.
    c. Parietal.
    d. Lacrimal.
    e. Occipital.
13. What is the foramen located between the occipital bone and petrous part of temporal bone?
    a. Condylar canal.
    b. Jugular foramen.
    c. Mastoid foramen.
    d. Foramen magnum.
    e. Hypoglossal canal.
14. The internal acoustic meatus is a canal within the ____ bone:
    a. Occipital.
    b. Maxillary.
    c. Temporal.
    d. Sphenoid.
    e. Zygomatic.
15. The acoustic labyrinth is situated within the ___ part of temporal bone:
    a. Petrous.
    b. Squamous.
    c. Tympanic.
    d. Styloid process.
    e. Zygomatic process.
16. The superior orbital fissure is a gap within the ___ bone:
    a. Frontal.
    b. Parietal.
    c. Occipital.
    d. Sphenoid.
    e. Temporal.
17. What is the foramen located between the sphenoid bone and petrous part of temporal bone?
    a. Foramen ovale.
    b. Jugular foramen.
    c. Foramen lacerum.
    d. Foramen rotundum.
    e. Foramen spinosum.
18. The internal carotid artery passes across the foramen ____ to the cranium:
    a. Ovale.
    b. Lacerum.
    c. Magnum.
    d. Spinosum.
    e. Rotundum.
19. Foramen spinosum is located in the __ of sphenoidal bone:
    a. Body.
    b. Lesser wing.
    c. Sella turcica.
    d. Greater wing.
    e. Pterygoid plates.
20. What is the nerve that passes through foramen rotundum?
    a. Olfactory.
    b. Maxillary.
    c. Mandibular.
    d. Ophthalmic.
    e. Vestibulocochlear.



Anatomy Q-Bank / Hosam Eldeen Elsadig Gasmalla / ISBN: 978 – 99942 – 4- 142 - 2

Scanned with CamScanner

21. The mandibular division of the trigeminal nerve exits the skull through:
a. Foramen ovale.
b. Foramen lacerum.
c. Foramen spinosum.
d. Foramen rotundum.
e. Superior orbital fissure.

22. Which part of sphenoid bone ossifies by intramembranous ossification?
a. Body.
b. Lesser wing.
c. Sella turcica.
d. Greater wing.
e. Pterygoid plates.

23. Scaphocephaly results from premature closure of the ___ suture:
a. Sagittal.
b. Coronal.
c. Lambdoid.
d. Parietomastoid.
e. Sphenosquamosal.

24. The Axis is an atypical cervical vertebra because of the:
a. Lack of body. Atlas → $C_1$ (neither body nor spine)
b. Presence of longest spine. - $C_7$
c. Presence of the odontoid process. - $C_2$
d. Absence of foramen transversarium ($C_7$)
e. Presence of lateral mass on each side.

25. The absence of a body makes the ___ vertebra atypical:
a. 1st cervical.
b. 2nd cervical.
c. 3rd thoracic.
d. 4th thoracic.
e. 5th lumbar.

26. The foramen transversarium of the _____ cervical vertebra does not transmit the vertebral artery.





a. 1st.
b. 2nd.
c. 4th.
d. 6th.
e. 7th. ($C_1 - C_7$)

27. What is the axis of the Atlantoaxial Joints during their movement?
a. Odontoid process.
b. Occipital condyles.
c. Lateral mass of C1.
d. Lateral mass of C2.
e. Anterior arch of C1.

28. The hyoid bone is connected to the scapula through ___ muscle:
a. Omohyoid.
b. Mylohyoid.
c. Stylohyoid.
d. Geniohyoid.
e. Sternohyoid.

26. The foramen transversarium of the _____ cervical vertebra does not transmit the vertebral artery.

## Neck

29. Which one of the following vessels is located just deep to platysma?
a. External jugular vein.
b. External carotid artery.
c. Common carotid artery.
d. Internal carotid artery.
e. Internal jugular vein.

30. The platysma is innervated by the ____ nerve:
a. Facial
b. Accessory
c. Trigeminal
d. Glossopharyngeal
e. Transverse cervical

31. Which one of the following ligaments is a thickened part of investing layer of the deep cervical fascia?
a. Thyrohyoid.
b. Spurspinous.
c. Cricothyroid.
d. Stylomandibular.
e. Ligamentum flaveum.

32. Sternocleidomastoid muscle is completely surrounded by the:
a. Carotid sheath.
b. Investing layer.
c. Superficial fascia.
d. Pretracheal fascia.
e. Prevertebral fascia.

33. The ligament that suspends the hyoid bone to digastric muscle is derived from _____?
a. Carotid sheath.
b. Superficial fascia.
c. Pretracheal fascia.
d. Prevertebral fascia.
e. Thyrohyoid membrane.

34. Which one of the following ligaments of the neck limits hyperextension?
a. Nuchal ligament.
b. Ligamentum flavum.
c. Interspinous ligaments.
d. Intertransverse ligaments.
e. Anterior longitudinal ligament.

35. What is the layer that overlies the phrenic nerve?
a. Platysma.
b. Superficial fascia.
c. Pretracheal fascia.
d. Prevertebral fascia.
e. Investing layer of deep fascia.

36. Pus in _____ pace of the neck passes as downward as the superior mediastinum.
a. Submental.
b. Prevertebral.
c. Submandibular.
d. Parapharyngeal.
e. Reteropharyngeal.

37. Moving the head so that the face looks upwards and to the left is a movement produced by the:
a. Platysma.
b. Left trapezius only.
c. Right geniohyoid only.
d. Right sternocleidomastoid.
e. Posterior belly of digastric.

38. The neck is divided Into anterior and posterior triangles using ____ muscle as land mark:
a. Trapezius.
b. Sternohyoid.
c. Scalenus anterior
d. Sternocleidomastoid.
e. Anterior belly of digastric.

Anatomy Q-Bank / Hosam Eldern Elsadig Gasmalla / ISBN: 978 - 99942 - 4- 142 - 2

Scanned with CamScanner

39. The _____ triangle of the neck lies between the anterior bellies of digastric muscles.
a. Occipital.
b. Posterior.
c. Muscular.
d. Submental.
e. Submandibular.

40. The posterior triangle is divided into two triangles by the ___ muscle:
a. Digastric.
b. Trapezius.
c. Omohyoid.
d. Mylohyoid.
e. Sternohyoid.

41. Stab wound directed into submandibular triangle may lead to:
a. Ptosis of the eye.
b. Drop of the shoulder.
c. Deviation of the mouth.
d. Deviation of the tongue.
e. Tilting of the head to the same side.

42. Roots of cervical plexus pass anterior to _____ muscle.
a. Scalenus medius.
b. Scalenus anterior.
c. Scalenus posterior.
d. Semispinalis capitis.
e. Sternocleidomastoid.

43. The nerve point of the neck is at the middle of the posterior border of the _____ muscle.
a. Trapezius.
b. Omohyoid.
c. Sternohyoid.
d. Sternothyroid.
e. Sternocleidomastoid.

44. What is the cutaneous branch of the cervical plexus that crosses over the sternocleidomastoid muscle?
a. Phrenic.
b. Great auricular.
c. Supraclavicular.
d. Lesser occipital.
e. Transverse cervical.

45. The branch from the hypoglossal nerve that supplies thyrohyoid muscle is actually derived from _____
a. C1.
b. C2.
c. C3.
d. C4.
e. C5.

46. Inferior root of Ansa hypoglossi is derived from:
a. C1.
b. C2 and C3.
c. C4 and C5.
d. C6 and C7.
e. C8.

47. The inferior root of ansa cervicalis spirals around the:
a. Subclavian artery.
b. Internal jugular vein.
c. External jugular vein.
d. Internal carotid artery.
e. External carotid artery.

48. Which one of the following branches from the cervical plexus is mixed (motor and sensory)?
a. Phrenic.
b. Great auricular.
c. Supraclavicular.
d. Lesser occipital.
e. Transverse cervical.

cutaneous



49. The _____ nerve from the cervical plexus supplies the skin over the acromion process of the scapula:
a. Phrenic.
b. Supraclavicular.
c. Great auricular.
d. Lesser occipital.
e. Transverse cervical.

50. A patient had lost cutaneous sensation over the angle of the mandible, which nerve is most likely damaged?
a. Mandibular.
b. Ophthalmic.
c. Great auricular.
d. Auriculotemporal.
e. Supra clavicular.

51. The thyroid gland moves with swallowing because it is tight by the ___ layer of deep cervical fascia:
a. Parotid.
b. Carotid.
c. Investing.
d. Pretracheal. (ligament of berry)
e. Prevertebral.

52. The superior thyroid artery is closely related to the ___ nerve:
a. Lingual.
b. Hypoglossal.
c. Internal laryngeal.
d. Glossopharyngeal.
e. Recurrent laryngeal.

53. Miosis in Horner's syndrome is due to lack of innervation from the _____ ganglion:
a. Otic.
b. Ciliary.
c. Cervical.
d. Submandibular.
e. Pterygopalatine.

SOT

54. Ptosis in Horner's syndrome is due to paralysis of the smooth muscle part of the _____ muscle:
a. Superior rectus.
b. Superior oblique.
c. Depressor anguli oris.
d. Levator palpebrae superioris.
e. Levator labii superioris alaeque nasii.

55. Damage to _____ ganglion will cause xerophthalmia:
a. Otic.
b. Ciliary.
c. Stellate.
d. Submandibular.
e. Pterygopalatine.

Ciliary
External

4 sensory cervical pl
great au
Lesser o
Transverse cervical
Supraclavicular
Phrenic → mixed

Scanned with CamScanner

## Face and SCALP

56. Cutaneous innervation of the forehead is provided by the ____ nerve:
    a. Facial.
    b. Infraorbital.
    c. External nasal.
    d. Supratrochlear.
    e. Zygomaticofacial.
57. Loss of sensation of skin over the tip of the nose is due to injury of _____ nerve.
    a. Facial
    b. Olfactory.
    c. Maxillary.
    d. Mandibular.
    e. Ophthalmic.
58. The cutaneous innervation of the anterior part of the SCALP is derived from branches of the _____ nerve:
    a. Facial.
    b. Maxillary.
    c. Ophthalmic.
    d. Great auricular.
    e. Lesser occipital.
59. Among the nerves of the face, which one of the following is a direct continuation of the maxillary nerve?
    a. Lacrimal
    b. Infraorbital.
    c. Supraorbital.
    d. Supratrochlear.
    e. Infratrochlear.
60. The ____ muscle compresses cheek against molar teeth:
    a. Risorius.
    b. Mentalis.
    c. Buccinator.
    d. Orbicularis oris.
    e. Zygomaticus major.
61. The _____ muscle draws a smile on the face:
    a. Risorius.
    b. Levator anguli oris.
    c. Zygomaticus minor.
    d. Corrugator supercilii.
    e. Levator labii superioris.



62. The _____ muscle helps to produce the expression of surprise on the face:
    a. Platysma.
    b. Buccinator.
    c. Occipitofrontalis.
    d. Orbicularis oculi.
    e. Corrugator supercilii.
63. Parasympathetic innervation to the parotid gland is provided through the _____ ganglion:
    a. Otic.
    b. Ciliary.
    c. Geniculate.
    d. Submandibular.
    e. Pterygopalatine.
64. Sensory innervation of the parotid sheath is carried by ___ nerve:
    a. Facial.
    b. Vidian
    c. Maxillary.
    d. Great auricular.
    e. Glossopharyngeal
65. Sensory innervation of the parotid gland is carried by _____ nerve:
    a. Vagus.
    b. Buccal.
    c. Great auricular.
    d. Auriculotemporal. (Post-ganglionic)
    e. Glossopharyngeal.
66. The preganglionic fibers to the otic ganglion reach it through the _____ nerve:
    a. Vidian.
    b. Deep petrosal.
    c. Lesser petrosal. (Glossopharyngeal)
    d. Greater petrosal.
    e. Lesser occipital.

## Temporal, infratemporal and pterygopalatine fossae

67. The _____ muscle retracts the mandible:
    a. Masseter.
    b. Temporalis.
    c. Buccinator.
    d. Medial pterygoid
    e. Lateral pterygoid.
68. Medial pterygoid muscle is inserted into the medial surface of the:
    a. Zygomatic bone.
    b. Medial pterygoid plate.
    c. Lateral pterygoid plate.
    d. Ramus of the mandible.
    e. Coronoid process of the mandible.
69. Which one of the following muscles of mastication opens the mouth?
    a. Masseter.
    b. Temporalis.
    c. Buccinator.
    d. Medial pterygoid
    e. Lateral pterygoid.
70. Masseter muscle is inserted into the _____ of the mandible:
    a. Neck.
    b. Angle.
    c. Mylohyoid line.
    d. Coronoid process.
    e. Medial surface of ramus.
71. Which one of the following branches of the mandibular nerve supplies the anterior belly of Digastric muscle:
    a. Buccal.
    b. Masseteric.
    c. Deep temporal.
    d. Nerve to Mylohyoid.
    e. Nerve to Lateral pterygoid.
72. The sensory supply of the chin is derived from the _____ nerve:
    a. Buccal.
    b. Lingual.
    c. Meningeal.
    d. Inferior alveolar.
    e. Auriculotemporal.



73. The pterygopalatine fossa is bounded superiorly by the:
    a. Zygomatic arch.
    b. Greater wing of the sphenoid.
    c. Posterior aspect of the maxilla
    d. Pterygoid process of the sphenoid.
    e. Perpendicular plate of the palatine bone.
74. The maxillary nerve enters the pterygopalatine fossa through ____:
    a. Foramen ovale.
    b. Foramen rotundum.
    c. Inferior orbital fissure.
    d. Pterygomaxillary fissure.
    e. Sphenopalatine foramen.
75. The maxillary nerve is connected to the Ophthalmic nerve through the ___ nerve:
    a. Ganglionic.
    b. Infraorbital.
    c. Inferior Alveolar.
    d. Zygomatico-temporal.
    e. Posterior Superior Alveolar.
76. The preganglionic parasympathetic root of the Pterygopalatine ganglion is the _____ nerve:
    a. Deep petrosal.
    b. Lesser occipital.
    c. Greater petrosal.
    d. Greater occipital.
    e. Superficial petrosal.

(Pterygopalatine fossa)
sup. Greater wing of sph.
ant. Maxilla (body of Maxilla)
Lat. Mandible (Ramus)
Med. Sphenoid (Pterygoid lateral Plate)
inferiorly. open hota

Scanned with CamScanner

## Oral Cavity and palate

77. Which one of the following muscles forms the diaphragm of the oral cavity?
    a. Digastric.
    b. Mylohyoid.
    c. Genohyoid.
    d. Stylohyoid.
    e. Genoglossus.

78. Which one of the following structures is located inferior to mylohyoid muscle?
    a. Facial artery.
    b. Lingual nerve.
    c. Sublingual gland.
    d. Geniohyoid muscle.
    e. Submandibular duct.

79. Regarding hypoglossal nerve injury (at one side), the deviation of the tongue is produced mainly due to the paralysis of _____ muscle:
    a. Stylohyoid.
    b. Hyoglossus.
    c. Mylohyoid.
    d. Genioglossus.
    e. Palatoglossus.

80. Which one of the following muscles of the tongue narrows it?
    a. Hyoglossus.
    b. Genioglossus.
    c. Palatoglossus.
    d. Intrinsic vertical.
    e. Intrinsic Transverse.

81. Special sensation from the tip of the tongue is carried by the _____ nerve:
    a. Vagus.
    b. Lingual.
    c. Hypoglossal.
    d. Chorda tympani.
    e. Glossopharyngeal.

82. The ____ muscle of the tongue is innervated by the pharyngeal plexus:
    a. Hyoglossus.
    b. Styloglossus.
    c. Palatoglossus.
    d. Intrinsic vertical.
    e. Intrinsic Transverse.

83. Which one of the following muscles raises the floor of the mouth in the first stage of swallowing?
    a. Digastric.
    b. Mylohyoid.
    c. Palatopharyngeus.
    d. Tensor veli palatine.
    e. Levator veli palatine.

84. In the voluntary stage of swallowing, the bolus is compressed against the palate mainly by the action of _____ muscle:
    a. Palatopharyngeus
    b. Middle constrictor.
    c. Tensor veli palatine.
    d. Superior constrictor.
    e. Levator veli palatine.

85. Which one of the following muscles of the palate is innervated by the trigeminal nerve?
    a. Palatoglossus.
    b. Musculus uvulae.
    c. Palatopharyngeus.
    d. Tensor veli palatine.
    e. Levator veli palatine.

86. Nasopalatine nerve block affects the alveolar bone of the:
    a. Upper molars.
    b. Upper incisors.
    c. Lower incisors.
    d. Lower canines.
    e. Lower premolars.

87. The palatine tonsil is normally found within the_____.
    a. Valeculla.
    b. Oro-pharynx.
    c. Piriform fossa.
    d. Naso-pharynx.
    e. Laryngo pharynx.



88. The lingual tonsils are normally found within the_____.
    a. Valeculla.
    b. Oro-pharynx.
    c. Piriform fossa.
    d. Naso-pharynx.
    e. Laryngo-pharynx.

## Nasal Cavity and Paranasal Sinuses

89. What is the bone that forms major parts of roof, lateral walls and septum of the nasal cavity, all at the same time?
    a. Vomer.
    b. Maxilla.
    c. Ethmoid.
    d. Sphenoid.
    e. Palatine bone.

90. The superior concha is part of_____ bones.
    a. Frontal.
    b. Palatine.
    c. Sphenoid.
    d. Maxillary.
    e. Ethmoidal.

91. The bone that forms parts of both lateral wall and floor of the nasal cavity is:
    a. Vomer.
    b. Frontal.
    c. Maxilla.
    d. Sphenoid.
    e. Inferior concha.

92. The nasal and cranial cavities are communicated together through:
    a. Pterygoid canal.
    b. Cribriform plate.
    c. Incisive foramen.
    d. Lesser palatine foramen.
    e. Pterygo-maxillary fissure.

93. Which one of the following opens into the inferior meatus of the nasal cavity?
    a. Maxillary sinus.
    b. Sphenoidal sinus.
    c. Naso-lacrimal Duct.
    d. Posterior ethmoidal.
    e. Middle ethmoidal cells.

94. The ____ artery is main artery of the nasal cavity:
    a. Inferior labial.
    b. Superior labial.
    c. Lesser palatine.
    d. Sphenopalatine.
    e. Greater palatine.



Scanned with CamScanner

95. The common site of nosebleed, (Little's area) is located at the_____of the nose.
a. Roof.
b. Floor.
c. Septum.
d. Left lateral wall.
e. Right lateral wall.

96. General sensation from the roof the nasal cavity is carried by_____nerve.
a. Facial
b. Olfactory.
c. Maxillary.
d. Mandibular.
e. Ophthalmic.

97. The special sensation from the upper part of the nasal septum is carried by_____ nerve.
a. Facial
b. Olfactory.
c. Maxillary.
d. Mandibular.
e. Ophthalmic.

98. The mucous membrane of the inferior concha is innervated by_____ nerve.
a. Facial.
b. Olfactory.
c. Maxillary.
d. Mandibular.
e. Ophthalmic.

99. The sphenoidal sinus is related laterally to the:
a. Pons.
b. Pituitary fossa.
c. Medulla oblongata.
d. Internal carotid artery. + Cavernous sinus
e. Roof of the Nasopharynx.



## Pharynx and Larynx

103. The pharynx extends lower down to the level of ____ vertebra:
a. 3rd cervical.
b. 4th cervical.
c. 6th cervical.
d. 1st thoracic.
e. 4th thoracic.

104. The hypopharynx is located posterior to the:
a. Larynx.
b. Soft palate.
c. Oral cavity.
d. Nasal cavity.
e. Nasal septum.

105. The roof of the nasopharynx is built up by two bones: the basilar part of the occipital bone and the _____ bone:
a. Palatine.
b. Lacrimal.
c. Maxillary.
d. Ethmoidal.
e. Sphenoidal.

106. The pharyngeal muscle that originates from the pterygomandibular raphe is the:
a. Stylo-pharyngeus.
b. Middle constrictor.
c. Palato-pharyngeus.
d. Superior constrictor.
e. Salpingo-pharyngeus.

107. The pharyngeal muscle that is innervated solely by the glossopharyngeal nerve is the _____ muscle:
a. Stylopharyngeus.
b. Palatopharyngeus
c. Middle constrictor.
d. Salpingopharyngeus.
e. Superior constrictor.

108. The upper border of thyroid cartilage lies at level with _____ vertebra:



100. Sensory innervation of the sphenoidal sinus is provided by the _____ nerve:
a. Nasociliary.
b. Supraorbital.
c. Posterior ethmoidal.
d. Anterior superior alveolar.
e. Posterior superior alveolar.

101. The Base of the maxillary sinus forms the:
a. Hard palate.
b. Floor of the orbit.
c. Floor of the nasal cavity.
d. Alveolar part of the maxilla.
e. Lateral wall of the nasal cavity.

102. The adenoid is located within the _____.
a. Valeculla
b. Oro-pharynx.
c. Naso-pharynx.
d. Piriform fossa.
e. Laryngo-pharynx.

Pharyngeal tonsil

- Pharynx end C6
- Upper thyroid start C4
- Cricoid C6

a. C 2.
b. C 4.
c. C6.
a. T2.
b. T4.

109. The cricoid cartilage lies at level with .... vertebra:
a. C 4.
b. C6.
c. T2.
d. T4.
e. T6

110. The type of joint in cricothyroid joint is:
a. Fibrous.
b. Typical synovial.
c. Atypical synovial.
d. Primary cartilaginous.
e. Secondary cartilaginous.

111. The laryngeal vestibule is located:
a. Below the vocal folds.
b. Above the vestibular folds.
c. Above the aryepiglottic fold.
d. Between the vestibular and vocal folds.
e. Lateral to the middle part of the laryngeal

112. Which one of the following muscles elevates the hyoid bone during swallowing?
a. Omohyoid.
b. Genohyoid.
c. Thyrohyoid.
d. Sternohyoid.
e. Sternothyroid.

113. Which one of the following muscles abducts the vocal folds?
a. Cricothyroid.
b. Thyroarytenoid.
c. Transverse arytenoids.
d. Lateral cricoarytenoid. (Adduct)
e. Posterior cricoarytenoid.

114. The cricothyroid muscle is innervated by the __ nerve:
a. Hypoglossal.



Scanned with CamScanner

e. Posterior cricoarytenoid.

114. The cricothyroid muscle is innervated by the __ nerve:
a. Hypoglossal.



b. Internal laryngeal.
c. Glossopharyngeal.
d. External laryngeal.
e. Recurrent laryngeal.



## Ear

115. Among the depressions of the auricle of the ear, the ------ is the deepest one:
a. Scapha.
b. Concha.
c. Scaphoid fossa.
d. Triangular fossa.
e. Intertragic notch.

116. The sensory innervation of the medial surface of the auricle (back of the ear) is provided by the _____ nerve:
a. Facial.
b. Ophthalmic
c. Great auricular.
d. Greater occipital.
e. Auriculotemporal

117. "Pars interna" of the external acoustic meatus is directed:
a. Anterolaterally.
b. Posterolaterally.
c. Anteromedially. (Pars ~~externa~~ interna)
d. Posteromedially. (Pars media)
e. Superiomedially.

Pars externa: mediallyanteriorly

118. The tympanic membrane is at the ….. wall of the middle ear:
a. Medial.
b. Lateral.
c. Inferior.
d. Anterior.
e. Posterior.

119. The roof of the middle ear is formed by the ….. component of the temporal bone:
a. Petrous part.
b. Squamous part.
c. Tympanic part.
d. Styloid process.
e. Zygomatic process.

120. The pharyngotympanic tube is at the ….. wall of the middle ear:
a. Medial.
b. Lateral.
c. Inferior.
d. Anterior.
e. Posterior.



121. The _____ is closely related to the floor of the middle ear:
a. Internal jugular vein.
b. Anterior jugular vein.
c. Retromandibular vein.
d. Internal carotid artery.
e. Common carotid artery.

122. The tensor tympani muscle is attached to the:
a. Incus.
b. Stapes. (Stapedius)
c. Malleus.
d. Round window.
e. Tympanic membrane.

123. The stapedius muscle pulls the stapes ……:
a. Laterally.
b. Medially. (tympanic)
c. Inferiorly.
d. Superiorly.
e. Posteriorly.



Lateral — tympanic memb
Medial — labyrinthine
Anterior — Pharyngotympanic tube
Roof — Tympanic part of temporal
floor — Internal jugular vein

## Orbital Region and Eyeball

124. The medial wall of the orbit is formed primarily by the ____ bone:
a. Palatine.
b. Maxillary.
c. Temporal.
d. Zygomatic.
e. Ethmoidal.

125. The lacrimal bone participates in the formation of the ____ of the orbit:
a. Roof.
b. Floor.
c. Apex.
d. Medial wall.
e. Lateral wall.

126. The apex of the orbit is in the ___ bone:
a. Orbital part of the frontal.
b. Squamous part of temporal.
c. Lesser wing of the sphenoid.
d. Perpendicular plate of palatine.
e. Frontal process of the zygomatic.

127. The thickest part of the orbital walls is the:
a. Roof.
b. Floor.
c. Apex.
d. Medial wall.
e. Lateral wall.

128. The medial angel of the eye is known as:
a. Tarsus.
b. Canthus.
c. Conjunctival sac.
d. Palpebral fissure.
e. Conjunctival fornix.

129. Post-synaptic parasympathetic innervation to the lacrimal gland is conveyed through ___ nerve:
a. Vidian.
b. Zygomatic.
c. Nasociliary.
d. Deep petrosal.
e. Greater petrosal.



Scanned with CamScanner

130. The _____ muscle opens the eye widely:
a. Medial rectus.
b. Lateral rectus.
c. Superior rectus.
d. Superior oblique.
e. Levator palpebrae superioris.

131. The Abducent nerve supplies the _____ muscle:
a. Lateral rectus.
b. Superior tarsal.
c. Superior rectus.
d. Inferior oblique.
e. Superior oblique.

132. The Superior oblique muscle is supplied by the ____ nerve:
a. Frontal.
b. Lacrimal.
c. Abducent.
d. Trochlear.
e. Oculomotor.

133. The medial rectus muscle _____ the eye:
a. Rotates.
b. Adducts.
c. Abducts. (lateral)
d. Elevates.
e. Depresses.

134. The pre-synaptic parasympathetic root of the ciliary ganglion arises from the _____ nerve:
a. Facial.
b. Frontal.
c. Abducent.
d. Nasociliary.
e. Oculomotor.

135. The anterior transparent part of the outer (fibrous) layer of the eye is named:
a. Sclera.
b. Retina.
c. Cornea.
d. Choroid
e. Ciliary body.



136. The "red eye" phenomenon that occurs in photography is due to flash reflection from the:
a. Iris.
b. Pupil.
c. Retina
d. Cornea.
e. Choroid (Red eye)

137. The dilator pupillae muscle receives postsynaptic sympathetic innervation through the _____ nerve:
a. Facial.
b. Frontal.
c. Lacrimal.
d. Oculomotor.
e. Long ciliary.


138. In pupillary light reflex, the efferent signal is carried by the ___ nerve:
a. Optic.
b. Facial.
c. Abducent.
d. Trochlear.
e. Oculomotor.

139. The "blind spot" in the retina refers to the _____:
a. Sclera.
b. Retina.
c. Optic disc.
d. Ciliary body.
e. Macula lutea.

140. The area of the most acute visual resolution is the:
a. Iris.
b. Pupil.
c. Retina.
d. Fovcola.
e. Optic disc.

high visual resolution

141. Macula lutea is located _____ to the optic disc:
a. Deep.
b. Lateral.
c. Medial.
d. Inferior.
e. Superior.





142. In the eye, the main site that is responsible for refraction of the light is:
a. Iris.
b. Lens.
c. Cornea.
d. Aqueous humor.
e. Vitreous humor

143. The lens is adjusted for near vision by its increased convexity due to the contraction of the _____ muscle:
a. Ciliary.
b. Superior rectus.
c. Dilator pupillae.
d. Sphincter pupillae.
e. Levator Palpebrae Superioris.

## Section Two: Scenario-Based MCQs

### Clinical Case Scenario (1)

Jaafar is a 24 years old university student who was involved in a fight, presented to the emergency room with lacerations on the side of the face over the mandible, the patient was feeling numbness in the lower lip and chin, examination and imaging studies revealed mandibular fracture and dislocation.

Answer the following questions:

144. Which part of the mandible participates in the formation of the temporomandibular joint (TMJ)?
a. Angle.
b. Ramus.
c. Genial tubercle.
d. Symphysis menti.
e. Condyloid process

145. What is the most common type of mandibular dislocation?
a. Lateral.
b. Superior.
c. Anterior.
d. Posterior.
e. Posterio-superior.

146. What is the most common site of fracture in the mandible?
a. Body.
b. Angle.
c. Symphysis menti.
d. Coronoid process.
e. Condyloid process.

147. What is the most commonly injured nerve following mandibular fracture?
a. Buccal.
b. Lingual.
c. Inferior alveolar.
d. Transverse cervical.
e. Marginal mandibular.



Scanned with CamScanner