Home | Neuro bcqs test.docx | Sign in | Upgrade now
Menu | Home | Insert | Page Layout | References | Review | View | Tools | WPS AI | Share
Table Of Contents | Update TOC | TOC Level | Insert Footnote | Previous Footnote | Next Footnote | Insert Endnote | Previous Endnote | Next Endnote | Footnote/Endnote Separator | Caption | Insert Table of Figures | Cross-reference

1.What lies directly above the tentorium?

A. The parietal lobes.
B. The temporal lobes.
C. The cerebellum
D. The frontal lobes.
E. The occipital lobes..

2.What are the major structural differences between dorsal root ganglia (DRG) and autonomic ganglia?

A. One is derived from neural crest and the other from neural tube.
B. The DRG contain the cell bodies of sensory neurons whereas the sympathetic ganglia contain the synaptic junctions between preganglionic and postganglionic neurons.
C. Both have synapses.
D. One belongs to the CNS and the other belongs to the PNS.

3.What vessel(s) serve(s) areas involved in speech in the majority of people?

A. Right middle cerebral artery.
B. Right and left middle cerebral arteries
C. Left middle cerebral artery.
D. Right and left posterior cerebral arteries

Words: 1518 | Spell Check | 130%
Type here to search | 34°C | 12:4 12/0

Scanned with CamScanner

Home | Neuro bcqs test.docx | Sign in | Upgrade now
Menu | Home | Insert | Page Layout | References | Review | View | Tools | WPS AI | Share
Table Of Contents | Update TOC | TOC Level | Insert Footnote | Previous Footnote | Next Footnote | Insert Endnote | Previous Endnote | Next Endnote | Footnote/Endnote Separator | Caption | Insert Table of Figures | Cross-reference

2.What are the major structural differences between dorsal root ganglia (DRG) and autonomic ganglia?

A. One is derived from neural crest and the other from neural tube.

B. The DRG contain the cell bodies of sensory neurons whereas the sympathetic ganglia contain the synaptic junctions between preganglionic and postganglionic neurons.

C. Both have synapses.

D. One belongs to the CNS and the other belongs to the PNS.

3.What vessel(s) serve(s) areas involved in speech in the majority of people?

A. Right middle cerebral artery.

B. Right and left middle cerebral arteries

C. Left middle cerebral artery.

D. Right and left posterior cerebral arteries.

E. Left middle and posterior cerebral arteries.

4.



1/7 | Words: 1518 | Spell Check | 130%
Type here to search | 34°C | 12:40

Scanned with CamScanner

Scanned with CamScanner

Obstruction of the cerebral aqueduct would result in dilation of which of the following?

A. Lateral ventricles
B. Third ventricle

Double-click to edit header

C. Fourth ventricle
D. A and B
E. A, B, and C

5. Which of the following structures does not border on the third ventricle?

A. Fourth ventricle

Scanned with CamScanner

5.Which of the following structures does not border on the third ventricle?

A. Fourth ventricle
B. Hypothalamus
C. Thalamus
D. Interventricular foramen
E. Aqueduct

6.The spinal nerves consist of ventral and dorsal roots. Where are the cell bodies of the axons in each root?

A. Ventral root cell bodies are in the gray matter of the spinal cord and dorsal root cell bodies are in ganglia.
B. Ventral and dorsal root cell bodies are in the gray matter of the cord.
C. Ventral and dorsal root cell bodies are in ganglia, (clusters of cell bodies outside the CNS).
D. Ventral root cell bodies are in ganglia and dorsal root cell bodies are in the gray matter of the spinal cord

7. How do the meninges cover the spinal cord?

A. The pia, arachnoid, and dura cover the spinal cord in tight, closely apposed layers.
B. The dura, arachnoid, and pia of the brain and spinal cord are continuous. All

Type here to search | 34°C | 12:41 PM 12/06/2024

Scanned with CamScanner

6.The spinal nerves consist of ventral and dorsal roots. Where are the cell bodies of the axons in each root?

A. Ventral root cell bodies are in the gray matter of the spinal cord and dorsal root cell bodies are in ganglia.

B. Ventral and dorsal root cell bodies are in the gray matter of the cord.

C. Ventral and dorsal root cell bodies are in ganglia, (clusters of cell bodies outside the CNS).

D. Ventral root cell bodies are in ganglia and dorsal root cell bodies are in the gray matter of the spinal cord

7. How do the meninges cover the spinal cord?

A. The pia, arachnoid, and dura cover the spinal cord in tight, closely apposed layers.

B. The dura, arachnoid, and pia of the brain and spinal cord are continuous. All three layers loosely cover the spinal cord.

C. Only the dura mater continues down from the brain to cover the spinal cord, protecting it from the surrounding bone

D. The dura, arachnoid, and pia all cover the spinal cord; the dura and arachnoid are tightly connected with each other..

8. The spinal cord is held in place by these structures?

Scanned with CamScanner

protecting it from the surrounding bone

D. The dura, arachnoid, and pia all cover the spinal cord; the dura and arachnoid are tightly connected with each other..

8. The spinal cord is held in place by these structures?
   A. Pia mater and choroid plexus
   B. Conus medullaris and cauda equina
   C. Denticulate ligament and Filum terminale
   D. Anterior and posterior spinal ligaments
   E. Ligamentum flavum

9. In the medulla, where is the anterolateral system located with respect to the nucleus ambiguus?
   A. It is dorsal to the nucleus ambiguus.
   B. It is lateral to the nucleus ambiguus.
   C. It is ventral to the nucleus ambiguus.
   D. It is medial to the nucleus ambiguus.

Scanned with CamScanner

10. Which of the following tracts carry proprioception fibres?
    A. Anterior spinothalamic tract
    B. Lateral corticospinal tract
    C. Rubrospinal tract
    D. Lateral spinothalamic tract
    E. Dorsal column-medial lemniscus pathway

11. Sensory dissociation is the loss of pain and temperature with sparing of vibration, joint position, and fine touch. By knowing the location of the two pathways that carry the information, at which level are the second order axons located:
    A. Thalamus.
    B. Midbrain.
    C. Pons.
    D. Hypothalamus
    E. Medulla.

12. A patient suffers loss of sensation from the face and upper body. The obstructed vessel is most likely:
    A. Anterior cerebral artery.
    B. Middle cerebral artery

Scanned with CamScanner

12. A patient suffers loss of sensation from the face and upper body. The obstructed vessel is most likely:

A. Anterior cerebral artery.
B. Middle cerebral artery.
C. Posterior cerebral artery.
D. Internal carotid artery.

13. The neostriatum consists of which of the following structures?

A. Lentiform and caudate nucleus
B. Putamen and caudate nucleus
C. Lentiform and globus pallidus
D. Putamen and lentiform nucleus
E. Putamen and globus pallidus

14. The suprachiasmatic nucleus of the hypothalamus is responsible for?

A. Regulation of the circadian rhythm
B. Control of feeding impulses
C. Regulation of osmolality
D. Thermoregulation
E. Oxytocin release

Scanned with CamScanner

15. How do axons from the VPM nucleus of thalamus reach the postcentral gyrus?

A. Internal capsule.
B. External capsule.
C. Extreme capsule.
D. Internal medullary lamina.

16. To what cortical region does the VPL and VPM nucleus of the thalamus project?

A. Transverse temporal lobe.

Scanned with CamScanner

16. To what cortical region does the VPL and VPM nucleus of the thalamus project?
    A. Transverse temporal lobe.
    B. Postcentral gyrus.
    C. Precentral gyrus.
    D. Prefrontal cortex

17. The area of the medulla containing the spinal nucleus of V may be supplied by which artery?
    A. Anterior inferior cerebellar artery (AICA).
    B. Superior cerebellar artery.
    C. Anterior spinal artery.
    D. Posterior inferior cerebellar artery (PICA).
    E. Posterior cerebral artery.

18. The cerebrospinal fluid (CSF) leaves the lateral ventricles to flow into the third ventricle through which of the following structures?
    A. Foramen of Magendie
    B. Cisterna Magna
    C. Foramen of Monro
    D. Cerebral aqueduct
    E. Foramen of Luschka

Scanned with CamScanner

E. Foramen of Luschka

19. The internal cerebral vein is formed by the union of which of the following?
    A. Superficial cerebral vein and the basal vein
    B. Anterior cerebral vein and striate vein
    C. Superior cerebral vein and the great cerebral vein
    D. Superficial and deep middle cerebral vein
    E. Thalamostriate and choroidal veins
20. Which of the following is the correct option regarding the cerebellum?
    A. Stellate and basket cells are present in the molecular layer
    B. Purkinje neurons are of bipolar variety
    C. Purkinje cells have scarce branching
    D. Superior cerebellar peduncle connects cerebellum to pons
    E. Molecular, granular, and pyramidal layers are present in cortex

21. Which of the following is a sensory relay nucleus of the thalamus?
    A. Pulvinar nucleus.
    B. Anterior nucleus.
    C. Ventral posteromedial nucleus.
    D. Intralaminar nuclei (i.e., centromedian).
    E. Dorsomedial nucleus.

Scanned with CamScanner

21. Which of the following is a sensory relay nucleus of the thalamus?
    A. Pulvinar nucleus.
    B. Anterior nucleus.
    C. Ventral posteromedial nucleus.
    D. Intralaminar nuclei (i.e., centromedian).
    E. Dorsomedial nucleus.
22. The role of the corpus callosum can best be defined as
    A. Association cortical connections from all areas.
    B. Intracortical connections between the same gyri on opposite sides of each hemisphere.
    C. A dorsal extension of the internal capsule.

Home | Neuro bcqs test.docx | Sign in | Upgrade now
Menu | Home | Insert | Page Layout | References | Review | View | Tools | WPS AI | Share
Table Of Contents | Update TOC | TOC Level | Insert Footnote | Previous Footnote | Next Footnote | Insert Endnote | Previous Endnote | Next Endnote | Footnote/Endnote Separator | Caption | Insert Table of Figures | Cross-reference

hemisphere.

E. A pathway to ensure that the same information goes to each hemisphere.

23. The red nucleus which gives off the rubrospinal tract, is located in which of the following?
    A. Pons
    B. Medulla
    C. Cerebellum
    D. Basal ganglia
    E. Midbrain
24. Crude touch sensation is lost due to damage to which of the following?
    A. Lateral corticospinal tract
    B. Lateral spinothalamic tract
    C. Anterior spinothalamic tract
    D. Dorsal column-medial lemniscus pathway
    E. Spinocerebellar tract
25. Which of the following structures is responsible for emotional memory?
    A. Hippocampus
    B. Midbrain
    C. Cerebellum
    D. Hypothalamus
    E. Amygdala

Words: 1518 | Spell Check | 130% | Type here to search | 34°C

Scanned with CamScanner

Scanned with CamScanner

26. What attaches the cerebellum to the brain stem?
    - A. Cerebral peduncles.
    - B. Vermis.
    - C. Cerebellar peduncles.
    - D. Tonsil.

27. What is the vascular supply of the pons?
    - A. Vertebral arteries.
    - B. Basilar artery.
    - C. Superior cerebellar artery.
    - D. Posterior inferior cerebellar artery.

28. What route do the thalamocortical fibers from the ventral lateral nucleus take?
    - A. Internal capsule
    - B. External capsule
    - C. Extreme capsule
    - D. Cerebral peduncle
    - E. Pyramid

29. Climbing fibers:
    - A. Excite Purkinje cells.
    - B. Have cells bodies on the opposite side of the medulla.

Home | Neuro bcqs test.docx | Sign in | Upgrade now
Menu | Home | Insert | Page Layout | References | Review | View | Tools | WPS AI | Share
Table Of Contents | Update TOC | TOC Level | Insert Footnote | Previous Footnote | Next Footnote | Insert Endnote | Previous Endnote | Next Endnote | Footnote/Endnote Separator | Caption | Insert Table of Figu | Cross-reference

E. Pyramid

29. Climbing fibers:

A. Excite Purkinje cells.
B. Have cells bodies on the opposite side of the medulla.
C. Contact only a few Purkinje cells.

D. Are axons of cells in the inferior olivary nucleus.
E. All of the above.

30.



5/7 | Words: 1518 | Spell Check | 130%
Type here to search | 34°C

Scanned with CamScanner

Home | Neuro bcqs test.docx | Sign in | Upgrade now
Menu | Home | Insert | Page Layout | References | Review | View | Tools | WPS AI | Share
Table Of Contents | Update TOC | TOC Level | Insert Footnote | Previous Footnote | Next Footnote | Insert Endnote | Previous Endnote | Next Endnote | Footnote/Endnote Separator | Caption | Insert Table of Fig | Cross-reference



The prominent caudate and putamen seen in this axial or horizontal MRI section are separate- by the:

A. Anterior limb of the internal capsule.
B. Genu of the internal capsule.
C. Posterior limb of the internal capsule.
D. All limbs of the internal capsule.

31. The basal ganglia exert their effects on motor behavior through the:

A. Rubrospinal tract.
B. Vestibulospinal tract.
C. Reticulospinal tract.
D. Corticospinal tract.
E. All of the above.

32. Pontocerebellar fibres continue as:

A. Middle cerebellar peduncle
B. Superior cerebellar peduncle
C. Inferior cerebellar peduncle
D. Posterior cerebral peduncle
E. Middle cerebral peduncle

5/7 | Words: 1518 | Spell Check | 130%

Scanned with CamScanner

Home
Neuro bcqs test.docx
Sign in
Upgrade now
Menu
Home Insert Page Layout References Review View Tools
WPS AI
Share
Table Of Contents
Update TOC
TOC Level
Insert Footnote
Previous Footnote
Next Footnote
Insert Endnote
Previous Endnote
Next Endnote
Footnote/Endnote Separator
Caption
Insert Table of Figures
Cross-reference

E. Middle cerebral peduncle

33. The appropriate statement regarding the medulla oblongata is?
    A. Olives are oval elevations produced by underlying superior olivary nuclei
    B. Cuneate tubercle lies lateral to the gracilis tubercle
    C. Middle cerebellar peduncle connects medulla to the cerebellum
    D. Nucleus ambiguous consists of large sensory neurons
    E. Nucleus of the spinal tract of trigeminal nerve lies medial to the corticospinal tract

Double-click to edit header
ords: 1518
Spell Check
130%

Scanned with CamScanner

Scanned with CamScanner

34. Tuberal nuclei are located in:
    A. Hypothalamus.
    B. Median eminence.
    C. Infundibulum.
    D. Anterior lobe of pituitary.

35. Preganglionic sympathetic cell bodies are located in:
    A. lateral horn (T1-L2).
    B. lateral horn (S2-5).
    C. Both A and B
    D. Edinger-Westphal, salivatory and dorsal motor nucleus of X.

36. Which set of cranial nerves contains preganglionic parasympathetic axons.
    A. V, VII, IX, X.
    B. IV, VI, XI, XII.
    C. III, VII, IX, X.
    D. I, II, VII, XI.

37. What is the relationship of the hippocampus to the inferior horn of the lateral ventr
    A. Its in the floor of the ventricle.

Scanned with CamScanner

37. What is the relationship of the hippocampus to the inferior horn of the lateral ventricle?
    - A. Its in the floor of the ventricle.
    - B. Its in the roof of the ventricle.
    - C. It is rostral to the ventricle.
    - D. It has no relationship to the ventricle.

38. Axons of hippocampal pyramidal cells travel in the
    - A. Ventral (amygdalofugal) pathway.
    - B. Stria terminalis.
    - C. Fornix.
    - D. Cingulum.

39. Regarding boundaries of the 4th ventricle, the caudal part of the lateral boundary is formed by?
    - A. Superior cerebellar peduncle
    - B. Middle cerebellar peduncle
    - C. Superior and middle cerebellar peduncle
    - D. Middle and inferior cerebellar peduncle
    - E. Inferior cerebellar peduncle

40. Which of the following is/are true regarding filum terminale?
    - A. It is attached to the conus medullaris.
    - B. It is a continuation of the pia and ependyma of the spinal cord.

Neuro bcqs test.docx
Sign in
Upgrade now
Home Insert Page Layout References Review View Tools
WPS AI
Share
Update TOC
TOC Level
Insert Footnote
Previous Footnote
Next Footnote
Insert Endnote
Previous Endnote
Next Endnote
Footnote/Endnote Separator
Caption
Insert Table of Figu
Cross-reference

C. Fornix.
D. Cingulum.

39. Regarding boundaries of the 4th ventricle, the caudal part of the lateral boundary is formed by?

A. Superior cerebellar peduncle
B. Middle cerebellar peduncle
C. Superior and middle cerebellar peduncle
D. Middle and inferior cerebellar peduncle
E. Inferior cerebellar peduncle

40. Which of the following is/are true regarding filum terminale?

A. It is attached to the conus medullaris.
B. It is a continuation of the pia and ependyma of the spinal cord.
C. It penetrates the dura at the end of the dural sac (vertebra S2).
D. It terminates as the coccygeal ligament fusing with the periosteum of the co
E. All of the above.

18
Spell Check
Double-click to edit footer
130%
to search
34°C
Scanned with CamScanner