c. Cytotrophoblasts
d. Trophoblasts
e. Blastomere

Q19. Maternal site of placenta is formed by

a. Decidua parietalis
b. Decidua basalis
c. Chorionic frondosum (villous)
d. Decidua capsularis
e. Chorionic laevae (smooth)

Q20. Hormone NOT synthesized by Placenta is;

a. Glycoprotein hCG.
b. Human chorionic somatomamotrophin.
c. Luteinizing hormone.
d. Human chorionic thyrotropin .
e. Human chorionic corticotrophin.

Q21. Within an osteon we cannot find:

a. Lamellae
b. Haversian canal
c. Canaliculi
d. Chondrocytes
e. Lacunae

Hyaline cartilage does not include the following:

b. Epiphysial plates
c. Nasal septum
d. Pinna of ear
e. Sternal ends of ribs
Articular surfaces of movable joints

Q25. Each muscle fiber has a sleeve of areolar connective tissue around

a. Deep fascia.
b. Superficial fascia.
c. Perimysium.
d. Epimysium.
e. Endomysium

Q26.Which one is the peculiar feature of a skeletal muscle fiber

a. It has a single nucleus located centrally
b. It has multiple nuclei which are located periphery of the fiber
c. Its ends are branching
d. It unites with its fellow fiber with the help of intercalated disc
e. Its contraction cannot be controlled voluntarily

Q27.Tendons are composed of _____connective tissue.

a. Skeletal
b. Areolar
c. Dense irregular
d. Yellow elastic

Q44. The most vascular layer of meninges is:
a. Meningeal layer of dura matter
b. Pia matter
c. Arachnoid layer of dura matter
d. Endosteleal layers of dura matter
e. Falxcerebri

Q45. Choose false statement regarding blood supply of brain
a. Anterior cerebral Artery
b. Basilar artery
c. Posterior cerebral
d. External carotid artery
e. Internal carotid artery

Q46. In sensory pathways, first order neurons are present
a. Anterior horn of spinal cord
b. Posterior horn of spinal cord
c. Lateral horn of spinal cord
d. Posterior root ganglia
e. In the thalamus

Q47. Regarding white mater of brain
a. Corona radiate has only the descending fibers
b. Internal capsule lies between the putamen and glob
c. Corpus callosum is the largest commissure of brai
d. Anterior commissure links two occipitals poles v
e. Habenular commissure is part of the hypothalam

b. Areolar
c. Dense irregular
d. Yellow elastic
e. Dense regular

Q28. Blood vessels are lined with _____epithelium.

a. Stratified squamous
b. Transitional
c. Simple columnar
d. Simple squamous
e. Stratified cuboidal

Q29. Keratinized stratified squamous epithelium is found in

a. The oral cavity .
b. The nasal cavity.
c. The epidermis
d. The pharynx
e. The vagina

Q30. Choose the true statement about the Skin.

a. Having thick dermis is called the thick skin.
b. Epidermis is devoid of nerve supply

b. Areolar
c. Dense irregular
d. Yellow elastic
e. Dense regular

Q28. Blood vessels are lined with _____epithelium.

a. Stratified squamous
b. Transitional
c. Simple columnar
d. Simple squamous
e. Stratified cuboidal

Q29. Keratinized stratified squamous epithelium is found in

a. The oral cavity .
b. The nasal cavity.
c. The epidermis
d. The pharynx
e. The vagina

Q30. Choose the true statement about the Skin.

a. Having thick dermis is called the thick skin.
b. Epidermis is devoid of nerve supply

b) The nose.
c) The outer ear.
d) The costal cartilage of the rib cage.
e) Intervertebral discs.

Q34. The junction between the diaphysis and epiphysis in a growing bone is called th

a) Endosteum
b) Articular plate
c) Periosteum
d) Epiphyseal plate
e) Epiphyseal line

Q35. Which of the following part of diencephalon controls endocrine hormonal s autonomic nervous system

a) Hypothalamus
b) Dorsal thalamus
c) Epithalamus
d) Subthalamus
e) Thalamus

Q36. Which of the following nuclei are associated with the planning of the n

a) Limbic system
b) Cerebral cortex
c) Thalamus
d) Basal nuclei
e) Hypothalamus

Q37. Which of the following is not the part of hypothalamus

a) Tubes cinereum
b) Median eminence
c) Mammillary body
d) Optic chiasma
e) Pineal gland

b) The nose.
c) The outer ear.
d) The costal cartilage of the rib cage.
e) Intervertebral discs.

Q34. The junction between the diaphysis and epiphysis in a growing bone is called th

a) Endosteum
b) Articular plate
c) Periosteum
d) Epiphyseal plate
e) Epiphyseal line

Q35. Which of the following part of diencephalon controls endocrine hormonal s autonomic nervous system

a) Hypothalamus
b) Dorsal thalamus
c) Epithalamus
d) Subthalamus
e) Thalamus

Q36. Which of the following nuclei are associated with the planning of the n

a) Limbic system
b) Cerebral cortex
c) Thalamus
d) Basal nuclei
e) Hypothalamus

Q37. Which of the following is not the part of hypothalamus

a) Tubes cinereum
b) Median eminence
c) Mammillary body
d) Optic chiasma
e) Pineal gland

Q33. Fibrocartilage is found in
a) The joint surfaces
b) The nose.
c) The outer ear.
d) The costal cartilage of the rib cage.
e) Intervertebral discs.

Q34. The junction between the diaphysis and epiphysis in a growing bone is called
a) Endosteum
b) Articular plate
c) Periosteum
d) Epiphyseal plate
e) Epiphyseal line

Q35. Which of the following part of diencephalon controls endocrine hormonal autonomic nervous system
a) Hypothalamus
b) Dorsal thalamus
c) Epithalamus
d) Subthalamus
e) Thalamus

Q36. Which of the following nuclei are associated with the planning of the

d) Fibroblasts
e) Osteoblasts

Q7. Soft connective tissue

a) An indication of microcephaly
b) Frontal sinuses
c) Epiphyseal plates
d) Cribriform plates
e) Fontanelles

Q8. Endochondral and intramembranous

a) Bone remodeling
b) Embryonic skeletal ossification
c) Controlling blood calcium levels
d) Cartilage synthesis

a) An indication of microcephaly
b) Frontal sinuses
c) Epiphyseal plates
d) Cribriform plates
e) Fontanelles

Q8.Endochondral and intramembranous are two mechanisms o

a) Bone remodeling
b) Embryonic skeletal ossification
c) Controlling blood calcium levels
d) Cartilage synthesis
e) Cartilage development

Q9. The Haversian (central) canal in each osteon contains

a) Chondroitin sulfate
b) Hydroxyapatite
c) Osteoblasts
d) Blood vessels
e) Synovial fluid

Q10.Which of the following is an event of 3rd week of

a) Development of Notochord
b) Webbing of digits

luteum abundantly produces which of the following hormone

a) Follicle stimulating hormone
b) Luteinizing hormone
c) Progesterone
d) Estrogen
e) Human chorionic gonadotropin hormone

Q14. Somites are derived from;

a. Extra embryonic somatic mesoderm
b. Extra embryonic splanchnic mesoderm.
c. Paraxial mesoderm.
d. Mesenchymal transformation.
e. Neural crest cells.

Q15. Neural crest cells give rise to;

a. Meninges.
b. Suprarenal medulla.
c. Pigment cell of dermis.
d. Enamel of teeth.
e. Muscle ,bone and connective tissue of pharyngeal arches origin.

Q16. The following changes do not occur in the period of 9th to 12th v

a. Primary ossification centers appear in the skeleton.
b. Liver is major site of erythropoiesis.
c. External genitalia of male and female appear.

luteum abundantly produces which of the following hormone

a) Follicle stimulating hormone
b) Luteinizing hormone
c) Progesterone
d) Estrogen
e) Human chorionic gonadotropin hormone

Q14. Somites are derived from;

a. Extra embryonic somatic mesoderm
b. Extra embryonic splanchnic mesoderm.
c. Paraxial mesoderm.
d. Mesenchymal transformation.
e. Neural crest cells.

Q15. Neural crest cells give rise to;

a. Meninges.
b. Suprarenal medulla.
c. Pigment cell of dermis.
d. Enamel of teeth.
e. Muscle ,bone and connective tissue of pharyngeal arches origin.

Q16. The following changes do not occur in the period of 9th to 12th v

a. Primary ossification centers appear in the skeleton.
b. Liver is major site of erythropoiesis.
c. External genitalia of male and female appear.

Pineal gland

Q38. The following statements concern the longitudinal cerebral fissure

a. The fissure contains the fold of dura mater, the falxcerebelli.
b. The fissure contais the middle cerebral arteries .
c. The superior sagittal sinus lies below it.
d. In the depths of the fissure, the corpus callosum crosses the midline.
e. The inferior sagittal sinus lies above it.

Q39. The following statements concern the central sulcus:

a. The central sulcus extends onto the medial surface of the cerebral he
b. The frontal lobe lies posterior to it.
c. The parietal lobe lies anterior to it.
d. The central sulcus is continuous inferiorly with the lateral sulcus.
e. The arachnoid mater extends into the central sulcus.

Q40.Structures found within the cerebellum does not include:

a. The festigeal nucleus
b. The dentate nucleus
c. The emboliform nucleus
d. The arcuate nucleus
e. The globose nucleus

Q41. Choose the true statement regarding the spinal cord:

a. It's the long cylindrical upper part of CNS.
b. It occupies the upper 2/3rd of vertebral canal
c. It gives rise 31 pairs of cranial nerve.
d. Superiorly it is continuous with filumterminale

structures are at the center of compact bone lame
the bone length:
a) Haversian canals
b) Canaliculi
c) Perforating canals
d) Osteocytes
e) Lacunae

Q6. The cell type that is responsible for maintaining bo

a) Osteoclasts
b) Chondrocytes
c) Osteocytes
d) Fibroblasts
e) Osteoblasts

Q7. Soft connective tissue membranes between th

a) An indication of microcephaly
b) Frontal sinuses
c) Epiphyseal plates
d) Cribriform plates

structures are at the center of compact bone lame
the bone length:
a) Haversian canals
b) Canaliculi
c) Perforating canals
d) Osteocytes
e) Lacunae

Q6. The cell type that is responsible for maintaining bo

a) Osteoclasts
b) Chondrocytes
c) Osteocytes
d) Fibroblasts
e) Osteoblasts

Q7. Soft connective tissue membranes between th

a) An indication of microcephaly
b) Frontal sinuses
c) Epiphyseal plates
d) Cribriform plates

Saving screenshot...

the best option:

BCQs

. Regarding Endochondral bone formation
) Secondary spongiosa is formed by the woven bone
) Secondary spongiosa is found only on diaphysis
) Secondary spongiosa is formed by lamellar bone
) Secondary spongiosa is found only on cartilage surface
) Bone develops directly within the soft connective tissues

2. Rhombomeres is the eight series of buldges formed from
a) Three layered folding
b) By the fusion of neural tube
c) By the fusion of neural folds
d) From the part of hind brain
e) From the rostal fold of embryo

3. Cervical sinus developed on the side of the neck by the growth of
a) Second groove
b) Third groove
c) Fourth groove
d) By over growth of second arch
e) By the growth of all above

doesn't include

Second brachial arch derivatives are
a) Stylohyoid ligament
b) Lesser horns of the hyoid bone
c) Upper part of the body of hyoid bone
d) Reichert's cartilage
e) Styloid process of temporal bone.

5. Primary function of Oral Mucosa is:

a) Three layered folding
b) By the fusion of neural tube
c) By the fusion of neural folds
d) From the part of hind brain
e) From the rostal fold of embryo

3. Cervical sinus developed on the side of the neck by the growth of
a) Second groove
b) Third groove
c) Fourth groove
d) By over growth of second arch
e) By the growth of all above

4. Second brachial arch derivatives are
a) Stylohyoid ligament
b) Lesser horns of the hyoid bone
c) Upper part of the body of hyoid bone
d) Reichert's cartilage
e) Styloid process of temporal bone.

Primary function of Oral Mucosa is:
Sensation
Secretion
Thermal regulation
Protection
Conductivity

egarding compact bone
The inner layer of periostium consists of dense, irregular
The outer layer of periostium consists of bone cells, their
supply
Periosteal surface of bone is more active in booe format
Endosteal surface of bone is more active in bone format
The inner layer of periosteum is fibrous layer.

11. Which is the derivative of 2nd pharyngeal arch:
    a. Malleus
    b. Incus
    c. Auditory tube
    d. Stapes
    e. Middle ear cavity

12. Which is the derivative of 1st pharyngeal arch:
    a. Stapedius muscle
    b. Tympanic membrane
    c. Malleus
    d. Auricle
    e. External auditory meatus

13. Utricle, semisulcular ducts, vestibular ganglion of VIII cranial nerve are derived from:
    a. Otic vesicle
    b. Auricular hillocks
    c. Pharyngeal groove
    d. 1st pharyngeal arch
    e. Pharyngeal membrane

14. Auditory tube is supplied by:
    a. Middle menengial artery
    b. Artery of pterigoid canal
    c. Ascending palatine artery
    d. A & B
    e. A & C

15. The saccule, cochlear duct, spiral ganglion of VIII cranial nerve are derived from:
    a. Utrical portion of otic vesicle
    b. Saccular portion of otic vesicle
    c. Auricular hillocks
    d. Pharyngeal groove
    e. Pharyngeal arch 1st

16. The blood supply of pinna comes from:
    a. External carotid artery
    b. Internal carotid artery
    c. Both A & B
    d. Facial artery
    e. Maxillary artery

17. The blood supply of Eustachian tube comes from all of the arteries except:
    a. Middle meningeal artery
    b. Ascending pharyngeal artery
    c. Artery of the pterigoid canal
    d. Deep auricular artery
    e. None of the above

18. The ear develops from the :
    a. 1st pharyngeal pouch
    b. 2nd pharyngeal pouch
    c. 3rd pharyngeal pouch
    d. 4th pharyngeal pouch
    e. 5th pharyngeal pouch

19. The vesicles begins to develop around:
    a. 20th day of development
    b. 25th day of development
    c. 33rd day of development
    d. 60th day of development
    e. 65th day of development

20. Pinna originates as a fusion of:
    a. 2 hillocks
    b. 3 hillocks
    c. 4 hillocks
    d. 5 hillocks
    e. 6 hillocks

Q39) Lower limb and anal region in cerebral hemisphere are supplied by:

a) Middle cerebral artery.
b) Posterior communicating branch.
c) Anterior cerebral artery.
d) Anterior communicating branch.
e) Posterior cerebral artery.

Q40) REGARDING FERTILIZATION FOLLOWING STATEMENT IS INCORRECT:

a) it occurs in the ampullary part of uterine tube.
b) disintegration of the zona pelucida begins.
c) secondary oocyte completes its second meiotic division.
d) genetic sex of the baby can be determined.
e) restores the diploid number of chromosomes of species.

Q41) 'Posterior cerebral artery' is a terminal branch of:

a) External carotid artery.
b) Vertebral aretery.
c) Basilar artery.

Second Year BDS Semester-III Examinations 2010

Paper "Anatomy"

Time Allowed: Two Hours

Max. Marks:

Q1) A muscle which elevates and retracts the mandible is
- a) Medial pterygoid
- b) Masseter
- c) Temporalis
- d) Digastric
- e) Lateral pterygoid

Q2) Unique features in the development of skull is
- a) It encloses a cavity
- b) It is contributed by paraxial mesoderm
- c) Neural crest cells forms membranous portion
- d) It is mesodermal in origin
- e) It ins two layer of bone

Q3) The best statement regarding the microscopic features of cerebellar cortex
- a) The afferent fibers terminate in granular layer
- b) It consists of six layers of cells
- c) In medullary layer cells are compactly arranged
- d) The Purkinje layer consists of flask shaped cells with highly developed dendritic tree
- e) In granuler layer the cells are dispersed

b) Medial
c) Superior
d) Anterior
e) Posterior

Q6) Regarding the tympanic cavity

a) Tensor tympani is the only muscle which is present in the tympanic cavity
b) It is the air containing part of ear
c) A cranial nerve is related to tympanic cavity
d) It contains small bones
e) Auditory tube extends from the anterior wall of tympanic cavity to naso pharynx

Q7) The roots of first cervical spinal nerve corresponds to the

a) level of formen magnum
b) atlas vertebra
c) upper edge of spinal cord
d) junction of medulla and spinal cord
e) lower edge of medulla oblongata

Q8) The following fibre lies immediately beneath the cortex and connect the adjacent gyri:

a) Long association fibres
b) Uncinate fasciculus
c) Inferior longitudinal fasciculus
d) Short association fibres
e) Fronto occipital fasciculus

Q9) During developments of tongue, the copula contributes to

a) Filiform papillae
b) posterior 1/3rd of the tongue
c) Incisive foramen
d) Frenulum of the tongue
e) Anterior 2/3rd of the tongue

Q10) Regarding Retina (point out the odd statement)

a) Nuclei of cones are dispersed near the limiting membrane

Select an odd statement regarding the structure of eye:

a) Ciliary epithelium consist of a single layer of columnar unpigmented cells.
b) Photo receptors of the retina are modified bipolar neurons.
c) Inner layers of retina are absent in the fovea centralis.
d) Descemet's membrane of cornea is highly refractile under the light microscope.
e) A single layer of cuboidal cells is present all over just under the lens capsule.

Q44) Which of the following nerve gives parasympathetic fibres to parotid gland

a) Facial
b) Trigeminal
c) Accessory nerve
d) Vagus
e) Glossopharyngeal

Q45) The importance of alar plates in the developing spinal cord is that

a) it is present bilaterally
b) it is separate from basal plate by sulcus limitans
c) it produces a longitudinal bulge
d) its cell bodies form the dorsal gray column of spinal cord
e) its enlargement on both the sides form dorsal median septum

Q46) Where are the vestibular nuclei located:

a) Beneath the lateral part of floor of fourth ventricle
b) In the olive
c) In the cerebellum
d) Beneath the medial part of floor of fourth ventricle
e) In the midbrain

Q47) The development of papilloedema in the case of

b) mitochondria
c) Golgi complex
d) nucleus
e) rough endoplasmic reticulum

Q46) Which of the layers of epidermis comprises keratinized dead cells

a) Papillary layer
b) Stratum spinosum
c) Stratum corneum
d) Stratum basale
e) Stratum granulolsm

Q47) THE SKELETAL MUSCLE FIBRES:

a) posses diads associated with contraction.
b) are multinucleated.
c) are cylindrical and branching.
d) have T-tubules located at Z-disks.
e) have gap junctions.

Q48) A ring shaped section of small intestine could be

a) Coronal Section
b) Coronal section
c) Transverse section
d) Median section
e) Sagittal section

Q49) The dural venous sinuses related to tentorium cerebelli are all except

a) Straight sinus
b) Superior petrosal sinus
c) Inferior petrosal sinus
d) Right transverse sins
e) Left transverse sinus

Q50) Cavenous sinus is situated in

a) Anterior cranial fossa
b) Middle cranial fossa
c) Posterior cranial fossa
d) Inferior cranial fossa
e) Superior cranial fossa

★★★★★★★★★★ THE END ★★★★★★★★★★

b) Masseter
c) Temporalis muscle
d) Lateral pterygoid
e) Medial pterygoid

Q39) Select a wrong statement regarding the filiform papillae of the tongue
a) Are distributed over the entire anterior two third dorsal surface of the tongue
b) Covered by stratified squamous cornified epithelium
c) Are finger like in shape - cone-shaped projection
d) Taste buds are located on their dorsal surface
e) Are the most numerous of the lingual papillae


Q40) The dorsal root ganglion contains cells bodies of
a) Dorsal column of spinal cord
b) Anterolateral tract
c) Anterior spino thalamic tract
d) Anterior spino vestibular tract
e) Lateral spino thalamic tract

Q41) Which of the following statement is incorrect about lateral Medullary syndrome
a) Ipsilateral loss of touch.
b) Ipsilateral Paralysis of vocal cord
c) Ipsilateral loss of pain sensation of the face
d) Dysphonia and dysphargia
e) Ipsilateral Horners syndrome

Q42) The correct statement regarding the microscopic features of cerebral cortex
a) Large pyramidal cells (Betz cells) are found in ganglionic layer
b) Molecular layer consists of stellate cells
c) The axons of Martinotti cells pass in deeper layers of the cortex
d) Size of the cells decreases from superficial to deep layer
e) Multiform layer is the most superficial layer

Q43) Select an odd statement regarding the structure of eye:
a) Ciliary epithelium consist of a single layer of columnar unpigmented cells.
b) Photo receptors of the retina are modified bipolar neurons
c) Inner layers of retina

WhatsApp 05:14 66%
drive.google.com
IMG-20170112-WA0031.jpg



Head & Neck Module Internal Evaluation

26. Surface ectoderm produces which of the following parts in development of eye:
(a) Lens
(b) Corneal stroma
(c) Sclera
(d) Extraocular muscles
(e) Endothelial lining of blood vessels of eye

27. An abscess was surgically removed from the middle of the posterior triangle on the right side. During recovery the patient noticed that her shoulder drooped and she could no longer raise her right hand above her head to brush her hair. Which nerve has been cut?
(a) Accessory (XI)
(b) Ansa cervicalis
(c) Facial (VII)
(d) Hypoglossal (XII)
(e) Suprascapular

28. Which of the following hyoid muscles is an important landmark in both the anterior and posterior triangles of the neck?
(a) geniohyoid
(b) mylohyoid
(c) omohyoid
(d) sternohyoid
(e) stylohyoid


29. The development of thyroid gland is marked by a depression called foramen cecum in the floor of the primitive pharynx. The foramen cecum gives rise to a diverticulum which later elongates and descends anterior to the hyoid bone and thyroid cartilage to assume the adult position.
What is this diverticulum called?
(a) Hyoglossal duct
(b) Laryngeal diverticulum
(c) Respiratory diverticulum
(d) Thyroglossal duct
(e) Thymus

30. cricothyroid muscle is supplied by
(a) recurrent laryngeal nerve
(b) external laryngeal nerve
(c) internal laryngeal nerve
(d) superior laryngeal nerve
(e) medial laryngeal nerve

31. Nerve of third pharyngeal arch is
(a) facial
(b) glossopharyngeal
(c) vagus
(d) hypoglossal
(e) trigeminal

IMG-20 ... 11.jpg
IMG-20 ... 12.jpg

WhatsApp 05:14 66%
drive.google.com
Bcqs



38. **During a gang fight, a 17 – year – old boy is punched, and his nassal septum is broken. Which of the following structure would be damaged?**
(a) Septal cartilage and nasal bone
(b) Inferior concha and vomer
(c) Vomer and perpendicular plate of ethmoid
(d) Cribriform plate and frontal bone
(e) Middle concha and vomer

39. **Xerostomia is a major feature of which of the most likely condition given below:**
(a) Down syndrome
(b) Tuberculosis
(c) Sjogren syndrome
(d) Ulcerative colits
(e) Crohn's diseade

40. **Peeriodontal disease can be a component of which of the following condition:**
(a) Dental caries
(b) Pleomorphic adenoma
(c) Gingivitis
(d) Systemic disease
(e) Mucoepidermoid

41. **Which one of the following is the most common antecedent of leukoplakia:**
(a) EBV
(b) HPV
(c) Tabacco
(d) Radiations
(e) Fungi.

42. **Rhodopsin is a light sensitive pigment. It contains protein called as**
(a) Opsin
(b) Keratin
(c) Collagen
(d) Casein
(e) Renin

43. **Aqueous humor is mainly formed by the process of**
(a) active secretion
(b) osmosis
(c) bulk transport
(d) passive diffusion
(e) facilitated diffusion

IMG-20 ... 11.jpg
IMG-20 ... 12.jpg

c) Myelencephalon
d) Rhombencephalon
e) Telencephalon

Q14) THE CARDIAC MUSCLE FIBRES:
a) have triads.
b) show absence of gap junctions.
c) are cylindrical and branching.
d) have regenerative ability.
e) contain nuclei in the periphery.

Q15) A PERSON IS UNABLE TO SEE THE FOLLOWING ST
MICROSCOPE:
a) mitochondria
b) cell membrane.
c) nucleus
d) Golgi complex.
e) rough endoplasmic reticulum.

Q16) A SPECIMEN OF LYMPHOID ORGAN UNDER MICRO
IT IS COVERED BY STRATIFIED SQUAMOUS EPITH
LIKELY TO BE:
a) palatine tonsil.
b) thymus
c) lymph node.
d) tubal tonsil
e) spleen

Q17) Axon Hillock is the site in neurons with no;
a) Cytoplasm
b) Granules
c) Cellular pale
d) Dendrites
e) Axon.

Q18) tendons are composed of _____ connective
a) Skeletal
b) Yellow elastic
c) Dense irregular
d) Dense regular
e) Areolar line

Q19) Blood vessels are lined with _____ epitheliu
a) Transitional
b) Stratified cuboidal
c) Stratified aquamous
d) Simple columnar
e) Simple Squamous

Q20) IN CASE OF PLACENTA, FOLLOWING OPTION IS
a) maternal surface is smooth and covered by amnion
b) fetal part is formed by chorionic frondosum.
c) contains cotyledons on it maternal surface.
d) umbilical cord is attached near its center.
e) it weighs 500 grams.

Q21) Conus Medullaris is the inferior apex of the:
a) Spinal cord
b) Cerebellum
c) Sympathetic chain
d) Cerebrum
e) Brain stain

Q22) THE MAJOR CHARACTERISTIC OF MEIOSIS-1 IS
a) producing the primordial germ cells

a) V cranial nerve
b) VI cranial nerve
c) III cranial nerve
d) II cranial nerve
e) IV cranial nerve

Q18) Select an incorrect statement regarding the structure of layer of the cerebral cortex

a) Inner band of Baillarger is present in the external pyramidal layer
b) External granular layer contains large number of small pyramidal cells
c) Martinotti cells are present in the multiform and ganglionic layers
d) Horizontal cells are present in molecular layer
e) Outer band of Baillarger is present in the internal granular layer

Q19) Herniation of the intervertebral disc between the fifth and sixth cervical vertebrae compress the

a) Fourth cervical nerve root
b) Sixth, seventh and eight cervical nerve roots
c) Seventh cervical nerve root
d) Fifth cervical nerve root
e) Sixth cervical nerve root

Q20) Select an odd statement regarding the structures of the oral cavity.

a) Superior surface of soft palate is lined by pseudostratified columnar ciliated epithelium.
b) Taste buds are located on the dorsal surface of the filiform papillae.
c) There are no sebaceous or sweat glands in the red area of lips.
d) Von Ebner glands of tongue are serous in nature.
e) In tongue, muscle bundles cross each other running in longitudinal, transverse and vertical

21) Regarding the glial cells

a) Fibrous astrocytes are found in gray matter
b) Microglia are found in 'PNS'
c) Protoplasmic astrocytes form scar tissue in CNS in case of any damage

c) Temporalis
d) Lateral pterygoid
e) Masseter

5) **The best statement regarding the microscopic features of cerebellar cortex**
a) The Purkinje layer consists of flask shaped cells with highly developed dendritic tree
b) It consists of six layers of cells
c) In granuler layer the cells are dispersed
d) In medullary layer cells are compactly arranged
e) The afferent fibers terminate in granular layer

6) **All the features develops during fourth week except**
a) Neural tube closes.
b) Bucco-pharyngeal membrane break down.
c) Most somites are formed.
d) Primary centre of ossification appears in the skull.
e) Optic vesicle first appears.

7) **Select an odd statement regarding the structure of eye:**
a) Ciliary epithelium consist of a single layer of columnar unpigmented cells.
b) A single layer of cuboidal cells is present all over just under the lens capsule.
c) Descemet's membrane of cornea is highly refractile under the light microscope.
d) Inner layers of retina are absent in the fovea centralis.
e) Photo receptors of the retina are modified bipolar neurons.

8) **Which of the following are derivatives from neural crest cells <u>except</u>?**
a) Schwann cells
b) Ganglion of sympathetic trunk
c) Melanocytes of skin
d) Cortex of adrenal gland
e) Dorsal root ganglion

9) **The correct statement regarding the microscopic features of cerebral cortex**
a) Size of the cells decreases from superficial to deep layer
b) Large pyramidal cells (Betz cells) are found in ganglionic layer
c) The axons of Martinotti cells pass in deeper layers of the cortex
d) Multiform layer is the most superficial layer
e) Molecular layer consists of stellate cells

0) **Foot plate of stapes is placed at**
a) Opening of pyramid
b) Tubal opening
c) Round window (fenestra cochlear)
d) Oval window (fenestra vestibuli)
e) Aditus

hown by Nissl staining method,

ctate or "molecular" appeara
The molecular layer is essentially
of the cerebral cortex. Neuroglia

d) It is common site of meningeal herniation
e) It is on the scalp

Q16) Spinous process is absent in which of the following vertebra

a) 2nd cervical vertebra
b) 5th cervical vertebra
c) 1st cervical vertebra
d) 3rd cervical vertebra
e) 7th cervical vertebra

Q17) Lateral rectus of eye is supplied by

a) V cranial nerve
b) VI cranial verve
c) III cranial nerve
d) II cranial nerve
e) IV cranial nerve

Q18) Select an incorrect statement regarding the structure of layer of the cerebral cortex

a) Inner band of Baillarger is present in the external pyramidal layer
b) External granular layer contains large number of small pyramidal cells
c) Martinotti cells are present in the multiform and ganglionic layers
d) Horizontal cells are present in molecular layer
e) Outer band of Baillarger is present in the internal granular layer

9) Herniation of the intervertebral disc between the fifth and sixth cervical vertebrae will compress the

a) Fourth cervical nerve root
b) Sixth, seventh and eight cervical nerve roots
c) Seventh cervical nerve root
d) Fifth cervical nerve root
e) Sixth cervical nerve root

Q4) The derivatives of alar plate of the neural tube are the
a) Basal ganglia of cerebrum
b) Pre vertebral sympathetic ganglia
c) Para vertebral sympathetic ganglia
d) Trigeminal ganglia
e) Dorsal root ganglia of spinal cord

Q5) The Thalamus forms the following boundary of 3RD ventricle:
a) Lateral
b) Medial
c) Superior
d) Anterior
e) Posterior

Q6) Regarding the tympanic cavity
a) Tensor tympani is the only muscle which is present in the tympanic cavity
b) It is the air containing part of ear
c) A cranial nerve is related to tympanic cavity
d) It contains small bones
e) Auditory tube extends from the anterior wall of tympanic cavity to naso pharynx

Q7) The roots of first cervical spinal nerve corresponds to the
a) level of formen magnum
b) atlas vertebra
c) upper edge of spinal cord
d) junction of medulla and spinal cord
e) lower edge of medulla oblongata

Q8) The following fibre lies immediately beneath the cortex and connect the adjacent gyri:
a) Long association fibres
b) Uncinate fasciculus
c) Inferior longitudinal fasciculus

Q7) ANATOMICAL POSITION OF THE BODY IN... EXCEPT THE:

a) feet are held together.
b) person is standing erect.
c) palms are facing towards body (medially). ✗
d) person is looking straight forward.
e) arms are hanging by sides.

Q8) ANTIBODIES ARE SYNTHESIZED AND SECRETED BY:

a) T cytotoxic cells.
b) T memory cells.
c) plasma cells.
d) T helper cells.
e) B memory cells.

Q9) Posterior root ganglia contains cell bodies of;

a) Pre-ganglionic neurons
b) Central motor neurons
c) Peripheral sensory neurons
d) Peripheral motor neurons
e) Central sensory neurons

Q10) The midbrain is a part of:

a) Cerebrum.
b) Diencephalon.
c) Pons.
d) Brain stem.
e) Cerebellum

1ST YEAR

Q78) The development of papilloedema in the case of increased intracranial pressure, is the following feature of optic nerve:

a) It is an extension of white matter of brain
b) It meets the sclera 3mm medial to posterior pole
c) It is surrounded by subarachnoid space as far as back of eye
d) It enters the orbit through optic canal
e) It receives the central artery of retina

Q79) Longest spinous process is the characteristics of

a) 1st thoracic vertebra
b) 7th cervical vertebra
c) 1st cervical vertebra
d) 12th thoracic vertebra
e) 2nd cervical vertebra

Q80) The dorsal roots of spinal nerves are derived from the following structure

a) The posterior gray column
b) Peripheral process of posterior root ganglia
c) The autonomic ganglia
d) White ramus communicans
e) Central process of posterior root ganglia

Q81) With swelling of mucous membrane of superior nasal meatus, which of the openin plugged?

a) Sphenoidal sinus
b) Frontal sinus
c) Anterior ethmoidal sinus
d) Superior Para nasal sinus
e) Middle ethmoidal sinus

Q82) The accurate statement about lacrimal gland is:

a) It curls around the medial margin of levator palpebrae superioris
b) Its orbital part is visible through superior fornix of conjunctiva
c) It is drained by ducts draining from palpebral part into superior fornix
d) Its palpebral part lies in lacrimal fossa
e) Secremotor fibers to the gland from inferior salivary nucleus travel in greater petrosal nerve

Q83) The tract which is concerned with reflex postural movements in response to visua stimuli is

a) Rubro spinal tract
b) Tecto spinal tract
c) Vestibulo spinal tract
d) Olivospinal tract
e) Corticospinal tract

Q84) A muscle which elevates and retracts the mandible is

a) Digastric
b) Medial pterygoid
c) Temporalis
d) Lateral pterygoid
e) Masseter

Q85) The best statement regarding the microscopic features of cerebellar cortex

a) The Purkinje layer consists of flask shaped cells with highly developed dendritic tree
b) It consists of six layers of cells
c) In granuler layer the cells are dispersed
d) In medullary layer cells are compactly arranged
e) The afferent fibers terminate in granular layer

Q86) All the features develops during fourth week except

a) Neural tube closes.
b) Bucco-pharyngeal membrane break down.
c) Most somites are formed.
d) Primary centre of ossification appears in the skull.
e) Optic vesicle first appears.

Q87) Select an odd statement regarding the structure of eye:

a) Ciliary epithelium consist of a single layer of columnar unpigmented cells.
b) A single layer of cuboidal cells is present all over just under the lens capsule.
c) Descemet's membrane of cornea is highly refractile under the light microscope.
d) Inner layers of retina are absent in the fovea centralis.
e) Photo receptors of the retina are modified bipolar neurons.

Q88) Which of the following are derivatives from neural crest cells <u>except</u>?

a) Schwann cells
b) Ganglion of sympathetic trunk
c) Melanocytes of skin
d) Cortex of adrenal gland
e) Dorsal root ganglion

Q89) The correct statement regarding the microscopic features of cerebral cortex

a) Size of the cells decreases from superficial to deep layer
b) Large pyramidal cells (Betz cells) are found in ganglionic layer
c) The axons of Martinotti cells pass in deeper layers of the cortex
d) Multiform layer is the most superficial layer
e) Molecular layer consists of stellate cells

Q90) Foot plate of stapes is place[illegible]

a) Opening of pyramid
b) Tubal opening

Q13) The forebrain is divided into cerebrum and:

a) Diencephalon
b) Mesencephalon
c) Myelencephalon
d) Rhombencephalon
e) Telencephalon

Q14) THE CARDIAC MUSCLE FIBRES:

a) have triads.
b) show absence of gap junctions.
c) are cylindrical and branching.
d) have regenerative ability.
e) contain nuclei in the periphery.

Q15) A PERSON IS UNABLE TO SEE THE FOLLOWING STRUCTURE UNDER L MICROSCOPE:

a) mitochondria
b) cell membrane.
c) nucleus
d) Golgi complex.
e) rough endoplasmic reticulum.

Q16) A SPECIMEN OF LYMPHOID ORGAN UNDER MICROSCOPE SHOWED T IT IS COVERED BY STRATIFIED SQUAMOUS EPITHELIUM. THIS ORGA LIKELY TO BE:

a) palatine tonsil.
b) thymus
c) lymph node.
d) tubal tonsil

Q26) A simplest structure considered to be alive are
a) Cells
b) Organs
c) Organelles
d) Tissues
e) Organisms

Q27) THE HYALINE CARTILAGE:
a) is devoid of perichondrium.
b) contains type-1 collagen.
c) is present on articular surfaces of synovial joints.
d) is the vascular tissue.
e) is capable of extensive turnover and repair.

Q28) THE MOST IMPORTANT EVENT FOR IMPLANTATION OF THE FETILIZED OVUM IS THE:
a) disintegration of zona pellucida.
b) entry of morula in uterine cavity.
c) formation of decidua.
d) formation of blastocyst
e) cleavage

Q29) THE MOST COMMON SITE FOR ABNORMAL IMPLANTATION IS THE:
a) uterine tube.
b) mesentery.
c) lower uterine segment
d) pouch of Douglas.
e) ovary.

Q30) Mid brain connects the forebrain to:
a) Pons.
b) Hind brain.
c) Spinal cord.
d) Cerebellum.
e) Brain stem.

Q31) Cerebral aqueduct passes through:
a) Medulla oblongatae
b) Cerebrum
c) Pons
d) All along brain stem
e) Mid brain.

Q32) PREGNANCY CAN BE DOCUMENTED IN A FEMALE IN SECOND WEEK OF DEVELOPMENT BY:
a) pelvic X-Ray (radiography)
b) assay of human chorionic gonadotropin in urine
c) C.T scan.
d) ultrasound
e) assay of serum progesterone in the blood.

Q33) Post ganglionic sympathic fibers innervate the following viscera except
a) Adrenal medulla
b) Lungs
c) Heart
d) Duodenum
e) Stomach

Q34) Cell - bodies of preganglionic sympathetic fibers all present in lateral gray born at the level between:
a) T_1 to L_2
b) S_2 S_3 S_4
c) T_4 to L_2
d) C_1 to C_4
e) S_1 S_2 S_3

Q35) Regarding the type of muscle contraction:
a) Muscle do not change its length during eccentric contraction
b) Muscle decrease in length during eccentric contraction
c) Muscle exerts force with changing the length in isometric contraction
d) Muscle exerts force without changing length in isometric contraction
e) Muscle increase in length in isotonic contraction

Q36) SIMPLE SQUAMOUS EPITHELIUM IS ABSENT IN THE FOLLOWING ORGANS:
a) serous membranes.
b) lung alveoli.
c) blood vessels.
d) loops of Henle of the kidney.
e) distal convoluted tubules of the kidney.

Q37) Which of the following is not an essential part of anatomical position
a) Forearms supine
b) Feet flat on the floor
c) Mouth closed
d) Arms down to the sides
e) Eyes facing forward

Q38) SECONDARY OOCYTE COMPLETES ITS SECOND MEIOTIC DIVISION TO BECOME A MATURE OVUM:
a) at puberty.
b) before ovulation
c) at ovulation.
d) at fertilization.
e) before fertilization.

Pearl Papers
HEALTH SCIENCES
P.U.H.S

tos

Zoom Rotate Draw Edit

true statement regarding the spinal cord:

a. It's the long cylindrical upper part of CNS.
b. It occupies the upper 2/3rd of vertebral canal
c. It gives rise 31 pairs of cranial nerve.
d. Superiorly it is continuous with filumterminale.
e. It ends at upper border of first lumber vertebrae.

Myelin of the peripheral nerves is produced by:

a) Fibroblast
b) Macrophage
c) Schwann cell
d) Oligodendrocyte
e) Muller cells

part of the brain-ster

Q13) The characteristic feature about the pericyte is
a) They have a contractile function
b) They are present in the basement membrane of vessels
c) These cells surround the arterioles
d) These cells are developed from ectoderm
e) They are spherical in shape

Q14) The anastomosis between external and internal carotid arteries include all of the following except
a) Dorsal nasal artery and facial artery
b) Greater palatine and superior labial artery
c) Anterior ethamoidal artery and facial artery
d) Lateral nasal artery and supratrochlear artery
e) Lacrimal artery and transverse facial artery

Q15) The clinical importance of anterior frontale is
a) It is the site of fusion of parietal bones with frontal bone
b) It bulges in case of raised intracranial pressure
c) Two parietal bone meet one another at this point
d) It is common site of meningeal herniation
e) It is on the scalp

Q16) Spinous process is absent in which of the following vertebra
a) 2nd cervical vertebra
b) 5th cervical vertebra
c) 1st cervical vertebra
d) 3rd cervical vertebra
e) 7th cervical vertebra

Q17) Lateral rectus of eye is supplied by
a) V cranial nerve
b) VI cranial nerve
c) III cranial nerve
d) II cranial nerve
e) IV cranial nerve

Q18) Select an incorrect statement regarding the structure of layer of the cerebral cortex
a) Inner band of Baillarger is present in the external pyramidal layer
b) External granular layer contains large number of small pyramidal cells
c) Martinotti cells are present in the multiform and ganglionic layers
d) Horizontal cells are present in molecular layer
e) Outer band of Baillarger is present in the internal granular layer

Q19) Herniation of the intervertebral disc between the fifth and sixth cervical vertebrae will compress the
a) Fourth cervical nerve root
b) Sixth, seventh and eight cervical nerve roots
c) Seventh cervical nerve root
d) Fifth cervical nerve root
e) Sixth cervical nerve root

Q20) Select an odd statement regarding the structures of the oral cavity.
a) Superior surface of soft palate is lined by pseudostratified columnar ciliated epithelium.
b) Taste buds are located on the dorsal surface of the filliform papillae.
c) There are no sebaceous or sweat glands in the red area of lips.
d) Von Ebner glands of tongue are serous in nature.
e) In tongue, muscle bundles cross each other running in longitudinal, transverse and vertical directions.

Q21) Regarding the glial cells
a) Fibrous astrocytes are found in gray matter
b) Microglia are found in 'PNS'
c) Protoplasmic astrocytes form scar tissue in CNS, in case of any damage
d) Ependymal cells are of simple squamous variety
e) Schawn cells form myelin in central nervous system

Q22) Occulomotor nucleus is situated at which of the following level
a) Medulla oblongata
b) Tegumentum pones
c) Superior colliculus
d) Cerebello potine angle
e) Inferior colliculus

Q23) Which of the following tracts does not cross the midline
a) Rubro-spinal tract
b) Pyramidal Tract
c) Dento thalamic tract
d) Spinothalamic tract
e) Dorsal spinocerebellar tract

Q24) Considering the myelination of nerve fiber
a) Unmyelinated nerve fibers are not covered by Schwann cells
b) The nucleus of Schwann cell is present at the node of Ranvier
c) Myelin is formed by the cytoplasm of Schwann cells
d) Myline is stained black in routine histological stains i.e H & E
e) Internode is the distance between two nodes

Q25) Regarding mid brain
a) Origin to 4th cranial nerve in its upper part
b) Cerebral peduncle lies posterior
c) Closely related to free border of falx cerebelli
d) Superior colliculi receive auditory impulses
e) Contains cerebral aqueduct

11. Which is the derivative of 2nd pharyngeal arch:
    a. Malleus
    b. Incus
    c. Auditory tube
    d. Stapes
    e. Middle ear cavity

12. Which is the derivative of 1st pharyngeal arch:
    a. Stapedius muscle
    b. Tympanic membrane
    c. Malleus
    d. Auricle
    e. External auditory meatus

13. Utricle, semisulcular ducts, vestibular ganglion of VIII cranial nerve are derived from:
    a. Otic vesicle
    b. Auricular hillocks
    c. Pharyngeal groove
    d. 1st pharyngeal arch
    e. Pharyngeal membrane

14. Auditory tube is supplied by:
    a. Middle menengial artery
    b. Artery of pterigoid canal
    c. Ascending palatine artery
    d. A & B
    e. A & C

15. The saccule, cochlear duct, spiral ganglion of VIII cranial nerve are derived from:
    a. Utrical portion of otic vesicle
    b. Saccular portion of otic vesicle
    c. Auricular hillocks
    d. Pharyngeal groove
    e. Pharyngeal arch 1st

16. The blood supply of pinna comes from:
    a. External carotid artery
    b. Internal carotid artery
    c. Both A & B
    d. Facial artery
    e. Maxillary artery

78) **The development of papilloedema in the case of increased intracranial pressure, is due to the following feature of optic nerve:**
a) It is an extension of white matter of brain
b) It meets the sclera 3mm medial to posterior pole
c) It is surrounded by subarachnoid space as far as back of eye
d) It enters the orbit through optic canal
e) It receives the central artery of retina

Q79) **Longest spinous process is the characteristics of**
a) 1st thoracic vertebra
b) 7th cervical vertebra
c) 1st cervical vertebra
d) 12th thoracic vertebra
e) 2nd cervical vertebra

Q80) **The dorsal roots of spinal nerves are derived from the following structure**
a) The posterior gray column
b) Peripheral process of posterior root ganglia
c) The autonomic ganglia
d) White ramus communicans
e) Central process of posterior root ganglia

Q81) **With swelling of mucous membrane of superior nasal meatus, which of the opening is plugged?**
a) Sphenoidal sinus
b) Frontal sinus
c) Anterior ethmoidal sinus
d) Superior Para nasal sinus
e) Middle ethmoidal sinus

Q82) **The accurate statement about lacrimal gland is:**
a) It curls around the medial margin of levator palpebrae superioris
b) Its orbital part is visible through superior fornix of conjunctiva
c) It is drained by ducts draining from palpebral part into superior fornix
d) Its palpebral part lies in lacrimal fossa
e) Secremotor fibers to the gland from inferior salivary nucleus travel in greater petrosal nerve

Q83) **The tract which is concerned with reflex postural movements in response to visual stimuli is**
a) Rubro spinal tract
b) Tecto spinal tract
c) Vestibulo spinal tract
d) Olivospinal tract
e) Corticospinal tract

Q84) **A muscle which elevates and retracts the mandible is**
a) Digastric
b) Medial pterygoid
c) Temporalis
d) Lateral pterygoid
e) Masseter


Q85) **The best statement regarding the microscopic features of cerebellar cortex**
a) The Purkinje layer consists of flask shaped cells with highly developed dendritic tree
b) It consists of six layers of cells
c) In granuler layer the cells are dispersed
d) In medullary layer cells are compactly arranged
e) The afferent fibers terminate in granular layer

Q86) **All the features develops during fourth week except**
a) Neural tube closes.
b) Bucco-pharyngeal membrane break down.
c) Most somites are formed.
d) Primary centre of ossification appears in the skull.
e) Optic vesicle first appears.

Q87) **Select an odd statement regarding the structure of eye:**
a) Ciliary epithelium consist of a single layer of columnar unpigmented cells.
b) A single layer of cuboidal cells is present all over just under the lens capsule.
c) Descemet's membrane of cornea is highly refractile under the light microscope.
d) Inner layers of retina are absent in the fovea centralis.
e) Photo receptors of the retina are modified bipolar neurons.

Q88) **Which of the following are derivatives from neural crest cells <u>except</u>?**
a) Schwann cells
b) Ganglion of sympathetic trunk
c) Melanocytes of skin
d) Cortex of adrenal gland

WhatsApp 05:12 69%
drive.google.com
Bcqs



15. **Brocas aarea is situated** in:
A. Superior temporal gyrus
B. Inferior temporal gyrus
C. Superior frontal gyrus
D. Inferior frontal gyrus

16. **Fine motor movement** is controlled by:
A. Cerebellum
B. Basal ganglia
C. Cerebrum
D. None

17. **Area of sound perception** is located in:
A. In the cortex in temporal lobe
B. Frontal lobe
C. Occipital lobe
D. None

18. **Which of the following** is not the part of brain stem:
A. Mid brain
B. Pons
C. Medulla
D. Cerebellum


19. **Patient presents with intention tremors.** The lesion is in:
A. Cerebellum
B. Basal ganglia
C. Cerebrum
D. Subthalamic nucleus

20. **Regarding diencephalon which statement is true?**
A. Forms the medial wall of third ventricle
B. Includes thalamus geniculate body,hypothalamus
C. Is apart of mesencephalon
D. None

The End

IMG-20 ... 11.jpg IMG-20 ... 12.jpg

WhatsApp 05:12 69%
drive.google.com



the correct statement regarding Filumterminale
enlargement of cervical segment
enlargement of Lumbar segment
cord is taper off
gation of Pia matter
gation of Dura matter

following nerve cell groups is NOT present in Posterior colu
a. Substantiagelatinosa
b. Nucleus Proprius
c. Nucleus dorsalis
d. Visceral afferent nucleus
e. Visceral efferent nucleus

Q42) Mid Brain is traversed by a narrow channel:
a. Interventricular Foramen
b. Cerebral aqueduct
c. Foramen Lushka
d. Foramen Magendie
e. Central canal

Q43) The Superior colliculi are centers for
a. Auditory reflexes
b. Visual reflexes
c. Motor reflexes
d. Medial lemniscus
e. Sensory reflexes

Q 44) The inferior colliculi are centers for
a. Auditory reflexes
b. Visual reflexes
c. Sensory reflexes
d. Motor reflexes
e. Medial lemniscus

Q 45) Which of the following is an unpaired venous sinus
a. Superior sinus
b. Sigmoid sinus
c. Cavernous sinus
d. Transverse sinus
e. Sphenoparietal sinus

Q 46) What nerve is found inferolateral to internal carotid artery within the cavernous sinus?
a. Frontal nerve
b. Lacrimal nerve
c. Trochlear nerve
d. Abducent nerve
e. Opthalmic nerve

Q 47) Which of the following structure is NOTincluded in Basal Ganglia?
a. Thalamus
b. Putamen
c. Pallidum
d. Claustrum
e. Lentiform nucleus

Q 48) The largest bundle of commissural fibers in brain is called?
a. Anterior Commissure
b. Posterior Commissure
c. Habenular Commissure
d. Hippocampal Commissure
e. Corpus callosum

The Primary motor area is located in which gyrus of brain?
a. Precentral gyrus
b. Postcentral gyrus
c. Superior frontal gyrus
d. Middle frontal gyrus
e. Superior temporal gyrus

Inferior horn of lateral ventricle extends into which lobe of cerebral hemisphere?

* * * * * * * * * THE END * * * * * * *

Share Bookmarks Tabs

a) Positive Quenkenstedt's sign when pressure rises (CSF)
b) In Prone position
c) For histopathological examination
d) Vertebral column is well extended
e) Normal pressure is about 60 to 150 mm of water

Q66) Select a wrong statement regarding the structure of exte

a) Ceruminous glands are found in the submucosa
b) External auditory meatus is lined by stratified squamous epithe
c) Ceruminous glands are a type of modified sebaceous gland
d) External surface of eardrum is lined with epidermis
e) Pinna is composed of a plate of elastic cartilage

Q67) Select an odd statement regarding the structure of nose:

a) Superior conchae are lined by olfactory mucosa.
b) Smell bodies are more prominent over the inferior conchae.
c) Poor population of goblet cells is present in the mucosa.
d) Vibrissae are present in the skin vestibule.
e) Many simple branched tubulo-alveolar glands are present in th

Q68) The important characteristic feature of mossy fibres in t

a) They are terminal fibers of all cerebral afferent tracts
b) They have multiple branches
c) They exert a much more diffuse excitatory effect
d) The deep cerebellar nuclei receive afferent information from
e) A single mossy fibre may stimulate thousands of Purkinje cel

Q69) All the structures contain fibers of corticospinal tracts e

a) Posterior limb of internal capsule
b) Tegmentum of medulla
c) Pyramids of medulla
d) Crus cerebri of mid brain
e) Lateral white column of spinal cord

Q70) Which of the following structure does not take part in f

a) Left maxillary Proces
b) Medial nasal proiminences
c) Lateral nasal proiminences
d) Right maxillary process
e) The upper jaw

Q71) Select an odd statement regarding the structure of cer

a) Unmyelinated axon of a basket cell ascends toward the mole
b) Mossy fibers originate from the inferior olivary nucleus of t
c) Four or more branching dendrites pass outward from the bas
d) Axons of Golgi type II cells end by making synapses with t
e) Cytoplasm of a granule cell contains large number of Nissl
concentrically around the nucleus.

Q72) Pulsation of facial artery can be easily felt at

a) Body of masseter
b) Posterior border of buccinator
c) Anterior border of temporalis
d) Posterior border of masseter
e) Anterior border of masseter

Q73) The pterion is the point

a) Is 1.5 inches above the mid point of zygomatic arch
b) It is marked by an eminence
c) It is ,marked by a depression
d) Where lesser wing of sphenoid bone meets with parietal b
e) Posterior branch of middle meningeal artery is related to i

Q74) Which of following level of spinal cord involve in par

a) S_1 and S_2
b) L_5 and S_1
c) S_3 , S_4 And S_5

Q15. Asim, a 5 years old boy falls from his bicycle and receives a cut over the knee. His tissue macrophages & fibroblasts will move to the site of injury for repair by:

a) Ciliary movement
b) Ameboid movement
c) Chemotaxis
d) Margination
e) Whip- like movement

Q16. Red blood cells can pass from bone marrow to blood capillaries by diapedesis during:

a) Basophil erythroblast stage
b) Mature erythrocyte stage
c) Reticulocyte stage
d) Pro-erythroblast stage
e) Orthochromatic erythroblast stage

Q17. Cell mediated immunity is caused by

a) Red blood cell
b) Plasma cell
c) T cell
d) B lymphocytes
e) Platelets

Q18. A blood transfusion reaction is likely to occur if a patient with blood group A+ is transfused of group:

a) A+
b) AB-
c) A-
d) O+
e) O-

Q19. Which of the following factor is a potent stimulator of RBC production:

a) Increased tissue oxygenation
b) Decreased tissue oxygenation
c) Barometric pressure at sea level
d) Normal pulmonary functions
e) Normal cardiac function

Q20. Intracellular negativity of charge is due to presence of

a) Intracellular Cl- ions
b) Intracellular HCO_3^- ions
c) Intracellular Proteins
d) Intracellular carbohydrates
e) Intracellular sodium ions

Q21. Different substances that make up the cell are collectively called

a) Ectoplasm
b) Endoplasm
c) Cytoplasm
d) Protoplasm
e) Nucleoplasm

Q22. The cell which contain leas[illegible]ty of water is

a) Cardiac cell
b) Nerve cell

32. Taste buds are present in following papillae except
(a) Fungiform
(b) Circumvallate
(c) Filliform
(d) Foliate
(e) Fulliform

33. The posteromedial surface of the parotid gland is related to:
(a) Masseter.
(b) Ramus of mandible.
(c) Temporomandibular joint.
(d) Styloid process.
(e) Branches of the facial nerve

34. Which of the following muscle assists in the elevation of the mandible?
(a) Lateral pterygoid.
(b) Medial pterygoid.
(c) Digastric.
(d) Genohyoid.
(e) Mylohyoid

35. The lateral wall of the infratemporal fossa is formed by:
(a) Lateral pterygoid plate.
(b) Tensor palatine muscle.
(c) Zygomatic arch.
(d) Ramus of mandible.
(e) Carotid sheath

36. A 45 year – old woman is suffering from numbness over the tip of her nose. Which of the following nerve is most likely to be damaged?
(a) Ophthalmic division of the trigeminal nerve
(b) Maxillary division of the trigeminal nerve
(c) Mandibular division of the trigeminal nerve
(d) Auriculotemporal nerve
(e) Auricular nerve

37. After having a tonsillectomy, a 57- year- old man with a long history of chewing tobacco use is unable to detect taste on the posterior one – third of his tongue. Which of the following nerve has most likely injured?
(a) Internal laryngeal nerve
(b) Lingual nerve
(c) Glossopharyngeal nerve
(d) Chorda nerve
(e) Facial nerve

26. Surface ectoderm produces which of the following parts in the development of eye:

(a) Lens
(b) Corneal stroma
(c) Sclera
(d) Extraocular muscles
(e) Endothelial lining of blood vessels of eye

27. An abscess was surgically removed from the middle of the posterior triangle on the right side. During recovery the patient noticed that her shoulder drooped and she could no longer raise her right hand above her head to brush her hair. Which nerve has been cut?

(a) Accessory (XI)
(b) Ansa cervicalis
(c) Facial (VII)
(d) Hypoglossal (XII)
(e) Suprascapular

28. Which of the following hyoid muscles is an important landmark in both the anterior and posterior triangles of the neck?

(a) geniohyoid
(b) mylohyoid
(c) omohyoid
(d) sternohyoid
(e) stylohyoid

29. The development of thyroid gland is marked by a depression called foramen cecum in the floor of the primitive pharynx. The foramen cecum gives rise to a diverticulum which later elongates and descends anterior to the hyoid bone and thyroid cartilage to assume the adult position.
What is this diverticulum called?

(a) Hyoglossal duct
(b) Laryngeal diverticulum
(c) Respiratory diverticulum
(d) Thyroglossal duct
(e) Thymus

30. cricothyroid muscle is supplied by

(a) recurrent laryngeal nerve
(b) external laryngeal nerve
(c) internal laryngeal nerve
(d) superior laryngeal nerve
(e) medial laryngeal nerve

31. Nerve of third pharyngeal arch is

(a) facial
(b) glossopharyngeal
(c) vagus
(d) hypoglossal
(e) trigeminal

the correct statement regarding Filumtermina
[illegible] enlargement of cervical segment
[illegible]um enlargement of Lumbar segment
[illegible]inal cord is taper off
[illegible]ntegation of Pia matter
[illegible]ntegation of Dura matter

following nerve cell groups is NOT present in Posterior colu
a. Substantiagelatinosa
b. Nucleus Proprius
c. Nucleus dorsalis
d. Visceral afferent nucleus
e. Visceral efferent nucleus

Q42) Mid Brain is traversed by a narrow channel:
a. Interventricular Foramen
b. Cerebral aqueduct
c. Foramen Lushka
d. Foramen Magende
e. Central canal

Q 43) The Superior colliculi are centers for:
a. Auditory reflexes
b. Visual reflexes
c. Motor reflexes
d. Medial leminiscus
e. Sensory reflexes

Q 44) The inferior colliculi are centers for
a. Auditory reflexes
b. Visual reflexes
c. Sensory reflexes
d. Motor reflexes
e. Medial leminiscus

Q 45) Which of the following is an unpaired venous sinus
a. Straight sinus
b. Sigmoid sinus
c. Cavernous sinus
d. Transverse sinus
e. Sphenopalatine sinus

Q 46) What nerve is found inferolateral to internal carotid artery within the cavernous sinus?
a. Frontal nerve
b. Lacrimal nerve
c. Trochlear nerve
d. Abducent nerve
e. Ophthalmic nerve

Q 47) Which of the following structure is NOTincluded in Basal Ganglia?
a. Thalamus
b. Striatum
c. Pallidum
d. Claustrum
e. Lentiform nucleus

Q 48) The largest bundle of commissural fibers in brain is called?
a. Anterior Commissure
b. Posterior Commissure
c. Habenular Commissure
d. Corpus callosum
e. Fornix

The Primary motor area is located in which gyrus of brain?
a. Precentralgyrus
b. Postcentralgyrus
c. Middle frontal gyrus
d. Superior temporal gyrus
e. Middle temporal gyros

Interior horn of lateral ventricle extends into which lobe of cerebral hemisphere?
frontal lobe
parietal lobe
occipital lobe
temporal lobe
insula

* * * * * * * * * THE END * * * * * *

15. **Brocas aarea is situated** in:
A. Superior temporal gyrus
B. Inferior temporal gyrus
C. Superior frontal gyrus
D. Inferior frontal gyrus

16. **Fine motor movement** is controlled by:
A. Cerebellum
B. Basal ganglia
C. Cerebrum
D. None

17. **Area of sound perception** is located in:
A. In the cortex in temporal lobe
B. Frontal lobe
C. Occipital lobe
D. None

18. **Which of the following** is not the part of brain stem:
A. Mid brain
B. Pons
C. Medulla
D. Cerebellum

19. **Patient presents with intention tremors.** The lesion is in:
A. Cerebellum
B. Basal ganglia
C. Cerebrum
D. Subthalamic nucleus

20. **Regarding diencephalon which statement** is true?
A. Forms the medial wall of third ventricle
B. Includes thalamus,geniculate body,hypothalamus
C. Is apart of mesencephalon
D. None

The End

Share
Edit
Delete

0 KB/s
66%
10:16 pm
MORE

11. Which is the derivative of 2nd pharyngeal arch:
    a. Malleus
    b. Incus
    c. Auditory tube
    d. Stapes
    e. Middle ear cavity

12. Which is the derivative of 1st pharyngeal arch:
    a. Stapedius muscle
    b. Tympanic membrane
    c. Malleus
    d. Auricle
    e. External auditory meatus

13. Utricle, semisulcular ducts, vestibular ganglion of VIII cranial nerve are derived from:
    a. Otic vesicle
    b. Auricular hillocks
    c. Pharyngeal groove
    d. 1st pharyngeal arch
    e. Pharyngeal membrane

14. Auditory tube is supplied by:
    a. Middle menengial artery
    b. Artery of pterigoid canal
    c. Ascending palatine artery
    d. A & B
    e. A & C

15. The saccule, cochlear duct, spiral ganglion of VIII cranial nerve are derived from:
    a. Utrical portion of otic vesicle
    b. Saccular portion of otic vesicle
    c. Auricular hillocks
    d. Pharyngeal groove
    e. Pharyngeal arch 1st

16. The blood supply of pinna comes from:
    a. External carotid artery
    b. Internal carotid artery
    c. Both A & B
    d. Facial artery
    e. Maxillary artery

54. The anterior fontanelle in fetal skull usually closes at?
a. Birth
b. One year after birth
c. One and half year after birth
d. Two years after birth

55. A malignant tumor is damaging the patient is glossopharyngeal nerve, they will experience
a. Loss of taste over the anterior two – third of tongue
b. Loss of somaesthetic sensation over anterior two third of the tongue
c. Loss of taste of sensation over posterior third of tongue
d. Paralysis of muscle of tongue

56. Examination of a patient indicates that he has a medially directed strabismus (squint). This could be due to damage to the.
a. Oculomotor nerve
b. Trochlear nerve
c. Ophthalmic
d. Abducent nerve

57. One of the following nucleus related to visual body relay ?
a. Lateral geniculate body
b. Edinger westpal nucleus
c. Pretectal nucleus
d. Superior colliculus

58. The sensory decussation is formed by?
a. Decussation of cortico spinal tract
b. Internal arcuate fibers
c. Descussation of superior cerebral peduncle
d. Descussation of rubrospinal tracts

59. All of the following are included in limbic system except:
a. Amygdaloid nucleus
b. Fornix
c. Parahippocampal gyrus
d. Hypothalamus

60. Regarding basal nuclei
a. They are composed of white matter
b. They are joined to cortex at posterior perforated substance
c. Caudate nucleus and putamen together forms neostriatum
d. Amygdale is part of lentiform nucleus

61. Vertebral artery
a. Is a branch of second part of subclavian artery
b. Gives off posterior cerebral artery
c. Passes through lower 6 central vertebral
d. Enters the cranium through foramen magnum

62. Dura mater is attached to:
a. Body of sphenoid
b. Margins of foramen magnum
c. Cribriform plate of ethmoid
d. petrous temporal bone

EACH OPTION CAN BE SELECTED ONCE, MORE THAN ONCE OR NOT AT ALL.

a	Head fold	f	philtrum
b	Nucleus pulposus	g	teratoma
c	Hydatidform mole	h	Upper incisor teeth
d	epiglottis	i	Phocomelia
e	Branchial cyst	j	Palatine tonsil

Lead in: Choose the correct option and write it against each stem

1. The structure derived from the intermaxillary segment of the embryonic face
2. Primitive streak fails to degenerate in fetal life
3. A normal remnant of an embryonic structure found in adult body
4. are found along the anterior border of the sternocleidomastoid muscle
5. The hypobranchial eminence contributes to the formation of

a	Pseudo stratified columnar epithelium	f	Thymus
b	Cuboidal epithelium	g	Palatine tonsil
c	Dentin	h	Osteon
d	Spinal cord	i	Osteoid
e	cerebellum	j	Cementum

Regarding histological features choose the best option for the following statements:

6. Unmineralized bone matrix
7. Crypts lined by stratified squamous Non- Keratinized epithelium
8. Covering of root surface of tooth
9. Gray commissure with central canal
10. Lining epithelium of thyroid follicles

a	Suture	f	Synovial Hinge joint
b	Syndesmosis	g	Synovial saddle joint
c	Gomphoses	h	Synovial pivot joint
d	Synchondrosis	i	Synovial bicondylar joint
e	Symphysis	j	Synovial ellipsoid joint

Choose the appropriate category of joints from the mentioned above for the following examples of joint with the given specific characteristics.

11. Between tooth and alveolar socket
12. Typically present in median plane
13. That permits rotation of thumb
14. Limited to skull bone
15. With fibrous membrane

37. Temporomandibular joint is innervated by:
a. Facial nerve
b. Mandibular nerve
c. Maxillary nerve
d. Vagus nerve

38. In the little area of Nasal cavity, all of following arteries anastomose EXCEPT:
a. Superior labial
b. Greater palatine
c. Spheno palatine
d. Posterior ethmoidal

39. A transient facial paralysis may result from entrance of anesthetic agent lingual nerve block into
a. Parotid capsule
b. Parotid fascia
c. Pterygomandibular raphe
d. Buccal fascia

40. The oral diaphragm is formed by which one of the following muscle:
a. Geniohyoid
b. Mylohyoid
c. Hyoglossus
d. Genioglossus

41. Anterior dislocation of temporomandibular joint is limited by
a. Sphenomandibular ligament
b. Stylomandibular ligament
c. Lateral ligament
d. Arcuate eminence

42. A young batsman received serve injury by a fast bouncer ball hitting his neck in posterior auricular region causing unconsciousness – injury to which structure had caused unconsciousness.
a. Occipital artery
b. Internal carotid artery
c. Basilar artery
d. Vertebral artery

43. A patient who had undergone the surgery to remove right lobe of thyroid gland, past operation his voice reduces to a whisper, which of following nerve was damaged?
a. Right external laryngeal
b. Right recurrent laryngeal
c. Left recurrent laryngeal
d. Left superior laryngeal

44. Which of the following is the dangerous layer of scalp
a. Pericranium
b. Loose areolar tissue
c. Galea aponeurosis
d. Skin

The area of cerebral cortex listed below receive their arterial supply as indicated:
a. the precentral gyrus is supplied by the middle cerebral artery
b. the post central gyrus (face area) is supplied by anterior cerebral artery
c. The cuneus is supplied by anterior cerebral artery
d. The Wernicke area is supplied by posterior cerebral artery

72. The motor speech area of Broca:
a. In most individual, important on left dominant hemisphere
b. Brings about the formation of words by its connections with secondary motor area
c. Connected to sensory speech area of wernicke
d. Brodmann area 34 and 35 represent motor speech area

73. The posterior surface of medulla oblongata shows.
a. Pyramids
b. Olives
c. Hypoglossal triangle
d. Middle cerebellar peduncle

74. The nerve emerging between pon and medulla oblongata is
a. Glossopharyngeal nerve
b. Facial nerve
c. Vagus nerve
d. Occulomotor nerve

75. 60 year old man came to ER with complain of paralysis on left side of lower limb with no difficulty in his speech , which artery is most likely to be involved.
a. Left anterior cerebral artery
b. Left middle cerebral artery
c. Right anterior cerebral artery
d. Right middle cerebral artery

EACH OPTION CAN BE SELECTED ONCE, MORE THAN ONCE OR NOT AT ALL.

a	Head fold	f	philtrum
b	Nucleus pulposus	g	teratoma
c	Hydatidform mole	h	Upper incisor teeth
d	epiglottis	i	Phocomelia
e	Branchial cyst	j	Palatine tonsil

Lead in: Choose the correct option and write it against each stem

1. The structure derived from the intermaxillary segment of the embryonic face
2. Primitive streak fails to degenerate in fetal life
3. A normal remnant of an embryonic structure found in adult body
4. are found along the anterior border of the sternocleidomastoid muscle
5. The hypobranchial eminence contributes to the formation of

a	Pseudo stratified columnar epithelium	f	Thymus
b	Cuboidal epithelium	g	Palatine tonsil
c	Dentin	h	Osteon
d	Spinal cord	i	Osteoid
e	cerebellum	j	Cementum

Regarding histological features choose the best option for the following statements:

6. Unmineralized bone matrix
7. Crypts lined by stratified squamous Non- Keratinized epithelium
8. Covering of root surface of tooth
9. Gray commissure with central canal
10. Lining epithelium of thyroid follicles

a	Suture	f	Synovial Hinge joint
b	Syndesmosis	g	Synovial saddle joint
c	Gomphoses	h	Synovial pivot joint
d	Synchondrosis	i	Synovial bicondylar joint
e	Symphysis	j	Synovial ellipsoid joint

Choose the appropriate category of joints from the mentioned above for the following examples of joint with the given specific characteristics.

11. Between tooth and alveolar socket
12. Typically present in median plane
13. That permits rotation of thumb
14. Limited to skull bone
15. With fibrous membrane

a	Parotid sheath	f	Auriculotemporal nerve
b	Retromandibular vein	g	Great auricular nerve
c	Facial nerve	h	Mastoid process
d	External carotid artery	i	Transverse cervical artery
e	Preauricular lymph nodes	J	Maxillary artery

Select the correct structure located in the parotid region from the above mentioned options which is most likely to be involved in the following conditions

16. In mumps, the parotid gland swells resulting in the excruciating pain during chewing.
17. During parotidectomy, the most superficial structure lying within the tissue of parotid gland.
18. During the dissection of the parotid region, the structure that indents the posteromedial surface of the parotid gland.

a	Medial pterygoid plate	f	Spine of sphenoid
b	Greater wing of sphenoid	g	Internal acoustic meatus
c	Lateral pterygoid plate	h	Pterygomaxillary fissure
d	Juglar foramen	i	Infraorbital fissure
e	Styloid process	J	Lesser wing of sphenoid

Select the most appropriate bony structure from the above mentioned list that is most likely to be related to the description given below.

19. Lying lateral to it is the auriculotemporal nerve and media to it is the chorda tympani nerve.
20. The bony opening has the fundus consisting of a bony plate called lamina cribrosa.

a	Occulomotor nerve	f	Cranial accessory nerve
b	Mandibular nerve	g	Hypoglossal nerve
c	Abducent nerve	h	Vagus nerve
d	Facial nerve	i	Opthalmic nerve
e	Spinal accessory nerve	J	Glossopharyngeal nerve

Select the correct cranial nerve from the above mentioned options that is most likely to be damaged resulting in the following clinical conditions

21. A Patient complains to the general physician that in the early morning when she tried to get up from the bed she is unable to move her head but can move her body.
22. A hypertensive patient in the emergency ward presents with double vision. On examination the pupils were dilated
23. During a routine physical examination, when the patient was asked to show his teeth, he was unable to do so.
24. A patient came to the general physician and complains that during chewing she is unable to touch the floor of the oral cavity.
25. When an examiner asks a child to pronounce "A", "O", "E", the child was unable to do so.

22. Disorders of the basal ganglia include all of the following except:

A. Chorea
B. Athetosis
C. Trigeminal neuralgia
D. Parkinsonism

23. Circle of willis is formed by the following except:

A. Anterior cerebral artery
B. Anterior choroidal artery
C. Posterior communicating artery
D. Posterior cerebral artery

24. The primary visual area of broadmann is located in the:

A. superior parietal lobule
B. Calcarine sulcus
C. Parieto-occipital sulcus
D. Frontal lobe

25. WHICH OF THE FOLLOWING STATEMENTS CHARACTERIZES OSTEOCLASTS:

A. Uni-nucleated cells
B. Secrete osteiod
C. Derived from osteoprogenitor cells
D. Occupy Howship lacunae

26. THE TYPE OF CARTILAGE IN EXTERNAL EAR:

A. Hyaline
B. Elastic
C. Fibrocartilage
D. All of the above

27. REGARDING OSTEON:

A. The central component is the Haversian canal
B. Composed of concentric lamellae of mineralized matrix
C. Primary cell type in osteon is the osteoclast
D. Blood vessels reach osteon from bone marrow via Volkmann's canals

28. BIPOLAR NEURONS ARE FOUND IN:

A. Dorsal root ganglia
B. Ventral horns of spinal cord
C. White matter of cerebrum
D. Olfactory epithelium

29. WHICH TYPE OF CELL PROVIDES STRUCTURAL AND NUTRITIONAL SUPPORT TO NEURONES IN THE CNS?

A. Microglia
B. Schwann cells
C. Astrocytes
D. Ependymal cells

EXTENDED MATCHING QUESTION

EACH OPTION CAN BE SELECTED ONCE, MORE THAN ONCE OR NOT AT ALL.

a	Head fold	f	philtrum
b	Nucleus pulposus	g	teratoma
c	Hydatidform mole	h	Upper incisor teeth
d	epiglottis	i	Phocomelia
e	Branchial cyst	j	Palatine tonsil

Lead in: Choose the correct option and write it against each stem

1. The structure derived from the intermaxillary segment of the embryonic face
2. Primitive streak fails to degenerate in fetal life
3. A normal remnant of an embryonic structure found in adult body
4. are found along the anterior border of the sternocleidomastoid muscle
5. The hypobranchial eminence contributes to the formation of

---

a	Pseudo stratified columnar epithelium	f	Thymus
b	Cuboidal epithelium	g	Palatine tonsil
c	Dentin	h	Osteon
d	Spinal cord	i	Osteoid
e	cerebellum	J	Cementum

Regarding histological features choose the best option for the following statements:

6. Unmineralized bone matrix
7. Crypts lined by stratified squamous Non- Keratinized epithelium
8. Covering of root surface of tooth
9. Gray commissure with central canal
10. Lining epithelium of thyroid follicles

---

a	Suture	f	Synovial Hinge joint
b	Syndesmosis	g	Synovial saddle joint
c	Gomphoses	h	Synovial pivot joint
d	Synchondrosis	i	Synovial bicondyiar joint
e	Symphysis	J	Synovial ellipsoid joint

Choose the appropriate category of joints from the mentioned above for the following examples of joint with the given specific characteristics.

11. Between tooth and alveolar socket
12. Typically present in median plane
13. That permits rotation of thumb
14. Limited to skull bone
15. With fibrous membrane

65.High endothelial venules are present in.

A. Thymus
B. Lymph node
C. Langerhan cell
D. Tonsil
E. Spleen

66.The pharyngeal plexus is situated at.

A.Posterolateral surface of superior constrictor
B. Posterolateral surface of middle constrictor
C.Posterolateral surface of inferior constrictor
D. In the piriform fossa

67.The following is not a content of cavernous sinus

A. Internal carotid artery
B. optic nerve .
C Trochlear nerve
D Abducent nerve

68.Excessive aphasia (inability to speak) results from damage to the

A. The motor speech area of bronca's
B. The sensory area of wernicke
C. The prefrontal cortex
D. The postcentral gyrus

69.Astereognosis results from damage to the


A. Primary sensory cortex
B. Sensory association cortex
C. Primary visual area
D. Primary auditory area

70.The facial colliculus overlies the nucleus of

A. Vagus nerve
B. Trochlear nerve
C. Oculomotor nerve
D. Abducent nerve

9:41

Talha Ahmed
Active 34 minutes ago



1st Year BDS/Pre-Prof 2017 (6)

...maxillary artery is supplied by
...part of maxillary artery
...part of maxillary artery
...part of maxillary artery
...dibular artery

46. A patient came to you with dry eyes due to decreased lacrimal gland secretion. This may be due to following ganglion.
a. Ciliary ganglion
b. Pterygopalatine ganglion
c. Otic ganglion
d. Submandibular ganglion

47. Infection from face to cavernous sinus can easily spread by:
a. Pterygoid venous plexus
b. External jugular vein
c. Internal jugular vein
d. Ophthalmic vein

48. Maxillary artery terminates as:
a. Infraorbital artery
b. Sphenopalatine artery
c. Palatine artery
d. Nasopalatine artery

49. In hanging, death occurs due to fracture of
a. Base of skull
b. Odontoid process of axis
c. Body of atlas
d. Arch of atlas

50. Regarding inferior alveolar nerve:
a. It is purely sensory
b. Purely motor
c. Mixed nerve
d. Parasympathetic nerve

51. Lower gums are innervated by:
a. Greater palatine nerve
b. Lingual nerve
c. Anterior superior alveolar nerve
d. Posterior superior alveolar nerve

52. The palatoglossal arch is formed by which of the following muscle:
a. Tensor veli palatine
b. Levator veli palatine
c. Palatoglossus muscle
d. Palate pharyngeus

53. Regarding left common carotid artery:
a. It arises from brachiocephalic trunk
b. It is embedded in carotid sheath
c. It gives two carotid and superior thyroid artery
d. It lies medial to internal jugular vein

Remix

a	Parotid sheath	f	Auriculotemporal nerve
b	Retromandibular vein	g	Great auricular nerve
c	Facial nerve	h	Mastoid process
d	External carotid artery	i	Transverse cervical artery
e	Preauricular lymph nodes	J	Maxillary artery

Select the correct structure located in the parotid region from the above mentioned options which is most likely to be involved in the following conditions

16. In mumps, the parotid gland swells resulting in the excruciating pain during chewing.
17. During parotidectomy, the most superficial structure lying within the tissue of parotid gland.
18. During the dissection of the parotid region, the structure that indents the posteromedial surface of the parotid gland.

a	Medial pterygoid plate	f	Spine of sphenoid
b	Greater wing of sphenoid	g	Internal acoustic meatus
c	Lateral pterygoid plate	h	Pterygomaxillary fissure
d	Juglar foramen	i	Infraorbital fissure
e	Styloid process	J	Lesser wing of sphenoid

Select the most appropriate bony structure from the above mentioned list that is most likely to be related to the description given below.

19. Lying lateral to it is the auriculotemporal nerve and media to it is the chorda tympani nerve.
20. The bony opening has the fundus consisting of a bony plate called lamina cribrosa.

a	Occulomotor nerve	f	Cranial accessory nerve
b	Mandibular nerve	g	Hypoglossal nerve
c	Abducent nerve	h	Vagus nerve
d	Facial nerve	i	Opthalmic nerve
e	Spinal accessory nerve	J	Glossopharyngeal nerve

Select the correct cranial nerve from the above mentioned options that is most likely to be damaged resulting in the following clinical conditions

21. A Patient complains to the general physician that in the early morning when she tried to get up from the bed she is unable to move her head but can move her body.
22. A hypertensive patient in the emergency ward presents with double vision. On examination the pupils were dilated
23. During a routine physical examination, when the patient was asked to show his teeth, he was unable to do so.
24. A patient came to the general physician and complains that during chewing she is unable to touch the floor of the oral cavity.
25. When an examiner asks a child to pronounce "A", "O", "E", the child was unable to do so.