a	Parotid sheath	f	Auriculotemporal nerve
b	Retromandibular vein	g	Great auricular nerve
c	Facial nerve	h	Mastoid process
d	External carotid artery	i	Transverse cervical artery
e	Preauricular lymph nodes	J	Maxillary artery

Select the correct structure located in the parotid region from the above mentioned options which is most likely to be involved in the following conditions

16. In mumps, the parotid gland swells resulting in the excruciating pain during chewing.
17. During parotidectomy, the most superficial structure lying within the tissue of parotid gland.
18. During the dissection of the parotid region, the structure that indents the posteromedial surface of the parotid gland.

a	Medial pterygoid plate	f	Spine of sphenoid
b	Greater wing of sphenoid	g	Internal acoustic meatus
c	Lateral pterygoid plate	h	Pterygomaxillary fissure
d	Juglar foramen	i	Infraorbital fissure
e	Styloid process	J	Lesser wing of sphenoid

Select the most appropriate bony structure from the above mentioned list that is most likely to be related to the description given below.

19. Lying lateral to it is the auriculotemporal nerve and media to it is the chorda tympani nerve.
20. The bony opening has the fundus consisting of a bony plate called lamina cribrosa.

a	Occulomotor nerve	f	Cranial accessory nerve
b	Mandibular nerve	g	Hypoglossal nerve
c	Abducent nerve	h	Vagus nerve
d	Facial nerve	i	Opthalmic nerve
e	Spinal accessory nerve	J	Glossopharyngeal nerve

Select the correct cranial nerve from the above mentioned options that is most likely to be damaged resulting in the following clinical conditions

21. A Patient complains to the general physician that in the early morning when she tried to get up from the bed she is unable to move her head but can move her body.
22. A hypertensive patient in the emergency ward presents with double vision. On examination the pupils were dilated
23. During a routine physical examination, when the patient was asked to show his teeth, he was unable to do so.
24. A patient came to the general physician and complains that during chewing she is unable to touch the floor of the oral cavity.
25. When an examiner asks a child to pronounce "A", "O", "E", the child was unable to do so.

47. All the nuclei in brain stem are parasympathetic **except**:

a. Superior salivatory
b. Dorsal vagal
c. Edinger Westpaul
d. Tractus solitarious

48. The white rami communicants :

a. Are demylinated fiber
b. Distributed to thoracic viscera
c. Are post synaptic to sympathetic chain
d. Courses in anterior root of spinal nerve

49. Sympathetic chain:

a. Consist of unilateral 31 ganglion
b. Consolidate cervical ganglion are three on each side
c. Chain terminates at stellate ganglion
d. Receive pre ganglion fiber from anterior horn of spinal cord

of growing bone:

22. Disorders of the basal ganglia include all of the following except:

A. Chorea
B. Athetosis
C. Trigeminal neuralgia
D. Parkinsonism

23. Circle of willis is formed by the following except:

A. Anterior cerebral artery
B. Anterior choroidal artery
C. Posterior communicating artery
D. Posterior cerebral artery

24. The primary visual area of broadmann is located in the:

A. superior parietal lobule
B. Calcarine sulcus
C. Parieto-occipital sulcus
D. Frontal lobe

25. WHICH OF THE FOLLOWING STATEMENTS CHARACTERIZES OSTEOCLASTS:

A. Uni-nucleated cells
B. Secrete osteiod
C. Derived from osteoprogenitor cells
D. Occupy Howship lacunae

26. THE TYPE OF CARTILAGE IN EXTERNAL EAR:

A. Hyaline
B. Elastic
C. Fibrocartilage
D. All of the above

27. REGARDING OSTEON:

A. The central component is the Haversian canal
B. Composed of concentric lamellae of mineralized matrix
C. Primary cell type in osteon is the osteoclast
D. Blood vessels reach osteon from bone marrow via Volkmann's canals

28. BIPOLAR NEURONS ARE FOUND IN:

A. Dorsal root ganglia
B. Ventral horns of spinal cord
C. White matter of cerebrum
D. Olfactory epithelium

29. WHICH TYPE OF CELL PROVIDES STRUCTURAL AND NUTRITIONAL SUPPORT TO NEURONES IN THE CNS?

A. Microglia
B. Schwann cells
C. Astrocytes
D. Ependymal cells

PRE PROF - 1st

1. The part of the sperm containing proteolytic enzymes to digest the zona pellucida is the:
   a. Capacitor
   b. Head
   c. Corona
   d. Acrosome (Hyaluronidase)

DONE

2. The ovulated mammalian oocyte is arrested at:
   a. Prophase of meiosis I
   b. Metaphase of meiosis I
   c. Prophase of meiosis II
   d. Metaphase of meiosis II

Somite → Sclerotome (vertebrae), Myotome (skeletal)

3. Somites:
   a. Differentiate into myotomes which give rise to skeletal muscle in trunk
   b. Differentiate into sclerotomes which give rise to vertebrae
   c. Arise from segmentation of the paraxial mesoderm
   d. All of the above are correct

4. When the amount of amniotic fluid exceeds two liters, the condition is called:
   a. Oligohydramnios
   b. Polyhydramnios
   c. Amniotitis
   d. Hydrogravida

5. After parturition a gynecologist identified the maternal surface of placenta due to 15-20 bulging areas called:

Fistula (maxillary Air Sinus)

   a. Cotyledons.
   b. Placental septa.
   c. Intervillous space.
   d. Cytotrophoblastic shell.

6. A 21 year old female present to the obstetrician at 16 weeks of prenatal care. The obstetrician finds increased amniotic fluid and orders alpha fetoprotein and HCG levels. One week later the called patient back to tell her that the alpha fetoprotein level is increased. The most probable fetal diagnosis is

   a. Anencephaly
   b. Jejunal atresia
   c. Renal agenesis
   d. Sacrococcgeal teratoma

Emgysl

7. Fetal membranes are as under EXCEPT
   a. Amnion
   b. Allantois
   c. Germ disc
   d. Chorion

21 - E
22 - b
23 - d
24 - g
25 - b

1 - f
2 - g
3 - b
4 - e
5 - d

8. At the time of ovulation there comes a surge of
   a. Estrogen
   b. Progesterone
   c. Follicle stimulating hormone
   d. Luteinizing hormone (Foatcaval).

9. In case of Dizygotic twins there is found to be
   a. One chorionic cavity
   b. Two amniotic cavities
   c. One umbilical cord
   d. One connecting stalk

a. Superior salivatory
b. Dorsal vagal
c. Edinger Westpaul
d. Tractus solitarious

48. The white rami communicants :

a. Are demylinated fiber
b. Distributed to thoracic viscera
c. Are post synaptic to sympathetic chain
d. Courses in anterior root of spinal nerve

49. Sympathetic chain:

a. Consist of unilateral 31 ganglion
b. Consolidate cervical ganglion are three on each side
c. Chain terminates at stellate ganglion
d. Receive pre ganglion fiber from anterior horn of spinal cord

EMQ No: 4

For each of the conditions described below match the nerve that is involved:

Options List

A. Peritoneum
B. Extra peritoneal fat
C. Fascia transversalis
D. Transverses oblique
E. Internal oblique
F. External oblique
G. Deep fascia
H. Membranous superficial fascia
I. Fatty superficial fascia
J. skin

Question No: 14. Forms mainly the entire posterior wall of inguinal canal

Question No: 15. Replaced by tunica vaginalis that surrounds the testes

Question No: 16. Replaced by cremasteric fascia within the spermatic cord.

Question No: 17. The layer that contains the dartos muscle

EMQ No: 5.

For each question below, chose a single option from the list

Options

A. Rathke's pouch
B. Foramen cecum
C. First pharyngeal cleft
D. Third pharyngeal pouch
E. Pars nervosa
F. Inter-maxillary segment
G. Second pharyngeal arch
H. First pharyngeal arch
I. Infundibulum
J. [illegible]

Question No; 18. Anterior pituitary is the derivative of

Question No; 19. Site of origin of thyroid gland in the adult is present between anterior 2/3 and posterior 1/3 of tongue

Question No; 20. Ear drum is the derivative of

Question No; 21. Muscles of mastication are the derivatives of

EMQ No: 1.

For each question below, chose a single option from the list

Options List

A. Internal cerebral vein
B. Anterior cerebral artery
C. Great cerebral vein
D. Posterior communicating artery
E. Basal vein
F. Posterior inferior cerebellar artery
G. Posterior cerebral artery
H. Basal artery
I. Inferior cerebral vein
J. Middle cerebral artery

Question No: 1. This is a superficial cerebral vein

Question No: 2. A 52-year old patient was admitted to the hospital with paralysis of right half of the trunk, right upper limb, and lower part of the right side of the face. It was also observed that there is an altered sensorium in the same area. This was diagnosed as an arterial lesion in the cranial cavity. The involved artery is:

Question No: 3. This is formed below the splenium of corpus callosum and terminates into anterior part of straight sinus

Question No: 4. Arises from internal carotid artery close to its bifurcation and joins the posterior cerebral artery.

Question No: 5. A 64- year old man is diagnosed with an acute stroke. His main deficit is a partial loss of his visual field. The neurologist diagnoses a lesion of the occipital lobe. Which of the arteries is likely to be involved?

EMQ No: 2.

For each question below, chose a single option from the list

A. Arachnoid mater
B. Cauda equina
C. Conus medullaris
D. Dura mater
E. Extradural space
F. L2
G. L3
H. L4
I. Pia mater
J. Subarachnoid space

Question No: 6. A lumbar puncture needle is inserted directly above or below this vertebra

Question No: 7. The lumbar cistern contains CSF and what structure?

Question No: 8. Separates the dura from the vertebral periosteum and is a site for injection of anaesthesia?

Question No: 9. Which is the last layer of the meninges to be penetrated when sampling CSF?

EACH OPTION CAN BE SELECTED ONCE, MORE THAN ONCE OR NOT AT ALL.

a	Head fold		
b	Nucleus pulposus	f	philtrum
c	Hydatidform mole	g	teratoma
d	epiglottis	h	Upper incisor teeth
e	Branchial cyst	i	Phocomelia
		j	Palatine tonsil

Lead in: Choose the correct option and write it against each stem

1. The structure derived from the intermaxillary segment of the embryonic face (f) philtrum
2. Primitive streak fails to degenerate in fetal life (g)
3. A normal remnant of an embryonic structure found in adult body. (e)
4. are found along the anterior border of the sternocleidomastoid muscle (e)
5. The hypobranchial eminence contributes to the formation of (d)

43. Inferior orbital fissure lies between :

a. Greater and lesser wing of sphenoid
b. Maxilla and greater wing
c. Maxilla and lesser wing
d. Lesser wing and body of sphenoid

(b)

44. A patient came in OPD with the complain of difficulty to open his eyes. Most probab
be:

a. Facial nerve
b. Occulomotor nerve
c. Trochlear nerve
d. Abducent nerve

45. In dissection of human body, a section right angle to long axis of the organs is:

a. Oblique section
b. Longitudinal section
c. Transverse section

54. The anterior fontanelle in fetal skull usually closes at?
a. Birth
b. One year after birth
c. One and half year after birth
d. Two years after birth

55. A malignant tumor is damaging the patient is glossopharyngeal nerve, they will experience
a. Loss of taste over the anterior two – third of tongue
b. Loss of somaesthetic sensation over anterior two third of the tongue
c. Loss of taste of sensation over posterior third of tongue
d. Paralysis of muscle of tongue

56. Examination of a patient indicates that he has a medially directed strabismus (squint). This could be due to damage to the.
a. Oculomotor nerve
b. Trochlear nerve
c. Ophthalmic
d. Abducent nerve

57. One of the following nucleus related to visual body relay ?
a. Lateral geniculate body
b. Edinger westpal nucleus
c. Pretectal nucleus
d. Superior colliculus

58. The sensory decussation is formed by?
a. Decussation of cortico spinal tract
b. Internal arcuate fibers
c. Descussation of superior cerebral peduncle
d. Descussation of rubrospinal tracts

59. All of the following are included in limbic system except:
a. Amygdaloid nucleus
b. Fornix
c. Parahippocampal gyrus
d. Hypothalamus

60. Regarding basal nuclei
a. They are composed of white matter
b. They are joined to cortex at posterior perforated substance
c. Caudate nucleus and putamen together forms neostriatum
d. Amygdale is part of lentiform nucleus

61. Vertebral artery
a. Is a branch of second part of subclavian artery
b. Gives off posterior cerebral artery
c. Passes through lower 6 central vertebral
d. Enters the cranium through foramen magnum

62. Dura mater is attached to:
a. Body of sphenoid
b. Margins of foramen magnum
c. Cribriform plate of ethmoid
d. petrous temporal bone

22. Disorders of the basal ganglia include all of the following except:

A. Chorea
B. Athetosis
C. Trigeminal neuralgia
D. Parkinsonism

23. Circle of willis is formed by the following except:

A. Anterior cerebral artery
B. Anterior choroidal artery
C. Posterior communicating artery
D. Posterior cerebral artery

24. The primary visual area of broadmann is located in the:

A. superior parietal lobule
B. Calcarine sulcus
C. Parieto-occipital sulcus
D. Frontal lobe

25. WHICH OF THE FOLLOWING STATEMENTS CHARACTERIZES OSTEOCLASTS:

A. Uni-nucleated cells
B. Secrete osteiod
C. Derived from osteoprogenitor cells
D. Occupy Howship lacunae

26. THE TYPE OF CARTILAGE IN EXTERNAL EAR:

A. Hyaline
B. Elastic
C. Fibrocartilage
D. All of the above

27. REGARDING OSTEON:

A. The central component is the Haversian canal
B. Composed of concentric lamellae of mineralized matrix
C. Primary cell type in osteon is the osteoclast
D. Blood vessels reach osteon from bone marrow via Volkmann's canals

28. BIPOLAR NEURONS ARE FOUND IN:

A. Dorsal root ganglia
B. Ventral horns of spinal cord
C. White matter of cerebrum
D. Olfactory epithelium

29. WHICH TYPE OF CELL PROVIDES STRUCTURAL AND NUTRITIONAL SUPPORT TO NEURONES IN THE CNS?

A. Microglia
B. Schwann cells
C. Astrocytes
D. Ependymal cells

49. Sympathetic chain:

a. Consist of unilateral 31 ganglion
b. Consolidate cervical ganglion are three on each side
c. Chain terminates at stellate ganglion
d. Receive pre ganglion fiber from anterior horn of spinal cord

growing bone:

pe here to search

a. Superior salivatory
b. Dorsal vagal
c. Edinger Westpaul
d. Tractus solitarious

48. The white rami communicants :

a. Are demylinated fiber
b. Distributed to thoracic viscera
c. Are post synaptic to sympathetic chain
d. Courses in anterior root of spinal nerve

49. Sympathetic chain:

a. Consist of unilateral 31 ganglion
b. Consolidate cervical ganglion are three on each side
c. Chain terminates at stellate ganglion
d. Receive pre ganglion fiber from anterior horn of spinal cord

EMQ No: 1.

For each question below, chose a single option from the list

Options List

A. Internal cerebral vein
B. Anterior cerebral artery
C. Great cerebral vein
D. Posterior communicating artery
E. Basal vein
F. Posterior inferior cerebellar artery
G. Posterior cerebral artery
H. Basal artery
I. Inferior cerebral vein
J. Middle cerebral artery

Question No: 1. This is a superficial cerebral vein

Question No: 2. A 52-year old patient was admitted to the hospital with paralysis of right half of the trunk, right upper limb, and lower part of the right side of the face. It was also observed that there is an altered sensorium in the same area. This was diagnosed as an arterial lesion in the cranial cavity. The involved artery is:

Question No: 3. This is formed below the splenium of corpus callosum and terminates into anterior part of straight sinus

Question No: 4. Arises from internal carotid artery close to its bifurcation and joins the posterior cerebral artery.

Question No: 5. A 64- year old man is diagnosed with an acute stroke. His main deficit is a partial loss of his visual field. The neurologist diagnoses a lesion of the occipital lobe. Which of the arteries is likely to be involved?

EMQ No: 2.

For each question below, chose a single option from the list

A. Arachnoid mater
B. Cauda equina
C. Conus medullaris
D. Dura mater
E. Extradural space
F. L2
G. L3
H. L4
I. Pia mater
J. Subarachnoid space

Question No: 6. A lumbar puncture needle is inserted directly above or below this vertebra

Question No: 7. The lumbar cistern contains CSF and what structure?

Question No: 8. Separates the dura from the vertebral periosteum and is a site for injection of anaesthesia?

Question No: 9. Which is the last layer of the meninges to be penetrated when sampling CSF?

EMQ No: 4

For each of the conditions described below match the nerve that is involved:

Options List

A. Peritoneum
B. Extra peritoneal fat
C. Fascia transversalis
D. Transverses oblique
E. Internal oblique
F. External oblique
G. Deep fascia
H. Membranous superficial fascia
I. Fatty superficial fascia
J. skin

Question No: 14. Forms mainly the entire posterior wall of inguinal canal

Question No: 15. Replaced by tunica vaginalis that surrounds the testes

Question No: 16. Replaced by cremasteric fascia within the spermatic cord.

Question No: 17. The layer that contains the dartos muscle

EMQ No: 5.

For each question below, chose a single option from the list

Options

A. Rathke's pouch
B. Foramen cecum
C. First pharyngeal cleft
D. Third pharyngeal pouch
E. Pars nervosa
F. Inter-maxillary segment
G. Second pharyngeal arch
H. First pharyngeal arch
I. Infundibulum
J. [illegible]

Question No; 18. Anterior pituitary is the derivative of

Question No; 19. Site of origin of thyroid gland in the adult is present between anterior 2/3 and posterior 1/3 of tongue

Question No; 20. Ear drum is the derivative of

Question No; 21. Muscles of mastication are the derivatives of

10. Stellate Reticulam is a part of developing
a. Enamel organ
b. Dental papilla
c. Dental follicle
d. Dental sac

11. Regarding nucleus of a cell all of the following are correct except:
a. Has porous membrane
b. Forms RNA of ribosomes
c. Continuous with mitochondria ER
d. Has nucleoli

12. Stratified squamous epithelium
a. Develops only from ectoderm
b. Found in pelvis of kidney
c. Continuously replaced by basal layer
d. Has superficial nucleated layer

13. Collagen fibres has following characteristic except:
a. Responsible for tensile strength
b. Unbranched fibres
c. Appear banded on microscopy
d. Produced by mast cells

14. Skeletal muscle fibres have
a. Peripheral single nuclei numerous
b. T system of tubules
c. Intercalated disc
d. Sarcomere between two I bands Z lines

15. Osteoblasts
a. Forms collagen
b. Remove bone during remodeling - osteoclast
c. Present in lacunae ← osteocytes
d. Derived from osteocytes ← osteoprogenitor cells

16. Lymph node has:
a. Connective tissue framework of collagen - No
b. Valveless single afferent lymph vessel — numerous Valves
c. T lymphocytes in cortex — Paracortex
d. B lymphocytes both in cortex and medulla

17. Tongue histology reveals that :
a. Filiform papillae has taste buds (No buds)
b. Lymphoid tissue is confined to tip
c. Mucous Gland numerous in anterior part (Posterior part)
d. Muscles fibres are predominantly skeletal

18. The salivary Glands:
a. Demilunes consist of mucous cells Serous
b. Serous cells are eosinophilic basophilic)
c. Striated ducts are lined by myoepithelial cells (intercalated ducts).
d. Lack adipose tissue

71. A patient complains of difficulty in swallowing , injury to which of the following nerve is cause of this complain:

a. Cranial part of accessory nerve
b. Spinal part of accessory nerve
c. Hypoglossal nerve
d. 1st cervical nerve

72. Nucleus ambigus is common for following nerves Except:

a. Vagus nerve
b. Accessory nerve
c. Hyopoglossal nerve
d. Glossopharyngeal nerve

73. Cavernous sinus coutains:

a. Sympathetic polexus
b. Mandibular nerve
c. Middle menengial artery
d. Pituitary stalk

74. With in the internal capsule, the sensory radiations from thalamus to Brodmann's area 3,2,1 run in the :

a. Anterior ½ of posterior limb
b. Posterior ½ of posterior limb
c. Whole of the posterior limb
d. Whole of the anterior limb

75. Forceps minor of corpus callosum are the fibers of :

a. Rostrum
b. Genu
c. Body
d. Splenium

1. Which of the following is not derived from neural crest cells:
a. "C" cells of the thyroid gland
b. Facial skeleton
c. Pia matter
d. Erector pili muscle

2. Which of the statement is true regarding primitive streak:
a. Completely regresses by the end of third week
b. Extends to the cranial end of embryonic disc
c. Gives rise to intraembryonic mesoderm
d. Its caudal termination is known as primitive node

3. Which of the following event during foetal period (3 month to birth) is true:
a. There is faster growth of head as compared to the rest of the body
b. During 3rd month the movements of the foetus are clearly felt by mother
c. By the 6th month the co ordination between respiratory system and nervous system is well established
d. At the end of 9th month the testes should be in scrotum

4. Which of the following virus is a confirmed teratogen
a. Measles
b. Cytomegalo virus
c. Influenza virus
d. Herpes simplex virus

5. Which of the statement regarding intraembryonic mesoderm is NOT true
a. It is formed during second week of development
b. It is continues with extra embryonic mesoderm at the margins of the disc
c. Intraembryonic blood vessels develop in it
d. Intraembryonic mesoderm form somites

6. Which of the statement regarding placenta is NOT true
a. Most of the maternal hormones cross placenta
b. The blood pressure is much higher in spiral arteries than veins
c. Umbilical arteries carry deoxygenated blood
d. Placenta previa can lead to massive hermorrhage

7. The otic vesicle forms from the following EXCEPT:
a. Cochlear duct
b. Semicircular canals
c. Membranous labyrinth
d. Tubotympanic recess

8. At birth the skin is covered by a whitish paste known as:
a. Ichthyosis layer
b. Cornium
c. Subcorium
d. Vernix caseosa

9. Cardiac muscle is differentiated from skeletal muscle by
a. T-tubules
b. Striations
c. Peripheral nucleus
d. Intercalated disc

10. The peculiar feature of thymus is
a. Lymphoid nodules
b. Hassal's corpuscles
c. Central artery
d. Paracortical zone

11. Regarding elastic cartilage which of the statement is true.
a. It is covered by periosteum
b. The chondrocytes lie within lacunae
c. It calcifies with age
d. Interwoven collagen fibers in matrix

53. Which of the following is the tributary of internal jugular vein?
a. Superior thyroid vein
b. Inferior thyroid vein
c. External jugular vein
d. Retro mandibular vein

54. Muscle of the soft palate which is supplied by mandubular nerve is
a. Palatoglossus
b. Palatopharyngeus
c. Tensor palati
d. Levator palati

55. Pterygopalatine fossa communicates with infratemporal fossa through
a. Foramen rotundum
b. Spheno-palatine foramen
c. Inferior orbital fissure
d. Pterygomaxillary fissure

56. Which of the following muscle forms oral diaphragm?
a. Stylohyoid
b. Mylohyoid
c. Geniohyoid
d. Diagastric

57. The sympathetic root of pterygopalatine ganglion is
a. Greater petrosal nerve
b. Lesser petrosal nerve
c. Deep petrosal nerve
d. Nerve to pterygoid canal

58. Maxillary artery terminates as
a. Infra orbital artery
b. Sphenopalatine artery
c. Posterior superior alveolar artery
d. Dorsal nasal artery

59. The infrahyoid muscle supplied by hypoglossal nerve is
a. Sternohyoid
b. Omohyoid
c. Thyrohyoid
d. Sternothyroid

60. In atlas, the spinous process is replaced by
a. Lateral mass
b. Anterior arch
c. Posterior arch
d. Posterior tubercle

61. The tract present in the anterior white coloumn of spinal cord is:
a. Rubrospinal tract
b. Olivospinal tract
c. Cuneocerebellar
d. Spinotectal tract

62. The auditory fibers relay in which of the following thalamic nucleus:
a. Lateral geniculate body
b. Medial geniculate body
c. Pulvinar
d. Dorsomedial nucleus

63. The internal capsule continuous below with the:
a. Corona radiata
b. Crus cerebri
c. Optic radiation
d. Auditory radiation

12. An oral tissue, which shows finger like processes and covered by stratified Squamous keratinized epithelium is
a. Circumvillate papillae
b. Foliate papillae
c. Fungiform papillae
d. Filiform papillae

13. The characteristic feature of mixed salivary gland is presence of
a. Large number of adipose cells
b. Small striated ducts
c. Serous demilunes
d. Small mucous acini

14. Which of the following feature is present in thyroid gland
a. Sinusoidal capillaries
b. Cuboidal follicular cells
c. Fenestrated capillaries
d. Ciliated parafollicular cells

15. Spongy bone can be differentiated from compact bone by the presence of
a. Bony trabeculae
b. Mineralized bone matrix
c. Osteoclasts
d. Bony lamellae

16. The epithelial tissue exhibits
a. Intertwining elongated cellular process
b. Elongated contractile cells
c. Aggregated polyhedral cells
d. Wandering and fixed cells

17. The spinal nerve:
a. Innervate the cardiac muscle
b. Contain ganglion in its posterior ramus
c. Supply the sturucture of head and face
d. Carries autonomic component in its anterior root

18. The diaphysis is:
a. Secondary ossification center
b. Present at the time of birth
c. Grows towards the shaft
d. More than one in each bone

19. The example of cruciate muscle is
a. Pectoralis major
b. Tibialis posterior
c. Diagastric
d. Masseter

20. Pelvic splancnic nerve
a. Are postganglionic fiber
b. Supply the glandular tissue
c. Synaps in visceral ganglion
d. Contains parasympathetic fibers

21. One of the following is the example of the pneumatic bone:
a. Nasal
b. Vomer
c. Palatine
d. Mastoid

22. Grey Rami communicants are:
a. Preganglionic fibers
b. Present in anterior root of spinal nerve
c. The supply of sweat gland
d. Content of vertebral canal

Options:

A. Growth hormone.
B. Prolactin.
C. ADH.
D. Oxytocin.
E. T3, T4.
F. Parathyroid hormone.*
G. Epinephrine.
H. Aldosterone.
I. Cortisol.
J. Insulin.

Lead in:

For each of the clinical states of hormonal dysfunction given below, select the most appropriate option. Each option may be used once, more than once or not at all.

1. A patient with resting blood pressure of 180/105 mmHg and a serum sodium level of 156mEq/L.
2. A 36 year old man complaining of palpitation with a heart rate of 105/ min and fine tremors.
3. A 16 year old girl whose height is 6 feet 3 inches and whose fasting blood sugar is 175mg/dl.
4. A 45 year old male complaining of palpitation who has an irregular pulse and a serum calcium level of 15mg/dl (normal 10mg/dl)
5. A 34 year old girl with history of delayed wound healing, rounded face and a fasting blood sugar level of 156mg/dl.

5.

Options:

A. Gastrin
B. Pancreatic polypeptide
C. Cholecystokinin
D. Leptin
E. Secretin
F. VIP
G. Motilin
H. Histamine
I. Somatostatin
J. GIP

Lead in:

For each of the following functions of GIT hormones given below, select the most appropriate option . Each option may be used once, more than once or not at all.

1. Hormone that inhibits gastric emptying.
2. Hormone that increases gastric H^+ ion secretion.
3. Hormone that stimulates contraction of gallbladder and relaxation of sphincter of Oddi.
4. Hormone that increases biliary HCO_3^- secretion.
5. Hormone that produces relaxation of GI smooth muscles.

EXTENDED MATCHING QUESTIONS

a	Anencephalous	f	Ankyloglossia
b	Microcephalous	g	Thyroid gland
c	Lacrimal fissure	h	Superior Parathyroid gland
d	Cleft lip	i	Dental pulp
e	Microglossia	j	Enamel

Select the nerve involved in the following conditions:

1. Incomplete merging of two medial nasal swellings.
2. Skull fails to close results in absence of brain.
3. Tongue fails to protrude because of frenulam.
4. Ectoderm germinal layer contributing tooth.
5. Endoderm of 4th pharyngeal pouch.

---

a	Fibroblast	f	Macrophage
b	Pericyte	g	Osteocyte
c	Reticular cell	h	Osteoblast
d	Adipocyte	i	Chondroblast
e	Lymphocytes	j	Chondrocyte

Select the correct type of cell from the list above that corresponds the statements given below:

6. Produces and maintain all extracellular components of connective tissue
7. In histologic sections the cells have a signet ring appearance because of the lipid that washed away during slide preparation.
8. Under light microscope, the cells appear large multinucleated with acidophilic cytoplasm having reffled border on their cell surface which faces the clear zone.
9. The cells constitute mononuclear phagocyte lysosomes.
10. The mesenchymal derived cells that scattered along capillaries with contractile filaments

---

a	Parallel	f	Unipennate
b	Multipennate	g	Parallel with intersection
c	Circumpennate	h	Bipennate
d	Twisted	i	Triangular
e	Crossed	j	Quadrate

Select the most appropriate muscle fasiculis mentioned above for the following example of the muscles given below:

11. Clavicular fiber of detoid
12. Rectus abdominus at interior abdominal wall.
13. Tibialis anterior of the leg

43. The free and attached margin of tentorium cerebelli cross each other at:
a. Anterior clinioid process
b. Apex of petrous temporal bone
c. Internal occipital protuberance
d. Opening of superior saggital sinus

44. Sphenopalatine foramen lies in
a. Horizontal plate of palatine bone
b. Perpendiculate plate of palatine bone
c. Orbital process of palatine bone
d. Pyramidal process of palatine bone

45. A 23 years old male represented with swelling of Parotid region. He has contracted Mumps five weeks back and he was suffering from excruciating pain in parotid region and ear, which increased while chewing. The nerve most likely to be mediating the pain is
a. Anterior Superior Alveolar
b. Middle Superior Alveolar
c. Glossopharyngeal
d. Auriculotemporal

46. Fracture of Pterion will cause injury to
a. Superficial temporal Artery
b. Middle cerebral Artery
c. Middle Meningeal Artery
d. Anterior Cerebral Artery

47. All the following nuclei are present beneath the floor of fourth ventricle except:
a. Dorsal nucleus of Vagus nerve
b. Vestibular nuclei
C. Hypoglossal nucleus
d. Facial nucleus

48. A patient came in emergency after gun shot injury in posterior triangle, the structure which most likely to be injured will be:
a. Trunks of brachial plexus
b. External juglar vein
c. Spinal accessory nerve
d. Supraclavicular lymph nodes

49. Nasopharynx is supplied by
a. Maxillary nerve
b. Glossopharyngeal nerve
c. Vagus nerve
d. Cranial accessory nerve

50. The nerve that does not pass through middle ear cavity is
a. Chorda tympanic
b. Tympanic
c. Greater petrosal
d. Lesser petrosal

51. All are the arterial sources of thyroid gland EXCEPT.
a. External carotid artery
b. Vertebral artery
c. thyrocervical artery
d. Brachiocephallic artery

52. On the lateral wall of nose the maxillary air sinus opens into
a. Sphenoidal recess
b. Superior meatus
c. Middle meatus
d. Inferior meatus

primary visual area
secondary visual area
occipital eyefield
frontal eyefield

65. Which of the following nucleus is present in cerebrum
a. Caudate
b. Cuneate
c. Dentate
d. Red

66. Nucleus ambigus is common for following nerve except
a. Vagus nerve
b. Accessory nerve
c. Hypoglossal nerve
d. Glossopharyngeal nerve

67. Dura matter is attached to
a. Body of sphenoid
b. Margins of foramen magnum
c. Ciribriform plate of ethemoid
d. Petrous temporal bone

68. Which one of the following is included in basal nuclei:
a. Habenular nucleus
b. Red nucleus
c. Tractus solitaries nucleus
d. Amygdaloid nucleus

69. Crus cerebri contain:
a. Nuclei of motor system
b. Nuclei of visual pathway
c. Descending tract
d. Ascending tract

70. In the pyramidal decussation the fibers that decussate are:
a. Anterior corticospinal tract fibers
b. Lateral corticospinal tract fibers
c. Lateral reticulospinal tract
d. Rubrospinal tract

71. on the anterolateral surface of the pons which of the following nerves emerge:
a. Abducent nerve
b. Trochlear nerve
c. Trigeminal nerve
d. Facial nerve

72. A patient came in emergency room on examination his visual area is effected which artery is susceptible:
a. Anterior cerebral artery
b. Middle cerebral artery
c. Posterior cerebral artery
d. Superior cerebral artery

73. Sylvian fissure separates:
a. Frontal lobe from parietal lobe
b. Frontal lobe from temporal lobe
c. Temporal lobe from occipital lobe
d. Occipital lobe from temporal lobe

d
d
d
1

Select the most correct cortical gyrus of the cerebral hemisphere according to the location/function given below:

14. Area 44,45 that produces speech
15. Primary motor area (area 4) produces the muscle movements on the opposite side of the body
16. Area 6, 8 and 9 that controls the voluntary scanning movements of the eye
17. Area 43 that receives the taste sensations
18. Area 17 represents the area where all the visual sensation reaches

a	Superior laryngeal nerve	f	Vagus nerve
b	Recurrent laryngeal nerve	g	Cranial accessory nerve
c	External laryngeal nerve	h	Spinal accessory nerve
d	Internal laryngeal nerve	i	Transverse cervical nerve
e	Glossopharyngeal nerve	j	Great auricular nerve

Select the most appropriate nerve from above mentioned options that is most likely to be damaged in the following clinical conditions:

19. A 30 years old surgeon operated upon a middle aged lady for enlarged thyroid and ligated superior and inferior arteries as close as possible to the poles. After two weeks the lady presented with hoarseness of voice.
20. A patient came in emergency with inhalation of a foreign body and severe respiratory obstruction. On examination the foregin body is found to lodged at the region of rima glottides and there is sensory loss in upper part of larynx.
21. A patient came in OPD with asymmetric neckline, drooping shoulder and weakness of forward elevation of shoulder.

a	Medial pterygoid plate	f	Spine of sphenoid
b	Greater wing of sphenoid	g	Internal acoustic meatus
c	Lateral pterygoid plate	h	Pterygomaxillary fissure
d	Jugular foramen	i	Infraorbital fissure
e	Styloid process	j	Lesser wing of sphenoid

Select the most appropriate bony structure from the above mentioned list that is most likely to be related to the description given below:

22. Lying lateral to it is the auriculotemporal nerve and medial to it is the chorda tympani nerve.
23. The bony opening has the fundus consisting of a bony plate called lamina cribrosa.

a	Posterior auricular artery	f	Lingual artery
b	Caroticotymapnic artery	g	Palatine artery
c	Vertebral artery	h	Superior thyroid artery
d	Middle meningeal artery	i	Inferior thyroid artery
e	Ascending pharyngeal artery	j	Facial artery

Select the artery which will rupture in the following condition from above mentioned list:

24. An unconscious man brought to emergency with history of fracture of neck. CT scan shows ischemia of medulla oblongata.
25. In a complicated case of surgery done for the removal of palatine tonsil, the tonsillar bed was penetrated leading to profuse bleeding.

c	Parotid gland	f	Auriculotemporal nerve
b	Retromandibular vein	g	Great auricular nerve
i	Facial nerve	h	Masseter muscle
a	External carotid artery	j	Transverse cervical artery
d	Preauricular lymph nodes	k	Maxillary artery

Select the correct structure located in the parotid region from the above mentioned options which is most likely to be involved in the following conditions

16. In mumps, the parotid gland swells resulting in the excruciating pain during chewing.
17. During parotidectomy, the most superficial structure lying within the tissue of parotid gland
18. During the dissection of the parotid region, the structure that indents the posteromedial surface of the parotid gland

---

a	Medial pterygoid plate	f	Spine of sphenoid
b	Greater wing of sphenoid	g	Internal acoustic meatus
c	Lateral pterygoid plate	h	Pterygomaxillary fissure
d	Jugular foramen	i	Infraorbital fissure
e	Styloid process	j	Lesser wing of sphenoid

Select the most appropriate bony structure from the above mentioned list that is most likely to be related to the description given below.

19. Lying lateral to it is the auriculotemporal nerve and media to it is the chorda tympani nerve.
20. The bony opening has the fundus consisting of a bony plate called lamina cribrosa.

---

71. The area of cerebral cortex listed below receive their arterial supply as indicated
a. The precentral gyrus is supplied by the middle cerebral artery
b. The post central gyrus (face area) is supplied by anterior cerebral artery
c. The cuneus is supplied by anterior cerebral artery
d. The Wernicke area is supplied by posterior cerebral artery

72. The motor speech area of Broca
a. In most individual, important on left dominant hemisphere
b. Brings about the formation of words by its connections with secondary motor area
c. Connected to sensory speech area of wernicke
d. Brodmann area 34 and 35 represent motor speech area

73. The posterior surface of medulla oblongata shows
a. Pyramids
b. Olives
c. Hypoglossal triangle
d. Middle cerebellar peduncle

74. The nerve emerging between pon and medulla oblongata is
a. Glossopharyngeal nerve
b. Facial nerve
c. Vagus nerve
d. Occulomotor nerve

75. 60 year old man came to ER with complain of paralysis on left side of lower limb with no difficulty in his speech, which artery is most likely to be involved
a. Left anterior cerebral artery
b. Left middle cerebral artery
c. Right anterior cerebral artery
d. Right middle cerebral artery

74. All of the following take part in circle of Willis except
a. Anterior cerebral artery
b. Middle cerebral artery
c. Posterior cerebral artery
d. Anterior communicating artery

75. The most medial intracerebellar nuclei is
a. Fastigial
b. Globose
c. Emboliform
d. Dentate

54. The anterior fontanelle in fetal skull usually closes at?
a. Birth
b. One year after birth
c. One and half year after birth
d. Two years after birth

55. A malignant tumor is damaging the patient's glossopharyngeal nerve, they will experience
a. Loss of taste over the anterior two – third of tongue
b. Loss of pain/thermo sensation over anterior two third of the tongue
c. Loss of taste of sensation over posterior third of tongue
d. Paralysis of muscle of tongue

56. Examination of a patient indicates that he has a medially directed strabismus (squint). This could be due to damage to the
a. Oculomotor nerve
b. Trochlear nerve
c. Ophthalmic
d. Abducent nerve

57. One of the following nucleus related to visual body relay ?
a. Lateral geniculate body
b. Edinger westphal nucleus
c. Pretectal nucleus
d. Superior colliculus

58. The sensory decussation is formed by?
a. Decussation of variable spinal tract
b. Internal arcuate fibers
c. Decussation of superior cerebral peduncle
d. Decussation of rubrospinal tracts

59. All of the following are included in limbic system except
a. Amygdaloid nucleus
b. Fornix
c. Parahippocampal gyrus
d. Hypothalamus

60. Regarding basal nuclei
a. They are composed of white matter
b. They are present to cortex at posterior perforated substance
c. Caudate nucleus and putamen together forms neostriatum
d. Amygdala is part of striatum nucleus

61. Vertebral artery
a. Is a branch of second part of subclavian artery
b. Gives off posterior meningeal artery
c. Passes through lower 6 cervical vertebra
d. Enters the cranium through foramen magnum

62. The brachial is enclosed in
a. Sheet of connective
b. Prolongation of foramen magnum
c. Prolongation of deep fascia
d. Posterior cutaneous nerve

Mandibular nerve
Middle meningeal artery
Pituitary stalk

64. All of the following nuclei are present beneath the floor of fourth ventricle except
a. Dorsal nucleus of vagus nerve
b. Vestibular nuclei
c. Hypoglossal nucleus
d. Facial nucleus

65. 4th ventricle communicates with all EXCEPT
a. Arachnoid cisterns
b. Arachnoid villi
c. Subarachnoid space
d. Spinal canal

66. All sensations reach cerebral cortex via
a. Thalamus
b. Internal capsule
c. Hypothalamus
d. Cerebellum

67. 20 years old girl present with sings of meningitis, during her investigation, lumber puncture block at $L_3 - L_4$ level alternate route for obtaining CSF will be
a. Between $L_2 - L_3$
b. Between $L_4 - S_1$
c. Of foramen magnum and C_1
d. Between $C_1 - C_2$

68. Forcep minor of corpus callosum are the fibers of
a. Rostrum
b. Genu
c. Body
d. Splenium

69. Great cerebral vein drain into:
a. Transverse sinus
b. Inferior sagittal sinus
c. Confluence of sinuses
d. Straight sinus

70. A 40 years old man present in causality of hospital with hemiplegic headache and difficulty in speaking. Finding of CT scan shows a large hematoma at the area of pterion those characteristic finding are due to.
a. Lesion at pyramidal decussation
b. Epidural hemorrhage
c. Lower moto- neuron lesion
d. Lesion at motor cortex

23. Syndesmosis is:
a. A cartilaginous joint
b. Present in skull
c. Formed by fibers and membrane
d. Biaxial in movement

24. Elastic cartilage is found in:
a. Ear ossicles
b. Nasal septum
c. Epiglottis
d. Intervertebral disc

25. Articular disc of temporomandibular joint is:
a. Made up of elastic cartilage
b. Divides joint cavity into anterior and posterior compartment
c. Receives insertion of medial pterygoid on its anterior margin
d. Slides forward during protraction

26. The carotid sheath and its contents may be safely retracted as a unit during surgical procedures of the neck. The contents of the carotid sheath include all the following structures EXCEPT:
a. Common carotid artery
b. Internal carotid artery
c. Sympathetic trunk
d. Vagus nerve

27. The vocal ligament is the thickened upper margin of the lateral portion of the
a. Quadrangular membrane
b. Hyoepiglottic ligament
c. Thyroepiglottic ligament
d. Cricothyroid ligament

28. Destruction of nerve cell bodies found in the pterygopalatine fossa would result in which of the following?
a. Dilated pupil
b. Dry eye
c. Loss of taste at the root of tongue
d. Dry mouth

29. The patient's left cornea is in abducted and depressed position and the right cornea appears to be in the normal position. Which nerve is injured on the left side?
a. Oculomotor
b. Trochlear
c. Abducent
d. Frontal

30. The skin over the angle of the mandible and over most of the parotid gland is supplied by
a. Mandibular branch of facial nerve
b. Great auricular nerve
c. Lesser occipital nerve
d. Greater occipital nerve

31. The motor branch of cervical plexus is:
a. Suboccipital nerve
b. Lesser occipital nerve
c. Great auricular nerve
d. Supraclavicular nerve

32. The layer of cervical fascia which forms a fascial sheath for the parotid gland is the
a. Prevertebral
b. Pretracheal
c. Investing layer
d. Retropharyngeal

19. Spleen has:
a. Loose connective tissue capsule
b. Lymphoid follicles around arterioles
c. Hassell's corpuscles
d. Crypts lined by stratified squamous epithelium

20. Hyaline cartilage:
a. Present in epiglottis
b. Forms inter vertebral disc
c. Has basophilic matrix with meta chromatic reaction
d. Always lined by perichondrium

21. The pneumatic bone
a. Are the type of flat bone
b. Act as paranasal sinuses
c. Contains serous fluid inside
d. Present at the skull cap

22. Example of secondary cartilaginous joint is
a. 1st costochondral joint
b. Joint between epiphysis and diaphysis
c. Manibrusternal joint
d. Sternoclavicular joint

23. In dissection of human body, a section right angle to long axis of the organ is
a. Oblique section
b. Longitudinal section
c. Transverse section
d. Horizontal section

24. In typical spinal nerve:
a. Posterior ramus contain autononic fibers
b. Anterior fiber root contains sensory fiber
c. Lateral column contain parasympathetic nuceli
d. Trunk contain sensory ganglion

25. Elastic cartilage is found in:
a. Ear ossicles
b. Nasal septum
c. Epiglottis
d. Intervertebral disc

26. Regarding blood supply of bones
a. Diaphysial arteries anastomose with metaphyseal
b. Epiphysial arteries are the branches of metaphyseal
c. Metaphyseal arteries are the branches of a nutrient artery
d. Nutrient artery supply epiphysis

27. Grey rami communicants
a. Preganglionic fibers
b. Arises from sympathetic chain
c. Terminate inside the organs
d. Covered by myline sheath

45. Muscle of mastication is supplied by
a. 1st part of maxillary artery
b. 2nd part of maxillary artery
c. 3rd part of maxillary artery
d. Facial artery

46. A patient came to you with dry eyes due to decreased lacrimal gland secretion. This may be due to following ganglion
a. Ciliary ganglion
b. Pterygopalatine ganglion
c. Otic ganglion
d. Submandibular ganglion

47. Infection from face to cavernous sinus can easily spread by:
a. Pterygoid venous plexus
b. External jugular vein
c. Internal jugular vein
d. Ophthalmic vein

48. Maxillary artery terminates as:
a. Infraorbital artery
b. Sphenopalatine artery
c. Palatine artery
d. Nasopalatine artery

49. In hanging , death occurs due to fracture of:
a. Base of skull
b. Odontoid process of axis
c. Body of atlas
d. Arch of atlas

50. Regarding inferior alveolar nerve:
a. It is purely sensory
b. Purely motor
c. Mixed nerve
d. Parasympathetic nerve

51. Lower gums are innervated by:
a. Greater palatine nerve
b. Lingual nerve
c. Anterior superior alveolar nerve
d. Posterior superior alveolar nerve

52. The palatoglossal arch is formed by which of the following muscle:
a. Tensor velli palatine
b. Levator velli palatine
c. Palatoglossus muscle
d. Palate pharygeus

53. Regarding left common carotid artery:
a. It arises from brachiocephalic trunk
b. It is embedded in carotid sheath
c. It gives two carotid and superior thyroid artery
d. It lies medial to internal jugular vein

Not sure

54. The anterior fontanelle in fetal skull usually closes at?
a. Birth
b. One year after birth
c. One and half year after birth (18 months)
d. Two years after birth

55. A malignant tumor is damaging the patient is glossopharyngeal nerve, they will experience
a. Loss of taste over the anterior two – third of tongue
b. Loss of somaesthetic sensation over anterior two third of the tongue
c. Loss of taste of sensation over posterior third of tongue
d. Paralysis of muscle of tongue

56. Examination of a patient indicates that he has a medially directed strabismus (squint). This could be due to damage to the.
a. Oculomotor nerve
b. Trochlear nerve
c. Ophthalmic
d. Abducent nerve

57. One of the following nucleus related to visual body relay ?
a. Lateral geniculate body
b. Edinger westpal nucleus
c. Pretectal nucleus
d. Superior colliculus

58. The sensory decussation is formed by?
a. Decussation of cortico spinal tract
b. Internal arcuate fibers
c. Descussation of superior cerebral peduncle
d. Descussation of rubrospinal tracts

59. All of the following are included in limbic system except:
a. Amygdaloid nucleus
b. Fornix
c. Parahippocampal gyrus
d. Hypothalamus

60. Regarding basal nuclei grey
a. They are composed of white matter
b. They are joined to cortex at posterior perforated substance
c. Caudate nucleus and putamen together forms neostriatum
d. Amygdale is part of lentiform nucleus
caudate

61. Vertebral artery
a. Is a branch of second part of subclavian artery
b. Gives off posterior cerebral artery
c. Passes through lower 6 central vertebral
d. Enters the cranium through foramen magnum

62. Dura matter is attached to:
a. Body of sphenoid
b. Margins of foramen magnum
c. Cribriform plate of ethmoid
d. petrous temporal bone

63. cavernous sinus contains
a. Sympathetic plexus
b. Mandibular nerve
c. Middle meningeal artery
d. Pituitary stalk

64. All of the following nuclei are present beneath the floor of fourth ventricle except:
a. Dorsal nucleus of vagus nerve
b. Vestibular nuclei
c. Hypoglossal nucleus
d. Facial nucleus

65. 4th ventricle communicates with all EXCEPT:
a. Arachnoid cisterns
b. Arachnoid villi
c. Subarachnoid space
d. Spinal canal

66. All sensations reach cerebral cortex via:
a. Thalamus
b. Internal capsule
c. Hypothalamus
d. Cerebellum

67. 20 years old girl present with sings of meningitis, during her investigation, lumber puncture block at $L_3 - L_4$ level alternate route for obtaining CSF will be
a. Between $L_2 - L_3$
b. Between $L_5 - S_1$
c. Of foramen magnum and C_1
d. Between $C_1 - C_2$

68. Forcep minor of corpus callosum are the fibers of
a. Rostrum
b. Genu
c. Body
d. Splenium

69. Great cerebral vein drain into:
a. Transverse sinus
b. Inferior sagittal sinus
c. Confluence of sinuses
d. Straight sinus

70. A 40 years old man present in causality of hospital with hemiplegic headache and difficulty in speaking Finding of CT scan shows a large hematoma at the area of pterion those characteristic finding are due to:
a. Lesion at pyramidal decussation
b. Epidural hemorrhage
c. Lower motor neuron lesion
d. Lesion at motor cortex

a. Ciliary ganglion
b. Pterygopalatine ganglion
c. Otic ganglion
d. Submandibular ganglion

47. Infection from face to cavernous sinus can easily spread by
a. Frontal sensory nerve
b. External nasal vein
c. Internal nasal vein
d. Ophthalmic vein

48. Maxillary artery terminates as
a. Infraorbital artery
b. Sphenopalatine artery
c. Palatine artery
d. Nasopalatine artery

49. In hanging , death occurs due to fracture of
a. Base of skull
b. Odontoid process of axis
c. Body of atlas
d. Arch of atlas

50. Regarding inferior alveolar nerve:
a. It is purely sensory
b. Purely motor
c. Mixed nerve
d. Parasympathetic nerve

51. Lower gums are innervated by:
a. Greater palatine nerve
b. Lingual nerve
c. Anterior superior alveolar nerve
d. Posterior superior alveolar nerve

52. The palatoglossal arch is formed by which of the following muscle
a. Tensor velli palatine
b. Levator velli palatine
c. Palatoglossus muscle
d. Palate pharygeus

53. Regarding left common carotid artery:
a. It arises from brachiocephalic trunk
b. It is embedded in carotid sheath
c. It gives two carotid and superior thyroid artery
d. Medial to internal jugular vein

Example of [illegible] muscle
a. Deltoid
b. Latissimus dorsi
c. Biceps brachii
d. Sternocleidomastoid

29. In quadrupeds a median plane is
a. Superiorly
b. Anteriorly
c. Posteriorly
d. Inferiorly

30. Regarding joint s which one of the following is not true?
a. Primary cartilaginous joints are termed as synchondroses
b. Synovial joint occurs between manubrium and body of sternum
c. Hinge joint is example of uniaxial type of synovial joint
d. Inferior tibiofibular joint is the example of syndesmosis

31. The posterior belly of the diagastric muscle is innervated by [illegible]
a. Trigeminal nerve
b. Facial nerve
c. Glossopharyngeal nerve
d. Accessory nerve

32. The cricoids cartilage lies at this vertebral level
a. Second cervical vertebrae
b. Fourth cervical vertebrae
c. Sixth cervical vertebrae
d. Eight cervical vertebrae

33. Medical student studying the venous drainage of brain, reveals that the sinus which is continous as internal jugular vein is:
a. Sigmoid sinus
b. Inferior petrosal sinus
c. Cavernous sinus
d. Superior petrosal sinus

34. The ganglion which lies in infratemporal fossa is

1. The part of the sperm containing proteolytic enzymes to digest the zona pellucida is the:
   a. Capacitor
   b. Head → contains nucleus with DNA
   c. Corona
   d. Acrosome → contains enzymes
   Middle piece → contains mitochondria → that provide energy

2. The ovulated mammalian oocyte is arrested at:
   a. Prophase of meiosis I
   b. Metaphase of meiosis I
   c. Prophase of meiosis II
   d. Metaphase of meiosis II

3. Somites:
   a. Differentiate into myotomes which give rise to skeletal muscle in trunk
   b. Differentiate into sclerotomes which give rise to vertebrae
   c. Arise from segmentation of the paraxial mesoderm
   d. All of the above are correct

4. When the amount of amniotic fluid exceeds two liters, the condition is called:
   a. Oligohydramnios decreased amniotic fluid (<300ml)
   b. Polyhydramnios increased amniotic fluid >2L
   c. Amniotitis infection in uterus
   d. Hydrogravida

5. After parturition a gynecologist identified the maternal surface of placenta due to 15-20 bulging areas called:
   a. Cotyledons.
   b. Placental septa.
   c. Intervillous space.
   d. Cytotrophoblastic shell.

6. A 21 year old female present to the obstetrician at 16 weeks of prenatal care. The obstetrician finds increased amniotic fluid and orders alpha fetoprotein and HCG levels. One week later the called patient back to tell her that the alpha fetoprotein level is increased. The most probable fetal diagnosis is

   a. Anencephaly
   b. Jejunal atresia
   c. Renal agenesis
   d. Sacrococcgeal teratoma

7. Fetal membranes are as under EXCEPT
   a. Amnion
   b. Allantois
   c. Germ disc
   d. Chorion

8. At the time of ovulation there comes a surge of
   a. Estrogen
   b. Progesterone
   c. Follicle stimulating hormone
   d. Luteinizing hormone

9. In case of Dizygotic twins there is found to be
   a. One chorionic cavity
   b. Two amniotic cavities
   c. One umbilical cord
   d. One connecting stalk

Dizygotic Twins → 2 amniotic cavities, 2 chorion, 2 placenta.
Monozygotic Twins → 2 amniotic cavities, 1 chorion, 1 placenta.

28. Example of twisted fasciculi (PB 137 GA)
a. Deltoid Oblique → Multipennate
b. Latismus dorsi
c. Bicep brachi Parallel → fusiform
d. Sternocleidomastoid twisted but crossed so cruciate

29. In quadrupeds a median places
a. Superiorly
b. Anteriorly
c. Posteriorly
d. Inferiorly

30. Regarding joint s which one of the following is not true?
a. Primary cartilaginous joints are termed as synchondroses
b. Synovial joint occurs between manubrium and body of sternum
c. Hinge joint is example of uniaxial type of synovial joint
d. Inferior tibiofibular joint is the example of syndesmosis

31. The posterior belly of the diagastric muscle is innervated by a branch of this cranial nerve.
a. Trigeminal nerve
b. Facial nerve
c. Glossopharyngeal nerve
d. Accessory nerve

32. The cricoids cartilage lies at this vertebral level.
a. Second cervical vertebrae
b. Fourth cervical vertebrae
c. Sixth cervical vertebrae
d. Eight cervical vertebrae

33. Medical student studying the venous drainage, of brain, reveals that the sinus which is continous as internal jugular vein is:
a. Sigmoid sinus
b. Inferior petrosal sinus
c. Cavernous sinus
d. Superior petrosal sinus

34. The ganglion which lies is infratemporal fossa is:
a. Otic ganglion
b. Ciliary ganglion
c. Pterygopalatine ganglion
d. Submandibular ganglion

35. A patient came in emergency with bleed from nose –X-Ray shows the fracture of floor of the nose. The most probably involved bone will be:
a. Vomer
b. Cribriform plate of ethmoid
c. Palatine process of maxilla
d. Perpendicular plate of palatine

36. The opening which has connection with middle meatus is
a. Sphenoid sinus
b. Frontal sinus
c. Lacrimal duct
d. Posterior ethmoidal sinus

a	Occulomotor nerve	f	Cranial accessory nerve
b	Mandibular nerve	g	Hypoglossal nerve
c	Abducent nerve	h	Vagus nerve
d	Facial nerve	i	Opthalmic nerve
e	Spinal accessory nerve	J	Glossopharyngeal nerve

Select the correct cranial nerve from the above mentioned options that is most likely to be damaged resulting in the following clinical conditions

21. A Patient complains to the general physician that in the early morning when she tried to get up from the bed she is unable to move her head but can move her body.
22. A hypertensive patient in the emergency ward presents with double vision. On examination the pupils were dilated
23. During a routine physical examination, when the patient was asked to show his teeth. he was unable to do so.
24. A patient came to the general physician and complains that during chewing she is unable to touc the floor of the oral cavity.
25. When an examiner asks a child to pronounce "A", "O", "E", the child was unable to do so.

EXTENDED MATCHING QUESTION

EACH OPTION CAN BE SELECTED ONCE, MORE THAN ONCE OR NOT AT ALL.

a	Head fold	f	philtrum
b	Nucleus pulposus	g	teratoma
c	Hydatidform mole	h	Upper incisor teeth
d	epiglottis	i	Phocomelia
e	Branchial cyst	j	Palatine tonsil

Lead in: Choose the correct option and write it against each stem

1. The structure derived from the intermaxillary segment of the embryonic face
2. Primitive streak fails to degenerate in fetal life
3. A normal remnant of an embryonic structure found in adult body
4. are found along the anterior border of the sternocleidomastoid muscle
5. The hypobranchial eminence contributes to the formation of

---

a	Pseudo stratified columnar epithelium	f	Thymus
b	Cuboidal epithelium	g	Palatine tonsil
c	Dentin	h	Osteon
d	Spinal cord	i	Osteoid
e	cerebellum	j	Cementum

Regarding histological features choose the best option for the following statements:

6. Unmineralized bone matrix
7. Crypts lined by stratified squamous Non- Keratinized epithelium
8. Covering of root surface of tooth
9. Gray commissure with central canal
10. Lining epithelium of thyroid follicles

---

a	Suture	f	Synovial Hinge joint
b	Syndesmosis	g	Synovial saddle joint
c	Gomphoses	h	Synovial pivot joint
d	Synchondrosis	i	Synovial bicondylar joint
e	Symphysis	j	Synovial ellipsoid joint

Choose the appropriate category of joints from the mentioned above for the following examples of joint with the given specific characteristics.

11. Between tooth and alveolar socket
12. Typically present in median plane
13. That permits rotation of thumb
14. Limited to skull bone
15. With fibrous membrane

71. A patient complains of difficulty in swallowing , injury to which of the following nerve is cause of this complain:

a. Cranial part of accessory nerve
b. Spinal part of accessory nerve
c. Hypoglossal nerve
d. 1st cervical nerve

72. Nucleus ambigus is common for following nerves Except:

a. Vagus nerve
b. Accessory nerve
c. Hyopoglossal nerve
d. Glossopharyngeal nerve

73. Cavernous sinus coutains:

a. Sympathetic polexus
b. Mandibular nerve
c. Middle menengial artery
d. Pituitary stalk

74. With in the internal capsule, the sensory radiations from thalamus to Brodmann's area 3,2,1 run in the :

a. Anterior ½ of posterior limb
b. Posterior ½ of posterior limb
c. Whole of the posterior limb
d. Whole of the anterior limb

75. Forceps minor of corpus callosum are the fibers of :

a. Rostrum
b. Genu
c. Body
d. Splenium

EACH OPTION CAN BE SELECTED ONCE, MORE THAN ONCE OR NOT AT ALL.

a	Head fold	f	philtrum
b	Nucleus pulposus	g	teratoma
c	Hydatidform mole	h	Upper incisor teeth
d	epiglottis	i	Phocomelia
e	Branchial cyst	j	Palatine tonsil

Lead in: Choose the correct option and write it against each stem

1. The structure derived from the intermaxillary segment of the embryonic face
2. Primitive streak fails to degenerate in fetal life
3. A normal remnant of an embryonic structure found in adult body
4. are found along the anterior border of the sternocleidomastoid muscle
5. The hypobranchial eminence contributes to the formation of

a	Pseudo stratified columnar epithelium	f	Thymus
b	Cuboidal epithelium	g	Palatine tonsil
c	Dentin	h	Osteon
d	Spinal cord	i	Osteoid
e	cerebellum	j	Cementum

Regarding histological features choose the best option for the following statements:

6. Unmineralized bone matrix
7. Crypts lined by stratified squamous Non- Keratinized epithelium
8. Covering of root surface of tooth
9. Gray commissure with central canal
10. Lining epithelium of thyroid follicles

a	Suture	f	Synovial Hinge joint
b	Syndesmosis	g	Synovial saddle joint
c	Gomphoses	h	Synovial pivot joint
d	Synchondrosis	i	Synovial bicondylar joint
e	Symphysis	j	Synovial ellipsoid joint

Choose the appropriate category of joints from the mentioned above for the following examples of joint with the given specific characteristics.

11. Between tooth and alveolar socket
12. Typically present in median plane
13. That permits rotation of thumb
14. Limited to skull bone
15. With fibrous membrane

d. Has nucleoli

12. Stratified squamous epithelium
a. Develops only from ectoderm
b. Found in pelvis of kidney
c. Continuously replaced by basal layer
d. Has superficial nucleated layer

13. Collagen fibres has following characteristic except:
a. Responsible for tensile strength
b. Unbranched fibres
c. Appear banded on microscopy
d. Produced by mast cells

14. Skeletal muscle fibres have
a. Peripheral single nuclei
b. T system of tubules
c. Intercalated disc
d. Sarcomere between two I bands

15. Osteoblasts
a. Forms collagen
b. Remove bone during remodeling
c. Present in lacunae
d. Derived from osteocytes

16. Lymph node has:
a. Connective tissue framework of collagen
b. Valveless single afferent lymph vessel
c. T lymphocytes in cortex
d. B lymphocytes both in cortex and medulla

17. Tongue histology reveals that :
a. Filiform papillae has taste buds
b. Lymphoid tissue is confined to tip
c. Mucous Gland numerous in anterior part
d. Muscles fibres are predominantly skeletal

18. The salivary Glands:
a. Demilunes consist of mucous cells
b. Serous cells are eosinophilic
c. Striated ducts are lined by myoepithelial cells
d. Lack adi pose tissue

37. Temporomandibular joint is innervated by:
a. Facial nerve
b. Mandibular nerve
c. Maxillary nerve
d. Vagus nerve

38. In the little area of Nasal cavity, all of following arteries anastomose EXCEPT:
a. Superior labial
b. Greater palatine
c. Spheno palatine
d. Posterior ethmoidal

39. A transient facial paralysis may result from entrance of anesthetic agent lingual nerve block into
a. Parotid capsule
b. Parotid fascia
c. Pterygomandibular raphe
d. Buccal fascia

40. The oral diaphragm is formed by which one of the following muscle:
a. Geniohyoid
b. Mylohyoid
c. Hyoglossus
d. Genioglossus

41. Anterior dislocation of temporomandibular joint is limited by
a. Sphenomandibular ligament
b. Stylomandibular ligament
c. Lateral ligament
d. Arcuate eminence

42. A young batsman received serve injury by a fast bouncer ball hitting his neck in posterior auricular region causing unconsciousness – injury to which structure had caused unconsciousness.
a. Occipital artery
b. Internal carotid artery
c. Basilar artery
d. Vertebral artery

43. A patient who had undergone the surgery to remove right lobe of thyroid gland, past operation his voice reduces to a whisper, which of following nerve was damaged?
a. Right external laryngeal
b. Right recurrent laryngeal
c. Left recurrent laryngeal
d. Left superior laryngeal

44. Which of the following is the dangerous layer of scalp
a. Pericranium
b. Loose areolar tissue
c. Galea aponeurosis
d. Skin

54. The anterior fontanelle in fetal skull usually closes at?
a. Birth
b. One year after birth
c. One and half year after birth
d. Two years after birth

55. A malignant tumor is damaging the patient is glossopharyngeal nerve, they will experience
a. Loss of taste over the anterior two – third of tongue
b. Loss of somaesthetic sensation over anterior two third of the tongue
c. Loss of taste of sensation over posterior third of tongue
d. Paralysis of muscle of tongue

56. Examination of a patient indicates that he has a medially directed strabismus (squint). This could be due to damage to the.
a. Oculomotor nerve
b. Trochlear nerve
c. Ophthalmic
d. Abducent nerve

57. One of the following nucleus related to visual body relay ?
a. Lateral geniculate body
b. Edinger westpal nucleus
c. Pretectal nucleus
d. Superior colliculus

58. The sensory decussation is formed by?
a. Decussation of cortico spinal tract
b. Internal arcuate fibers
c. Descussation of superior cerebral peduncle
d. Descussation of rubrospinal tracts

59. All of the following are included in limbic system except:
a. Amygdaloid nucleus
b. Fornix
c. Parahippocampal gyrus
d. Hypothalamus

60. Regarding basal nuclei
a. They are composed of white matter
b. They are joined to cortex at posterior perforated substance
c. Caudate nucleus and putamen together forms neostriatum
d. Amygdale is part of lentiform nucleus

61. Vertebral artery
a. Is a branch of second part of subclavian artery
b. Gives off posterior cerebral artery
c. Passes through lower 6 central vertebral
d. Enters the cranium through foramen magnum

62. Dura mater is attached to:
a. Body of sphenoid
b. Margins of foramen magnum
c. Cribriform plate of ethmoid
d. petrous temporal bone

cavernous sinus contains
Sympathetic plexus
Mandibular nerve
Middle meningeal artery
Pituitary stalk

64. All of the following nuclei are present beneath the floor of fourth ventricle except:
a. Dorsal nucleus of vagus nerve
b. Vestibular nuclei
c. Hypoglossal nucleus
d. Facial nucleus

65. 4th ventricle communicates with all EXCEPT:
a. Arachnoid cisterns
b. Arachnoid villi
c. Subarachnoid space
d. Spinal canal

66. All sensations reach cerebral cortex via:
a. Thalamus
b. Internal capsule
c. Hypothalamus
d. Cerebellum

67. 20 years old girl present with sings of meningitis, during her investigation, lumber puncture block at $L_3 - L_4$ level alternate route for obtaining CSF will be
a. Between $L_2 - L_3$
b. Between $L_5 - S_1$
c. Of foramen magnum and C_1
d. Between $C_1 - C_2$

8. Forcep minor of corpus callosum are the fibers of
Rostrum
Genu

EACH OPTION CAN BE SELECTED ONCE, MORE THAN ONCE OR NOT AT ALL.

a	Head fold	f	philtrum
b	Nucleus pulposus	g	teratoma
c	Hydatidform mole	h	Upper incisor teeth
d	epiglottis	i	Phocomelia
e	Branchial cyst	j	Palatine tonsil

Lead in: Choose the correct option and write it against each stem

1. The structure derived from the intermaxillary segment of the embryonic face
2. Primitive streak fails to degenerate in fetal life
3. A normal remnant of an embryonic structure found in adult body
4. are found along the anterior border of the sternocleidomastoid muscle
5. The hypobranchial eminence contributes to the formation of

a	Pseudo stratified columnar epithelium	f	Thymus
b	Cuboidal epithelium	g	Palatine tonsil
c	Dentin	h	Osteon
d	Spinal cord	i	Osteoid
e	cerebellum	j	Cementum

Regarding histological features choose the best option for the following statements:

6. Unmineralized bone matrix
7. Crypts lined by stratified squamous Non- Keratinized epithelium
8. Covering of root surface of tooth
9. Gray commissure with central canal
10. Lining epithelium of thyroid follicles

a	Suture	f	Synovial Hinge joint
b	Syndesmosis	g	Synovial saddle joint
c	Gomphoses	h	Synovial pivot joint
d	Synchondrosis	i	Synovial bicondylar joint
e	Symphysis	j	Synovial ellipsoid joint

Choose the appropriate category of joints from the mentioned above for the following examples of joint with the given specific characteristics.

11. Between tooth and alveolar socket
12. Typically present in median plane
13. That permits rotation of thumb
14. Limited to skull bone
15. With fibrous membrane

47. All the nuclei in brain stem are parasympathetic **except**:

a. Superior salivatory
b. Dorsal vagal
c. Edinger Westpaul
d. Tractus solitarious

48. The white rami communicants :

a. Are demylinated fiber
b. Distributed to thoracic viscera
c. Are post synaptic to sympathetic chain
d. Courses in anterior root of spinal nerve

49. Sympathetic chain:

a. Consist of unilateral 31 ganglion
b. Consolidate cervical ganglion are three on each side
c. Chain terminates at stellate ganglion
d. Receive pre ganglion fiber from anterior horn of spinal cord

of growing bone:

The area of cerebral cortex listed below receive their arterial supply as indicated:
a. The precentral gyrus is supplied by the middle cerebral artery
b. The post central gyrus (face area) is supplied by anterior cerebral artery
c. The cuneus is supplied by anterior cerebral artery
d. The Wernicke area is supplied by posterior cerebral artery

72. The motor speech area of Broca:
a. In most individual, important on left dominant hemisphere
b. Brings about the formation of words by its connections with secondary motor area
c. Connected to sensory speech area of wernicke
d. Brodmann area 34 and 35 represent motor speech area

73. The posterior surface of medulla oblongata shows.
a. Pyramids
b. Olives
c. Hypoglossal triangle
d. Middle cerebellar peduncle

74. The nerve emerging between pon and medulla oblongata is
a. Glossopharyngeal nerve
b. Facial nerve
c. Vagus nerve
d. Occulomotor nerve

75. 60 year old man came to ER with complain of paralysis on left side of lower limb with no difficulty in his speech , which artery is most likely to be involved.
a. Left anterior cerebral artery
b. Left middle cerebral artery
c. Right anterior cerebral artery
d. Right middle cerebral artery

a	Parotid sheath	f	Auriculotemporal nerve
b	Retromandibular vein	g	Great auricular nerve
c	Facial nerve	h	Mastoid process
d	External carotid artery	i	Transverse cervical artery
e	Preauricular lymph nodes	j	Maxillary artery

Select the correct structure located in the parotid region from the above mentioned options which is most likely to be involved in the following conditions

16. In mumps, the parotid gland swells resulting in the excruciating pain during chewing.
17. During parotidectomy, the most superficial structure lying within the tissue of parotid gland.
18. During the dissection of the parotid region, the structure that indents the posteromedial surface of the parotid gland.

a	Medial pterygoid plate	f	Spine of sphenoid
b	Greater wing of sphenoid	g	Internal acoustic meatus
c	Lateral pterygoid plate	h	Pterygomaxillary fissure
d	Juglar foramen	i	Infraorbital fissure
e	Styloid process	j	Lesser wing of sphenoid

Select the most appropriate bony structure from the above mentioned list that is most likely to be related to the description given below.

19. Lying lateral to it is the auriculotemporal nerve and media to it is the chorda tympani nerve.
20. The bony opening has the fundus consisting of a bony plate called lamina cribrosa.

a	Occulomotor nerve	f	Cranial accessory nerve
b	Mandibular nerve	g	Hypoglossal nerve
c	Abducent nerve	h	Vagus nerve
d	Facial nerve	i	Opthalmic nerve
e	Spinal accessory nerve	j	Glossopharyngeal nerve

Select the correct cranial nerve from the above mentioned options that is most likely to be damaged resulting in the following clinical conditions

21. A Patient complains to the general physician that in the early morning when she tried to get up from the bed she is unable to move her head but can move her body.
22. A hypertensive patient in the emergency ward presents with double vision. On examination the pupils were dilated
23. During a routine physical examination, when the patient was asked to show his teeth, he was unable to do so.
24. A patient came to the general physician and complains that during chewing she is unable to touc the floor of the oral cavity.
25. When an examiner asks a child to pronounce "A", "O", "E", the child was unable to do so.

a	Parotid sheath	f	Auriculotemporal nerve
b	Retromandibular vein	g	Great auricular nerve
c	Facial nerve	h	Mastoid process
d	External carotid artery	i	Transverse cervical artery
e	Preauricular lymph nodes	J	Maxillary artery

Select the correct structure located in the parotid region from the above mentioned options which is most likely to be involved in the following conditions

16. In mumps, the parotid gland swells resulting in the excruciating pain during chewing.
17. During parotidectomy, the most superficial structure lying within the tissue of parotid gland.
18. During the dissection of the parotid region, the structure that indents the posteromedial surface of the parotid gland.

a	Medial pterygoid plate	f	Spine of sphenoid
b	Greater wing of sphenoid	g	Internal acoustic meatus
c	Lateral pterygoid plate	h	Pterygomaxillary fissure
d	Juglar foramen	i	Infraorbital fissure
e	Styloid process	J	Lesser wing of sphenoid

Select the most appropriate bony structure from the above mentioned list that is most likely to be related to the description given below.

19. Lying lateral to it is the auriculotemporal nerve and media to it is the chorda tympani nerve.
20. The bony opening has the fundus consisting of a bony plate called lamina cribrosa.

a	Occulomotor nerve	f	Cranial accessory nerve
b	Mandibular nerve	g	Hypoglossal nerve
c	Abducent nerve	h	Vagus nerve
d	Facial nerve	i	Opthalmic nerve
e	Spinal accessory nerve	J	Glossopharyngeal nerve

Select the correct cranial nerve from the above mentioned options that is most likely to be damaged resulting in the following clinical conditions

21. A Patient complains to the general physician that in the early morning when she tried to get up from the bed she is unable to move her head but can move her body.
22. A hypertensive patient in the emergency ward presents with double vision. On examination the pupils were dilated
23. During a routine physical examination, when the patient was asked to show his teeth, he was unable to do so.
24. A patient came to the general physician and complains that during chewing she is unable to touch the floor of the oral cavity.
25. When an examiner asks a child to pronounce "A", "O", "E", the child was unable to do so.

22. Disorders of the basal ganglia include all of the following except:

A. Chorea
B. Athetosis
C. Trigeminal neuralgia
D. Parkinsonism

23. Circle of willis is formed by the following except:

A. Anterior cerebral artery
B. Anterior choroidal artery
C. Posterior communicating artery
D. Posterior cerebral artery

24. The primary visual area of broadmann is located in the:

A. superior parietal lobule
B. Calcarine sulcus
C. Parieto-occipital sulcus
D. Frontal lobe

25. WHICH OF THE FOLLOWING STATEMENTS CHARACTERIZES OSTEOCLASTS:

A. Uni-nucleated cells
B. Secrete osteiod
C. Derived from osteoprogenitor cells
D. Occupy Howship lacunae

26. THE TYPE OF CARTILAGE IN EXTERNAL EAR:

A. Hyaline
B. Elastic
C. Fibrocartilage
D. All of the above

27. REGARDING OSTEON:

A. The central component is the Haversian canal
B. Composed of concentric lamellae of mineralized matrix
C. Primary cell type in osteon is the osteoclast
D. Blood vessels reach osteon from bone marrow via Volkmann's canals

28. BIPOLAR NEURONS ARE FOUND IN:

A. Dorsal root ganglia
B. Ventral horns of spinal cord
C. White matter of cerebrum
D. Olfactory epithelium

29. WHICH TYPE OF CELL PROVIDES STRUCTURAL AND NUTRITIONAL SUPPORT TO NEURONES IN THE CNS?

A. Microglia
B. Schwann cells
C. Astrocytes
D. Ependymal cells

71. The area of cerebral cortex listed below receive their arterial supply as indicated:
a. The precentral gyrus is supplied by the middle cerebral artery
b. The post central gyrus (face area) is supplied by anterior cerebral artery
c. The cuneus is supplied by anterior cerebral artery
d. The Wernicke area is supplied by posterior cerebral artery

72. The motor speech area of Broca:
a. In most individual, important on left dominant hemisphere
b. Brings about the formation of words by its connections with secondary motor area primary
c. Connected to sensory speech area of wernicke
d. Brodmann area 34 and 35 represent motor speech area
44 45

73. The posterior surface of medulla oblongata shows.
a. Pyramids
b. Olives
c. Hypoglossal triangle
d. Middle cerebellar peduncle

74. The nerve emerging between pon and medulla oblongata is
a. Glossopharyngeal nerve
b. Facial nerve
c. Vagus nerve
d. Occulomotor nerve

75. 60 year old man came to ER with complain of paralysis on left side of lower limb with no difficulty in his speech , which artery is most likely to be involved.
a. Left anterior cerebral artery
b. Left middle cerebral artery
c. Right anterior cerebral artery
d. Right middle cerebral artery

65.High endothelial venules are present in.

A. Thymus
B. Lymph node
C. Langerhan cell
D. Tonsil
E. Spleen

66.The pharyngeal plexus is situated at.

A.Posterolateral surface of superior constrictor
B. Posterolateral surface of middle constrictor
C.Posterolateral surface of inferior constrictor
D. In the piriform fossa

67.The following is not a content of cavernous sinus

A. Internal carotid artery
B. optic nerve .
C Trochlear nerve
D Abducent nerve

68.Excessive aphasia (inability to speak) results from damage to the

A. The motor speech area of bronca's
B. The sensory area of wernicke
C. The prefrontal cortex
D. The postcentral gyrus

69.Astereognosis results from damage to the

A. Primary sensory cortex
B. Sensory association cortex
C. Primary visual area
D. Primary auditory area

70.The facial colliculus overlies the nucleus of

A. Vagus nerve
B. Trochlear nerve
C. Oculomotor nerve
D. Abducent nerve

EXTENDED MATCHING QUESTION

EACH OPTION CAN BE SELECTED ONCE, MORE THAN ONCE OR NOT AT ALL.

a	Head fold	f	philtrum
b	Nucleus pulposus	g	teratoma
c	Hydatidform mole	h	Upper incisor teeth
d	epiglottis	i	Phocomelia
e	Branchial cyst	j	Palatine tonsil

Lead in: Choose the correct option and write it against each stem

1. The structure derived from the intermaxillary segment of the embryonic face
2. Primitive streak fails to degenerate in fetal life
3. A normal remnant of an embryonic structure found in adult body
4. are found along the anterior border of the sternocleidomastoid muscle
5. The hypobranchial eminence contributes to the formation of

---

a	Pseudo stratified columnar epithelium	f	Thymus
b	Cuboidal epithelium	g	Palatine tonsil
c	Dentin	h	Osteon
d	Spinal cord	i	Osteoid
e	cerebellum	J	Cementum

Regarding histological features choose the best option for the following statements:

6. Unmineralized bone matrix
7. Crypts lined by stratified squamous Non- Keratinized epithelium
8. Covering of root surface of tooth
9. Gray commissure with central canal
10. Lining epithelium of thyroid follicles

---

a	Suture	f	Synovial Hinge joint
b	Syndesmosis	g	Synovial saddle joint
c	Gomphoses	h	Synovial pivot joint
d	Synchondrosis	i	Synovial bicondylar joint
e	Symphysis	j	Synovial ellipsoid joint

Choose the appropriate category of joints from the mentioned above for the following examples of joint with the given specific characteristics.

11. Between tooth and alveolar socket
12. Typically present in median plane
13. That permits rotation of thumb
14. Limited to skull bone
15. With fibrous membrane

37. Temporomandibular joint is innervated by:
a. Facial nerve
b. Mandibular nerve (Masseteric + Auriculotemporal N).
c. Maxillary nerve
d. Vagus nerve

38. In the little area of Nasal cavity, all of following arteries anastomose EXCEPT:
a. Superior labial
b. Greater palatine
c. Spheno palatine
d. Posterior ethmoidal Ant ethmoidal

39. A transient facial paralysis may result from entrance of anesthetic agent lingual nerve block into
a. Parotid capsule
b. Parotid fascia
c. Pterygomandibular raphe
d. Buccal fascia

40. The oral diaphragm is formed by which one of the following muscle:
a. Geniohyoid
b. Mylohyoid
c. Hyoglossus
d. Genioglossus

41. Anterior dislocation of temporomandibular joint is limited by
a. Sphenomandibular ligament
b. Stylomandibular ligament
c. Lateral ligament
d. Arcuate eminence

42. A young batsman received serve injury by a fast bouncer ball hitting his neck in posterior auricular region causing unconsciousness – injury to which structure had caused unconsciousness.
a. Occipital artery
b. Internal carotid artery
c. Basilar artery
d. Vertebral artery

43. A patient who had undergone the surgery to remove right lobe of thyroid gland, past operation his voice reduces to a whisper, which of following nerve was damaged?
a. Right external laryngeal
b. Right recurrent laryngeal
c. Left recurrent laryngeal
d. Left superior laryngeal

44. Which of the following is the dangerous layer of scalp
a. Pericranium
b. Loose areolar tissue
c. Galea aponeurosis
d. Skin

a	Parotid sheath	f	Auriculotemporal nerve
b	Retromandibular vein	g	Great auricular nerve
c	Facial nerve	h	Mastoid process
d	External carotid artery	i	Transverse cervical artery
e	Preauricular lymph nodes	J	Maxillary artery

Select the correct structure located in the parotid region from the above mentioned options which is most likely to be involved in the following conditions

16. In mumps, the parotid gland swells resulting in the excruciating pain during chewing.
17. During parotidectomy, the most superficial structure lying within the tissue of parotid gland.
18. During the dissection of the parotid region, the structure that indents the posteromedial surface of the parotid gland.

a	Medial pterygoid plate	f	Spine of sphenoid
b	Greater wing of sphenoid	g	Internal acoustic meatus
c	Lateral pterygoid plate	h	Pterygomaxillary fissure
d	Juglar foramen	i	Infraorbital fissure
e	Styloid process	J	Lesser wing of sphenoid

Select the most appropriate bony structure from the above mentioned list that is most likely to be related to the description given below.

19. Lying lateral to it is the auriculotemporal nerve and media to it is the chorda tympani nerve.
20. The bony opening has the fundus consisting of a bony plate called lamina cribrosa.


a	Occulomotor nerve	f	Cranial accessory nerve
b	Mandibular nerve	g	Hypoglossal nerve
c	Abducent nerve	h	Vagus nerve
d	Facial nerve	i	Opthalmic nerve
e	Spinal accessory nerve	J	Glossopharyngeal nerve

Select the correct cranial nerve from the above mentioned options that is most likely to be damaged resulting in the following clinical conditions

21. A Patient complains to the general physician that in the early morning when she tried to get up from the bed she is unable to move her head but can move her body.
22. A hypertensive patient in the emergency ward presents with double vision. On examination the pupils were dilated
23. During a routine physical examination, when the patient was asked to show his teeth, he was unable to do so.
24. A patient came to the general physician and complains that during chewing she is unable to touch the floor of the oral cavity.
25. When an examiner asks a child to pronounce "A", "O", "E", the child was unable to do so.

9:41

Talha Ahmed
Active 34 minutes ago



1st Year BDS/Pre-Prof 2017 (6)

...maxillation is supplied by
...of maxillary artery
...k of maxillary artery
...rt of maxillary artery
...ial artery

46. A patient came to you with dry eyes due to decreased lacrimal gland secretion. This may be due to following ganglion.
a. Ciliary ganglion
b. Pterygopalatine ganglion
c. Otic ganglion
d. Submandibular ganglion

47. Infection from face to cavernous sinus can easily spread by:
a. Pterygoid venous plexus
b. External jugular vein
c. Internal jugular vein
d. Ophthalmic vein

48. Maxillary artery terminates as:
a. Infraorbital artery
b. Sphenopalatine artery
c. Palatine artery
d. Nasopalatine artery

49. In hanging, death occurs due to fracture of:
a. Base of skull
b. Odontoid process of axis
c. Body of atlas
d. Arch of atlas

50. Regarding inferior alveolar nerve:
a. It is purely sensory
b. Purely motor
c. Mixed nerve
d. Parasympathetic nerve

51. Lower gums are innervated by:
a. Greater palatine nerve
b. Lingual nerve
c. Anterior superior alveolar nerve
d. Posterior superior alveolar nerve

52. The palatoglossal arch is formed by which of the following muscle:
a. Tensor veli palatine
b. Levator veli palatine
c. Palatoglossus muscle
d. Palate pharyngeus

53. Regarding left common carotid artery:
a. It arises from brachiocephalic trunk
b. It is embedded in carotid sheath
c. It gives two carotid and superior thyroid artery
d. It lies medial to internal jugular vein

Remix

a	Parotid sheath	f	Auriculotemporal nerve
b	Retromandibular vein	g	Great auricular nerve
c	Facial nerve	h	Mastoid process
d	External carotid artery	i	Transverse cervical artery
e	Preauricular lymph nodes	J	Maxillary artery

Select the correct structure located in the parotid region from the above mentioned options which is most likely to be involved in the following conditions

16. In mumps, the parotid gland swells resulting in the excruciating pain during chewing.
17. During parotidectomy, the most superficial structure lying within the tissue of parotid gland.
18. During the dissection of the parotid region, the structure that indents the posteromedial surface of the parotid gland.

a	Medial pterygoid plate	f	Spine of sphenoid
b	Greater wing of sphenoid	g	Internal acoustic meatus
c	Lateral pterygoid plate	h	Pterygomaxillary fissure
d	Juglar foramen	i	Infraorbital fissure
e	Styloid process	J	Lesser wing of sphenoid

Select the most appropriate bony structure from the above mentioned list that is most likely to be related to the description given below.

19. Lying lateral to it is the auriculotemporal nerve and media to it is the chorda tympani nerve.
20. The bony opening has the fundus consisting of a bony plate called lamina cribrosa.

a	Occulomotor nerve	f	Cranial accessory nerve
b	Mandibular nerve	g	Hypoglossal nerve
c	Abducent nerve	h	Vagus nerve
d	Facial nerve	i	Opthalmic nerve
e	Spinal accessory nerve	J	Glossopharyngeal nerve

Select the correct cranial nerve from the above mentioned options that is most likely to be damaged resulting in the following clinical conditions

21. A Patient complains to the general physician that in the early morning when she tried to get up from the bed she is unable to move her head but can move her body.
22. A hypertensive patient in the emergency ward presents with double vision. On examination the pupils were dilated
23. During a routine physical examination, when the patient was asked to show his teeth, he was unable to do so.
24. A patient came to the general physician and complains that during chewing she is unable to touch the floor of the oral cavity.
25. When an examiner asks a child to pronounce "A", "O", "E", the child was unable to do so.