45. Muscle of mastication is supplied by
a. 1st part of maxillary artery
b. 2nd part of maxillary artery
c. 3rd part of maxillary artery
d. Facial artery

46. A patient came to you with dry eyes due to decreased lacrimal gland secretion. This may be due to following ganglion
a. Ciliary ganglion
b. Pterygopalatine ganglion
c. Otic ganglion
d. Submandibular ganglion

47. Infection from face to cavernous sinus can easily spread by:
a. Pterygoid venous plexus
b. External jugular vein
c. Internal jugular vein
d. Ophthalmic vein

48. Maxillary artery terminates as:
a. Infraorbital artery
b. Sphenopalatine artery
c. Palatine artery
d. Nasopalatine artery

49. In hanging , death occurs due to fracture of:
a. Base of skull
b. Odontoid process of axis
c. Body of atlas
d. Arch of atlas

50. Regarding inferior alveolar nerve:
a. It is purely sensory
b. Purely motor
c. Mixed nerve
d. Parasympathetic nerve

51. Lower gums are innervated by:
a. Greater palatine nerve
b. Lingual nerve
c. Anterior superior alveolar nerve
d. Posterior superior alveolar nerve

52. The palatoglossal arch is formed by which of the following muscle:
a. Tensor velli palatine
b. Levator velli palatine
c. Palatoglossus muscle
d. Palate pharygeus

53. Regarding left common carotid artery:
a. It arises from brachiocephalic trunk
b. It is embedded in carotid sheath
c. It gives two carotid and superior thyroid artery
d. It lies medial to internal jugular vein

Not sure

19. Spleen has:
a. Loose connective tissue capsule
b. Lymphoid follicles around arterioles
c. Hassell's corpuscles
d. Crypts lined by stratified squamous epithelium

20. Hyaline cartilage:
a. Present in epiglottis
b. Forms inter vertebral disc
c. Has basophilic matrix with meta chromatic reaction
d. Always lined by perichondrium

21. The pneumatic bone
a. Are the type of flat bone
b. Act as paranasal sinuses
c. Contains serous fluid inside
d. Present at the skull cap

22. Example of secondary cartilaginous joint is
a. 1st costochondral joint
b. Joint between epiphysis and diaphysis
c. Manibrusternal joint
d. Sternoclavicular joint

23. In dissection of human body, a section right angle to long axis of the organ is
a. Oblique section
b. Longitudinal section
c. Transverse section
d. Horizontal section

24. In typical spinal nerve:
a. Posterior ramus contain autononic fibers
b. Anterior fiber root contains sensory fiber
c. Lateral column contain parasympathetic nuceli
d. Trunk contain sensory ganglion

25. Elastic cartilage is found in:
a. Ear ossicles
b. Nasal septum
c. Epiglottis
d. Intervertebral disc

26. Regarding blood supply of bones
a. Diaphysial arteries anastomose with metaphyseal
b. Epiphysial arteries are the branches of metaphyseal
c. Metaphyseal arteries are the branches of a nutrient artery
d. Nutrient artery supply epiphysis

27. Grey rami communicants
a. Preganglionic fibers
b. Arises from sympathetic chain
c. Terminate inside the organs
d. Covered by myline sheath

28. Example of twisted fasciculi (PB 137 GA)
a. Deltoid Oblique → Multipennate
b. Latismus dorsi
c. Bicep brachi Parallel → fusiform
d. Sternocleidomastoid twisted but crossed so cruciate

29. In quadrupeds a median places
a. Superiorly
b. Anteriorly
c. Posteriorly
d. Inferiorly

30. Regarding joint s which one of the following is not true?
a. Primary cartilaginous joints are termed as synchondroses
b. Synovial joint occurs between manubrium and body of sternum
c. Hinge joint is example of uniaxial type of synovial joint
d. Inferior tibiofibular joint is the example of syndesmosis

31. The posterior belly of the diagastric muscle is innervated by a branch of this cranial nerve.
a. Trigeminal nerve
b. Facial nerve
c. Glossopharyngeal nerve
d. Accessory nerve

32. The cricoids cartilage lies at this vertebral level.
a. Second cervical vertebrae
b. Fourth cervical vertebrae
c. Sixth cervical vertebrae
d. Eight cervical vertebrae

33. Medical student studying the venous drainage, of brain, reveals that the sinus which is continous as internal jugular vein is:
a. Sigmoid sinus
b. Inferior petrosal sinus
c. Cavernous sinus
d. Superior petrosal sinus

34. The ganglion which lies is infratemporal fossa is:
a. Otic ganglion
b. Ciliary ganglion
c. Pterygopalatine ganglion
d. Submandibular ganglion

35. A patient came in emergency with bleed from nose –X-Ray shows the fracture of floor of the nose. The most probably involved bone will be:
a. Vomer
b. Cribriform plate of ethmoid
c. Palatine process of maxilla
d. Perpendicular plate of palatine

36. The opening which has connection with middle meatus is
a. Sphenoid sinus
b. Frontal sinus
c. Lacrimal duct
d. Posterior ethmoidal sinus

63. cavernous sinus contains
a. Sympathetic plexus
b. Mandibular nerve
c. Middle meningeal artery
d. Pituitary stalk

64. All of the following nuclei are present beneath the floor of fourth ventricle except:
a. Dorsal nucleus of vagus nerve
b. Vestibular nuclei
c. Hypoglossal nucleus
d. Facial nucleus

65. 4th ventricle communicates with all EXCEPT:
a. Arachnoid cisterns
b. Arachnoid villi
c. Subarachnoid space
d. Spinal canal

66. All sensations reach cerebral cortex via:
a. Thalamus
b. Internal capsule
c. Hypothalamus
d. Cerebellum

67. 20 years old girl present with sings of meningitis, during her investigation, lumber puncture block at $L_3 - L_4$ level alternate route for obtaining CSF will be
a. Between $L_2 - L_3$
b. Between $L_5 - S_1$
c. Of foramen magnum and C_1
d. Between $C_1 - C_2$

68. Forcep minor of corpus callosum are the fibers of
a. Rostrum
b. Genu
c. Body
d. Splenium

69. Great cerebral vein drain into:
a. Transverse sinus
b. Inferior sagittal sinus
c. Confluence of sinuses
d. Straight sinus

70. A 40 years old man present in causality of hospital with hemiplegic headache and difficulty in speaking Finding of CT scan shows a large hematoma at the area of pterion those characteristic finding are due to:
a. Lesion at pyramidal decussation
b. Epidural hemorrhage
c. Lower motor neuron lesion
d. Lesion at motor cortex

a. Ciliary ganglion
b. Pterygopalatine ganglion
c. Otic ganglion
d. Submandibular ganglion

47. Infection from face to cavernous sinus can easily spread by
a. Frontal sensory nerve
b. External nasal vein
c. Internal nasal vein
d. Ophthalmic vein

48. Maxillary artery terminates as
a. Infraorbital artery
b. Sphenopalatine artery
c. Palatine artery
d. Nasopalatine artery

49. In hanging , death occurs due to fracture of
a. Base of skull
b. Odontoid process of axis
c. Body of atlas
d. Arch of atlas

50. Regarding inferior alveolar nerve:
a. It is purely sensory
b. Purely motor
c. Mixed nerve
d. Parasympathetic nerve

51. Lower gums are innervated by:
a. Greater palatine nerve
b. Lingual nerve
c. Anterior superior alveolar nerve
d. Posterior superior alveolar nerve

52. The palatoglossal arch is formed by which of the following muscle
a. Tensor velli palatine
b. Levator velli palatine
c. Palatoglossus muscle
d. Palate pharygeus

53. Regarding left common carotid artery:
a. It arises from brachiocephalic trunk
b. It is embedded in carotid sheath
c. It gives two carotid and superior thyroid artery
d. Medial to internal jugular vein

Example of [illegible] muscle
a. Deltoid
b. Latissimus dorsi
c. Biceps brachii
d. Sternocleidomastoid

29. In quadrupeds a median plane is
a. Superiorly
b. Anteriorly
c. Posteriorly
d. Inferiorly

30. Regarding joint s which one of the following is not true?
a. Primary cartilaginous joints are termed as synchondroses
b. Synovial joint occurs between manubrium and body of sternum
c. Hinge joint is example of uniaxial type of synovial joint
d. Inferior tibiofibular joint is the example of syndesmosis

31. The posterior belly of the diagastric muscle is innervated by [illegible]
a. Trigeminal nerve
b. Facial nerve
c. Glossopharyngeal nerve
d. Accessory nerve

32. The cricoids cartilage lies at this vertebral level
a. Second cervical vertebrae
b. Fourth cervical vertebrae
c. Sixth cervical vertebrae
d. Eight cervical vertebrae

33. Medical student studying the venous drainage of brain, reveals that the sinus which is continous as internal jugular vein is:
a. Sigmoid sinus
b. Inferior petrosal sinus
c. Cavernous sinus
d. Superior petrosal sinus

34. The ganglion which lies in infratemporal fossa is

cavernous sinus contains
Sympathetic plexus
Mandibular nerve
Middle meningeal artery
Pituitary stalk

64. All of the following nuclei are present beneath the floor of fourth ventricle except:
a. Dorsal nucleus of vagus nerve
b. Vestibular nuclei
c. Hypoglossal nucleus
d. Facial nucleus

65. 4th ventricle communicates with all EXCEPT:
a. Arachnoid cisterns
b. Arachnoid villi
c. Subarachnoid space
d. Spinal canal

66. All sensations reach cerebral cortex via:
a. Thalamus
b. Internal capsule
c. Hypothalamus
d. Cerebellum

67. 20 years old girl present with sings of meningitis, during her investigation, lumber puncture block at $L_3 - L_4$ level alternate route for obtaining CSF will be
a. Between $L_2 - L_3$
b. Between $L_5 - S_1$
c. Of foramen magnum and C_1
d. Between $C_1 - C_2$

8. Forcep minor of corpus callosum are the fibers of
Rostrum
Genu

d. Has nucleoli

12. Stratified squamous epithelium
a. Develops only from ectoderm
b. Found in pelvis of kidney
c. Continuously replaced by basal layer
d. Has superficial nucleated layer

13. Collagen fibres has following characteristic except:
a. Responsible for tensile strength
b. Unbranched fibres
c. Appear banded on microscopy
d. Produced by mast cells

14. Skeletal muscle fibres have
a. Peripheral single nuclei
b. T system of tubules
c. Intercalated disc
d. Sarcomere between two I bands

15. Osteoblasts
a. Forms collagen
b. Remove bone during remodeling
c. Present in lacunae
d. Derived from osteocytes

16. Lymph node has:
a. Connective tissue framework of collagen
b. Valveless single afferent lymph vessel
c. T lymphocytes in cortex
d. B lymphocytes both in cortex and medulla

17. Tongue histology reveals that :
a. Filiform papillae has taste buds
b. Lymphoid tissue is confined to tip
c. Mucous Gland numerous in anterior part
d. Muscles fibres are predominantly skeletal

18. The salivary Glands:
a. Demilunes consist of mucous cells
b. Serous cells are eosinophilic
c. Striated ducts are lined by myoepithelial cells
d. Lack adi pose tissue

1. The part of the sperm containing proteolytic enzymes to digest the zona pellucida is the:
   a. Capacitor
   b. Head → contains nucleus with DNA
   c. Corona
   d. Acrosome → contains enzymes
   Middle piece → contains mitochondria → that provide energy

2. The ovulated mammalian oocyte is arrested at:
   a. Prophase of meiosis I
   b. Metaphase of meiosis I
   c. Prophase of meiosis II
   d. Metaphase of meiosis II

3. Somites:
   a. Differentiate into myotomes which give rise to skeletal muscle in trunk
   b. Differentiate into sclerotomes which give rise to vertebrae
   c. Arise from segmentation of the paraxial mesoderm
   d. All of the above are correct

4. When the amount of amniotic fluid exceeds two liters, the condition is called:
   a. Oligohydramnios decreased amniotic fluid (<300ml)
   b. Polyhydramnios increased amniotic fluid >2L
   c. Amniotitis infection in uterus
   d. Hydrogravida

5. After parturition a gynecologist identified the maternal surface of placenta due to 15-20 bulging areas called:
   a. Cotyledons.
   b. Placental septa.
   c. Intervillous space.
   d. Cytotrophoblastic shell.

6. A 21 year old female present to the obstetrician at 16 weeks of prenatal care. The obstetrician finds increased amniotic fluid and orders alpha fetoprotein and HCG levels. One week later the called patient back to tell her that the alpha fetoprotein level is increased. The most probable fetal diagnosis is

   a. Anencephaly
   b. Jejunal atresia
   c. Renal agenesis
   d. Sacrococcgeal teratoma

7. Fetal membranes are as under EXCEPT
   a. Amnion
   b. Allantois
   c. Germ disc
   d. Chorion

8. At the time of ovulation there comes a surge of
   a. Estrogen
   b. Progesterone
   c. Follicle stimulating hormone
   d. Luteinizing hormone

9. In case of Dizygotic twins there is found to be
   a. One chorionic cavity
   b. Two amniotic cavities
   c. One umbilical cord
   d. One connecting stalk

Dizygotic Twins → 2 amniotic cavities, 2 chorion, 2 placenta.
Monozygotic Twins → 2 amniotic cavities, 1 chorion, 1 placenta.

c. Epiphysis usually one in each other
d. Metaphysic is the growing part

65. Event occurring in the third week of development is:
a. Appearance of amniotic cavity
b. Lying down of extra embryonic mesoderm
c. The development of blastocoele
d. Formation of tertiary villi

66. Coronal plane
a. Divides the body into right and left halves
b. Parallel to sagital plane
c. Crosses the body from front and back
d. Right angle to vertical plane

67. Which one of the following is a normal content of umbilical cord at full term
a. Meckles diverticulum
b. Wharton's jelly
c. Two umbilical veins
d. Loops of intestine

68. A 21 year old female to the observation at 16 weeks prenatal core. The observation finds increased amniotic fluid and orders alpha fetoprotein HCG levels. One week later the called patient back to tell her that the alpha fetoprotein level is increased. The most probable fetal diagonsis is:
a.
b. G
c. J
d. J

69. Kinefelters syndrome is the defect of
a. Autosomes
b. Lusosomes
c. Sex chromosomes
d. Monosomes

d. Supported by temporomandibular ligment medially

58. The fascia forming the floor of posterior triangle of neck is:
    a. Investing layer of deep cervical fascia
    b. None of above
    c. Pretracheal fascia
    d. Prevertebral of deep cervical fascia
59. Examination of patient indicate that he has a medially directed strabismus (squint). This could be due to damage to the
    a. Adbucent nerve
    b. Oeculomotor nerve
    c. Ophthalmic nerve
    d. Trochlear nerve
60. Smooth muscle:
    a. Has a distinct sarcomere
    b. Passes diads
    c. Passes triads
    d. Caveolae store release calcium ions
61. Regarding the nuclei of cerebellum, the most medial nucleus is
    a. Dentate
    b. Emboli form
    c. Fastigial
    d. Globose
62. Muscles of mastication is supplied by:
    a. 1st part of maxillary artery
    b. 2nd part of maxillary artery
    c. 3rd part of maxillary artery
    d. Facial artery
63. The whit rami communicants:
    a. Are demylinated fyber
    b. Distributed to thoracic viscera
    c. Are post synaptic to sympathetic chain
    d. Courses in anterior root of spinal nerve
64. Regarding the part of growing bone
    a. Diaphysis grows away from the bone ends
    b. Epiphyseal cartilage represents the secondary ossification centre

32. An elastic artery has:
    a. Collagen fibers in tunica adventitia
    b. Inner endothelium in tunica intima
    c. Smooth muscle fibers in tunica media
    d. Internal elastic membrane in tunica intima
33. Cilia are found in the epithelium of:
    a. Gall bladder
    b. Stomach
    c. Uterine tube
    d. Alveoli of lung
34. A tooth development sequentially has last stage as
    a. Cap stage
    b. Enamel sac
    c. Dental root
    d. Eruption
35. A muscle with twisted arrangement of fibers is:
    a. Clavicular part of pectoralis major
    b. Latissimus dorsi
    c. Serratus anterior
    d. Trapezius
36. A great power of contraction of a muscle is due to its :
    a. Bipennate fiber
    b. Fusiform fibers
    c. Multipennate fibers
    d. Strap like fibers
37. A young singer after the local music theater visit her physician and complain of vocal difficulties. One examination, she is unable to abduct her vocal cord during quite breathing. Which of the following muscle most likely paralyzed.
    a. Cricoarytenoid
    b. Posterior Cricoarytenoid
    c. Thyroarytenoid
    d. Vocalis

44. The sensory decussation is formed by
    a. Decussation of corticospinal
    b. Decussation of internal arcuate fiber
    c. Decussation of superior cerebellar peduncle
    d. Decussation of rubrospinal tract
45. A vertical plane at right angle to median plane is
    a. Coronal plane
    b. Fontal plane
    c. Horizontal plane
    d. Sagittal plane
46. Regarding thymus:
    a. Epitherial reticular cells from "Hassal's Corpuscle"
    b. Lymphoid follicle are marked in the cortex
    c. Lobules are completely separated from each other
    d. Medulla contains "Cords of billroth"
47. Which of the following is the cervical branch of facial artery.
    a. Lateral nasal artery
    b. Superior labial artery
    c. Tonsilar artery
    d. Transverse facial artery
48. In a cervical vertebra, the:
    a. Body possess costal facets
    b. Identified by foramina transversaria
    c. Spinous process is blunt
    d. Vertical arch shows maxillary process
49. Injury on post triangle may damage which of the following structure:
    a. Cranial part of accessory nerve
    b. Spinal part of accessory nerve
    c. Branches of brachial plexus
    d. Branches of cervical plexus
50. Regarding the development of eye?
    a. Optic vesicle are out pounching from rhombencephalon
    b. Inner & outer layer of optic cup are firmly adherent to each other
    c. Coloboma develops due to complete fusion of choroid tissue
    d. The anterior one fifth of inner layer is regarded as pars ceca retina

8. Within carotid sheath
   a. Internal juglar vein is medial to atery
   b. Vagus nerve is between artery and vein
   c. Ansa cervicals is present in post wall
   d. Internal carotid artery in outside the sheath
9. Spinal nerve
   a. Anterior root contain automatic fibers
   b. Posterior ramus form plexus
   c. Ramities in vertebral canal
   d. Root exit through invertebral foramen
10. Which one is the following is a normal content of umbilical cord at full term.
   a. Meckles diverticuum
   b. Wharton's jelly
   c. Two umbilical veins
   d. Loops of intestine
11. The following structures are involved in formation of left lip.
   a. The fronto – nasal prominence
   b. Maxillary prominence
   c. Lateral nasal prominence
   d. Medical nasal prominence
12. Lymphatic from the lower lip and chin drain into which of the following lymph node?
   a. Post auricular
   b. Pre auricular
   c. Submandibular
   d. Submental
13. Sudden onset of stabbing pain on the left side of face in the infraorbital region , cheek and upper lip areas can occur due to increase stimulation of left:
   a. Facial nerve
   b. Mandibular nerve
   c. Maxillary nerve
   d. Opithalmic nerve

1. The morula
   a. Contains cells called blastomeres
   b. Has three distinctive cell types
   c. At birth in 5000 gm on an average
   d. Is similar to gastrula
2. Event occurring in the third week of development is
   a. Appearance of amniotic cavity
   b. Laying down of extra embryonic mesoderm
   c. The development of blastocoele
   d. Formation of tertiary with
3. Khnefelters syndrome is the defects of
   a. Autosomes
   b. Lysosomes
   c. Sex chromosomes
   d. Monosomes
4. Which is following is called the dangerous layer of scalp
   a. Galeaaponeurotica
   b. Loose areolar tissue
   c. Pencaranium
   d. Skin
5. Circumpennate fibers found in
   a. Deltoid
   b. Rectus abdommis
   c. Brachialis
   d.
6. Spinal nerve
   a. It is a part of A.N.S
   b. Contain post synaptic automatic fiber in their rimi
   c. Receive sensory innervations in its interior roots
   d. Supplying the visceral fibers to the thoraco abdominal organ
7. A patient came in emergency room , on examination his visual area is affected which artery is susceptible.
   a. Anterior cerebellar artery
   b. Middle cerebellar artery
   c. Post cerebellar artery
   d.

a	Parotid sheath	f	Auriculotemporal nerve
b	Retromandibular vein	g	Great auricular nerve
c	Facial nerve	h	Mastoid process
d	External carotid artery	i	Transverse cervical artery
e	Preauricular lymph nodes	j	Maxillary artery

Select the correct structure located in the parotid region from the above mentioned options which is most likely to be involved in the following conditions

16. In mumps, the parotid gland swells resulting in the excruciating pain during chewing.
17. During parotidectomy, the most superficial structure lying within the tissue of parotid gland.
18. During the dissection of the parotid region, the structure that indents the posteromedial surface of the parotid gland.

---

a	Medial pterygoid plate	f	Spine of sphenoid
b	Greater wing of sphenoid	g	Internal acoustic meatus
c	Lateral pterygoid plate	h	Pterygomaxillary fissure
d	Juglar foramen	i	Infraorbital fissure
e	Styloid process	j	Lesser wing of sphenoid

Select the most appropriate bony structure from the above mentioned list that is most likely to be related to the description given below.

19. Lying lateral to it is the auriculotemporal nerve and media to it is the chorda tympani nerve.
20. The bony opening has the fundus consisting of a bony plate called lamina cribrosa.

---

a	Occulomotor nerve	f	Cranial accessory nerve
b	Mandibular nerve	g	Hypoglossal nerve
c	Abducent nerve	h	Vagus nerve
d	Facial nerve	i	Opthalmic nerve
e	Spinal accessory nerve	j	Glossopharyngeal nerve

Select the correct cranial nerve from the above mentioned options that is most likely to be damaged resulting in the following clinical conditions

21. A Patient complains to the general physician that in the early morning when she tried to get up from the bed she is unable to move her head but can move her body.
22. A hypertensive patient in the emergency ward presents with double vision. On examination the pupils were dilated
23. During a routine physical examination, when the patient was asked to show his teeth, he was unable to do so.
24. A patient came to the general physician and complains that during chewing she is unable to touch the floor of the oral cavity.
25. When an examiner asks a child to pronounce "A", "O", "E", the child was unable to do so.

OPTION CAN BE SELECTED ONCE, MORE TH

a	Head fold
b	Nucleus pulposus
c	Hydatidform mole
d	epiglottis
e	Branchial cyst

Lead in: Choose the correct option and write it against e

1. The structure derived from the intermaxillary se
2. Primitive streak fails to degenerate in fetal life
3. A normal remnant of an embryonic structure foun
4. are found along the anterior border of the sternoc
5. The hypobranchial eminence contributes to the for

Pseudo stratified columnar epithelium		f	Thyr
uboidal epithelium		g	Palat
entin		h	Osteo
inal cord		i	Cysto
ebellum			

a. Stylomastoid foramen
b. Mandibular foramen
c. Mental foramen
d. Foramen ovale
e. Foramen spinosum
f. Optic canal
g. Internal acoustic meatus
h. Jugular foramen
i. Carotid canal
j. Hypoglossal canal

Lead in: Select the structure which exit through the skull from above mentioned foramen

1. Facial nerve
2. Maxillary nerve
3. Mandibular nerve
4. Glossophargngeal nerve
5. Hypoglossal nerve

14. Mental spine give attachment to:
    a. Genohyoid
    b. Mylohyoid
    c. Omohyoid
    d. Stylohyoid
15. Following thyroidectomy hoarseness of voice may occur, the condition is caused by injury to the :
    a. External laryngeal nerve
    b. Internal laryngeal nerve
    c. Recurrent laryngeal nerve
    d. Superior laryngeal nerve
16. Interior cerebellar peduncles carries the following tracts EXCEPT.
    a. Olivoeare bellar
    b. Derso spino cerebellar
    c. Ventro spinocerebellar
    d. Arculate fiber
17. Adenoid is the enlargement of
    a. Cerebellar tonsil
    b. Lingual tonsil
    c. Palatine tonsil
    d. Pharyngeal tonsil
18. Regarding the parts of growing bone
    a. Epiphyseal cartilage fuse with the disphysis
    b. Diaphysis is one on each side
    c. Metaphysis grows away from diaphysis
    d. Epiphysis represent the primary ossification centre
19. All the nuclei in brain stem are parasympathic EXCEPT.
    a. Superior salivatory
    b. Dorsal vagal
    c. Edinger westphal
    d. Tractus sulitarius

20. Which cells are responsible for the stability and integrity of bone:
    a. Osteobiasts
    b. Osteoclasts
    c. Osteocytes
    d. Osteoprogetor cells
21. Which of the following fontanelle is closed in the last?
    a. Anterior fontanelle
    b. Posterior fontanelle
    c. Lateral fontanelle
    d. Sphenoidal fontanelle
22. Structure passes through internal acoustic meatus IS:
    a. Internal jugular vein
    b. Facial nerve
    c. Infra petrosal
    d. Internal carotid artery
23. Red pulp of spleen contains:
    a. Venous sinusoids
    b. Central artery
    c. Germinal centre lymphoid follicle
    d. Hassel,s corpuscles
24. Epithelium adipose
    a. Is highly vascularized
    b. Is incapable of regeneration
    c. Rests on basement membrane
    d. Has abdundant intercellular space
25. The components of definitive chorion are:
    a. Extraembryonic somatic mesoderm, cytotrophoblast and syncytitotrophoblast
    b. Extraembryonic mesoderm and epiblast
    c. Extraembryonic visceral mesoderm
    d. Intraembyonic mesoderm syncytitotrophoblast and cytotrophoblast

26. A 43 year old man has onset of difficulty with speaking. Examination by ENT surgeon reveals problem in elevating the hyoid bone and floor of the mouth.
    a. Mylohyoid
    b. Omohyoid
    c. Sternothyroid
    d. Thyrohyoid
27. The example of the end arteries is:
    a. Celiac artery
    b. Central artery of the retina
    c. Coronary artery
    d. Posterior cerebral artery
28. Pseudotratified epithelium
    a. More than two layers of cells lying over basement membrane
    b. Cuboidal shaped cells are present
    c. Found in trachea
    d. Do not contain goblet cells
29. A principle difference between spongy bone and compact bone is the:
    a. Composition of amorphous and ground substance
    b. Amount of collagen in matrix
    c. Presence and absence of osseous lamellae
    d. Ratio of bone matrix to tissue matrix
30. The poster division of retomandibular vein join with posterior auricular vein to form :
    a. Angular vein
    b. Common facial vein
    c. External jugular vein
    d. Facial vein
31. Aponeuroses
    a. Encloses a muscle tendon
    b. Forms inter muscular scpta
    c. Is cord like in appearance
    d. Provides attachment for muscles fibers

51. Middle meiningeal artery is a branch of :
    a. 1st part of maxillary artery
    b. 2nd part of maxillary artery
    c. 3rd part of maxillary artery
    d. External carotid artery
52. A patient came in O.P.D with the complain of difficulty in elevating the lower jaw an examination physician reveals the loss of innervations of following muscles:
    a. Anterior belly of digastrics
    b. Lateral pterygoid
    c. Medial pterygoid
    d. Mylohyoid
53. Fetal membranes are as under EXCEPT:
    a. Amnion
    b. Allantois
    c. Germ disc
    d. Chorion
54. The fetal part of placenta is
    a. The wall of yolk sac
    b. Notochordal process
    c. Chorionic leave
    d. Chorionic frondosum
55. Opening of parotid duct lies :
    a. Lower moral tooth
    b. Lower pre moral tooth
    c. Upper moral tooth
    d. Upper pre moral tooth
56. A patient came to you with dry eyes due to decreased lacramial gland secretion This may be due to damage of the following ganglion:
    a. Cillary ganglion
    b. Otic ganglion
    c. Pterygopalatin ganglion
    d. Submandibular ganglion
57. Temporomandibular joint is
    a. Is deficient of its capsule anteriorly
    b. Present between temporal and mandibular bone
    c. Supplied by buccal branch of facial nerve

Q 11) [illegible]

Q 12) Primary ossification center appear in long bones are seen by:

Q 13) Organ which develops from intermediate mesoderm:
a) Liver
b) Spleen
c) Kidney
d) Gall bladder
e) Urinary bladder

Q 14) Site of fertilization is:
a) Ampulla of uterine tube
b) Isthmus of uterine tube
c) Uterus
d) Cervix
e) Ovary

Q 15) False statement regarding events of 2nd week of development is:
a) Formation of primary villi
b) Establishment of uteroplacental circulation
c) Formation of extraembryonic mesoderm
d) Bilaminar germ disc is formed
e) Implantation begins

Q 16) Layer of multinucleated zons without distinct cell boundaries:
a) Cytotrophoblast
b) Epiblast
c) Hypoblast
d) Syncytiotrophoblast
e) Amnioblast

Q 17) Finger-like cellular columns with syncytial covering in trophoblast is known as:
a) Free terminal villi
b) Secondary villi
c) Anchoring villi
d) Primary villi
e) Tertiary villi

Q 18) Dentin producing cells, odontoblasts develop from?
a) Endoderm
b) Paraxial mesoderm
c) Intermediate mesoderm
d) Neural crest cells
e) Ectoderm

Q 19) Cardiac muscles develop from:
a) Somatic layer of lateral plate mesoderm
b) Intra embryonic mesoderm
c) Endoderm
d) Neural crest cells
e) Splanchnic mesoderm

Q 20) Fate of notochord:
a) Spinal cord
b) Vertebral column
c) Vertebral arch only
d) Intervertebral disc
e) Nucleus pulposus

Q 21) Remnant of Primitive streak in sacrococcygeal region:
a) Sacrococcygeal Teratoma
b) Sirengiomelia
c) Situs inversus
d) Holoprosencephaly
e) Spina bifida

Q 22) Maternal portion of placenta is formed by:
a) Chorion frondosum
b) Decidua peritalis
c) Decidua Basalis
d) Chorion leave

Q 1) **Choose the false statement regarding sesamoid bone.**
a) Patella is largest one.
b) They are covered by periosteum.
c) Found under tendons.
d) They minimize the friction.
e) Maintain local circulation.

Q 2) **Length wise growth of long bones occurs by.**
a) Diaphysis.
b) Epiphysis.
c) Epiphysis plate of cartilage.
d) Metaphysis.
e) Number of nutrient foramen.

Q 3) **Choose the best example regarding pneumatic bone.**
a) Mandible.
b) Nasal.
c) lacrimal.
d) Inferior nasal concha.
e) Maxilla.

Q 4) **Regarding ossification of bones choose odd statement:**
a) Primary ossification center is usually one in number.
b) Primary ossification center appear before birth.
c) Secondary ossification center appear in diaphysis.
d) Secondary ossification centers are multiple.
e) Long bones ossify through endochondral ossification.

Q 5) **Select incorrect statement regarding architecture of skeletal muscles.**
a) Sartorius is strap like muscles.
b) Oblique fibers are more power full.
c) Biceps is fusiform in shape.
d) Temporalis is example of triangular muscle.
e) Parallel fibers have less range of movement.

Q 6) **choose correct statement regarding skeletal muscles.**
a) Fixed end of muscle is called insertion.
b) Origin is movable end of muscle.
c) Fleshy part of muscle is called tendon.
d) Flattened end of muscle is called aponeurosis.
e) Origin is always proximal to insertion.

Q 7) **Tooth and its socket is example of which joint.**
a) Suture.
b) Symphysis.
c) Syndesmosis.
d) Gomphosis.
e) Synchondrosis.

Q 8) **Choose the type of synovial joint in which articular surface is reciprocally co convex**
a) Hinge joint
b) Pivot joint.
c) Saddle joint
d) Ball and socket joint.
e) Plane joint

Q 9) **Choose incorrect statement regarding neural crest cell derivatives:**
a) Adrenal cortex
b) Melanocytes
c) Odontoblasts
d) Dorsal root ganglia.
e) Bones of face

Q 10) **Fetal portion of placenta is formed by:**
a) Chorión frondosum
b) Chorion leave
c) Decidua Basalis
d) Decidua caspularis
e) Decidua peritalis

Q 48) Which of the following statements describes correctly Actin?
a. Predominant Actin type in the epithelium of adult Uro Actin
b. Also known as cytoplasmic Actin
c. Characterized by the presence of myosin
d. Lines the diplon in cranial Actin
e. Forms the diplon in cranial Actin

Q 49) What term applies to collections of neuronal cell bodies (somata) in the central nervous system?
a. Ganglia
b. Neuroglia
c. Nodes
d. White matter
e. Nuclei

Q 50) Which of the following statements regarding the adblast is CORRECT?
a. It is composed of small chromaffin cells
b. It forms the roof of the sneak
c. It forms the primitive mouth
d. It forms the primitive streak
e. Notochordal cells incorporate into

★★★★★★★★★★ THE END ★★★★★★★★★★

[redacted]

[redacted]

[redacted]

Q 11) During 14th week which maternal antibodies are transported from mo...
a) IgA
b) IgE
c) IgM
d) IgD
e) IgG

Q 12) Primary ossification center appear in long bones and skull by:
a) 3rd week
b) 12th week
c) 16th week
d) 26th week.
e) 38th week

Q 13) Organ which develops from intermediate mesoderm:
a) Liver
b) Spleen.
c) Kidney
d) Gall bladder
e) Urinary bladder

Q 14) Site of fertilization is:
a) Ampulla of uterine tube
b) Isthmus of uterine tube
c) Uterus
d) Cervix
e) Ovary

Q 15) False statement regarding events of 2nd week of development is:
a) Formation of primary villi
b) Establishment of uteroplacental circulation
c) Formation of extraembryonic mesoderm
d) Bilaminar germ disc is formed
e) Implantation begins

Q 16) Layer of multinucleated zone without distinct cell boundaries:
a) Cytotrophoblast
b) Epiblast
c) Hypoblast
d) Syncytiotrophoblast
e) Amnioblast

Q 17) Finger-like cellular columns with syncytial covering in trophoblast is known as:
a) Free terminal villi
b) Secondary villi
c) Anchoring villi
d) Primary villi
e) Tertiary villi

Q 18) Dentin producing cells, odontoblasts develop from?
a) Endoderm
b) Paraxial mesoderm
c) Intermediate mesoderm
d) Neural crest cells
e) Ectoderm

Q 19) Cardiac muscles develop from:
a) Somatic layer of lateral plate mesoderm
b) Intra embryonic mesoderm
c) Endoderm
d) Neural crest cells
e) Splanchnic mesoderm

Q 20) Fate of notochord:
a) Spinal cord
b) Vertebral column
c) Vertebral arch only
d) Intervertebral disc
e) Nucleus pulposus

Q 21) Remnant of Primitive streak in sacrococcygeal region:
a) Sacrococcygeal Teratoma
b) Sirengiomelia
c) Situs inversus
d) Holoprosencephaly
e) Spina-bifida

Q 22) Maternal portion of placenta is formed by:
a) Chorion frondosum
b) Decidua peritalis
c) Decidua Basalis
d) Chorion leave
e) Decidua capsularis

Q 23) **Choose correct answer regarding oogenesis:**
a) Primary oocyte remain active until puberty
b) Graffian follicle is formed in preantral stage
c) Zona pellucida is formed by granulosa cells only
d) Theca externa is secretory layer
e) Oocyte maturation inhibitor secreted by follicular cells

Q 24) **Select the name of Ascending tract of lateral white matter responsible for carrying pain and temperature sensation:**
a) Anterior spinocerebellar tract
b) Posterior spinocerebellar tract
c) Lateral spinothalamic tract
d) Spino tectal tract
e) Spino olivary tract

Q 25) **Second order neuron for fascicular gracilis is:**
a) Substantia gelatinosa group
b) Nucleus gracilis
c) Dorsal root ganglia
d) Ventral posterolateral nucleus of thalamus
e) Nucleus dorsalis

Q 26) **Part of the visual reflex nucleus of Midbrain is:**
a) Lateral Geniculate body
b) Medial Geniculate body
c) VPL
d) Inferior colliculus
e) Superior colliculus

27) **Facial colliculus, a rounded swelling in floor of 4th ventricle is formed by:**
a) Abducent nucleus
b) Facial nucleus
c) Cochlear nucleus
d) Dorsal vagal nucleus
e) Superior Salivary nucleus

Q 28) **Select the false statement regarding deep intracerebellar nuclei:**
a) Dentate nucleus
b) Globus pallidus nucleus
c) Emboliform nucleus
d) Fastigeal nucleus
e) Globos nucleus

Q 29) **Choose false statement regarding the structure seen on transverse section of mid brain passing through inferior colliculus level:**
a) Red nucleus
b) Trochlear nucleus
c) Substantia nigra
d) Inferior collicular nucleus
e) Crus cerebri

30) **CSF Of lateral ventricle circulates into 3rd ventricle through which of the following openings:**
a) Cerebral aqueduct of sylvius
b) Foramen of magendie
c) Foramen of lushka
d) Foramen of Monro
e) Foramen Cecum

Q 31) **Choose the true statement regarding meninges of brain and spinal cord**
a) Endosteal layer of dura matter extends uptil second sacral vertebra
b) Pia matter is inner avascular layer
c) Arachanoid matter is outermost layer
d) Ligamentum divericulatum are triangular extensions of pia matter
e) Meningeal layer of dura matter stops at foramen magnum

Q 32) **Branches of basilar artery do not include:**
a. Anterior cerebral artery
b. Labyrinthine artery
c. Pontine arteries
d. Posterior cerebral artery
e. Superior cerebellar artery

Q 33) **Which of the following is a branch of vertebral artery?**
a. Anterior cerebral artery
b. B. Anterior inferior cerebellar artery
c. Middle cerebral artery
d. Ophthalmic arterys
e. Posterior inferior cerebellar artery

Time Allowed: Two Hours

Max. Marks: 100

Q1) A muscle which elevates and retracts the mandible is
a) Medial pterygoid
b) Masseter
c) Temporalis
d) Digastric
e) Lateral pterygoid

Q2) Unique features in the development of skull is
a) It encloses a cavity
b) It is contributed by paraxial mesoderm
c) Neural crest cells forms membranous portion
d) It is mesodermal in origin
e) It has two layer of bone

Q3) The best statement regarding the microscopic features of cerebellar cortex
a) The afferent fibers terminate in granular layer
b) It consists of six layers of cells
c) In medullary layer cells are compactly arranged
d) The Purkinje layer consists of flask shaped cells with highly developed dendritic tree
e) In granuler layer the cells are dispersed

Q4) The characteristic feature about the pericyte is
a) These cells are developed from ectoderm
b) These cells surround the arterioles
c) They are present in the basement membrane of vessels
d) They are spherical in shape
e) They have a contractile function

Q5) Which one of the following is best statement about scalp
a) Scalp has very poor nerve supply
b) Wounds of the scalp bleed profusely because the vessels do not retract
c) There are abundant emissary veins in frontal region
d) Epicranial muscles are supplied by mandibular nerve
e) Spread of infection is a common phenomena

Q6) Regarding Retina (point out the odd statement)
a) Nuclei of cones are dispersed near the limiting membrane
b) The rods and cones are polarized by neurons
c) Basal region of pigment epithelium rests firmly on Bruch's membrane
d) External plexiform layer is present between layer of rods and cones and bipolar cells
e) The anterior part of retina is photo sensitive

Q7) Mitosis of cells occurs in which layer of developing neural tube
a) Ependymal and mantle layers
b) Mantle layer
c) Ependymal (neuroepithelial) layer
d) Mantle and marginal layers
e) Marginal layer

Q8) Select an odd statement regarding the structures of the oral cavity.
a) Taste buds are located on the dorsal surface of the filiform papillae.
b) In tongue, muscle bundles cross each other running in longitudinal, transverse and vertical directions.
c) Superior surface of soft palate is lined by pseudostratified columnar ciliated epithelium.
d) Von Ebner glands of tongue are serous in nature.
e) There are no sebaceous or sweat glands in the red area of lips.

Q9) The most common site for spina bifida occulta is
a) Lower thoracic region
b) Mid thoracic region
c) Upper thoracic region
d) Lumbosacral region
e) Cervical region

Q10) Select an odd statement regarding the structure of cerebellum.
a) Cytoplasm of a granule cell contains large number of Nissl granules, which are arranged concentrically around the nucleus.
b) Unmyelinated axon of a basket cell ascends toward the molecular layer.
c) Mossy fibers originate from the inferior olivary nucleus of the medulla oblongata.
d) Axons of Golgi type II cells end by making synapses with terminal branches of the mossy fibers.
e) Four or more branching dendrites pass outward from the base of the perikaryon of the Purkinje cell.

Q11) Regarding the congenital anomalies of development of nose , which one of the following statement is incorrect
a) Nasal bridge is bifid
b) Single nostril results when only one nasal placode forms
c) Bifid nose means development of in separate noses
d) The nostrils are widely separated
e) Bifid nose results when medial nasal prominences do not merge completely

Time Allowed: One Hour

Q1) REGARDING AMNIOTIC FLUID, ALL OF THE FOLLOWING ARE TRUE EXCEPT THAT IT
a) contains the fetal cells and fetal urine.
b) allows mobility to fetus.
c) protects fetus from external injury.
d) helps in parturition.
e) amount is 500 ml at full term.

Q2) FOLLOWING STRUCTURE IS FORMED DURING THIRD WEEK OF DEVELOPMENT:
a) trilaminar germ disc.
b) neural tube.
c) genital ridge.
d) limb buds.
e) branchial arches.

Q3) 'Pterion' is located at
a) Pterygopalatine fossa
b) Occipital region
c) Zygomatic arch
d) Temporal region
e) Digastric fossa

Q4) THE BEST EXAMPLE OF LOOSE CONNECTIVE TISSUE IS THE:
a) dermis.
b) suprficial fascia.
c) tendon.
d) aponeurosis.
e) deep fascia.

Q5) IN CASE OF SKELETAL MUSCLES OF THE BODY:
a) sartorius is an example of bipennate muscles.
b) sternohyoid is an example of strap muscles.
c) triceps brachi is a multipennate muscle.
d) range of movement is greater than power in pennate muscles.
e) fibres run parallel to the line of pull in pennate muscles.

Q6) 'Visual area' on cortex is supplied by:
a) Vertebral atery.
b) Basilar artery.
c) Middle cerebral artery.
d) Posterior cerebral artery.
e) Anterior cerebral artery.

Q7) ANATOMICAL POSITION OF THE BODY INCLUDES ALL THE FOLLOWING STATEMENTS EXCEPT THE:
a) feet are held together.
b) person is standing erect.
c) palms are facing towards body (medially).
d) person is looking straight forward.
e) arms are hanging by sides.

Q8) ANTIBODIES ARE SYNTHESIZED AND SECRETED BY:
a) T cytotoxic cells.
b) T memory cells.
c) plasma cells.
d) T helper cells.
e) B memory cells.

Q9) Posterior root ganglia contains cell bodies of;
a) Pre-ganglionic neurons
b) Central motor neurons
c) Peripheral sensory neurons
d) Peripheral motor neurons
e) Central sensory neurons

Q10) The midbrain is a part of:
a) Cerebrum.
b) Diencephalon.
c) Pons.
d) Brain stem.
e) Cerebellum.

Q11) The following skin cells alert the immune system to pathogens
a) Langerhans
b) Melanocytes
c) Keratinocytes
d) Fibroblast
e) Homey cells

Q12) Skeletal muscles:
a) Do not form the main bulk of body
b) Are supplied by somatic nerves
c) Cells have one nuclei
d) Are capable of powerful contraction hence not fatigued easily
e) Form the main bulk of GIT muscles

Q26) A simplest structure considered to be alive are
a) Cells
b) Organs
c) Organelles
d) Tissues
e) Organisms

Q27) THE HYALINE CARTILAGE:
a) is devoid of perichondrium.
b) contains type-1 collagen.
c) is present on articular surfaces of synovial joints.
d) is the vascular tissue.
e) is capable of extensive turnover and repair.

Q28) THE MOST IMPORTANT EVENT FOR IMPLANTATION OF THE FETILIZED OVUM IS THE:
a) disintegration of zona pellucida.
b) entry of morula in uterine cavity.
c) formation of decidua.
d) formation of blastocyst
e) cleavage

Q29) THE MOST COMMON SITE FOR ABNORMAL IMPLANTATION IS THE:
a) uterine tube.
b) mesentery.
c) lower uterine segment.
d) pouch of Douglas
e) ovary.

Q30) Mid brain connects the forebrain to:
a) Pons.
b) Hind brain.
c) Spinal cord.
d) Cerebellum.
e) Brain stem.

Q31) Cerebral aqueduct passes through:
a) Medulla oblongatae
b) Cerebrum
c) Pons
d) All along brain stem
e) Mid brain.

Q32) PREGNANCY CAN BE DOCUMENTED IN A FEMALE IN SECOND WEEK OF DEVELOPMENT BY:
a) pelvic X-Ray (radiography).
b) assay of human chorionic gonadotropin in urine.
c) C.T scan.
d) ultrasound
e) assay of serum progesterone in the blood.

Q33) Post ganglionic sympathic fibers innervate the following viscera except
a) Adrenal medulla
b) Lungs
c) Heart
d) Duodenum
e) Stomach

Q34) Cell - bodies of preganglionic sympathetic fibers all present in lateral gray born at the level between:
a) T_1 to L_2
b) S_2 S_3 S_4
c) T_4 to L_5
d) C_1 to C_4
e) S_1 S_2 S_3

Q35) Regarding the type of muscle contraction:
a) Muscle do not change its length during eccentric contraction
b) Muscle decrease in length during eccentric contraction
c) Muscle exerts force with changing the length in isometric contraction
d) Muscle exerts force without changing length in isometric contraction
e) Muscle increase in length in isotonic contraction

Q36) SIMPLE SQUAMOUS EPITHELIUM IS ABSENT IN THE FOLLOWING ORGANS:
a) serous membranes.
b) lung alveoli.
c) blood vessels.
d) loops of Henle of the kidney.
e) distal convoluted tubules of the kidney.

Q37) Which of the following is not an essential part of anatomical position
a) Forearms supine
b) Feet flat on the floor
c) Mouth closed
d) Arms down to the sides
e) Eyes facing forward

Q38) SECONDARY OOCYTE COMPLETES ITS SECOND MEIOTIC DIVISION TO BECOME A MATURE OVUM:
a) at puberty.
b) before ovulation.
c) at ovulation.
d) at fertilization.
e) before fertilization.

Q13) The characteristic feature about the pericyte is
a) They have a contractile function
b) They are present in the basement membrane of vessels
c) These cells surround the arterioles
d) These cells are developed from ectoderm
e) They are spherical in shape

Q14) The anastomosis between external and internal carotid arteries include all of the following except
a) Dorsal nasal artery and facial artery
b) Greater palatine and superior labial artery
c) Anterior ethamoidal artery and facial artery
d) Lateral nasal artery and supratrochlear artery
e) Lacrimal artery and transverse facial artery

Q15) The clinical importance of anterior frontale is
a) It is the site of fusion of parietal bones with frontal bone
b) It bulges in case of raised intracranial pressure
c) Two parietal bone meet one another at this point
d) It is common site of meningeal herniation
e) It is on the scalp

Q16) Spinous process is absent in which of the following vertebra
a) 2nd cervical vertebra
b) 5th cervical vertebra
c) 1st cervical vertebra
d) 3rd cervical vertebra
e) 7th cervical vertebra

Q17) Lateral rectus of eye is supplied by
a) V cranial nerve
b) VI cranial verve
c) III cranial nerve
d) II cranial nerve
e) IV cranial nerve

Q18) Select an incorrect statement regarding the structure of layer of the cerebral cortex
a) Inner band of Baillarger is present in the external pyramidal layer
b) External granular layer contains large number of small pyramidal cells
c) Martinotti cells are present in the multiform and ganglionic layers
d) Horizontal cells are present in molecular layer
e) Outer band of Baillarger is present in the internal granular layer

Q19) Herniation of the intervertebral disc between the fifth and sixth cervical vertebrae will compress the
a) Fourth cervical nerve root
b) Sixth, seventh and eight cervical nerve roots
c) Seventh cervical nerve root
d) Fifth cervical nerve root
e) Sixth cervical nerve root

Q20) Select an odd statement regarding the structures of the oral cavity.
a) Superior surface of soft palate is lined by pseudostratified columnar ciliated epithelium.
b) Taste buds are located on the dorsal surface of the filiform papillae.
c) There are no sebaceous or sweat glands in the red area of lips.
d) Von Ebner glands of tongue are serous in nature.
e) In tongue, muscle bundles cross each other running in longitudinal, transverse and vertical directions.

Q21) Regarding the glial cells
a) Fibrous astrocytes are found in gray matter
b) Microglia are found in 'PNS'
c) Protoplasmic astrocytes form scar tissue in CNS, in case of any damage
d) Ependymal cells are of simple squamous variety
e) Schawn cells form myelin in central nervous system

Q22) Occulomotor nucleus is situated at which of the following level
a) Medulla oblongata
b) Tegumentum pones
c) Superior colliculus
d) Cerebello potine angle
e) Inferior colliculus

Q23) Which of the following tracts does not cross the midline
a) Rubro-spinal tract
b) Pyramidal Tract
c) Dento thalamic tract
d) Spinothalamic tract
e) Dorsal spinocerebellar tract

Q24) Considering the myelination of nerve fiber
a) Unmyelinated nerve fibers are not covered by Schwann cells
b) The nucleus of Schwann cell is present at the node of Ranvier
c) Myelin is formed by the cytoplasm of Schwann cells
d) Myline is stained black in routine histological stains i.e H & E
e) Internode is the distance between two nodes

Q25) Regarding mid brain
a) Origin to 4th cranial nerve in its upper part
b) Cerebral peduncle lies posterior
c) Closely related to free border of falx cerebelli
d) Superior colliculi receive auditory impulses
e) Contains cerebral aqueduct

Paper "Anatomy"

Time Allowed: Two Hours | Max. Marks: 100

Q1) Where are the vestibular nuclei located:
a) Beneath the lateral part of floor of fourth ventricle
b) In the cerebellum
c) In the olive
d) Beneath the medial part of floor of fourth ventricle
e) In the midbrain

Q2) Ostium of sphenoidal sinus opens into
a) Middle meatus
b) Superior meatus
c) Spheno - ethmoidal recess
d) External auditory meatus
e) Inferior meatus

Q3) The dorsal root ganglion contains cells bodies of
a) Dorsal column of spinal cord
b) Anterior spino vestibular tract
c) Anterolateral tract
d) Lateral spino thalamic tract
e) Anterior spino thalamic tract

Q4) The derivatives of alar plate of the neural tube are the
a) Basal ganglia of cerebrum
b) Pre vertebral sympathetic ganglia
c) Para vertebral sympathetic ganglia
d) Trigeminal ganglia
e) Dorsal root ganglia of spinal cord

Q5) The Thalamus forms the following boundary of 3RD ventricle:
a) Lateral
b) Medial
c) Superior
d) Anterior
e) Posterior

Q6) Regarding the tympanic cavity
a) Tensor tympani is the only muscle which is present in the tympanic cavity
b) It is the air containing part of ear
c) A cranial nerve is related to tympanic cavity
d) It contains small bones
e) Auditory tube extends from the anterior wall of tympanic cavity to naso pharynx

Q7) The roots of first cervical spinal nerve corresponds to the
a) level of formen magnum
b) atlas vertebra
c) upper edge of spinal cord
d) junction of medulla and spinal cord
e) lower edge of medulla oblongata

Q8) The following fibre lies immediately beneath the cortex and connect the adjacent gyri:
a) Long association fibres
b) Uncinate fasciculus
c) Inferior longitudinal fasciculus
d) Short association fibres
e) Fronto occipital fasciculus

Q9) During developments of tongue, the copula contributes to
a) Filiform papillae
b) posterior 1/3rd of the tongue
c) Incisive foramen
d) Frenulum of the tongue
e) Anterior 2/3rd of the tongue

Q10) Regarding Retina (point out the odd statement)
a) Nuclei of cones are dispersed near the limiting membrane
b) The anterior part of retina is photo sensitive
c) External plexiform layer is present between layer of rods and cones and bipolar cells
d) Basal region of pigment epithelium rests firmly on Bruch's membrane
e) The rods and cones are polarized by neurons

Q11) Concerning nerve fibers pick out the correct statement:
a) Unmyelinated nerve fibers posses nodes of ranvier.
b) The contral nervous system is rich in unmyiliated axons.
c) In adult nervous system single layer of mydin sheath is present.
d) Axons of small diaimeties are usually myelinated.
e) Group of nerve fibres are seldomly present in CNS.

Q12) Atresia of external auditory meatus is always associated with:
a) Absence of the middle ear.
b) Absence of the auricle.
c) None of these.
d) Absence of the inner ear.
e) 8th cranial nerve deafness.

Page 1 of 8

Q13) The forebrain is divided into cerebrum and:
a) Diencephalon
b) Mesencephalon
c) Myelencephalon
d) Rhombencephalon
e) Telencephalon

Q14) THE CARDIAC MUSCLE FIBRES:
a) have triads
b) show absence of gap junctions.
c) are cylindrical and branching.
d) have regenerative ability.
e) contain nuclei in the periphery.

Q15) A PERSON IS UNABLE TO SEE THE FOLLOWING STRUCTURE UNDER LIGHT MICROSCOPE:
a) mitochondria
b) cell membrane.
c) nucleus
d) Golgi complex
e) rough endoplasmic reticulum.

Q16) A SPECIMEN OF LYMPHOID ORGAN UNDER MICROSCOPE SHOWED THAT A PART OF IT IS COVERED BY STRATIFIED SQUAMOUS EPITHELIUM. THIS ORGAN IS MOST LIKELY TO BE:
a) palatine tonsil.
b) thymus
c) lymph node.
d) tubal tonsil.
e) spleen

Q17) Axon Hillock is the site in neurons with no;
a) Cytoplasm
b) Granules
c) Cellular pale
d) Dendrites
e) Axon.

Q18) tendons are composed of _____ connective tissue
a) Skeletal
b) Yellow elastic
c) Dense irregular
d) Dense regular
e) Areolar line

Q19) Blood vessels are lined with _____ epithelium
a) Transitional
b) Stratified cuboidal
c) Stratified aquamous
d) Simple columnar
e) Simple Squamous

Q20) IN CASE OF PLACENTA, FOLLOWING OPTION IS INCORRECT:
a) maternal surface is smooth and covered by amnion
b) fetal part is formed by chorionic frondosum.
c) contains cotyledons on it maternal surface.
d) umbilical cord is attached near its center.
e) it weighs 500 grams.

Q21) Conus Medullaris is the inferior apex of the:
a) Spinal cord
b) Cerebellum
c) Sympathetic chain
d) Cerebrum
e) Brain stain

Q22) THE MAJOR CHARACTERISTIC OF MEIOSIS-1 IS THE:
a) producing the primordial germ cells.
b) pairing of the homologous chromosomes.
c) reducing the amount of DNA to 1N.
d) acheiving the diploid number of chromosomes.
e) splitting of the centromere.

Q23) The following skin cells have a sensory function ?
a) Dendritic cells
b) Merkel's cells
c) Keratinocytes
d) Melanocytes
e) Prickle cells

Q24) Lateral wall of orbit is formed by frontal and
a) Mandible
b) Occipital bone
c) Nasal bone
d) Maxilla
e) Zygomatic bone

Q25) The following type of joints is most moveable
a) A synarthrosis
b) A condyloid joint
c) A symphysis
d) A diarthrosis
e) A synovial joint

Q39) The structure of tongue reveal all the following features except
a) The mucous membrane is smooth on its lower surface
b) Filli form papillae epithelium contain taste buds
c) Is a mass of striated muscles
d) Groups of glands are gathered in between muscle fibers
e) Circum uafrate papillae are present near sulcus terminalis

Q40) The corticospinal fibres are absent in the following regions of white matter
a) Lateral white column of the spinal cord
b) Pyramids of medulla oblongata
c) Cerebral peduncle of the midbrain
d) Anterior limb of the internal capsule
e) Corona radiata

Q41) Regarding the skin of face
a) There is no fascia on the face
b) The skin of face has a rich blood supply
c) Boil on the face can be dangerous and lead to cavernous sinus thrombosis
d) Sensory supply of face is from cranial nerves
e) It possesses many glands

Q42) It is untrue to say about middle ear that
a) Oval and round windows are related to its medial bony wall
b) Posteriorly it communicates with mastoid air cells
c) Tympanic cavity is lined by simple squamous cells
d) The auditory tube connect it to pharynx
e) Is an irregular space in interior of temporal bone

Q43) The following paranasal air sinus develops during late fetal life
a) Frontal sinus
b) Posterior ethmoid sinus
c) Maxillary sinus
d) Sphenoid sinus
e) Anterior ethmoid sinus

Q44) The best statement about parotid gland is
a) It is pierced by facial artery
b) Glossopharyngeal nerve supply secretomotor fibers to it
c) Gland is covered only by superficial fascia
d) The duct of the gland opens in the oral cavity at the level of 3rd upper molar tooth
e) It is a mixed type of gland

Q45) In comparison to the amount of blood supplied by the capillaries to the gray matter and white matter:
a) Gray matter receives 70% amount of blood supply to the brain
b) White matter receives more supply than gray matter
c) Both white and gray matter receive equal amount of blood
d) Gray matter receives 40% amount of blood supply to the brain
e) Gray matter receives more supply than white matter

Q46) A player in the hockey ground was struck by a ball on head. After remaining unconscious for a while he regained consciousness, and become unconscious again .A diagnosis of extradural hemorrhage was made. Which of the following blood vessels is most commonly involved:
a) Middle meningeal artery
b) Superior longitudinal sinus
c) Transverse sinus
d) Anterior meningeal artery
e) Internal maxillary artery

Q47) Regarding the infra temporal fossa
a) Its medial boundary is the pterygoid plate
b) It is related to carotid sheath
c) It is a space in the skull
d) Maxillary artery and the temporalis muscles are the only contents
e) It communicates with the pterygo palatine fossa through pterygo maxillary fissure

Q48) Which one of the following is best statement about scalp
a) Spread of infection is a common phenomena
b) Scalp has very poor nerve supply
c) There are abundant emissary veins in frontal region
d) Epicranial muscles are supplied by mandibular nerve
e) Wounds of the scalp bleed profusely because the vessels do not retract

Q49) Ascending pathways leading to consciousness consist of
a) Three neurons
b) Four neurons
c) Five neurons
d) One neuron
e) Two neurons

Q50) The chorda tympani nerve courses through tympanic cavity, lying on its:
a) Roof
b) Anterior wall
c) Floor
d) Medial wall
e) Lateral wall

Q51) Clark's column extends between the following
a) Only the cervical segment
b) Only the thoracic segment
c) 1st thoracic segment to 9th thoracic segment
d) 10th thoracic segment to 1st sacral segment
e) 8th cervical segment to 3rd lumbar segment

Q52) Select a wrong statement regarding the filiform papillae of the tongue
a) Are finger like in shape
b) Are distributed over the entire anterior two third dorsal surface of the tongue
c) Taste buds are located on their dorsal surface
d) Are the most numerous of the lingual papillae
e) Covered by stratified squamous cornified epithelium

Q53) The importance of alar plates in the developing spinal cord is that
a) it is separate from basal plate by sulcus limitans
b) it is present bilaterally
c) it produces a longitudinal bulge
d) its cell bodies form the dorsal gray column of spinal cord
e) its enlargement on both the sides form dorsal median septum

Q54) Which of the following muscle plays major role in opening the Jaw
a) Temporalis muscle
b) Masseter
c) Buccinator
d) Medial pterygoid
e) Lateral pterygoid

Q55) The Passavant's ridge is formed when the posterior wall of pharynx is drawn forward during swallowing, due to the contraction of
a) Palatoglossus
b) Salpingopharyngeus
c) Levator veli palatini
d) Palatopharyngeus
e) Tensor veli palatini

Q56) Regarding the muscle of face:
a) Levator palpebrae superioris is supplied by facial nerve.
b) Muscles of facial expression are supplied by the maxillary nerve.
c) Buccinator is an accessory muscle of mastication so supplied by the mandibular division of trigeminal nerve.
d) Embryologically the muscles of facial expression are developed from the mesoderm of the second pharyngeal arch.
e) Occipitofrontalis helps in the movement of skin of face.

Q57) The most common site for spina bifida occulta is
a) Upper thoracic region
b) Lumbosacral region
c) Cervical region
d) Mid thoracic region
e) Lower thoracic region

Q58) It is true to say about the structure of nasal cavity that
a) The inner surface of nares are devoid of sebaccous and sweat glands
b) Superior concha are covered by respiratory epithelium
c) The chemo receptor of smell are located with in inferior corcha
d) The epithelium with in the vestibule remain stratified squamous epithelium
e) Concha has large swell bodies in its mucous membrane

Q59) The important feature of Arnold chiari malformation through advance imaging is
a) Herniation of tonsils of cerebellum in to vertebral canal
b) Spina bifida
c) Blockage of cerebrospinal fluid
d) Internal hydrocephalus
e) Cranio vertebral anomaly

Q60) The vallate papillae in the oral part of tongue are innervated by nerve of third pharyngeal arch because
a) The mucosa of posterior third of tongue is pulled slightly anterior as tongue develops
b) Papillae appear towards the end of 8th week
c) Vallate papillae are the first to appear
d) The merged distal tongue buds form the anterior 2/3rd of tongue
e) The median tongue bud forms no recognizable part of adult tongue

Q61) Which of the following statement is correct about lingual artery
a) It is crossed superficially by hypoglossal nerve
b) It runs through substance of submandibular salivary gland
c) It is a branch of subclavian artery
d) It arises at the level of cricoid cartilage
e) It runs superficial to hyoglossus muscle

Q62) The striking feature of oligodendrocytes is that
a) They may surround nerve cell bodies
b) There are no fiaments in their Cytoplasm
c) They are frequently found in rows
d) They are responsible for formation of myelin sheath in the central nervous system
e) They have small cell bodies

Q63) The cervical spinal nerves lack
a) Connective tissue sheaths
b) Preganglionic sympathetic fibres
c) Motor fibres to smooth muscles
d) Post ganglionic sympathetic fibres
e) Motor fibres to skeletal muscles

Q64) Internal structure of spinal cord shows
a) Anterior Grey horn which contains mainly sensory neurons
b) Lateral Grey horn in thoracic region which contain sympathetic neurons
c) Lateral spinothalamic tracts carrying fibers for the sensation of temperature
d) Spinocerebellar tracts are located in posterior white column
e) Fasciculus cuneatus lies in anterior white column

Q26) Which of the statement concerning thalamus is incorrect

a) Joins to other side by thalamic adhesion
b) Forms anterior boundary of interventricular foramen
c) Seperated from lenitform nucleus by internal capsule
d) Serve as main relay station for all sensory impulse except olfactory
e) Largest part of Diencephlon

Q27) Neurons in the human body can be: (Choose the correct statement)

a) Golgi type one, if its axon is long some time up to or even more than a meter
b) Multipolar if it has many axons and one dendrite
c) Unipolar if it posses only one dendrite
d) Bipolar if it posses two dendrites
e) Golgi type two if it has short dendrons Less than few centimeters

Q28) Unique features in the development of skull is

a) It is mesodermal in origin
b) It is contributed by paraxial mesoderm
c) It encloses a cavity
d) Neural crest cells forms membranous portion
e) It has two layer of bone

Q29) The leptomeninges are derived from the following structure

a) Notochord
b) Neural crest cells
c) Denticulate ligament
d) Endoderm
e) Mesenchyme surrounding spinal cord

Q30) Regarding the nerve supply of extra ocular muscle:

a) Levotor palpeberae superioris is supplied by inferior division of the oculomotor nerve.
b) Superior oblique is supplied by the contra-lateral trochlear nerve.
c) Lateral rectus is supplied by trochlear nerve.
d) Medial rectus is supplied by the ophthalmic nerve.
e) Inferior oblique is supplied by the abducent nerve.

Q31) Following statements regarding the motor speech area of Broca are true except

a) Is connected to the sensory speech area of wernicke
b) Its damage will cause receptive aphasia
c) In most of the individuals it is present in left cerebral hemisphere
d) Is located in the inferior frontal gyrus
e) Is concerned with formation of words by its connections with the primary motor area

Q32) The vertebral artery arises from which of the following

a) 1st part of subclavian artery
b) 2nd part of subclavian artery
c) Thyrocervical trunk
d) 3rd part of subclavian artery
e) Costocervical trunk

Q33) Which statement is incorrect about cutaneous innervation of face

a) Parotid gland ---------mandibular nerve
b) Upper lip ---------- Maxillary nerve
c) Anterior part of temple --------mandibular nerve
d) Tip of nose ---------- ophthalmic nerve
e) Forhead ----------ophthalmic nerve

Q34) In the fourth ventricle

a) Nucleus of accessory nerve lies in its floor
b) Choroid plexus projects from the lateral walls
c) Roof contains one median and two lateral apertures (openings)
d) Lateral boundries are formed by inferior cerbellar peduncles only
e) Floor is formed by posterior surface of midbrain

Q35) Conus medullaris is the

a) Tapers off spinal cord
b) Cervical enlargement of spinal cord
c) Thickening of diameter in spinal cord
d) Lumbar enlargement of spinal cord
e) Enlargement of subarachnoid space

Q36) Mitosis of cells occurs in which layer of developing neural tube

a) Mantle layer
b) Marginal layer
c) Ependymal (neuroepithelial) layer
d) Mantle and marginal layers
e) Ependymal and mantle layers

Q37) Regarding the nasal cavity

a) Para nasal sinuses can not be visualized on the x-rays
b) Olfactory nerves are present in nasal cavity
c) The linning of nasal cavity and vestibule consists of keratinized stratified squamous epithelium .
d) It receives it's blood supply only from facial artery
e) Para nasal sinuses can normally be opaque on the x-rays

Q38) Regarding the congenital anomalies of development of nose , which one of the following statement is incorrect

a) Bifid nose means development of in separate noses
b) Single nostril results when only one nasal placode forms
c) Bifid nose results when medial nasal prominences do not merge completely
d) Nasal bridge is bifid
e) The nostrils are widely separated

Paper "Anatomy"

Time Allowed: Two Hours

Max. Marks: 100

Q1) Where are the vestibular nuclei located:
a) Beneath the lateral part of floor of fourth ventricle
b) In the cerebellum
c) In the olive
d) Beneath the medial part of floor of fourth ventricle
e) In the midbrain

Q2) Ostium of sphenoidal sinus opens into
a) Middle meatus
b) Superior meatus
c) Sphenio - ethmoidal recess
d) External auditory meatus
e) Inferior meatus

Q3) The dorsal root ganglion contains cells bodies of
a) Dorsal column of spinal cord
b) Anterior spino vestibular tract
c) Anterolateral tract
d) Lateral spino thalamic tract
e) Anterior spino thalamic tract

Q4) The derivatives of alar plate of the neural tube are the
a) Basal ganglia of cerebrum
b) Pre vertebral sympathetic ganglia
c) Para vertebral sympathetic ganglia
d) Trigeminal ganglia
e) Dorsal root ganglia of spinal cord

Q5) The Thalamus forms the following boundary of 3rd ventricle:
a) Lateral
b) Medial
c) Superior
d) Anterior
e) Posterior

Q6) Regarding the tympanic cavity
a) Tensor tympani is the only muscle which is present in the tympanic cavity
b) It is the air containing part of ear
c) A cranial nerve is related to tympanic cavity
d) It contains small bones
e) Auditory tube extends from the anterior wall of tympanic cavity to naso pharynx

Q7) The roots of first cervical spinal nerve corresponds to the
a) level of formen magnum
b) atlas vertebra
c) upper edge of spinal cord
d) junction of medulla and spinal cord
e) lower edge of medulla oblongata

Q8) The following fibre lies immediately beneath the cortex and connect the adjacent gyri:
a) Long association fibres
b) Uncinate fasciculus
c) Inferior longitudinal fasciculus
d) Short association fibres
e) Fronto occipital fasciculus

Q9) During developments of tongue, the copula contributes to
a) Filiform papillae
b) posterior 1/3rd of the tongue
c) Incisive foramen
d) Frenulum of the tongue
e) Anterior 2/3rd of the tongue

Q10) Regarding Retina (point out the odd statement)
a) Nuclei of cones are dispersed near the limiting membrane
b) The anterior part of retina is photo sensitive
c) External plexiform layer is present between layer of rods and cones and bipolar cells
d) Basal region of pigment epithelium rests firmly on Bruch's membrane
e) The rods and cones are polarized by neurons

Q11) Concerning nerve fibers pick out the correct statement:
a) Unmyelinated nerve fibers posses nodes of ranvier.
b) The contral nervous system is rich in unmyiliated axons.
c) In adult nervous system single layer of mydin sheath is present.
d) Axons of small diaimeties are usually myelinated.
e) Group of nerve fibres are seldomly present in CNS.

Q12) Atresia of external auditory meatus is always associated with:
a) Absence of the middle ear.
b) Absence of the auricle.
c) None of these.
d) Absence of the inner ear.
e) 8th cranial nerve deafness.

Q78) The development of papilloedema in the case of increased intracranial pressure, is due to the following feature of optic nerve:
a) It is an extension of white matter of brain
b) It meets the sclera 3mm medial to posterior pole
c) It is surrounded by subarachnoid space as far as back of eye
d) It enters the orbit through optic canal
e) It receives the central artery of retina

Q79) Longest spinous process is the characteristics of
a) 1st thoracic vertebra
b) 7th cervical vertebra
c) 1st cervical vertebra
d) 12th thoracic vertebra
e) 2nd cervical vertebra

Q80) The dorsal roots of spinal nerves are derived from the following structure
a) The posterior gray column
b) Peripheral process of posterior root ganglia
c) The autonomic ganglia
d) White ramus communicans
e) Central process of posterior root ganglia

Q81) With swelling of mucous membrane of superior nasal meatus, which of the opening is plugged?
a) Sphenoidal sinus
b) Frontal sinus
c) Anterior ethmoidal sinus
d) Superior Para nasal sinus
e) Middle ethmoidal sinus

Q82) The accurate statement about lacrimal gland is:
a) It curls around the medial margin of levator palpebrae superioris
b) Its orbital part is visible through superior fornix of conjunctiva
c) It is drained by ducts draining from palpebral part into superior fornix
d) Its palpebral part lies in lacrimal fossa
e) Secremotor fibers to the gland from inferior salivary nucleus travel in greater petrosal nerve

Q83) The tract which is concerned with reflex postural movements in response to visual stimuli is
a) Rubro spinal tract
b) Tecto spinal tract
c) Vestibulo spinal tract
d) Olivospinal tract
e) Corticospinal tract

Q84) A muscle which elevates and retracts the mandible is
a) Digastric
b) Medial pterygoid
c) Temporalis
d) Lateral pterygoid
e) Masseter

Q85) The best statement regarding the microscopic features of cerebellar cortex
a) The Purkinje layer consists of flask shaped cells with highly developed dendritic tree
b) It consists of six layers of cells
c) In granuler layer the cells are dispersed
d) In medullary layer cells are compactly arranged
e) The afferent fibers terminate in granular layer

Q86) All the features develops during fourth week except
a) Neural tube closes.
b) Bucco-pharyngeal membrane break down.
c) Most somites are formed.
d) Primary centre of ossification appears in the skull.
e) Optic vesicle first appears.

Q87) Select an odd statement regarding the structure of eye:
a) Ciliary epithelium consist of a single layer of columnar unpigmented cells.
b) A single layer of cuboidal cells is present all over just under the lens capsule.
c) Descemet's membrane of cornea is highly refractile under the light microscope.
d) Inner layers of retina are absent in the fovea centralis.
e) Photo receptors of the retina are modified bipolar neurons.

Q88) Which of the following are derivatives from neural crest cells except?
a) Schwann cells
b) Ganglion of sympathetic trunk
c) Melanocytes of skin
d) Cortex of adrenal gland
e) Dorsal root ganglion

Q89) The correct statement regarding the microscopic features of cerebral cortex
a) Size of the cells decreases from superficial to deep layer
b) Large pyramidal cells (Betz cells) are found in ganglionic layer
c) The axons of Martinotti cells pass in deeper layers of the cortex
d) Multiform layer is the most superficial layer
e) Molecular layer consists of stellate cells

Q90) Foot plate of stapes is placed at
a) Opening of pyramid
b) Tubal opening
c) Round window (fenestra cochlear)
d) Oval window (fenestra vestibuli)
e) Aditus

Q65) Select odd statement regarding lumbar puncture (CSF)
a) Positive Quenkenstedt's sign when pressure rises
b) In Prone position
c) For histopathological examination
d) Vertebral column is well extended
e) Normal pressure is about 60 to 150 mm of water

Q66) Select a wrong statement regarding the structure of external ear
a) Ceruminous glands are found in the submucosa
b) External auditory meatus is lined by stratified squamous epithelium
c) Ceruminous glands are a type of modified sebaceous gland
d) External surface of eardrum is lined with epidermis
e) Pinna is composed of a plate of elastic cartilage

Q67) Select an odd statement regarding the structure of nose:
a) Superior conchae are lined by olfactory mucosa.
b) Smell bodies are more prominent over the inferior conchae.
c) Poor population of goblet cells is present in the mucosa.
d) Vibrissae are present in the skin vestibule.
e) Many simple branched tubulo-alveolar glands are present in the lamina propria of mucosa.

Q68) The important characteristic feature of mossy fibres in the cerebellum is
a) They are terminal fibers of all cerebral afferent tracts
b) They have multiple branches
c) They exert a much more diffuse excitatory effect
d) The deep cerebellar nuclei receive afferent information from them
e) A single mossy fibre may stimulate thousands of Purkinje cells

Q69) All the structures contain fibers of corticospinal tracts except
a) Posterior limb of internal capsule
b) Tegmentum of medulla
c) Pyramids of medulla
d) Crus cerebri of mid brain
e) Lateral white column of spinal cord

Q70) Which of the following structure does not take part in formation of cleft lip
a) Left maxillary Proces
b) Medial nasal proiminences
c) Lateral nasal proiminences
d) Right maxillary process
e) The upper jaw

Q71) Select an odd statement regarding the structure of cerebellum.
a) Unmyelinated axon of a basket cell ascends toward the molecular layer.
b) Mossy fibers originate from the inferior olivary nucleus of the medulla oblongata.
c) Four or more branching dendrites pass outward from the base of the perikaryon of the Purkinje cell.
d) Axons of Golgi type II cells end by making synapses with terminal branches of the mossy fibers.
e) Cytoplasm of a granule cell contains large number of Nissl granules, which are arranged concentrically around the nucleus.

Q72) Pulsation of facial artery can be easily felt at
a) Body of masseter
b) Posterior border of buccinator
c) Anterior border of temporalis
d) Posterior border of masseter
e) Anterior border of masseter

Q73) The pterion is the point
a) Is 1.5 inches above the mid point of zygomatic arch
b) It is marked by an eminence
c) It is ,marked by a depression
d) Where lesser wing of sphenoid bone meets with parietal bone
e) Posterior branch of middle meningeal artery is related to it

Q74) Which of following level of spinal cord involve in parasympathetic out flow
a) S_1 and S_2
b) L_1 and S_1
c) S_3 , S_4 And S_5
d) S_1 , S_2 And S_3
e) S_2, S_3 And S_4

Q75) Which type of cells is not present in the structure of cerebellum?
a) Pyramidal cells
b) Golgi type II cells
c) Basket cells
d) Granule cells
e) Stellate cells

Q76) Select the odd statement regarding structure of larynx
a) Vocal segments are elastic in nature
b) The number of cartilages lie at out side the mucus membrane
c) The upper fold of mucosa constitute false vocal cords
d) The lingual surface of epithelium of epiglottis is stratified squamous epithelium
e) Vocalis muscles are skeletal in nature

Q77) Which of the following nerve gives parasympathetic fibres to parotid gland
a) Facial
b) Glossopharyngeal
c) Trigeminal
d) Vagus
e) Accessory nerve

Q91) Select the correct statement regarding neuropores.
a) Rostral neuropore closed after the closure of caudal neuropore
b) Closed prior to three months of developments
c) Neuropores must be closed prior to one month of development
d) Closed prior to two months of developments
e) Remains opened

Q92) Occulusion of middle cerebral artery may produce contra lateral homonymous hemianopia because of damage to
a) Visual cortex (area 17)
b) Optic chiasma
c) Lateral geniculate body
d) Optic radiation
e) Isual association cortex (areas 18 & 19)

Q93) Which of the following statement is incorrect about lateral Medullary syndrome
a) Dysphonia and dysphargia
b) Ipsilateral loss of pain sensation of the face
c) Ipsilateral loss of touch
d) Ipsilateral Paralysis of vocal cord
e) Ipsilateral Horners syndrome

Q94) The following cell lines the ventricles of brain and the central canal of spinal cord:
a) Tanycytes.
b) Astrocytes.
c) Choroidal epithelial cell.
d) Oligodendrocytes.
e) Ependymocytes.

Q95) The neural layer of retina is derived from the following
a) Mesenchyme surroundings the optic cup
b) Optic stalk
c) Outer thinner layer of optic cup
d) Inner thickened layer of optic cup
e) Lens placode

Q96) Sulcus limitans is the groove which separates the
a) Alar and roof plate
b) Basal and roof plate
c) Basal and floor plate
d) Alar and floor plate
e) Basal and alar laminae

Q97) In the development of eye, the structures which develop from optic cup include:
a) Retina, Iris and ciary body.
b) All layers of retina and ciliary muscles.
c) Cornea, choroid and retina.
d) Lense, cornea and retina.
e) Ciliary muscles, Iris and retina.

Q98) Select an odd statement regarding spinal nerve
a) Anterior root contain only motor axon
b) They are formed by union of anterior and posterior nerve root
c) Posterior root ganglion contain unipolar neurons
d) There are 31 pairs
e) Posterior ramus contain one sensory axon

Q99) Malformations of the auricle is most commonly associated with:
a) Congenital deafness.
b) Maternal X-irridiation.
c) Defects of the lower jaw.
d) Thalidomide administration.
e) Anomalies of the sex chromosomes.

Q100) Regarding blood supply of brain, the best statement is
a) Medial and lateral central striate arteries supply the internal capsule
b) Cortical branches are the common site of thrombosis
c) Anastomosis frequently occur in central arteries
d) Thalamus is supplied by anterior cerebral arteries
e) Circle of willis is formed by two internal carotid and two spinal arteries

★★★★★★★★★★ THE END ★★★★★★★★★★

3. moves inferiorly
4. compresses thoracic viscera
5. does not change position

368. The olfactory region of the nasal cavity is located on the

1. cribriform plate
2. nasal septum and lateral wall above the superior nasal concha
3. posterior nasal walls
4. nasal floor and lateral wall above the inferior nasal concha
5. lateral nasal walls

369. The nasolacrimal duct opens into the

1. inferior meatus
2. middle meatus
3. superior meatus
4. sphenoethmoidal recess

370. Blood is supplied to the nasal cavity by branches of the

1. ophthalmic artery
2. maxillary artery
3. facial artery
4. all of the above

371. Which of the following drain into the superior meatus of nose?

1. Frontal sinus
2. Sphenoidal sinus
3. Nasolacrimal duct

5. Carpal bones of wrist are which type of bone:

a) short bones
b) short long bones
c) long bones
d) long short bones
e) Flat bones

6. Intervertebral discs of vertebrae are composed of:

a) hyaline cartilage
b) fibrocartilage
c) elastic cartilage
d) Fibrohyaline cartilage
e) no cartilage

7. Interphalangeal joints are which kind of joints:

a) condylar
b) plane
c) hinge
d) ball and socket
e) pivot

8. During a dissection class of anatomy, students were observing the muscles of limb, on further inspection of the skeletal muscle, they observed that a mem was covering a bundle of muscle fibres, this membrane is called:

a) Endomysium
b) Perimysium
c) Epimysium
d) Superficial fascia
e) Skin

9. What is most appropriate about a motor unit:

a) A site where motor nerve enters a muscle
b) A single motor neuron only
c) A single motor neuron together with the muscle fibres supplied by i
d) Two motor neurons supplying one muscle
e) A group of motor neurons

15. A young patient developed conjunctivitis on exposure to perfume. Which epithelium was infected?

a) Simple columnar epithelium
b) Simple squamous epithelium
c) Stratified squamous non-keratinized
d) Pseudo stratified
e) Stratified columnar epithelium

16. At puberty, most of us face problems of acne which is due to overstimulation of sebaceous glands. How you classify these glands according to mode of secretion?

a) Merocrine secretion
b) Paracrine secretion
c) Autocrine secretion
d) Holocrine secretion
e) Apocrine secretion

17. *Transitional epithelium is not found in:*

a) *Ureter*
b) *Major calyx*
c) *Urinary bladder*
d) *Membranous urethra*
e) *Proximal part of urethra in males as far as the ejaculatory ducts*

18. *Most commonly used dye for tissue preparation for microscopy is:*

a) *Methylene blue*
b) *Hematoxylin and eosin*
c) *Orange g*
d) *Acid Fuscin*
e) *Silver stain*

19. *Euchromatin found in the nucleus is:*

a) *Folded, coiled, or condensed chromatin*
b) *Straight, uncoiled or extended chromatin*
c) *Coiling of long strand of nucleosomes*
d) *Degenerated form of DNA*
e) *Double coiled genetic material*

29. Hematopoietic tissue in healthy adults is

a) Spleeen
b) Liver
c) Red bone marrow
d) Kidney
e) Lymph node

30. The first blood cell progenitor in embryo is:

a) Paraxial mesoderm
b) Intermediate mesoderm
c) Lateral mesoderm
d) Extra embryonic mesoderm
e) Angioblast cell

31. The Ultrasound report revealed excessive amount of amniotic fluid in a pregnant woman (Polyhydramnios) What is normal amount of amniotic fluid in full term pregnancy?

a) 50ml
b) 400ml
c) 1000ml
d) 2000ml
e) 3000ml

32. A physician told a pregnant woman that she had Oligohydramnios. Following condition can cause oligohydramnios:

a) Anencephaly
b) Esophageal atresia
c) Maternal diabetes
d) Renal agenesis
e) Spina bifida

33. The placenta membrane (blood-placenta barrier) is formed by from out to inside in sequence:

a) Cytotrophoblast - syncytotrophoblast connective tissue, endothelium.
b) Syncytotrophoblast - cytotrophoblast - endothelium, connective tissue.
c) Syncytotrophoblast, cytotrophoblast, connective tissue, endothelium.
d) Endothelium - connective tissue - cytotrophoblast syncytotrophoblast.
e) Syncytotrophobast, connective tissue, cytotrophoblast, endothelium

44. The primary visual cortex is located in which lobe?

a) Temporal
b) Occipital
c) Parietal
d) Frontal
e) Insula

45. Injury to the medial lemniscus causes;

a) Pain deficits on the same side
b) Tactile deficits on the opposite side
c) Temperature deficits on the same side
d) Proprioceptive deficits on the same side
e) vibration deficits on the same side

46. The middle 2/3 of crus cerebri contain:

a) corticospinal fibre
b) frontopontine fibers
c) temporopontine fiber
d) medial longitudinal fasciculus
e) lateral longitudinal fasciculus

47. A patient presents with vertical diplopia, head tilting and also difficulty in coming downstairs. The likely nerve damaged is:

a) Oculomotor nerve
b) Trochlear nerve
c) Optic nerve
d) Abducent nerve
e) Trigeminal nerve

48. Patient has increased knee jerk due to

a) Corticospinal tract lesion
b) Lmn lesion
c) Spinothalamic lesion
d) Spinal nerve l2
e) None of the above

69. The sensory nerves supplying the scalp are all except

a) Supratrochlear nerve
b) Supra orbital nerve
c) Infra orbital nerve
d) Greater occipital nerve
e) Lesser occipital nerve

70. The neck is divided into anterior and posterior triangles using ____ muscle as land mark:

a) Trapezius.
b) Sternohyoid.
c) Scalenus anterior
d) Sternocleidomastoid
e) Anterior belly of digastric

71. Which of the following nerve gives parasympathetic fibres to parotid gland

a) Vagus
b) Facial
c) Trigeminal
d) Glossopharyngeal
e) Accessory nerve

72. Regarding the Thyroid gland, point out the incorrect statement

a) Is surrounded by sheath of pretracheal fascia
b) Has isthmus covering 2nd -4th tracheal ring
c) Receives blood from a branch of costocervical trunk
d) Receives blood from external carotid artery
e) Receives blood from aorta

73. Best statement regarding carotid vessels is

a) Common carotid arteries arise from arch of aorta on both sides
b) Internal juglar vein lies on medial side of common carotid artery within carotid sheath
c) External carotid arteries arise at the level of upper border of cricoid cartilage
d) Superior thyroid artery is the first branch of external carotid
e) Internal carotid artery gives two branches before entering the skull

84. In the anterior view of the viscerocranium, which one of the following bones is single?

a) Nasal.
b) Frontal.
c) Lacrimal.
d) Maxillary.
e) Zygomatic.

85. What is the strongest bone among the bones of the face?

a) Nasal.
b) Frontal.
c) Maxillary.
d) Zygomatic.
e) Mandibular.

86. The superior orbital margin is formed by the ____ bone:

a) Frontal.
b) Maxillary.
c) Zygomatic.
d) Ethmoidal.
e) Sphenoidal.

87. What part of the anterior pituitary gland is mainly responsible for hormone secretion?

a) Hypothalamus
b) Pars intermedia
c) Pars tuberalis
d) Pars anterior
e) Pars nervosa

88. Neurovascular bundle in chest wall is located at:

a) Inferior border of ribs
b) uperior border of ribs
c) Anterior to ribs
d) Lower part of intercostal space
e) Middle part of intercostal space

89. During respiration, Muscle which increases the anteroposterior and transverse diameter of thoraxic cavity is:

a) Diaphragm
b) External intercostal muscles
c) Internal intercostal muscles
d) Inner most intercostal muscles
e) Internal & Innermost intercostal muscles

90. Right border of heart is mainly formed by:

a) Right ventricle
b) Right atrium
c) Left ventricle
d) Left atrium
e) Left auricle

91. Left Anterior descending artery is a branch of:

a) Right coronary artery
b) Left coronary artery
c) Marginal artery
d) Circumflex artery
e) Posterior interventricular artery

92. Which among the following structures is present in posterior mediastenum:

a) Trachea
b) Ascending aorta
c) Descending thoracic aorta
d) Superior vana cava
e) Thymus

93. A little boy while playing engulfed a tiny peanut which got stuck in airway, where will it go:

a) Right bronchus
b) Left bronchus
c) Esophagus
d) Larynx
e) Pharynx

94. Which lobe of lung is inferior to horizontal fissure:

a) Inferior lobe of right lung
b) Middle lobe of right lung
c) Superior lobe of left lung
d) Middle lobe of left lung
e) Inferior lobe of left lung

95. Which of the following structures of Gastrointestinal tract is lined by stratified squamous non-keratinized epithelium:

a) Esophagus
b) Stomach
c) Ascending colon
d) Descending colon
e) Small intestine

96. Which structure is located immediately distal to the jejunum in the gastrointestinal tract:

a) Duodenum
b) Ileum
c) Cecum
d) Ascending colon
e) Transverse colon

97. Where in the abdomen is the liver predominantly located:

a) Epigastrium
b) Left hypochondrium
c) Right hypochondrium
d) Right flank
e) Subhepatic space

98. Implantation of embryo occurs in which part of female reproductive system:

a) Fundus of uterus
b) Body of uterus
c) Cervix of uterus
d) Isthmus of fallopian tube
e) Ampulla of fallopuan tube

99. Sertoli cells and leydig cell are present in:

a) Prostate
b) Testes
c) Vas deferens
d) Epididymis
e) Ejaculatory duct

100. Lymphatic drainage of supra-renal gland is carried via:

a) axillary nodes
b) Inguinal nodes
c) lumbar nodes
d) Messenteric nodes
e) Thoracic nodes

The Bell's palsy is characterized by lesion of:

a) Upper part of facial nerve nucleus
b) Lower part of facial nerve nucleus
c) Facial nerve in facial canal above the origin of nerve to stapedius
d) Facial nerve in facial canal above the origin of chorda tympani
e) Facial nerve in facial canal just before it emerges from stylomastoid foramen

80. In the occulomotor nerve lesion, pupil does not react to accommodation due to the damage to:

a) Postganglionic fibers from ciliary ganglion to sphincter pupillae
b) Post ganglionic fibers from ciliary ganglion to ciliary muscles
c) Sympathetic fibers to visceral part of levator palpabrace superiors
d) Motor fibers in superior division supplying superior rectus
e) Motor fibers in inferior division supplying medical rectus

81. Which statement is incorrect about cutaneous innervation of face

a) Forhead ----------ophthalmic nerve
b) Upper lip ---------- Maxillary nerve
c) Tip of nose ---------- ophthalmic nerve
d) Parotid gland ----------mandibular nerve
e) Anterior part of temple --------mandibular nerve

82. Regarding the nerves of orbit all are true except

a) Optic nerve
b) Zygomatic nerve
c) Lacrimal nerve
d) Inferior alveolar nerve
e) Nasociliary

83. Blood supply of trachea in neck is mainly derived from

a) Superior thyroid artery
b) Subclavian artey
c) Inferior thyroid artery
d) Internal carotid artery
e) Common carotid artery

49. The only abductor of vocal cord is

a) Posterior crico-arytenoid
b) Laterla crico-aryteniod
c) Cricothyroid
d) Thyroarytenoid
e) None of the above

50. Inferior thyroid artery ligation may cause injury to

a) External laryngeal nerve
b) Vagus nerve
c) Superior laryngeal nerve
d) Recurrent laryngeal nerve
e) Internal laryngeal nerve

51. Following muscle elevate the soft palate during swallowing

a) Levator velipalatini
b) Palato glossus
c) Palato pharyngeus
d) Salpingo pharyngeus
e) Tensor Veli palatini

52. Tensor tympani muscle inserted on

a) Neck of malleus
b) Long process of incus
c) Neck of stapes
d) Handle of malleus
e) Anterior process of malleus

53. Investing layer enclose which of the following structure

a) Thyroid gland
b) Parathyroid gland
c) Infrahyoid muscles
d) Trapezius
e) Subclavian vessels

74. Little's area of nose consists of

a) Arterial plexus
b) Venous plexus
c) Nerve plexus
d) Lymphatic plexus
e) Arterio-venous plexus

75. Regarding the blood supply of face

a) Facial artery is the posterior branch of external carotid artery
b) Skin flaps in the plastic surgery of face usually necrose because of profuse blood supply to face
c) Superficial temporal artery is a branch of facial artery
d) Superficial temporal branch is a terminal branch of external carotid artery
e) Sub mental artery is a branch of external carotid artery

76. Regarding the infra temporal fossa

a) It is a space in the skull
b) It is related to carotid sheath
c) Maxillary artery and the temporalis muscles are the only contents
d) Its medial boundary is the pterygoid plate
e) It communicates with the pterygo palatine fossa through pterygo maxillary fissure

77. With swelling of mucous membrane of superior nasal meatus, which of the opening is plugged?

a) Superior Para nasal sinus
b) Anterior ethmoidal sinus
c) Middle ethmoidal sinus
d) Sphenoidal sinus
e) Frontal sinus

78. Foot plate of stapes is placed at

a) Opening of pyramid
b) Oval window (fenestra vestibuli)
c) Round window (fenestra cochlear)
d) Aditus
e) Epitympanic recesses

64. Select incorrect statement regarding components of anterior triangle

a) Myohyoid muscle
b) Digastric muscle
c) Sternocleiod mastoid muscle
d) Thyrohyoid muscle
e) Sternohyoid muscle

65. All of the following structures pass through foramen ovale except

a) Lesser petrosal nerve
b) Middle meningeal artery
c) Emissary vein
d) Accessory meningeal artery
e) Mandibular nerve

66. The characteristics feature of second cervical vertebra is

a) Presence of odontoid process
b) Heart shaped body
c) Presence of condyles
d) Trifid spine
e) Massive body

67. The cutaneous nerves to neck carrying fibers of dorsal rami of cervical spinal nerves is

a) Greater auricular nerve
b) Greater occipital nerve
c) Lesser occipital nerve
d) Transverse cervical nerve
e) Supraclavicular nerve

68. Which of the following structure is not supplied by Hypoglossal nerve

a) Genioglossus
b) Hyoglossus
c) Stylo glossus
d) Superfical longitudinal muscle of tongue
e) Palatoglossus

54. Regarding the sclaneus anterior muscle, choose the best statement

a) It is attached to second rib
b) It lies posterior to subclavian vein
c) It is supplied by phrenic nerve
d) It lies behind roots of Brachial plexus
e) It runs behind the subclavain artery

55. A muscle which elevates and retracts the mandible is

a) Masseter
b) Temporalis
c) Medial pterygoid
d) Lateral pterygoid
e) Digastric

56. Stylo pharyngeus muscle is supplied by

a) Pharyngeal plexus
b) Accessory nerve
c) Vagus nerve
d) Hypoglossal nerve
e) Glossopharyngeal nerve

57. Cricopharyngeus muscle is part of

a) Upper constrictor
b) Lower constrictor
c) Middle constrictor
d) Salpingo pharyngeous
e) Stylo phyaryngeous

58. The only suprahyoid muscle which is supplied by cervical plexus is

a) Digastric
b) Stylohyoid
c) Mylohyoid
d) Thyroid
e) Geniohyoid

39. The passage between the third ventricle and lateral ventricle of brain is called the:

a) Foramen of Magendie
b) Aqueduct of sylvius
c) Interventricular foramen(Monro)
d) Foramen Luschka
e) Central canal

40. The part of motor cortex innervating the leg area is supplied by which artery?

a) Anterior cerebral
b) Middle cerebral
c) Posterior cerebral
d) Basilar artery
e) Vertebral artery

41. Which of the following nerve arises from floor of interpeduncular fossa?

a) Optic nerve
b) Trochlear
c) Oculomotor
d) Trigeminal
e) Facial

42. The opening present in the roof of the 4th ventricle is

a) Foramen Monro
b) Foramen Magenda
c) Foramen Lushka
d) Aqueduct of sylvius
e) Central canal

43. The main motor area is located in:

a) Insula
b) Superior frontal gyrus
c) Post central gyrus
d) Pre central gyrus
e) Superior temporal gyrus

20. Cell apoptosis (cell death) is carried out by:

a) Peroxisomes
b) Smooth Endoplasmic Reticulum
c) Lysosomes
d) Golgi apparatus
e) Mitochondria

21. A student observing histological section of skin on microscope. Lining epithelium identified will be:

a) Simple cuboidal
b) Simple squamous
c) Stratified squamous keratinized
d) Stratified squamous non-keratinized
e) Simple columnar

22. While studying respiratory epithelium on electron microscope, small hair like processes with rapid to-and-fro oscillatory movement is observed on its surface, these are called:

a) Flagellum
b) Cilia
c) Micro-villi
d) Villi
e) Stero-villi

23. During examination, histological section of active thyroid follicle has been placed. The identified epithelium will be:

a) Stratified squamous
b) Simple cuboidal
c) Simple columnar
d) Transitional
e) Stratified columnar

24. A 20-year old woman presents at the ER with severe abdominal pain on the right side with signs of abdominal bleeding. She indicated that she has been sexually active without contraception and missed her last menstrual period. Based on this information, which of the following disorders must be included as an option in the diagnosis?

a) Ovarian cancer
b) Appendicitis

c) Normal pregnancy
d) Ectopic tubal pregnancy
e) Toxemia of pregnancy

25. Mesoderm is formed from

a) Hypoblast cells
b) Extraembryonic ectoderm
c) Notochordal cells
d) Epiblast cells(ans)
e) Amnioblasts

26. Which of the following statements regarding the uteroplacental circulation is CORRECT?

a) Lacunae form within the cytotrophoblast
b) Hydrolytic enzymes produced by the trophoblast degrade the endothelium of embryonic capillaries
c) It is established during the second week of development(ans)
d) It is a transient nutrient supply for the blastocyst during implantation
e) It only forms in ectopic implantation

27. Angiogenesis is characterized by formation of :

a) New vascular channel
b) New vessels by budding and branching from preexisting vessels.
c) Angioblast cell
d) Mesenchymal cell
e) Mesodermal cell

28. Angioblast cell which aggregate to form blood island derived from:

a) Allantois
b) Umblical vesicle
c) Connecting stalk
d) Amnion
e) Extraembryonic Mesoderm

10. The right lymphatic duct drains which part of the body?

a) Upper left quadrant
b) Lower left quadrant
c) Upper right quadrant
d) Lower right quadrant
e) Left side of the body

11. Most commonly used dye for tissue preparation for microscopy is:

a) Methylene blue
b) Hematoxylin and eosin
c) Orange g
d) Acid Fuscin
e) Silver stain

12. Simple columnar epithelium

a) Choroidal plexus of ventricles
b) Ovary
c) Epidermis
d) testis
e) lips

13. Epithelium of ovary is

a) Simple cuboidal
b) Simple columnar
c) Simple squamous
d) stratified cuboidal
e) statified columnar

14. Compact bone contains:

a) Osteocytes in the lacunar space
b) large number of medullary cavities
c) Covered by spongy bone
d) does not have haversion canal system
e) By periostium internally