- Phase of Amelogenesis
- Dental Modification of Salivary gland
- Cap Stage (very imp)
- Diff b/w masticatory and lining mucosa
- Key hole appearance of Enamel
- Hertwig sheath makes what?
- Types and Location of papilla
- Development of Tongue
- Salivary gland and their secretion
- Tongue k muscles kahan se bante hain?
- Life cycle of Ameloblast
- Components of enamel organ.
- Function of stratum intermedium
- Define Tuft, Lamella, Spindle
- Gingival Fibres.
- Aprismatic enamel?
- Enamel protein?
- Tooth pattern?
- Keratinized & non keratinized k diff
- Diff b/w rods and interods & their strength
- Q where rods are not present

Ans. At the initial & final points (interods are present there).

MID-TERM VIVA QUESTIONS

- Reciprocal Induction
- Secretory Phase of Amelogenesis
- Ductal Modification of Salivary gland
- Cap Stage (very imp)
- Dif b/w masticatory and lining mucosa
- Key hole appearance of Enamel
- Hertwig sheath makes what?
- Types and location of papilla
- Development of Tongue
- Salivary gland and their secretion
- Tongue k muscles kahan se kahan se hain?
- Life cycle of Ameloblast
- Components of enamel organ.
- Function of stratum intermedium
- Define Tuft, Lamella, Spindle
- Gingival fibres.
- Aprismatic enamel?
- Enamel protein?
- Tooth pattern?
- Keratinized & non keratinized k dif

Identification Points

Maxillary Central Incisors -
1. Mesial side longer
2. Mesio-incisal angle at sharp 90°
3. Mesial side straight, distal side convex

Maxillary Lateral Incisors -
1. Mesial - side longer
2. Mesio-incisal angle at sharp 90°
3. Mesial side straight, distal side convex

Mandibular Central Incisor -
1. Most symmetrical tooth
2. Mesial edge sharper than distal
3. Smaller than lateral - most smallest tooth

Mandibular Lateral Incisor :-
1. Larger than central
2. Mesial edge sharper
3. Cingulum deviated towards mesial side

Maxillary Canine :-
1. Mesial slope shorter than distal
2. Cingulum well-developed compared to mandibular canine

Mandibular Canine :-
1. Distal slope shorter than mesial
2. Cingulum less developed
3. Shorter than maxillary

Maxillary 1st Premolar :-
1. Mesial developmental groove
2. Mesial Slope shorter than distal
3. Buccal Cusp 1 mm longer than lingual.

2nd Maxillary Premolar:-

1. Mesial developmental groove, less prominent than 1st Premolar
2. Buccal cusp longer
3. Mesial side longer than distal slope

Mandibular 1st Premolar:-

1. Tooth is inclined towards lingual side.
2. Buccal cusp longer than lingual
3. Mesial slope shorter
4. Lingual lobe present on distal side

Mandibular 2nd Premolar:-

1. Tooth inclined towards lingual side.
2. Buccal cusp longer.
3. Mesial slope shorter
4. Lingual lobe more prominent than 1st Premolar sometime may develop into 2nd Lingual cusp

Maxillary 1st Molar:-

1. 4 cusps
2. Oblique ridge. (mesiolingual to disto-buccal)
3. Mesiolingual largest cusp.
4. Lingual side more convex
5. Buccal side more in height than lingual side

Maxillary 2nd Molar:-

1. 4 cusps
2. Oblique ridge (mesiolingual to disto-buccal cusp)
3. Mesiobuccal largest cusp.
4. Lingual side more convex
5. Buccal side height more than lingual side.

Maxillary Teeth

1) Max Central Incisor: Triangular from Proximal & Incisal Surface.
Crown shape: Trapezoidal.

2) Max lateral Incisor. Triangular Proximal Surface
Incisal Surface Ovoid.
Crown shape Trapezoid.

3) Max Canine.
Proximal Surface: Triangular
Incisal Surface: Diamond
Crown Shape: Trapezoidal.

(4) Max 1st premolar:
Proximal Surface: Trapezoidal
Occlusal table: Trapezoidal
Occlusal Outline: Hexagonal
Crown Shape: Trapezoidal

Maxillary Teeth.

(5) Max 2nd Premolar:
Proximal Surface: Trapezoidal
Occlusal table: rectangular.
Occlusal Outline: Ovoid
Crown shape: Trapezoidal

(6) Max 1st Molar: Proximal Surface: Trapezoidal
Trapezoidal to roughly Rhomboidal: Occlusal Surface: Trapezoidal
Crown shape: Trapezoidal

(7) Max 2nd Molar: Proximal Surface: Trapezoidal
Occlusal Surface: Rhomboidal
Crown shape: ??

(8) Max 3rd Molar: Proximal Surface: Trapezoidal.
Occlusal surface: Heart shaped
Crown shape: ??

(25) How many tooth buds are formed At birth: 52 Confirmed answer

(26) Which fiber join two teeth:
Transseptal fiber.

(27) Function of Myoepithelial cells:
Contraction.

(28) Mums caused by Inflamation of what gland: Parotid gland.

(29) The layer above basal layer shrink away ..... : called Prickle layer.

(30) Basal lamna contain collagen: Type IV

(31) Example of Cartilaginous Joint:

(32) When the Jaw is at rest which Nerve Ending still work: Search Answer.

ERUPTION TIME

Primary Dentition

Mandible

CI = 8 months
LI = 13 months
1M = 16 months
C = 20 months
2M = 27 months

Maxillary

CI = 10 months
LI = 11 months
1M = 16 months
C = 19 months
2M = 29 months

Permanent Dentition

Mandible

1M, CI } 6-7 years
LI = 7-8 years
C = 9-10 years
P1 = 10-12 years
P2 = 11-12 years
2M = 11-13 years
3M = 17-21 years

Maxillary

1M = 6-7 years
CI = 7-8 years
LI = 8-9 years
P1 = 10-11 years
P2 = 10-12 years
C = 11-12 years
2M = 12-13 years
3M = 17-21 years

12- Muscle of mastication are derived from the _____ arch.

13- Muscles of Facial expression are derived from the _____ arch.

14- The branchial arches are covered by _____ from outside and lined by _____ from inside. except the first arch.

15- The Mandibular arch send out a prominence called _____ at its proximal end before extension, while its distal extension is called _____.

- During the _____ period, there is a rapid increase in the overall size of fetus.

Stomodeum is seperated from the blind end of foregut by the _____

All the tooth structures are derived from neural crest cells except _____

The _____ begins by fertilization of ovum till the development of 3 germ layer.

10- Soft palate develop from.

a) Primary palate.
b) Palatal processes fused with the nasal septum.
c) Medial nasal process.
d) Palatal process which are not fused with nasal septum.
) Frontal process.

o.) Inferior parathyroid gland originates from _____

oo) Superior parathyroid gland originate from _____

oo) Extrinsic muscles of Tongue Develop from the _____

.) The palatal shelves (palatine process)

a) covered by endoderm
b) are derived from primary palate
) Must assume a vertical position for palate closure to occur.
are derived from the first branchial arch
~~Must~~ Must undergo epithelio-mesenchymal transformation for seamless union.

Chapter - 2 and 3 -

1- Cervical sinus developed on the side of the neck by the growth of _____

2- Second branchial arch derivatives are: _____

3- Branchial arches sharing in Tongue development are: _____

4. Ossification center of the developing mandible arises between: _____

5- First branchial arch is supplied by _____

Second branchial arch is supplied by _____

. Except the mandibular condyle most of mandibular growth is.

a) Intramembranous. c) sutural. e) N. of abv.
b) Endochondral d) Both a & b

Which of the following is not a correct pairing of embryonic structure and adult derivative.

a) Maxillary process: secondary palate
b) primary palate: Upper incisor teeth
c) second pharyngeal arch: Mandible
d) Lateral lingual swelling: anterior 2/3 of Tongue
e) Meckels cartilage: Maxilla

25- Somites develop from the _____

26 Example of trisomic condition is _____

27- ~~Which~~ of the fate of the most posterior extremity of the meckel's Cartilage:

a) sphenomandibular ligament
b) sphenomaxillary ligament.
c) Mandibular condyle.
d) Coronoid process.
e) Incus & malleus.

28- Rhombomeres is the eight series of bulges formed from:

a) Three layered folding.
b) By the fusion of Neural tube
c) By the fusion of neural folds.
d) From the part of hind brain
e) from the Rostal fold of embryo.

fusion of hemimandibles at symphysis occurs at age of

a) 5 year
b) 1 year of age.
c) 5 month in utero
d) Donot fuse.
e) 10 year of age.

Lower lip developed by the median fusion of the two _____ prominence

Branchial arches are seperated by external shallow depression called _____

_____ arch is supplied by the internal and common carotid arteries.

- Maxillay Molar 3 root → largest root → Palat

- Fibre which Joint Two Root Together. → Transeptal fibre

- Sclerotic Dentin

- Formation of Secondary Dentin → of Pulp Root canal

1- In Slow Fibis, which of the following neurotransmitter is not used? Glutamate.

12- The following structure divides TMJ into two.

a- Right & left condylar head
b- Articular disk
c- Capsule & ligament
d- Temporal fossa
e- Mylohoid muscle

13- The following is responsible for the dynamic mechano reception & movement audendo in TMJ.

a- Ruffini's corpuscle
b- Pacini's corpuscle.
c- Golgi Tendon.
d- Free Nerve Organ
e- Nerve transmitter.

7Q. Which of the following is an anatomical feature of articular disk.

a- Thick posterior & anterior band.
b- Thick central zone
c- Thin posterior band
d- Thin bilaminar zone.
e- Thin perforated zone.

8- TMJ formed at which month.
Ans.- 3 month.

9- Example of Cartilaginous Joint

10- When the Jaw is at rest which Nerve Ending still work-

11- Gomphosis is the type of fibrous joint.

1- Golgi Tendon organ remain completely inactive in immobile Joints, becoming active only when the Joint is at extremes of its range of movement -

2- Articular disk is divided by capsule.

3- Ossification begin first in the Temporal blastema.

4- The human temporomandibular joint is an example of ginglymo-arthrodial articulation.

The Joint is capable of which of the following movements.

a- Gliding or translator movement only.
b- Hinge or ~~rotation~~ Rotational movement only.
c- Anterior & Superior movement only.
d- Combined rotational & translatory movement
e- Combined anterior & gliding movement

5. Basic structure of the TMJ consist of.

a- condyle, mandibular fossa, disc, articular eminence.
b- Condyle, fossa of sphenoid bone, articular eminence.
c- Condyle, fossa of ethmoid bone, articular eminence.

6- Which one of the following is considered to be accessary ligament of masticatory apparatus? TMJ.

a. stylomandibular ligament
b Lateral ligament.
c- Temporo-mandibular ligament
d- Zygomatico mandibular ligament.

- Chapter: 4 -

1. Tight Junction belong to Occludin, ____ Family.

2. Zonula Adherons belongs to Cadherin family.

3. In Collagen synthesis, 3rd part is glycine

Intermediate mesoderm derivative.
Urogenital system.

Secondary Palate development 7-8week

Maxilla ossification center.

∴ Congenital defect major factor.
Tekogen-

Mitossis- diploid no of chromosome.
~~trap~~

33- Have lip is median cleft

34- Primitive yolk sac is developed in 2nd Week

35- Fetal Stage is Prenatal development of 3rd stage

36- Autosomal Dominant Condition.
Achondroplasia.

37- Monosomy condition include what chromosomes

38- Primary Palate is formed by the fusion of:

39- secondary palate formed at what week _____

40- Patterning is a spatial & temporal event.

Regarding Endochondral bone formation:

condary spongiosa is formed by Woven bone
only on diaphysis
by Lamellar bone
only on cartilage surface.
Bone develop directly within the
soft connective tissue

- Blastocyst is formed in 2nd Week.

- Taste bud is supplied by ——
nerve

- Larynx is formed by ——
pharyngeal arch

27- ~~Which of~~ The fate of the most posterior extremity of the meckel's cartilage:

a) Sphenomandibular ligament
b) Sphenomaxillary ligament
c) Mandibular condyle.
d) Coronoid process
e) Incus & malleus.

28- Rhombomeres is the eight series of bulges formed from:

a) Three layered folding.
b) By the fusion of Neural tube
c) By the fusion of neural folds.
d) From the part of hind brain
e) from the Rostral fold of embryo.

Which of the following is not a correct pairing of embryonic structure and adult derivative.

a) Maxillary process: Secondary palate
b) Primary palate: Upper incisor teeth.
c) Second pharyngeal arch: Mandible.
d) Lateral lingual swelling: ant 2/3 of Tongue.

e) Meckels cartilage: Maxilla

f- Somites develop from the.

---

g) Example of trisomic condition is _____.

c) Inferior parathyroid gland originates from _____

d) Superior parathyroid gland originate from _____

e) Extrinsic muscles of Tongue Develop from the _____

f) The palatal shelves (palatine process)

a) covered by endoderm

b) are derived from primary palate Must assume a vertical position for palate closure to occur.

are derived from the first branchial arch Must Must undergo epithelio-mesenchymal transformation for seamless union.

Stomodeum is seperated from the blind end of foregut by the

- All the tooth structures are derived from neural crest cells except ____

- The ____ begins by fertilization of ovum till the development of 3 germ layers.

- Soft palate develop from.

a) Primary palate
b) Palatal processes fused with the nasal septum.
c) Medial nasal process.
d) Palatal process which are not fused with nasal septum.
e) Frontal process.

Fusion of hemimandibles at symphysis occur at age of

a) 5 year
b) 1 year of age.
c) 5 month in utero
d) Donot fuse.
e) 10 year of age.

Lower lip developed by the median fusion of the two _____ prominence

Branchial arches are seperated by external shallow depression called _____

_____ arch is supplied by the internal and common carotid arteries.

12- Muscle of mastication are derived from the _____ arch

13- Muscles of Facial expression are derived from the _____ arch

14- The branchial arches are covered by _____ from outside and lined by _____ from inside. except the first arch.

15- The Mandibular arch send out a prominence called _____ at its proximal end before extension, while its distal extension is called _____.

16- During the _____ period, there is a rapid increase in the overall size of fetus.

Chapter - 2 and 3 -

1- Cervical sinus developed on the side of the neck by the growth of.

_____

2- Second branchial arch derivatives are:- _____

3- Branchial arches sharing in Tongue development are:- _____

4- Ossification center of the developing mandible arises between _____

5- First branchial arch is supplied by.

_____

6- Second branchial arch is supplied by

_____

7- Except the mandibular condyle most of mandibular growth is

a) Intramembranous. c) Sutural. e) None of above.
b) Endochondral d) Both a & c

Primary function of dental pulp

1) Assure root & closure
2) Provide nutrition.
3) Provide sensation.
4) Form Dentin
5) Form Enamel.

1- ~~OBJECTIVE~~ Dentition.

Chapter 1.

- Teeth constitute about ___ % of total surface area of mouth.

2- Main constituent of tooth.
- Dentine,
  Pulp
  Enamel
  Cementum.

3- Main function of pulp
  a) Nutritive
  b) reperative
  c) Formation of dentine

1- Tooth proper consist of

enamel, dentin, pulp
(soft connective tissue).

103- The greatest no. of cell present on the upper surface of pulp are odontoblast
104- Pulp stones are frequently found in orifices
105- Contour lines of owen are found in dentine
06 Lingual groove present in ____
107- Mesiolingual groove present in ____
108- Mesial marginal groove present in ____
109→ In 1st premolar lingual cusp deviate from ____
110→ On occlusal aspect small concavity irregular depression is called
111- A tooth which have prominent transverse ridge and no central groove ____
112 Y Shape on occlusal aspect present in mand 2nd 3 cusp type
113- In maxillary 1st Pm buccal cusp is longer than lingual cusp