


Q.11: Right ward shift of the curve indicate

a) Haldane effect
b) Bohr's effect
c) Oxygen binds with haemoglobin
d) Low PCO2
e) Decreased 2,3,DPG

Q.12: Normal venous blood returning from the peripheral tissues, the P_{O_2} is about 40 mm Hg, and the saturation of haemoglobin averages

a) 95 %
b) 85 %
c) 75 %
d) 65 %
e) 55%

Q.13: If an adult man has 15gm haemoglobin / dl of blood the oxygen binding capacity in 100 ml of blood is


a) 1.34 ml of oxygen
b) 10.34 ml of oxygen
c) 15.01 ml of oxygen
d) 20.1 ml of oxygen
e) 21.1 ml of oxygen

Q.14: left ward shift of oxygen haemoglobin dissociation curve in

a) Increased body temperature
b) Decreased PO2
c) Increased 2,3 DPG
d) Acidosis
e) Alkalosis

Q.15: Normal partial pressure of oxygen in systemic arterial blood is

a) 40 mmHg
b) 50 mmHg
c) 87 mmHg
d) 95mmHg
e) 104 mmHg



Q16: In above mentioned diagram number "1" is

a) Submandibular gland
b) Sub lingual gland
c) Parotid gland
d) Otic ganglion
e) Pharyngeal salivary gland

Q17: Parotid gland supplied by post ganglionic fibers from

a) Otic ganglion
b) Petrosal ganglion
c) Submandibular ganglion
d) Ciliary ganglion
e) Geniculate ganglion

Q18: Preganglionic fibers to the parotid gland runs in which of the following cranial nerve

a) III
b) IV
c) VII
d) IX
e) X

Q19; Daily average secretion of saliva is


a) 1L
b) 2 L
c) 2.5 L
d) 2.6 L
e) 3L

Q20: Which of the following enzyme in saliva is concerned starch digestion

a) Lingual lipase
b) Trypsin
c) Chymotrypsin
d) Ptyalin
e) Carboxypeptidase

11:42

Facebook



Hall: Guyton and Hall Textbook of Medical Physiology, 12th Edition
Copyright © 2011 by Saunders, an imprint of Elsevier, Inc. All rights reserved.


Clinical Anatomy & Operative Surgery - Salivary Glands innervation ...

Visit ›

Images may be subject to copyright. Learn More

Share

Save



Vetscraft

Nerve supply of salivary gl...

From this image




Submandibular ganglion



Chorda tympani

superior inferior olivary nucleus tract otic gangliom



Q1. Phase 0 represents?

a) Influx of sodium
b) Closer of sodium channel
c) Influx of potassium channel
d) Closer of potassium channel
e) Opening of calcium channel

Q2. Phase 1 represents?

a) Influx of sodium
b) Partial repolarization
c) Influx of potassium channel
d) Closer of potassium channel
e) Opening of calcium channel

Q3.Phase 2 represents?

a) Influx of sodium
b) Partial repolarization
c) Influx of potassium channel
d) Closer of potassium channel
e) Opening of calcium channel

Q4.Phase 3 represents?

a) Influx of sodium
b) Partial repolarization
c) Influx of potassium channel
d) Closer of potassium channel
e) Opening of potassium channel

Q5.Phase 4 represent?

a) Influx of sodium
b) Partial repolarization
c) Influx of potassium channel
d) Closer of potassium channel
e) Resting membrane potential

Trousseau's sign

Trousseau sign of latent tetany is a medical sign observed in patients with low calcium.

