

Item Response Theory - Final Essay

Marius Keute

September 23, 2022

Contents

1	Introduction	3
2	Preparing the Data	4
3	Descriptive Analyses and Dichotomization	5
4	IRT modeling	7
4.1	Rasch model estimation	7
4.2	Model analysis	8
4.3	Differential Item Functioning	10
5	Higher-parameterized IRT models	10
6	Polytomous IRT model	13
7	Factor models	13
8	Reliability and Unidimensionality	15
9	Measurement Invariance	16
10	Theoretical Part: Key differences between IRT and CTT	17
10.1	Introduction	17
10.2	Core Ideas and Terminology	18
10.3	Strenghts	18
10.4	Limitations	18
10.5	Conclusion and Application	18
11	Analysis code	19
12	References	65

submitted to:

Dr. Stefano Noventa

University of Tübingen

submitted by:

Marius Keute (QDS, 5991873)

Statutory Declaration: I hereby declare that I composed the present paper independently and that I have used no other resources than those indicated. The text passages which are taken from other works in wording or meaning have been identified as such. I also declare that this work has not been partly or completely used in another examination.

The full R and Markdown code used for generating this essay is available on Github:

<https://github.com/mkeute/IRT-essay>

1 Introduction

Understanding sexual habits and behavior can be important for, e.g., improving sex education for adolescents, preventing sexually transmitted diseases (STDs), and identifying high-risk populations for sexual misconduct. The Sexual Compulsivity Scale (SCS) is a 10-item questionnaire constructed to measure hypersexuality and high libido in a given person (Kalichman and Rompa (1995), Kalichman and Rompa (2001)). Each of the 10 items is a statement about sexual habits, feelings, or experiences, and the test-taker can indicate how much they can relate to each statement on a four-level scale ranging from 1 (Not at all like me) to 4 (Very much like me).

The 10 items are (Kalichman and Rompa (2001)):

1. My sexual appetite has gotten in the way of my relationships.
2. My sexual thoughts and behaviors are causing problems in my life.
3. My desires to have sex have disrupted my daily life.
4. I sometimes fail to meet my commitments and responsibilities because of my sexual behaviors.
5. I sometimes get so horny I could lose control.
6. I find myself thinking about sex while at work.
7. I feel that sexual thoughts and feelings are stronger than I am.
8. I have to struggle to control my sexual thoughts and behavior.
9. I think about sex more than I would like to.
10. It has been difficult for me to find sex partners who desire having sex as much as I want to.

In this essay, using data from the original validation cohort (Kalichman and Rompa (2001)), I will provide a thorough analysis of the SCS, using methods derived from Item Response Theory (IRT), and to a lesser extent from Classical Test Theory (CTT). In the final section, I will give an overview over both theories and their key differences.

2 Preparing the Data

The dataset (Kalichman and Rompa (1995)) consists of 3376 observations, the variables being the ten items of the SCS, the sum score, gender and age. From the age variable, three cases where the reported age was 100 years or higher appeared implausible and therefore set to missing values. The remaining cases had a mean age of 30.9 years (median 28 years, range [14, 85]). From the gender variable, 13 values were missing and 15 cases where the reported gender was “3” (other) were set to missing values. Of the remaining cases, 2295 (68.5%) reported male gender (“1”) and 1053 (31.4%) reported female gender (“2”). In the dataset, 133 cases contained at least one missing value.

The pattern of missing SCS items is shown in Figure 1. It can be seen that item Q9 was missing most often, though not by a large margin (Q9: 27 missing values, Q5: 13 missing values). It can be seen that the majority of cases with missing values (118 cases / 88.7%) had only a single missing item, while there were no prominent patterns of items that tended to be jointly missing. Eight cases where more than two SCS items were missing were excluded from all further analyses. For the remaining 3368 cases, the probability of missing values at each SCS variable was modeled as a function of the values in *all other* SCS variables using a logistic regression model:

$$P(M_{i,q} = 1 | X_{i,q}) = \sigma(X_{i,q} \hat{\beta}),$$

where $M_{i,q}$ is 1 if the i^{th} person has a missing value at item $q \in \{Q1, Q2, \dots, Q10\}$, $X_{i,q}$ denotes the item values of all other items except item q , σ is the logistic function $\sigma(x) = \frac{1}{1+e^{-x}}$, and $\hat{\beta}$ are the estimated regression weights (Guan and Yusoff (2011)). Note that each variable’s pattern of missing values could only be predicted based on the observations without missing values in any other variable, since those cases were excluded by the logistic model by default of the implementation. Since the majority of cases had either no or only one variable missing, however, this should not bias the overall picture very much.




Figure 1: Pattern of missing SCS values.

3 Descriptive Analyses and Dichotomization

The distribution of responses for each item before dichotomization can be seen in Figure 2. All item categories show reasonable coverage of the range of responses (1-4), and there are no obvious flooring or ceiling effects, except for a slight tendency to a flooring effect with item Q6 (few cases with response 1).




Figure 2: Distribution of non-dichotomized responses per item

For dichotomization of the item data, I considered two options, namely, thresholding each of the 10 items at its own median, to ensure an even distribution of observations into both categories for each item, or finding a common threshold for all items. Since the items have only four levels each, a median split would not necessarily lead to a very balanced dichotomization. Furthermore, the item levels are designed to have the same meaning across all items, therefore I decided to dichotomize at a common threshold of 2, i.e., the dichotomous items $D_q \in \{D_1, D_2, \dots, D_{10}\}$ were defined such that

$$D_{i,q} = \begin{cases} 0 & \text{if } Q_{i,q} \in \{1, 2\}, \\ 1 & \text{if } Q_{i,q} \in \{3, 4\}, \end{cases}$$

Of note, simple models in IRT such as the Rasch model (see below) assume that all item responses are either correct or incorrect (or solved / unsolved, respectively). Since a personality test such as the SCS does not have right or wrong responses, it is common to dichotomize the values, as described above, and henceforth treat one of the dichotomous response options as the ‘correct’ one, in this case, responses greater than 2. This is, however, purely for compliance with IRT terminology and does not imply that the ‘correct’ dichotomous responses are better than the ‘incorrect’ ones in any way.

Descriptive characteristics of the 10 SCS items are shown in Table 1, the proportions of correct responses are shown in Figure 2. Since most variables’ median was 2, this was not much different from an item-wise median threshold (see Table 1).

Moreover, I calculated item discrimination, i.e., each items ability to discriminate between high- and low-scoring individuals, using the adjusted item-total correlation method (Reynolds and Livingston (2021)), i.e., by calculating biserial correlation coefficients between each (dichotomized) item’s scores and the sum of all other (dichotomized) items.

Tetrachoric item intercorrelations are shown in Figure 3. It can be seen that all pairs of items show moderate to high positive correlations, indicating that all items measure similar information yet are not redundant (see below for further scrutiny of factorial structure). Item easiness (i.e., proportion of correct responses) was between 27% (item Q4) and 71.9% (item Q6), item discrimination was

Table 1: Descriptive item statistics (mean, median and range *before* dichotomization)

X	stat	Q1	Q2	Q3	Q4	Q5	Q6	Q7	Q8	Q9	Q10
1	max	4.0	4.0	4.0	4.0	4.0	4.0	4.0	4.0	4.0	4.0
2	mean	2.3	2.2	2.2	1.9	2.2	3.1	2.2	2.3	2.5	2.5
3	median	2.0	2.0	2.0	2.0	2.0	3.0	2.0	2.0	2.0	3.0
4	min	1.0	1.0	1.0	1.0	1.0	1.0	1.0	1.0	1.0	1.0

Table 2: Distribution and discrimination of dichotomized items

X	Q1	Q2	Q3	Q4	Q5	Q6	Q7	
item easiness (percent in category 1)	40.60	37.90	37.30	27.10	38.20	71.90	37.80	4
number of cases in category 1	1366.00	1275.00	1257.00	912.00	1285.00	2420.00	1273.00	139
discrimination	0.45	0.46	0.44	0.34	0.29	0.26	0.42	

between .26 (item Q6) and .45 (items Q1, Q2), i.e., there was no item with a trivial response pattern (e.g., all or no responses correct), and no item was a good representation of the entire scale, since all item discriminations were only moderate in size.




Figure 3: Tetrachoric intercorrelations between items.

4 IRT modeling

After analyzing the SCS data using descriptive statistics and concepts derived from CTT, in the following I will fit and discuss different IRT models to the data.

4.1 Rasch model estimation

Next, I estimated a Rasch model for the SCS data, also known as either the one-parameter logistic model or one-parameter normal ogive model, depending on the parameterization.

It models a given person's chances of solving a given item as a logistic function of the difference between the q^{th} item's difficulty β_q and the i^{th} person's ability θ_i , where β_q and θ_i are latent (unobserved) quantities that are estimated from the dichotomous (solved vs. not solved) item data.

The probability for a given person can then be expressed by the logistic function: $P(D_{i,q} = 1 | \beta_q, \theta_i) = \sigma(\theta_i - \beta_q)$, where σ is the logistic function as specified above. That is to say, it is purely the difference between item difficulty and person ability that explains the correctness of item responses within the model.

Crucially, the model assumes that this relationship is identical for all items, i.e., the logistic function can only be shifted but not changed in slope across items with different difficulty. Item difficulty is, therefore, the only free parameter of the Rasch model, whereas alternative models (see below) also estimate additional parameters.

To obtain a comprehensive picture, I fitted Rasch models using three different software implementations in R 4.1.

The first method was the one implemented in the R package **eRm** (Mair and Hatzinger (2007)). The **eRm::RM** function estimates a Rasch model using conditional maximum likelihood estimation. To make the model identifiable, the user can choose between two model constraints, namely that the model parameters must sum to 0 or that the first item's parameter is fixed to 0. I chose the first (default) option, i.e., forcing item difficulties to sum to 0. Item discriminativity, i.e., the steepest slope of the logistic functions (at $\beta_q = \theta_i$), is fixed to 1 for all items in this implementation.

The second method was the one implemented in the R package **ltm** (Rizopoulos (2006)). The **ltm::rasch** function estimates a Rasch model using approximate marginal maximum likelihood estimation. This package provides the user with more flexibility to impose constraints on the model than **eRm**, I fixed item discriminativity to 1 for all items, to maximize comparability with the **eRm** parameters.

The third method was a structural equation model as implemented in **lavaan** (Rosseel (2012)). Unlike the two previous implementations, **lavaan** requires a more explicitly user-defined model specification, as it does not provide any ready-made function or syntax for Rasch models.

I used a modified copy of the syntax presented in Templin (2022):

```
SCS =~ 1*Q1 + 1*Q2 + 1*Q3 + 1*Q4 + 1*Q5 + 1*Q6 + 1*Q7 + 1*Q8 + 1*Q9 + 1*Q10
```

```
Q1 | t1; Q2 | t1; Q3 | t1; Q4 | t1; Q5 | t1; Q6 | t1; Q7 | t1;  
Q8 | t1; Q9 | t1; Q10 | t1;
```

```
SCS ~ 0;
```

SCS $\sim 1 \cdot \text{SCS}$

Again, I fixed item discriminativities to 1 for all items. The item parameters Q_1, \dots, Q_{10} were subjected to a common threshold τ_1 , and the sum of all item parameters (corresponding to the latent variable SCS) was fixed to 0, as with **eRm**. Moreover, its variance was fixed to unit. Of note, due to limitations of the implementation, **lavaan** is not able to estimate Rasch models using maximum likelihood estimation, but only using mean- and variance-adjusted weighted least squares (WLSMV) estimation, which limits model fit comparisons. **TODO describe conversion to IRT params**

4.2 Model analysis

The item difficulty parameters of the three models are shown in Figure 4, along with the item difficulty derived from CTT (i.e., the proportion of incorrect responses per item in the data). While the parameters differed between the different models, it is important to note that the parameters from all four models (including CTT) were perfectly correlated for all pairs of models (all $r > .999$), which indicates that the parameters of one model are simply affine linear transformations of the parameters of any other model, i.e., while numerically different, the models incorporated identical information about the items. The corresponding item-characteristic curves (ICC) are shown in Figure 5. ICCs are generated by calculating the function graph of the item-wise logistic functions parameterized by item difficulty, across a range of possible person ability values on the x-axis.




Figure 4: Item difficulties in comparison.




Figure 5: Item-characteristic curves for the three Rasch models.

The overall model fit indices are shown in Table3. Of note, log-likelihood and information criteria

Table 3: Fit indices for Rasch models

X	loglik	npar	AIC	BIC	cAIC
eRm	-17891.63	9	35801.26	35856.36	35865.36
ltm	-18567.86	10	37155.72	37216.95	NA

can only be reported for those models fitted using `ltm` and `eRm`, while the `lavaan` model’s fit indices are not comparable, as it was not fitted using maximum likelihood estimation. For brevity, I skip the discussion of the `lavaan` model fit indices. Comparing the fit indices for the `ltm` and `eRm` Rasch models, it can be seen that `eRm` had an overall higher log-likelihood. Since it also had one free parameter less (because of the sum constraint, see above), it was, overall, the preferred model also according to the Akaike and Bayes-Schwarz information criteria.

Finally, to get an impression of how the three models perform for each item, I calculated the mean 0-1-loss per item as:

$$\mathcal{L}_q = \frac{1}{n} \sum_{i=1}^n |f(\theta_i, \beta_q) - D_{i,q}|$$

that is to say, I used the item difficulty parameters β_q and person ability scores θ_i estimated by the models to make predictions for each person and item:

$$f(\theta_i, \beta_q) =: \begin{cases} 1 & \text{if } \sigma(\theta_i - \beta_q) > 0.5, \\ 0 & \text{otherwise} \end{cases}$$

The mean loss is then calculated as the proportion of incorrectly predicted cases for each item. It is shown in Figure 6. It can be seen that the three models, despite differences in parameterization, performed very similarly, with the `eRm` and `lavaan` models performing almost identically, whereas the `ltm` model tended to incur a slightly higher loss, except for items Q6 and Q10. Interestingly, these are the most difficult items, and the fact `ltm` outperformed `eRm` especially for those items might be related to the differences between conditional vs. marginal maximum likelihood estimation, which have the strongest effect in cases where either all or no responses are correct, i.e., for particularly easy or difficult items.




Figure 6: Mean 0-1-loss (proportion of incorrectly predicted responses) per item.

4.3 Differential Item Functioning

I tested for differential item functioning (DIF) using the package `difR` and the procedure outlined in the companion paper (Magis et al. (2010)). DIF is a disadvantageous property of a Rasch model, meaning that item responses differ between subjects from different participant groups, even given the same estimated ability level. The presence of DIF indicates a lack of measurement invariance of the model. The results are displayed in 7. I used the `difLord` method to investigate DIF, but obtained essentially the same results with the `difRaju` method. I discarded `difLRT`, the third recommended IRT-related method, due to its high computational demand. I tested for DIF across genders (male vs. female) and age groups (above median age vs. smaller or equal to median age). Significant DIF (FDR-corrected p -value $< .05$) across the gender groups was detected for items Q5 and Q10, and across the age groups for item Q1, Q5, and Q10. However, the effect sizes were in the negligible range for all but item Q5 in the gender group comparison, where a moderate effect was detected ($\Delta_{Lord} = -1.14$). Since I have only been considering Rasch (one-parameter) models so far, this analysis only tested for uniform DIF.




Figure 7: p -values (FDR corrected, negative log-transformed) from DIF analysis. Black line: significance threshold ($p < .05$)

5 Higher-parameterized IRT models

There are several extensions and alternatives to the Rasch model with its restrictive assumptions that differences between items can be described by just one parameter, namely item difficulty, while the Birnbaum model or 2-parameters logistic model also takes into account item discriminativity (corresponding to varying slopes of the item-characteristic curves of different items), and other possible models additionally include a guessing probability term (corresponding to various vertical offsets of the item-characteristic curves of different items) or ceiling probability term (corresponding to clipping the item-characteristic curves from above). For the given dataset, it appears reasonable to estimate a Birnbaum model, whereas 3-PL or 4-PL models seem difficult, since guessing and ceiling probabilities are not easy to operationalize for the dataset, considering that there is no ground truth to the items and we found no prominent ceiling or flooring in any item.

For fitting the 2-PL model, I used the `ltm::ltm` function, and the Rasch model fitted using `ltm::rasch` served as a baseline model for comparison. ICCs and estimated item difficulty parameters

are shown in Figure 8 and 9, respectively. It can clearly be seen that item discriminativities, and with them the slopes of the ICC curves, vary considerably between items. A side-effect of two-parameter modeling is that the ICCs now cross each other, i.e., there is no clear order of difficulty between items anymore, but whether one item is more difficult than another can now be dependent on the person ability.

Comparing the two models using a Likelihood Ratio Test, I found the 2-PL model to fit the data significantly better than the Rasch model (log-LR = 1741.55, $p < .001$).




Figure 8: Item-Characteristic Curves from 2-PL model

For further model comparison, I calculated infit and outfit indices for each item and model. Infit and outfit indices are based on model residuals. While outfit (short for outlier-sensitive fit statistic) is particularly sensitive to unexpected responses in cases where item difficulty and person ability are far apart (e.g., a low-ability person unexpectedly solves several very difficult items), infit (short for information-weighted fit statistic) is particularly sensitive to unexpected responses in cases where item difficulty and person ability match (e.g., a person solves far less or far more than half of the items whose difficulty equals their ability). Both are based on the normalized residuals $Z_{iq} = \frac{D_{iq} - \mathbb{E}(D_{iq})}{\sqrt{\text{var}(D_{iq})}}$, where D_{iq} is the actual response of person i to item q , $\mathbb{E}(D_{iq}) = P(D_{iq} = 1 | \beta_q, \theta_i)$ is the conditional expectation for this person's response to the item, given the model parameters, calculated using the logistic function as shown above. $\text{var}(D_{iq})$ can be calculated as $P(D_{iq} = 1 | \beta_q, \theta_i)(1 - P(D_{iq} = 1 | \beta_q, \theta_i))$. The infit index for item q is then defined as: $\text{Infit}_q = \sum_{i=1}^n \left(\frac{\text{var}(D_{iq})}{\sum_{i=1}^n \text{var}(D_{iq})} Z_{iq}^2 \right)$, the outfit index is defined as: $\text{Outfit}_q = \sum_{i=1}^n \frac{Z_{iq}^2}{n}$. For both indices, values close to 1 indicate good fit, whereas higher values indicate under- and lower values overfitting.

The infit and outfit values for both models are, for the most part, in the acceptable (>0.7 and <1.3) range (see Figure 10). For infit, the 2-PL model consistently outperforms the Rasch model, whereas the outfit values tend to be lower overall, and both models are much closer to each other.




Figure 9: Estimated item difficulties and discriminativities based on CTT, Rasch model and 2-PL model. For both IRT models, error bars indicate standard errors.



Figure 10: Infit and outfit indices for each item in the Rasch and 2-PL models

6 Polytomous IRT model

Following the procedure outlined in Smyth (2022), I calculated a polytomous IRT model as a more data-rich alternative to the Rasch model, that is to say, using the original data with four-level responses instead of the dichotomized data.

7 Factor models

Following the IRT analysis of the dichotomized data, I went back to the original, non-dichotomized data, to investigate its factorial structure. It has been suggested that the SCS can best be described by two latent factors, one comprising items Q1, Q2, Q3, Q4, and Q10, being related to consequences of sexual behavior and compulsivity to one's lifestyle, and a second one comprising items Q5, Q6, Q7, Q8, and Q9, being related to the compulsivity of one's sexual thoughts without necessarily affecting actual behavior. Using `lavaan::cfa`, I fitted several confirmatory factor analysis (CFA) models to the data to find the latent structure that describes the data best. I specified four candidate latent structures:

The first candidate structure was a unidimensional model, i.e., the data can be explained by a single underlying latent factor.

The second candidate structure was a two-factor correlated-traits model, i.e., two latent factors were specified with item loadings as described above, and correlations between the two latent factors were allowed.

The third candidate structure was a bifactor model, i.e., two latent factors were specified with item loadings as described above, with an additional general factor on which all items load. The general factor was constrained to be orthogonal to the subfactors.

The final candidate structure was a hierarchical factor model, which specifies the two item-specific factors, and additionally has them load on a shared second-order factor.

Models were fitted with standardized latent variables, i.e., the variance of all latent factors was fixed to unit. The models were specified in `lavaan` syntax as follows:

Unidimensional model:

```
xi1 =~ Q1+Q2+Q3+Q4+Q5+Q6+Q7+Q8+Q9+Q10
```

Correlated-traits model:

```
xi1 =~ Q1+Q2+Q3+Q4+Q10
xi2 =~ Q5+Q6+Q7+Q8+Q9
```

Bifactor model:

```
G =~ Q1+Q2+Q3+Q4+Q5+Q6+Q7+Q8+Q9+Q10
xi1 =~ Q1+Q2+Q3+Q4+Q10
xi2 =~ Q5+Q6+Q7+Q8+Q9
G ~~ 0*xi1
G ~~ 0*xi2
```

Hierarchical model:

```
xi1 =~ Q1+Q2+Q3+Q4+Q10
xi2 =~ Q5+Q6+Q7+Q8+Q9
```

Table 4: Model comparison between CFA models

X	model	Df	AIC	BIC
1	hierarchical	33	85576.75	85711.44
2	bifactor	24	85167.74	85357.52
3	correlated traits	34	85574.75	85703.32
4	unidimensional	35	86552.60	86675.04

$G \sim \xi_1 + \xi_2$

The comparison of fits between the four factor models is shown in 4. Models were compared with respect to the Akaike (AIC) and Bayes-Schwarz (BIC) information criteria. Among the four candidate models, the bifactor model was clearly the preferred one among the four candidate models according to AIC as well as BIC.

The ‘winning’ bifactor model is illustrated in Figure 11. Obviously, the fact that the bifactor model is the preferred option among the four candidate models does not mean that it is necessarily a good description of the data in an absolute sense. To understand the absolute goodness-of-fit (not just compared to other models), there is a range of fit indices that we can consider. In particular, the root mean squared error of approximation (RMSEA), standardized root mean squared residual (SRMR), comparative fit index (CFI), and Tucker-Lewis index (TLI) are informative. For the bifactor model, RMSEA was at 0.073 (RMSEA < 0.08 indicating acceptable, RMSEA < 0.05 indicating good fit by convention), SRMR was at 0.028 (SRMR < 0.05 indicating good fit by convention), CFI was at 0.973 (CFI > 0.95 indicating good fit by convention), and TLI was at 0.95 (TLI > 0.95 indicating acceptable, TLI > 0.97 indicating good fit by convention). Overall, the bifactor model was, therefore, an acceptable to good fit for the SCS data.




Figure 11: Factor structure and loadings of bifactor model

An open question with respect to the factorial structure is to which subscale item Q10 should belong.

While it has been assigned to the first subfactor, its loading on the factor is low (0.18, around half as high as the second-lowest loading item, Q4). To investigate the issue, I fitted two alternative bifactor models, one (Alternative 1) where item Q10 belonged to the second subfactor, together with items Q5 - Q9, and one (Alternative 2) where item Q10 constituted its own, third subfactor. Additionally, I used the `psych::omega` function to fit a bifactor model with automatized identification of the factor structure. This procedure also returned a three-factor structure, where the first four items loaded on one subfactor, items Q5 - Q9 loaded on another subfactor, and items Q1, Q6, and Q10 loaded on a third subfactor (see Figure 12). I refitted this model in `lavaan` to facilitate comparisons.




Figure 12: Factor structure and loadings of automatically identified bifactor model

Considering all fit indices reported above (see Table 5), the original factor structure was clearly preferred over the Alternative 1 bifactor model, but the Alternative 2 bifactor model was a close competitor to the original bifactor model, that is to say, item Q10 arguably constitutes a third subfactor on its own. The automatically identified factor structure, however, was clearly preferred over the original model by all fit indices.

Looking at the content of the items (see Introduction), we can see that item Q10 is the only item that explicitly involves sex partners and difficulties to find sex partners, while item Q1 mentions relationships, item Q6 mentions the work environment, but all other items do not explicitly refer to relationships with other persons. It could, therefore, be, that responses to those items are not only influenced by sexual compulsivity, but also by a range of social and communicative abilities that might influence whether someone has difficulties finding sex partners and getting along with relationship partners and coworkers or not. Therefore, it appears plausible that those three items apparently form a third subfactor.

8 Reliability and Unidimensionality

As outlined above, the SCS is better described by a bifactor model with two or three subfactors than by a fully unidimensional model. It can, therefore, be concluded, that the scale is not unidimensional.

Table 5: Model comparison between different bifactor models

X	npar	aic	bic	cfi	tli	srmr	rmsea
Bifactor (original)	31	85167.74	85357.52	0.9733127	0.9499613	0.0278664	0.0732799
Bifactor (Alternative 1)	31	85184.35	85374.13	0.9722914	0.9480464	0.0309848	0.0746688
Bifactor (Alternative 2)	33	85168.28	85370.30	0.9734025	0.9455960	0.0285013	0.0764095
Bifactor (Automatized)	35	84924.43	85138.70	0.9885177	0.9741647	0.0188950	0.0526548

Table 6: Composite reliability scores for the bifactor models with two and three subfactors

score	G	ξ_1	ξ_2	ξ_3	model
α	0.90	0.83	0.84	NA	two-factor
ω	0.87	0.71	0.71	NA	two-factor
ω_2	0.58	0.36	0.36	NA	two-factor
ω_3	0.58	0.36	0.36	NA	two-factor
α	0.90	0.84	0.84	0.66	three-factor
ω	0.81	0.83	0.78	0.52	three-factor
ω_2	0.32	0.57	0.54	0.29	three-factor
ω_3	0.32	0.57	0.54	0.29	three-factor

To determine the composite reliability of the scales with respect to its subscales determined by confirmatory factor analysis, I calculated composite reliability using `semTools::reliability`. Composite reliability is a measure of internal consistency and is related to the question of whether item sum scores are a good representation of the underlying, latent quantity.

The function gives back Cronbach’s α as well as the ω_1 , ω_2 , and ω_3 coefficients both for the original bifactor model with two subfactors (see Figure 11) and for the better-fitting, but more complex bifactor structure with three subfactors (see Figure 12). The results are shown in Table 6.

It can be seen that the estimated reliability differs substantially between scores, and between models. Unsurprisingly, the third subfactor ξ_3 from the three-factor model has relatively low reliability, which might be related to the fact that it is based on fewer items. The remaining factors G, ξ_1 and ξ_2 have acceptable to good composite reliability in both models according to Cronbach’s α and ω , ranging from 0.71 to 0.90. Differences between scores are related to the way those coefficients are calculated. All three variants of ω are calculated as a fraction, with the squared sum of loadings of a given factor, multiplied with the factor variance (which is 1 in our models) in the numerator. What differs is the denominator. Different variants of ω are calculated with denominators based on either the residual variance, empirical covariance, or model-implied covariance. The fact that ω_3 is consistently smaller than α could point to relatively equal general factor loadings.

9 Measurement Invariance

As a last step in data analysis, I investigated measurement invariance of the SCS data. Measurement invariance means that the factorial structure is independent of other variables that are not part of the structure. In particular, I tested whether or not the three-subfactor bifactorial structure was identical across genders and age groups, similar to the analysis of DIF (see above). Following the procedure outlined in Xu (2012) and Van de Schoot, Lugtig, and Hox (2012), I compared four models

Table 7: Model comparison for measurement invariance across genders

X	DF	AIC	BIC	Chisq	Chisq_diff	DF_diff	p
configural MI	48	85154.28	85656.29	522.8565	NA	NA	NA
weak MI	65	85142.56	85540.49	545.1302	22.273646	17	0.1744230
strong MI	72	85188.67	85543.76	605.2472	60.116972	7	0.0000000
strict MI	82	85174.69	85468.55	611.2630	6.015823	10	0.8139321

Table 8: Model comparison for measurement invariance across age groups

X	DF	AIC	BIC	Chisq	Chisq_diff	DF_diff	p
configural MI	34	84853.92	85441.64	238.6647	NA	NA	NA
weak MI	52	84971.18	85448.70	391.9201	153.25539	18	0e+00
strong MI	58	85028.49	85469.28	461.2282	69.30814	6	0e+00
strict MI	62	85057.70	85474.00	498.4420	37.21382	4	2e-07

with increasingly strict measurement invariance assumptions. The *first* model assumes *configural invariance*, i.e., model parameters can vary freely in each group, and no assumption about invariance between the groups is made. The *second* model assumes *weak invariance* or *metric invariance*, i.e., equal factor loadings across groups. The *third* model assumes *strong invariance* or *scalar invariance*, where factor loadings and item intercepts are assumed to be equal across groups. The final model assumes *strict invariance*, where, additionally, item residual variances are assumed to be equal across groups. Of note, I had to use the two-subfactor bifactorial structure to test for gender measurement invariance, and the three-subfactor bifactorial structure to test for age group measurement invariance, since otherwise, there would have been always one or several models that did not converge.

For gender, the overall preferred model was the one with weak invariance (see Table 7), while and Table 8).

Looking at modification indices of the weak invariance models for age and gender, I found that the suggested modifications for the gender model included allowing factor loadings for items Q1 and Q5 on the latent factors ξ_1 and ξ_2 , respectively, to differ between groups, while for the age model, freeing factor loadings for item Q1, Q5, and Q10 between groups was suggested. This is roughly in agreement with what I found with respect to DIF (see above).

10 Theoretical Part: Key differences between IRT and CTT

10.1 Introduction

Unlike some physical quantities, many of the variables of interest in psychology, economics, and other human-centric fields, are latent, i.e., not directly observable. Researchers often try to reconstruct such latent variables by combining several observable variables. In particular, for psychological concepts such as personality traits, a person's score will often be estimated as a combination of item responses in a psychological test. Even though forerunners of psychological tests have been around for centuries, a comprehensive theory of psychological testing only emerged roughly half a century ago. Commonly, Melvin Novick is considered the first author to present a comprehensive account of Classical Test Theory (CTT) (Novick (1965)). Around the same time, a probabilistic view of psychological testing began to emerge, which are now referred to as Item Response Theory

(IRT) (Rasch (1960)). It is interesting to note that *Classical* Test Theory does, therefore, not refer to the theory itself being older, but that it rather describes the ‘classical’ way authors thought about psychological testing from the early 20th century onward, whereas probabilistic approaches became popular only later, when increasing computational capacity made them practical. In the following, I will describe some of the core ideas underlying CTT and ITT, their respective strengths and limitations, and practical applications.

10.2 Core Ideas and Terminology

A test in the sense the CTT as well as IRT use the term is designed to measure a defined *trait* or *state* of a unit (often a person). The trait/state itself is assumed to be unobservable and is captured by combining several *items* that are thought to be reflective of the trait/state. Items in the test-theoretic sense have defined response categories, which can be either dichotomous (e.g., yes/no or solved/unsolved), or ordered (e.g. I . . . do not agree / agree a little / fully agree), or multinomial (e.g. my favorite color is . . . red/blue/green). Multinomial items where the responses can not be ordered, nor dichotomized (e.g., into correct and wrong) are more involved from a theoretical point of view and will not be further discussed here. In the end, all items of a test are usually combined by a linear function, i.e., a weighted sum, to form the test *score*.

10.2.1 CTT

The traditional view of psychological tests (Classical Test Theory, CTT) conceived a given person’s total score across all items of a test as an additive combination of the person’s true score and a testing error: $X_i = \tau_i + \epsilon_i$ (Van der Linden and Hambleton (1997)). (Crocker and Algina (2008))
Reliability, Consistency, Discrimination, Difficulty

10.2.2 IRT

10.3 Strengths

10.4 Limitations

10.5 Conclusion and Application

11 Analysis code

In the following, the complete analysis code and its output are shown.

```
library(ggplot2)
library(ggthemes)
library(reshape2)
library(readxl)
library(VIM)

## Loading required package: colorspace
## Loading required package: grid
## VIM is ready to use.
## Suggestions and bug-reports can be submitted at: https://github.com/statistikat/VIM/issues
##
## Attaching package: 'VIM'
## The following object is masked from 'package:datasets':
##
##      sleep
library(mice)

##
## Attaching package: 'mice'
## The following object is masked from 'package:stats':
##
##      filter
## The following objects are masked from 'package:base':
##
##      cbind, rbind
library(dplyr)

##
## Attaching package: 'dplyr'
## The following objects are masked from 'package:stats':
##
##      filter, lag
## The following objects are masked from 'package:base':
##
##      intersect, setdiff, setequal, union
library(tidyr)

##
## Attaching package: 'tidyr'
```

```

## The following object is masked from 'package:reshape2':
##
##      smiths
library(psych)

##
## Attaching package: 'psych'

## The following objects are masked from 'package:ggplot2':
##
##      %+%, alpha
library(ggcorrplot)
library(eRm)

##
## Attaching package: 'eRm'

## The following object is masked from 'package:psych':
##
##      sim.rasch
library(ltm)

## Loading required package: MASS
##
## Attaching package: 'MASS'

## The following object is masked from 'package:dplyr':
##
##      select
## Loading required package: msm
## Loading required package: polycor
##
## Attaching package: 'polycor'

## The following object is masked from 'package:psych':
##
##      polyserial
##
## Attaching package: 'ltm'

## The following object is masked from 'package:psych':
##
##      factor.scores
library(patchwork)

##
## Attaching package: 'patchwork'

```

```

## The following object is masked from 'package:MASS':
##
##      area
library(difR)
library(semPlot)
library(lavaan)

## This is lavaan 0.6-11
## lavaan is FREE software! Please report any bugs.
##
## Attaching package: 'lavaan'

## The following object is masked from 'package:psych':
##
##      cor2cov
library(semTools)

##
## #####
## This is semTools 0.5-6
## All users of R (or SEM) are invited to submit functions or ideas for functions.
## #####
##
## Attaching package: 'semTools'

## The following objects are masked from 'package:psych':
##
##      reliability, skew
#####
#part 1: data preparation, descriptive analyses
#####
{
df = read_xlsx("SCS_data.xlsx")
SCS_vars = names(df)[1:10]
#set missing values
print(table(df$gender))
df$gender[df$gender == 3] = NA
df[df==0] = NA

print(unique(df$age))
df$age[df$age >= 100] = NA
mean(df$age,na.rm=T)
median(df$age,na.rm=T)
min(df$age,na.rm=T)
max(df$age,na.rm=T)

```

```

sprintf("%i cases are incomplete",sum(!complete.cases(df)))
sprintf("%i cases have incomplete SCS data",sum(!complete.cases(df[,SCS_vars])))

#missing data motifs
# and missing proportion per item
pdf("missingplot.pdf",width = 8, height = 4)
aggr(df[!complete.cases(df[,SCS_vars]),SCS_vars],
     numbers=TRUE, sortVars=TRUE,prop=FALSE,
     labels=SCS_vars,
     ylab=c("#Cases Missing","Pattern"))
box(which = "figure",lwd=2)
dev.off()

nmissing = rowSums(is.na(df[,SCS_vars]))
table(nmissing[nmissing!=0])
prop.table(table(nmissing[nmissing!=0]))

#missing-at-random analysis
#(check whether missing data points in each variable
#can be jointly predicted by all the other variables)
pvals = data.frame(matrix(ncol = length(SCS_vars), nrow=0))
colnames(pvals) = SCS_vars
for (var in SCS_vars){
  formula = sprintf("I(is.na(%s)) ~ .", var)
  formula0 = sprintf("I(is.na(%s)) ~ 1", var)
  m = summary(glm(formula, data=df[,1:10]))$coefficients
  pvals[var,rownames(m)[2:10]] = m[2:10,"Pr(>|t|)"]
}
min(p.adjust(unlist(pvals), method="fdr"),na.rm=T)

#-> missing at random can be assumed
#remove cases where more than two SCS variables are missing

#15 cases removed
df_clean = df[rowSums(is.na(df[,SCS_vars])) <= 2,]

#use multiple imputation for remaining data
df_clean = complete(mice(df_clean))

#descriptives
df_clean[,1:10] %>% summarise_all(list(mean=mean, median = median,
                                       min = min, max = max)) %>%
  round(1) %>%
  gather(variable, value) %>%

```

```

separate(variable, c("var", "stat"), sep = "\\_") %>%
spread(var, value) -> descriptives

#fix order of columns in descriptives table
descriptives = descriptives[,c("stat",SCS_vars)]
write.csv(descriptives, "descriptives.csv")
#re-calculate sum score
df_clean$score = rowSums(df_clean[,1:10])

#distribution plot before dichotomization
tmp = melt(
  cbind(data.frame(id=1:nrow(df_clean)),df_clean[,SCS_vars]),
  id.vars="id")
tmp2 = data.frame(table(tmp$variable,tmp$value))
colnames(tmp2) = c("item","response","Freq")
ggplot(tmp2,aes(x=item, y=Freq, fill=response))+geom_col()+theme_clean()
ggsave("distroplot.pdf",width = 4,height = 2)

#dichotomization
dich = df_clean
dich[,1:10] = data.frame(lapply(df_clean[,1:10],
                                function (x) as.numeric(x > 2)))
dich$score = rowSums(dich[,1:10])
}

```

```

##
##      0      1      2      3
##    13 2295 1053    15
## [1]  41  50  23  42  36  29  24  35  26  43  21  39  37  64  28  46  34  31  47
## [20]  22  61  16  40  33  30  56  49  51  18  20  45  32  15  27  25  59  58  19
## [39]  14  38  48  44  55 100  65  17  77  57  60  52  53  62  71  78  54  63  67
## [58]  68  72 999  85  69  70  66  84 123  73

```

```

## Warning in plot.aggr(res, ...): not enough vertical space to display frequencies
## (too many combinations)

```

```

##
## Variables sorted by number of missings:
## Variable Count
##      Q9      27
##      Q3      22
##      Q4      22
##      Q7      22
##      Q8      21
##      Q6      20
##     Q10      17
##      Q2      16
##      Q1      15

```

```
##          Q5      13
##
## iter imp variable
##  1  1  Q1  Q2  Q3  Q4  Q5  Q6  Q7  Q8  Q9  Q10  gender  age
##  1  2  Q1  Q2  Q3  Q4  Q5  Q6  Q7  Q8  Q9  Q10  gender  age
##  1  3  Q1  Q2  Q3  Q4  Q5  Q6  Q7  Q8  Q9  Q10  gender  age
##  1  4  Q1  Q2  Q3  Q4  Q5  Q6  Q7  Q8  Q9  Q10  gender  age
##  1  5  Q1  Q2  Q3  Q4  Q5  Q6  Q7  Q8  Q9  Q10  gender  age
##  2  1  Q1  Q2  Q3  Q4  Q5  Q6  Q7  Q8  Q9  Q10  gender  age
##  2  2  Q1  Q2  Q3  Q4  Q5  Q6  Q7  Q8  Q9  Q10  gender  age
##  2  3  Q1  Q2  Q3  Q4  Q5  Q6  Q7  Q8  Q9  Q10  gender  age
##  2  4  Q1  Q2  Q3  Q4  Q5  Q6  Q7  Q8  Q9  Q10  gender  age
##  2  5  Q1  Q2  Q3  Q4  Q5  Q6  Q7  Q8  Q9  Q10  gender  age
##  3  1  Q1  Q2  Q3  Q4  Q5  Q6  Q7  Q8  Q9  Q10  gender  age
##  3  2  Q1  Q2  Q3  Q4  Q5  Q6  Q7  Q8  Q9  Q10  gender  age
##  3  3  Q1  Q2  Q3  Q4  Q5  Q6  Q7  Q8  Q9  Q10  gender  age
##  3  4  Q1  Q2  Q3  Q4  Q5  Q6  Q7  Q8  Q9  Q10  gender  age
##  3  5  Q1  Q2  Q3  Q4  Q5  Q6  Q7  Q8  Q9  Q10  gender  age
##  4  1  Q1  Q2  Q3  Q4  Q5  Q6  Q7  Q8  Q9  Q10  gender  age
##  4  2  Q1  Q2  Q3  Q4  Q5  Q6  Q7  Q8  Q9  Q10  gender  age
##  4  3  Q1  Q2  Q3  Q4  Q5  Q6  Q7  Q8  Q9  Q10  gender  age
##  4  4  Q1  Q2  Q3  Q4  Q5  Q6  Q7  Q8  Q9  Q10  gender  age
##  4  5  Q1  Q2  Q3  Q4  Q5  Q6  Q7  Q8  Q9  Q10  gender  age
##  5  1  Q1  Q2  Q3  Q4  Q5  Q6  Q7  Q8  Q9  Q10  gender  age
##  5  2  Q1  Q2  Q3  Q4  Q5  Q6  Q7  Q8  Q9  Q10  gender  age
##  5  3  Q1  Q2  Q3  Q4  Q5  Q6  Q7  Q8  Q9  Q10  gender  age
##  5  4  Q1  Q2  Q3  Q4  Q5  Q6  Q7  Q8  Q9  Q10  gender  age
##  5  5  Q1  Q2  Q3  Q4  Q5  Q6  Q7  Q8  Q9  Q10  gender  age
```

```
#####
#part 2: CTT-style item analysis
#####
{

  #tetrachoric correlations
  tetra_cor = tetrachoric(dich[,SCS_vars])
  ggcorrplot(tetra_cor$rho, type = "lower", lab = TRUE)+theme_clean()
  ggsave("tetrachoric_cor_mat.pdf",width = 6, height = 6)
  #dichotomous item statistics (percent and N correct, discriminativity)
  dich.distro = rbind(as.character(round(100*unlist(lapply(dich[,SCS_vars],
                                                         mean)),1)),
                     as.character(as.integer(unlist(lapply(dich[,SCS_vars],
                                                         sum))))))
  rownames(dich.distro) = c("item easiness\n(percent in category 1)",
                           "number of cases in category 1")

  discrimination = c()
  for (item in 1:10){
```



```

    itemname = SCS_vars[item]
    discrimination[itemname] = as.character(round(biserial(
      rowSums(dich[, -item]), dich[, item]), 2))
  }

  dich.stats = rbind(dich.distro, discrimination)
  write.csv(dich.stats, "dich_stats.csv")
}

#####
#part 3: estimate and analyze Rasch model
#####

#model fitting
{
  #approach 1: eRm
  #prepare data for eRm estimation
  #(just item data in wide format)
  rasch_model_eRm = RM(dich[, SCS_vars])

  #approach 2: ltm
  #constraint fixes item discriminativity to 1
  rasch_model_ltm = rasch(dich[, SCS_vars],
    constraint = cbind(length(SCS_vars) + 1, 1))
  smr_ltm = summary(rasch_model_ltm)

  #TODO check syntax
  #approach 3: lavaan
  #modified copy from https://jonathantemplin.com/wp-content/uploads/2022/02/EPsy906\_Example05\_Binary\_IFA-IRT\_Models.nb.html
  lavaansyntax = "

  # loadings/discrimination parameters:
  SCS =~ 1*Q1 + 1*Q2 + 1*Q3 + 1*Q4 + 1*Q5 + 1*Q6 + 1*Q7 + 1*Q8 + 1*Q9 + 1*Q10

  # thresholds use the | operator and start at value 1 after t:
  Q1 | t1; Q2 | t1; Q3 | t1; Q4 | t1; Q5 | t1; Q6 | t1; Q7 | t1;
  Q8 | t1; Q9 | t1; Q10 | t1;

  # factor mean:
  SCS ~ 0;

  # factor variance:
  SCS ~~ 1*SCS

  "

```

```

rasch_model_lavaan = sem(model = lavaansyntax, data = dich[,SCS_vars],
                        ordered = SCS_vars, mimic = "Mplus",
                        estimator = "WLSMV", std.lv = TRUE,
                        parameterization = "theta")
smr_lavaan = summary(rasch_model_lavaan, fit.measures = TRUE)

convertTheta2IRT = function(lavObject){
  #modified copy from
  #https://jonathantemplin.com/wp-content/uploads/2022/02/
  #EPSY906_Example05_Binary_IFA-IRT_Models.nb.html

  if (!lavObject@Options$parameterization == "theta") {
    stop("your model is not estimated with parameterization='theta'")
  }

  output = inspect(object = lavObject, what = "est")
  if (ncol(output$lambda)>1) { stop("IRT conversion is only valid
    for one dimensional factor models.
    Your model has more than one dimension.")
  }
  a = output$lambda
  b = output$tau/output$lambda
  return(list(a = a, b=b))
}

#make ICC plot function
ICC_plot = function(difficulty, discriminativity = 1){
  if (length(discriminativity)==1){
    discriminativity = rep(discriminativity, length(difficulty))
  }
  df = data.frame(x=seq(-6,6,.01))
  for (i in 1:length(difficulty)){
    df[[SCS_vars[i]]] = logistic(x=df$x, d=difficulty[i],
                                a=discriminativity[i])
  }

  df = melt(df, id.vars = "x")
  colnames(df)[2] = "item"
  plt=ggplot(df, aes(x = x, y = value, color = item, label = item)) +
    geom_line() + theme_clean() + xlab("Person parameter") +
    ylab("P(item solved)")
  return(directlabels::direct.label(plt, "last.qp"))
}

#make ICC plots

```

```

difficulties_eRm = -rasch_model_eRm$betapar
iccplot_eRm=ICC_plot(difficulties_eRm)+ggtitle("eRm")

#lme4 difficulties are shifted by .42 from eRm difficulties, why?

difficulties_ltm = smr_ltm$coefficients[1:10,"value"]
iccplot_ltm = ICC_plot(difficulties_ltm)+ggtitle("ltm")

difficulties_lavaan = convertTheta2IRT(lavObject = rasch_model_lavaan)$b

iccplot_lavaan=ICC_plot(difficulties_lavaan)+ggtitle("lavaan")

difficulties = rbind( data.frame(model="eRm",
                                item=factor(SCS_vars),
                                difficulty=as.numeric(difficulties_eRm)),
  data.frame(model="ltm",
            item=factor(SCS_vars),
            difficulty=as.numeric(difficulties_ltm)),
  data.frame(model="lavaan",
            item=factor(SCS_vars),
            difficulty=as.numeric(difficulties_lavaan)),
  data.frame(model="CTT",
            item=factor(SCS_vars),
            difficulty=1-as.numeric(dich.distro[1,])/100))
difficulties_plot = ggplot(difficulties,aes(x=item,y=difficulty,
                                           color=model,group=model)) +
  geom_point() + geom_line() + theme_clean() + ggtitle("model comparison")+
  scale_x_discrete(breaks=SCS_vars,limits=SCS_vars)

difficulties_plot
ggsave("diffcfig.pdf",width = 4,height = 3)

#arrange plots vertically and save
iccplot_eRm|iccplot_ltm|iccplot_lavaan

ggsave("iccfig.pdf",width = 12,height = 3)

#compare fits
#select second line of output (corresponding to marginal MLE)
eRm_fit = IC(person.parameter(rasch_model_eRm))[[1]][2,]

```

```

ltm_fit = c()
ltm_fit['value'] = smr_ltm$logLik
ltm_fit['npar'] = 10
ltm_fit['AIC'] = smr_ltm$AIC
ltm_fit['BIC'] = smr_ltm$BIC
ltm_fit['cAIC'] = NA

rasch_model_fits = rbind(eRm_fit, ltm_fit)
rownames(rasch_model_fits) = c("eRm", "ltm")
colnames(rasch_model_fits)[1] = "loglik"
write.csv(rasch_model_fits, "rasch_model_fits.csv")

#calculate loss per item

predict_responses = function(item_dffc, person_params, item_discr=1){
  item_dffc = as.numeric(item_dffc)
  if (length(item_discr)==1) item_discr = rep(item_discr, length(item_dffc))
  person_params = as.numeric(person_params)
  preds = matrix(nrow=length(person_params), ncol=length(item_dffc))
  for (p in 1:length(person_params)){
    for(i in 1:length(item_dffc)){
      preds[p,i] = logistic(x=person_params[p], d = item_dffc[i], a = item_discr[i])
    }
  }
  return(preds)
}

#extract latent person abilities
person_params_eRm = person.parameter(rasch_model_eRm)$theta.table[, "Person Parameter"]
person_params_ltm = factor.scores(rasch_model_ltm, dich[, SCS_vars])[1][, "z1"]
person_params_lavaan = as.numeric(predict(rasch_model_lavaan))

#make predictions for individual persons and items
preds_ltm = predict_responses(difficulties_ltm, person_params_ltm) > .5
preds_eRm = predict_responses(difficulties_eRm, person_params_eRm) > .5
preds_lavaan = predict_responses(difficulties_lavaan, person_params_lavaan) > .5

#calculate and plot mean 0-1-loss per item
itemloss_eRm = colMeans(dich[, SCS_vars] != preds_eRm)
itemloss_ltm = colMeans(dich[, SCS_vars] != preds_ltm)
itemloss_lavaan = colMeans(dich[, SCS_vars] != preds_lavaan)
itemloss = rbind(data.frame(model="eRm", item=SCS_vars, loss=itemloss_eRm),
                 data.frame(model="ltm", item=SCS_vars, loss=itemloss_ltm),
                 data.frame(model="lavaan", item=SCS_vars, loss=itemloss_lavaan))

```

```

ggplot(itemloss,aes(x=item,y=loss,
                    color=model,group=model)) +
  geom_point() + geom_line() + theme_clean() + ggtitle("0-1-loss\nmodel comparison")+
  scale_x_discrete(breaks=SCS_vars,limits=SCS_vars)

ggsave("itemlossfig.pdf",width = 4,height = 3)

}

```

```

## lavaan 0.6-11 ended normally after 7 iterations
##
##      Estimator                      DWLS
##      Optimization method            NLMINB
##      Number of model parameters      10
##
##      Number of observations          3368
##
## Model Test User Model:
##
##              Standard      Robust
##      Test Statistic      2427.602    1525.137
##      Degrees of freedom           45         45
##      P-value (Chi-square)        0.000        0.000
##      Scaling correction factor          1.610
##      Shift parameter              17.053
##      simple second-order correction (WLSMV)
##
## Model Test Baseline Model:
##
##      Test statistic      39123.644    24321.590
##      Degrees of freedom           45         45
##      P-value              0.000        0.000
##      Scaling correction factor          1.610
##
## User Model versus Baseline Model:
##
##      Comparative Fit Index (CFI)      0.939      0.939
##      Tucker-Lewis Index (TLI)        0.939      0.939
##
##      Robust Comparative Fit Index (CFI)          NA
##      Robust Tucker-Lewis Index (TLI)            NA
##
## Root Mean Square Error of Approximation:
##
##      RMSEA      0.125      0.099
##      90 Percent confidence interval - lower      0.121      0.095
##      90 Percent confidence interval - upper      0.130      0.103
##      P-value RMSEA <= 0.05      0.000      0.000

```

```

##
## Robust RMSEA NA
## 90 Percent confidence interval - lower NA
## 90 Percent confidence interval - upper NA
##
## Standardized Root Mean Square Residual:
##
## SRMR 0.109 0.109
##
## Weighted Root Mean Square Residual:
##
## WRMR 6.644 6.644
##
## Parameter Estimates:
##
## Standard errors Robust.sem
## Information Expected
## Information saturated (h1) model Unstructured
##
## Latent Variables:
## Estimate Std.Err z-value P(>|z|)
## SCS =~
## Q1 1.000
## Q2 1.000
## Q3 1.000
## Q4 1.000
## Q5 1.000
## Q6 1.000
## Q7 1.000
## Q8 1.000
## Q9 1.000
## Q10 1.000
##
## Intercepts:
## Estimate Std.Err z-value P(>|z|)
## SCS 0.000
## .Q1 0.000
## .Q2 0.000
## .Q3 0.000
## .Q4 0.000
## .Q5 0.000
## .Q6 0.000
## .Q7 0.000
## .Q8 0.000
## .Q9 0.000
## .Q10 0.000
##
## Thresholds:

```

	Estimate	Std.Err	z-value	P(> z)
##				
## Q1 t1	0.339	0.031	10.980	0.000
## Q2 t1	0.436	0.031	14.034	0.000
## Q3 t1	0.458	0.031	14.720	0.000
## Q4 t1	0.862	0.033	26.357	0.000
## Q5 t1	0.425	0.031	13.692	0.000
## Q6 t1	-0.819	0.032	-25.219	0.000
## Q7 t1	0.435	0.031	14.000	0.000
## Q8 t1	0.311	0.031	10.087	0.000
## Q9 t1	0.022	0.031	0.724	0.469
## Q10 t1	-0.027	0.031	-0.896	0.370

##

Variances:

	Estimate	Std.Err	z-value	P(> z)
##				
## SCS	1.000			
## .Q1	1.000			
## .Q2	1.000			
## .Q3	1.000			
## .Q4	1.000			
## .Q5	1.000			
## .Q6	1.000			
## .Q7	1.000			
## .Q8	1.000			
## .Q9	1.000			
## .Q10	1.000			

##

Scales y*:

	Estimate	Std.Err	z-value	P(> z)
##				
## Q1	0.707			
## Q2	0.707			
## Q3	0.707			
## Q4	0.707			
## Q5	0.707			
## Q6	0.707			
## Q7	0.707			
## Q8	0.707			
## Q9	0.707			
## Q10	0.707			

#DIF

```
{
  data_dif_age = dich[,SCS_vars]
  data_dif_age$age = dich$age > median(dich$age)
  dif_ageL = difLord(data_dif_age,"age",FALSE,"1PL")
  dif_ageR = difRaju(data_dif_age,"age",FALSE, "1PL")

  data_dif_gender= dich[,c(SCS_vars,"gender")]
}
```

```

dif_genderL = difLord(data_dif_gender,"gender",1, "1PL")
dif_genderR = difRaju(data_dif_gender,"gender",1, "1PL")

difstats=data.frame(
  p=-log10(p.adjust(
    c(dif_genderL$p.value,dif_ageL$p.value),method="fdr")),
  item = c(dif_genderL$names,dif_ageL$name),
  groups = c(rep("gender",10),rep("age",10))
)

ggplot(difstats, aes(x=item, y=p, group=groups,col=groups)) +
  geom_point() + geom_line() + theme_clean() + ylab("-log10(p)") +
  geom_hline(yintercept=-log10(.05))+scale_x_discrete(breaks=SCS_vars,
                                                    limits=SCS_vars)

ggsave("DIF_pvals.pdf",width = 4,height = 3)
}

#alternative model: 2PL
{

  #fit 1PL and 2PL, compare fit
  twoPL_model = ltm(dich[,SCS_vars] ~ z1, IRT.param = TRUE)
  difficulties_2PL = coef(twoPL_model)[,"Dffclt"]
  discriminativities_2PL = coef(twoPL_model)[,"Dscrmn"]
  ICC_2PL = ICC_plot(difficulty = difficulties_2PL,
                     discriminativity = discriminativities_2PL)

  Rasch_vs_twoPL_comparison = anova(rasch_model_ltm, twoPL_model)

  difficulties_1vs2PL = rbind( data.frame(model="Rasch (1-PL)",
                                         item=factor(SCS_vars),
                                         difficulty=as.numeric(difficulties_ltm)),
                              data.frame(model="Birnbbaum (2-PL)",
                                         item=factor(SCS_vars),
                                         difficulty=as.numeric(difficulties_2PL)),
                              data.frame(model="CTT",
                                         item=factor(SCS_vars),
                                         difficulty=as.numeric(dich.distro[1,]/100))
  ggplot(difficulties_1vs2PL,aes(x=item,y=difficulty,
                                color=model,group=model)) +
    geom_point() + geom_line() + theme_clean() + ggtitle("model comparison")+
    scale_x_discrete(breaks=SCS_vars,limits=SCS_vars)
  ggsave("difficulties_plot_2PL.pdf",width = 4,height = 3)
}

```



```

#calculate item-wise infit and outfit

get_outfit = function(ltm_model){
  X=ltm_model$X
  personscores = factor.scores(ltm_model,X)[[1]][,"z1"]
  dffc = coef(ltm_model)[,"Dffc1t"]
  discr = coef(ltm_model)[,"Dscrmn"]
  expected = predict_responses(dffc, personscores, discr)
  var_X = expected * (1-expected)
  Z_ij = (X-expected)/sqrt(var_X)
  chisq = colSums(Z_ij**2)
  #divide chisq by n
  return(chisq/nrow(ltm_model$X))
}

get_infit = function(ltm_model){
  X=ltm_model$X
  personscores = factor.scores(ltm_model,X)[[1]][,"z1"]
  dffc = coef(ltm_model)[,"Dffc1t"]
  discr = coef(ltm_model)[,"Dscrmn"]
  expected = predict_responses(dffc, personscores, discr)
  var_X = expected * (1-expected)
  Z_ij = (X-expected)/sqrt(var_X)
  infit = c()
  for (i in 1:length(dffc)){
    infit[i] = sum((var_X[,i] * (Z_ij[,i]**2))/sum(var_X[,i]))}
  return(infit)
}

outfit_rasch = get_outfit(rasch_model_ltm)
outfit_2PL = get_outfit(twoPL_model)
infit_rasch = get_infit(rasch_model_ltm)
infit_2PL = get_infit(twoPL_model)

inoutfit = rbind(data.frame(model="Rasch",fit="outfit",
                             item=names(outfit_rasch), value=outfit_rasch),
                 data.frame(model="2-PL",fit="outfit",
                             item=names(outfit_2PL), value=outfit_2PL),
                 data.frame(model="Rasch",fit="infit",
                             item=names(outfit_rasch),value=infit_rasch),
                 data.frame(model="2-PL",fit="infit",
                             item=names(outfit_2PL),value=infit_2PL))

ggplot(inoutfit,aes(x=item,y=value,
                    color=model,group=model)) + facet_wrap(~fit)+
  geom_point() + geom_line() + theme_clean() + ggtitle("model comparison")+
  scale_x_discrete(breaks=SCS_vars,limits=SCS_vars)

```

```


ggsave("inoutfit_plot_2PL.pdf",width = 8,height = 3)

}


#polytomous IRT model
{
  grm_model = grm(df_clean[,SCS_vars],constrained=T)
  plot(grm_model, ask=FALSE)
}

```


Item Response Category Characteristic Curves – Item: Q1




Item Response Category Characteristic Curves – Item: Q2




Item Response Category Characteristic Curves – Item: Q3




Item Response Category Characteristic Curves – Item: Q4




Item Response Category Characteristic Curves – Item: Q5




Item Response Category Characteristic Curves – Item: Q6




Item Response Category Characteristic Curves – Item: Q7




Item Response Category Characteristic Curves – Item: Q8



Item Response Category Characteristic Curves – Item: Q9



Item Response Category Characteristic Curves – Item: Q10



```
#####
#part 4: factorial structure of the data
#####

#factor analyses
{

  covdat = cov(df_clean[,SCS_vars])
  N=nrow(df_clean)

  #unidimensional model
  unidimensional_model <- '
xi1 =~ Q1+Q2+Q3+Q4+Q5+Q6+Q7+Q8+Q9+Q10

  '

  unidimensional_cfa <- cfa(unidimensional_model,
                           sample.cov=covdat,
                           sample.nobs=N,
                           std.lv=T)
```

```

#correlated traits
correlated_traits_model <- '
xi1 =~ Q1+Q2+Q3+Q4+Q10
xi2 =~ Q5+Q6+Q7+Q8+Q9
xi1 ~~ xi2
'

correlated_traits_cfa <- cfa(correlated_traits_model,
                             sample.cov=covdat,
                             sample.nobs=N,
                             std.lv=T)

#bifactor model (general factor and two item-specific factors)
bifactor_model <- '
G =~ Q1+Q2+Q3+Q4+Q5+Q6+Q7+Q8+Q9+Q10
xi1 =~ Q1+Q2+Q3+Q4+Q10
xi2 =~ Q5+Q6+Q7+Q8+Q9
G ~~ 0*xi1
G ~~ 0*xi2
'

bifactor_cfa <- cfa(bifactor_model,
                    sample.cov=covdat,
                    sample.nobs=N,
                    std.lv=T)

#hierarchical model

hierarchical_model <- '
xi1 =~ Q1+Q2+Q3+Q4+Q10
xi2 =~ Q5+Q6+Q7+Q8+Q9
G =~ xi1+xi2
'

hierarchical_cfa <- cfa(hierarchical_model,
                        sample.cov=covdat,
                        sample.nobs=N,
                        std.lv=T)

#
smr_hierarchical = summary(hierarchical_cfa, fit=T)$FIT
smr_bifactor = summary(bifactor_cfa, fit=T)$FIT

```



```

smr_correlated_traits = summary(correlated_traits_cfa, fit=T)$FIT
smr_unidimensional = summary(unidimensional_cfa, fit=T)$FIT

cfaaov_df = data.frame(model=c("hierarchical","bifactor","correlated traits",
                                "unidimensional"),
                        Df = c(smr_hierarchical["df"],
                                smr_bifactor["df"],
                                smr_correlated_traits["df"],
                                smr_unidimensional["df"]),
                        AIC = c(smr_hierarchical["aic"],
                                smr_bifactor["aic"],
                                smr_correlated_traits["aic"],
                                smr_unidimensional["aic"]),
                        BIC = c(smr_hierarchical["bic"],
                                smr_bifactor["bic"],
                                smr_correlated_traits["bic"],
                                smr_unidimensional["bic"]))

write.csv(cfaaov_df,"cfaaov_df.csv")

pdf("semplot_bifactor.pdf", width = 8,height = 4)
semPaths(bifactor_cfa, "std")
dev.off()

#fit alternative bifactor model (item Q10 belongs to subscale 2)
bifactor_model_2 <- '
G =~ Q1+Q2+Q3+Q4+Q5+Q6+Q7+Q8+Q9+Q10
xi1 =~ Q1+Q2+Q3+Q4
xi2 =~ Q5+Q6+Q7+Q8+Q9+Q10
G ~~ 0*xi1
G ~~ 0*xi2
'

bifactor_cfa_2 <- cfa(bifactor_model_2,
                      sample.cov=covdat,
                      sample.nobs=N,
                      std.lv=T)

#fit alternative bifactor model (item Q10 is its own subscale)
#-> does not converge

```

```

bifactor_model_3 <- '
G =~ Q1+Q2+Q3+Q4+Q5+Q6+Q7+Q8+Q9+Q10
xi1 =~ Q1+Q2+Q3+Q4
xi2 =~ Q5+Q6+Q7+Q8+Q9
xi3 =~ Q10
G ~~ 0*xi1
G ~~ 0*xi2
G ~~ 0*xi3
'

bifactor_cfa_3 <- cfa(bifactor_model_3,
                      sample.cov=covdat,
                      sample.nobs=N,
                      std.lv=T)

bifactor_model_4 <- '
G =~ Q1+Q2+Q3+Q4+Q5+Q6+Q7+Q8+Q9+Q10
xi1 =~ Q1+Q2+Q3+Q4
xi2 =~ Q5+Q6+Q7+Q8+Q9
xi3 =~ Q10+Q6+Q1
G ~~ 0*xi1
G ~~ 0*xi2
G ~~ 0*xi3
'

bifactor_cfa_4<- cfa(bifactor_model_4,
                    sample.cov=covdat,
                    sample.nobs=N,
                    std.lv=T)

pdf("semplot_bifactor_automatic.pdf", width = 8,height = 4)
semPaths(bifactor_cfa_4, "std")
dev.off()

smr_bif1=summary(bifactor_cfa,fit=T)$FIT
smr_bif2=summary(bifactor_cfa_2,fit=T)$FIT
smr_bif3=summary(bifactor_cfa_3,fit=T)$FIT
smr_bif4=summary(bifactor_cfa_4,fit=T)$FIT

bif_comparison=rbind(smr_bif1,smr_bif2,smr_bif3,smr_bif4)[,c("npar","aic",
                                                             "bic","cfi",
                                                             "tli","srmr",
                                                             "rmsea")]

```

```

rownames(bif_comparison) = c("Bifactor (original)",
                             "Bifactor (Alternative 1)",
                             "Bifactor (Alternative 2)",
                             "Bifactor (Automatized)")
write.csv(bif_comparison, "bifactor_comparison.csv")

}

```

```

## Warning in lav_model_vcov(lavmodel = lavmodel, lavsamplestats = lavsamplestats, : lavaan WAF
##      Could not compute standard errors! The information matrix could
##      not be inverted. This may be a symptom that the model is not
##      identified.

## lavaan 0.6-11 ended normally after 31 iterations
##
##      Estimator                      ML
##      Optimization method          NLMINB
##      Number of model parameters    22
##
##      Number of observations        3368
##
## Model Test User Model:
##
##      Test statistic                881.568
##      Degrees of freedom            33
##      P-value (Chi-square)          0.000
##
## Model Test Baseline Model:
##
##      Test statistic                16315.763
##      Degrees of freedom            45
##      P-value                        0.000
##
## User Model versus Baseline Model:
##
##      Comparative Fit Index (CFI)    0.948
##      Tucker-Lewis Index (TLI)      0.929
##
## Loglikelihood and Information Criteria:
##
##      Loglikelihood user model (H0)    -42759.769
##      Loglikelihood unrestricted model (H1) -42318.985
##
##      Akaike (AIC)                    85563.538
##      Bayesian (BIC)                   85698.224
##      Sample-size adjusted Bayesian (BIC) 85628.320
##
## Root Mean Square Error of Approximation:

```

```

##
## RMSEA 0.087
## 90 Percent confidence interval - lower 0.082
## 90 Percent confidence interval - upper 0.092
## P-value RMSEA <= 0.05 0.000
##
## Standardized Root Mean Square Residual:
##
## SRMR 0.041
##
## Parameter Estimates:
##
## Standard errors Standard
## Information Expected
## Information saturated (h1) model Structured
##
## Latent Variables:
## Estimate Std.Err z-value P(>|z|)
## xi1 =~
## Q1 0.315 NA
## Q2 0.363 NA
## Q3 0.380 NA
## Q4 0.306 NA
## Q10 0.248 NA
## xi2 =~
## Q5 0.335 NA
## Q6 0.224 NA
## Q7 0.366 NA
## Q8 0.394 NA
## Q9 0.312 NA
## G =~
## xi1 2.161 NA
## xi2 2.158 NA
##
## Variances:
## Estimate Std.Err z-value P(>|z|)
## .Q1 0.613 NA
## .Q2 0.398 NA
## .Q3 0.347 NA
## .Q4 0.542 NA
## .Q10 1.059 NA
## .Q5 0.610 NA
## .Q6 0.649 NA
## .Q7 0.380 NA
## .Q8 0.300 NA
## .Q9 0.783 NA
## .xi1 1.000
## .xi2 1.000

```

```

##      G              1.000
##
## lavaan 0.6-11 ended normally after 36 iterations
##
##      Estimator              ML
##      Optimization method    NLMINB
##      Number of model parameters    31
##
##      Number of observations    3368
##
## Model Test User Model:
##
##      Test statistic          457.923
##      Degrees of freedom      24
##      P-value (Chi-square)    0.000
##
## Model Test Baseline Model:
##
##      Test statistic          16315.763
##      Degrees of freedom      45
##      P-value                  0.000
##
## User Model versus Baseline Model:
##
##      Comparative Fit Index (CFI)    0.973
##      Tucker-Lewis Index (TLI)      0.950
##
## Loglikelihood and Information Criteria:
##
##      Loglikelihood user model (H0)    -42547.947
##      Loglikelihood unrestricted model (H1) -42318.985
##
##      Akaike (AIC)                  85157.894
##      Bayesian (BIC)                 85347.678
##      Sample-size adjusted Bayesian (BIC) 85249.177
##
## Root Mean Square Error of Approximation:
##
##      RMSEA                          0.073
##      90 Percent confidence interval - lower    0.067
##      90 Percent confidence interval - upper    0.079
##      P-value RMSEA <= 0.05                0.000
##
## Standardized Root Mean Square Residual:
##
##      SRMR                          0.028
##
## Parameter Estimates:

```

```

##
## Standard errors
## Information
## Information saturated (h1) model
##
## Latent Variables:
## Estimate Std.Err z-value P(>|z|)
## G =~
## Q1 0.552 0.037 14.940 0.000
## Q2 0.424 0.044 9.578 0.000
## Q3 0.751 0.033 23.095 0.000
## Q4 0.646 0.026 24.975 0.000
## Q5 0.707 0.024 28.962 0.000
## Q6 0.544 0.020 27.752 0.000
## Q7 0.604 0.031 19.189 0.000
## Q8 0.624 0.035 17.789 0.000
## Q9 0.509 0.031 16.359 0.000
## Q10 0.630 0.023 27.127 0.000
## xi1 =~
## Q1 0.504 0.042 11.853 0.000
## Q2 0.969 0.036 26.849 0.000
## Q3 0.481 0.046 10.460 0.000
## Q4 0.338 0.038 8.926 0.000
## Q10 0.177 0.034 5.219 0.000
## xi2 =~
## Q5 0.403 0.033 12.196 0.000
## Q6 0.187 0.028 6.785 0.000
## Q7 0.626 0.030 21.008 0.000
## Q8 0.728 0.030 24.409 0.000
## Q9 0.545 0.030 18.344 0.000
##
## Covariances:
## Estimate Std.Err z-value P(>|z|)
## G ~~
## xi1 0.000
## xi2 0.000
## xi1 ~~
## xi2 0.592 0.039 15.009 0.000
##
## Variances:
## Estimate Std.Err z-value P(>|z|)
## .Q1 0.618 0.017 35.612 0.000
## .Q2 0.029 0.068 0.429 0.668
## .Q3 0.368 0.013 28.113 0.000
## .Q4 0.541 0.016 33.910 0.000
## .Q5 0.583 0.017 34.146 0.000
## .Q6 0.601 0.017 34.903 0.000
## .Q7 0.381 0.013 29.836 0.000

```

```

##      .Q8                0.259    0.013    19.423    0.000
##      .Q9                0.779    0.021    37.237    0.000
##      .Q10               0.980    0.027    35.945    0.000
##      G                  1.000
##      xi1                1.000
##      xi2                1.000
##
## lavaan 0.6-11 ended normally after 20 iterations
##
##      Estimator                      ML
##      Optimization method          NLMINB
##      Number of model parameters    21
##
##      Number of observations        3368
##
## Model Test User Model:
##
##      Test statistic                881.568
##      Degrees of freedom             34
##      P-value (Chi-square)          0.000
##
## Model Test Baseline Model:
##
##      Test statistic                16315.763
##      Degrees of freedom             45
##      P-value                       0.000
##
## User Model versus Baseline Model:
##
##      Comparative Fit Index (CFI)    0.948
##      Tucker-Lewis Index (TLI)       0.931
##
## Loglikelihood and Information Criteria:
##
##      Loglikelihood user model (H0)   -42759.769
##      Loglikelihood unrestricted model (H1) -42318.985
##
##      Akaike (AIC)                   85561.538
##      Bayesian (BIC)                  85690.102
##      Sample-size adjusted Bayesian (BIC) 85623.375
##
## Root Mean Square Error of Approximation:
##
##      RMSEA                          0.086
##      90 Percent confidence interval - lower 0.081
##      90 Percent confidence interval - upper 0.091
##      P-value RMSEA <= 0.05           0.000
##

```

```

## Standardized Root Mean Square Residual:
##
##   SRMR                                0.041
##
## Parameter Estimates:
##
##   Standard errors                    Standard
##   Information                        Expected
##   Information saturated (h1) model    Structured
##
## Latent Variables:
##           Estimate   Std.Err   z-value   P(>|z|)
##   xi1 =~
##     Q1             0.750     0.017    43.760     0.000
##     Q2             0.865     0.016    54.405     0.000
##     Q3             0.904     0.016    57.392     0.000
##     Q4             0.729     0.016    44.757     0.000
##     Q10            0.590     0.020    29.185     0.000
##   xi2 =~
##     Q5             0.797     0.017    45.862     0.000
##     Q6             0.532     0.016    33.006     0.000
##     Q7             0.871     0.016    55.416     0.000
##     Q8             0.938     0.016    60.383     0.000
##     Q9             0.743     0.019    39.955     0.000
##
## Covariances:
##           Estimate   Std.Err   z-value   P(>|z|)
##   xi1 ~~
##     xi2            0.823     0.008    97.148     0.000
##
## Variances:
##           Estimate   Std.Err   z-value   P(>|z|)
##     .Q1            0.613     0.017    36.199     0.000
##     .Q2            0.398     0.013    30.925     0.000
##     .Q3            0.347     0.012    28.418     0.000
##     .Q4            0.542     0.015    35.864     0.000
##     .Q10           1.059     0.027    39.325     0.000
##     .Q5            0.610     0.017    35.904     0.000
##     .Q6            0.649     0.017    38.922     0.000
##     .Q7            0.380     0.012    30.993     0.000
##     .Q8            0.300     0.011    26.375     0.000
##     .Q9            0.783     0.021    37.587     0.000
##     xi1            1.000
##     xi2            1.000
##
## lavaan 0.6-11 ended normally after 16 iterations
##
##   Estimator                                ML

```



```

## Optimization method NLMINB
## Number of model parameters 20
##
## Number of observations 3368
##
## Model Test User Model:
##
## Test statistic 1860.670
## Degrees of freedom 35
## P-value (Chi-square) 0.000
##
## Model Test Baseline Model:
##
## Test statistic 16315.763
## Degrees of freedom 45
## P-value 0.000
##
## User Model versus Baseline Model:
##
## Comparative Fit Index (CFI) 0.888
## Tucker-Lewis Index (TLI) 0.856
##
## Loglikelihood and Information Criteria:
##
## Loglikelihood user model (H0) -43249.320
## Loglikelihood unrestricted model (H1) -42318.985
##
## Akaike (AIC) 86538.641
## Bayesian (BIC) 86661.082
## Sample-size adjusted Bayesian (BIC) 86597.533
##
## Root Mean Square Error of Approximation:
##
## RMSEA 0.124
## 90 Percent confidence interval - lower 0.120
## 90 Percent confidence interval - upper 0.129
## P-value RMSEA <= 0.05 0.000
##
## Standardized Root Mean Square Residual:
##
## SRMR 0.052
##
## Parameter Estimates:
##
## Standard errors Standard
## Information Expected
## Information saturated (h1) model Structured
##

```

```

## Latent Variables:
##           Estimate Std.Err z-value P(>|z|)
##   xi1 =~
##     Q1           0.705   0.017  40.855   0.000
##     Q2           0.808   0.016  49.971   0.000
##     Q3           0.848   0.016  53.034   0.000
##     Q4           0.694   0.016  42.503   0.000
##     Q5           0.779   0.017  44.860   0.000
##     Q6           0.530   0.016  33.130   0.000
##     Q7           0.821   0.016  51.377   0.000
##     Q8           0.876   0.016  55.077   0.000
##     Q9           0.715   0.019  38.349   0.000
##     Q10          0.599   0.020  30.087   0.000
##
## Variances:
##           Estimate Std.Err z-value P(>|z|)
##     .Q1           0.679   0.018  38.060   0.000
##     .Q2           0.493   0.014  35.632   0.000
##     .Q3           0.444   0.013  34.408   0.000
##     .Q4           0.591   0.016  37.718   0.000
##     .Q5           0.638   0.017  37.166   0.000
##     .Q6           0.652   0.017  39.290   0.000
##     .Q7           0.464   0.013  35.105   0.000
##     .Q8           0.412   0.012  33.415   0.000
##     .Q9           0.824   0.021  38.520   0.000
##     .Q10          1.048   0.026  39.647   0.000
##     xi1           1.000
##
## Warning in lav_object_post_check(object): lavaan WARNING: some estimated ov
## variances are negative
##
## Warning in lav_model_vcov(lavmodel = lavmodel, lavsamplestats = lavsamplestats, : lavaan WAF
## Could not compute standard errors! The information matrix could
## not be inverted. This may be a symptom that the model is not
## identified.
##
## Warning in lav_object_post_check(object): lavaan WARNING: some estimated ov
## variances are negative
##
## lavaan 0.6-11 ended normally after 36 iterations
##
## Estimator ML
## Optimization method NLMINB
## Number of model parameters 31
##
## Number of observations 3368
##
## Model Test User Model:
##
## Test statistic 457.923

```

```

## Degrees of freedom                24
## P-value (Chi-square)              0.000
##
## Model Test Baseline Model:
##
## Test statistic                    16315.763
## Degrees of freedom                45
## P-value                          0.000
##
## User Model versus Baseline Model:
##
## Comparative Fit Index (CFI)        0.973
## Tucker-Lewis Index (TLI)          0.950
##
## Loglikelihood and Information Criteria:
##
## Loglikelihood user model (H0)      -42547.947
## Loglikelihood unrestricted model (H1) -42318.985
##
## Akaike (AIC)                      85157.894
## Bayesian (BIC)                    85347.678
## Sample-size adjusted Bayesian (BIC) 85249.177
##
## Root Mean Square Error of Approximation:
##
## RMSEA                            0.073
## 90 Percent confidence interval - lower 0.067
## 90 Percent confidence interval - upper 0.079
## P-value RMSEA <= 0.05              0.000
##
## Standardized Root Mean Square Residual:
##
## SRMR                            0.028
##
## Parameter Estimates:
##
## Standard errors                    Standard
## Information                        Expected
## Information saturated (h1) model    Structured
##
## Latent Variables:
##
## Estimate Std.Err z-value P(>|z|)
## G =~
## Q1        0.552   0.037  14.940   0.000
## Q2        0.424   0.044   9.578   0.000
## Q3        0.751   0.033  23.095   0.000
## Q4        0.646   0.026  24.975   0.000
## Q5        0.707   0.024  28.962   0.000

```

```

##      Q6      0.544    0.020   27.752    0.000
##      Q7      0.604    0.031   19.189    0.000
##      Q8      0.624    0.035   17.789    0.000
##      Q9      0.509    0.031   16.359    0.000
##      Q10     0.630    0.023   27.127    0.000
##      xi1 =~
##      Q1      0.504    0.042   11.853    0.000
##      Q2      0.969    0.036   26.849    0.000
##      Q3      0.481    0.046   10.460    0.000
##      Q4      0.338    0.038    8.926    0.000
##      Q10     0.177    0.034    5.219    0.000
##      xi2 =~
##      Q5      0.403    0.033   12.196    0.000
##      Q6      0.187    0.028    6.785    0.000
##      Q7      0.626    0.030   21.008    0.000
##      Q8      0.728    0.030   24.409    0.000
##      Q9      0.545    0.030   18.344    0.000
##
## Covariances:
##      Estimate Std.Err z-value P(>|z|)
##      G ~~
##      xi1      0.000
##      xi2      0.000
##      xi1 ~~
##      xi2      0.592    0.039   15.009    0.000
##
## Variances:
##      Estimate Std.Err z-value P(>|z|)
##      .Q1      0.618    0.017   35.612    0.000
##      .Q2      0.029    0.068    0.429    0.668
##      .Q3      0.368    0.013   28.113    0.000
##      .Q4      0.541    0.016   33.910    0.000
##      .Q5      0.583    0.017   34.146    0.000
##      .Q6      0.601    0.017   34.903    0.000
##      .Q7      0.381    0.013   29.836    0.000
##      .Q8      0.259    0.013   19.423    0.000
##      .Q9      0.779    0.021   37.237    0.000
##      .Q10     0.980    0.027   35.945    0.000
##      G      1.000
##      xi1     1.000
##      xi2     1.000
##
## lavaan 0.6-11 ended normally after 2995 iterations
##
##      Estimator      ML
##      Optimization method      NLMINB
##      Number of model parameters      31
##

```

```

##      Number of observations                3368
##
## Model Test User Model:
##
##      Test statistic                473.665
##      Degrees of freedom                24
##      P-value (Chi-square)            0.000
##
## Model Test Baseline Model:
##
##      Test statistic                16315.763
##      Degrees of freedom                45
##      P-value                0.000
##
## User Model versus Baseline Model:
##
##      Comparative Fit Index (CFI)                0.972
##      Tucker-Lewis Index (TLI)                0.948
##
## Loglikelihood and Information Criteria:
##
##      Loglikelihood user model (H0)                -42555.818
##      Loglikelihood unrestricted model (H1)        -42318.985
##
##      Akaike (AIC)                85173.636
##      Bayesian (BIC)                85363.420
##      Sample-size adjusted Bayesian (BIC)        85264.919
##
## Root Mean Square Error of Approximation:
##
##      RMSEA                0.075
##      90 Percent confidence interval - lower        0.069
##      90 Percent confidence interval - upper        0.081
##      P-value RMSEA <= 0.05                0.000
##
## Standardized Root Mean Square Residual:
##
##      SRMR                0.031
##
## Parameter Estimates:
##
##      Standard errors                Standard
##      Information                Expected
##      Information saturated (h1) model        Structured
##
## Latent Variables:
##      Estimate Std.Err z-value P(>|z|)
##      G =~

```

```

##      Q1                0.709    0.032    22.375    0.000
##      Q2                0.600    0.031    19.222    0.000
##      Q3                0.906    0.023    39.581    0.000
##      Q4                0.747    0.018    42.135    0.000
##      Q5                0.719    0.019    37.865    0.000
##      Q6                0.514    0.017    31.097    0.000
##      Q7                0.702    0.022    31.712    0.000
##      Q8                0.745    0.024    30.485    0.000
##      Q9                0.585    0.024    24.419    0.000
##      Q10               0.599    0.021    28.118    0.000
##      xi1 =~
##      Q1                0.042    0.580    0.072    0.943
##      Q2                7.417   102.506    0.072    0.942
##      Q3                0.032    0.444    0.072    0.943
##      Q4                0.020    0.275    0.072    0.943
##      xi2 =~
##      Q5                0.339    0.023    14.464    0.000
##      Q6                0.169    0.020    8.531    0.000
##      Q7                0.512    0.025    20.682    0.000
##      Q8                0.597    0.027    21.877    0.000
##      Q9                0.470    0.026    17.878    0.000
##      Q10               0.090    0.024    3.691    0.000
##
## Covariances:
##      Estimate Std.Err z-value P(>|z|)
##      G ~~
##      xi1      0.000
##      xi2      0.000
##      xi1 ~~
##      xi2      0.050    0.696    0.072    0.943
##
## Variances:
##      Estimate Std.Err z-value P(>|z|)
##      .Q1      0.673    0.022    30.757    0.000
##      .Q2     -54.224  1520.571   -0.036    0.972
##      .Q3      0.342    0.014    23.661    0.000
##      .Q4      0.515    0.015    33.251    0.000
##      .Q5      0.613    0.017    36.728    0.000
##      .Q6      0.640    0.016    38.876    0.000
##      .Q7      0.383    0.013    29.834    0.000
##      .Q8      0.267    0.013    20.706    0.000
##      .Q9      0.772    0.021    36.838    0.000
##      .Q10     1.040    0.028    37.806    0.000
##      G        1.000
##      xi1      1.000
##      xi2      1.000
##
## lavaan 0.6-11 ended normally after 42 iterations

```

```

##
## Estimator ML
## Optimization method NLMINB
## Number of model parameters 33
##
## Number of observations 3368
##
## Model Test User Model:
##
## Test statistic 454.085
## Degrees of freedom 22
## P-value (Chi-square) 0.000
##
## Model Test Baseline Model:
##
## Test statistic 16315.763
## Degrees of freedom 45
## P-value 0.000
##
## User Model versus Baseline Model:
##
## Comparative Fit Index (CFI) 0.973
## Tucker-Lewis Index (TLI) 0.946
##
## Loglikelihood and Information Criteria:
##
## Loglikelihood user model (H0) -42546.028
## Loglikelihood unrestricted model (H1) -42318.985
##
## Akaike (AIC) 85158.056
## Bayesian (BIC) 85360.084
## Sample-size adjusted Bayesian (BIC) 85255.228
##
## Root Mean Square Error of Approximation:
##
## RMSEA 0.076
## 90 Percent confidence interval - lower 0.070
## 90 Percent confidence interval - upper 0.083
## P-value RMSEA <= 0.05 0.000
##
## Standardized Root Mean Square Residual:
##
## SRMR 0.029
##
## Parameter Estimates:
##
## Standard errors Standard
## Information Expected

```

```

## Information saturated (h1) model          Structured
##
## Latent Variables:
##      Estimate   Std.Err   z-value   P(>|z|)
##      G =~
##      Q1          0.561      NA
##      Q2          0.411      NA
##      Q3          0.780      NA
##      Q4          0.671      NA
##      Q5          0.689      NA
##      Q6          0.519      NA
##      Q7          0.603      NA
##      Q8          0.626      NA
##      Q9          0.498      NA
##      Q10         0.601      NA
##      xi1 =~
##      Q1          0.482      NA
##      Q2          1.027      NA
##      Q3          0.451      NA
##      Q4          0.314      NA
##      xi2 =~
##      Q5          0.424      NA
##      Q6          0.216      NA
##      Q7          0.628      NA
##      Q8          0.722      NA
##      Q9          0.558      NA
##      xi3 =~
##      Q10         0.702      NA
##
## Covariances:
##      Estimate   Std.Err   z-value   P(>|z|)
##      G ~~
##      xi1          0.000
##      xi2          0.000
##      xi3          0.000
##      xi1 ~~
##      xi2          0.572      NA
##      xi3          0.264      NA
##      xi2 ~~
##      xi3          0.224      NA
##
## Variances:
##      Estimate   Std.Err   z-value   P(>|z|)
##      .Q1          0.630      NA
##      .Q2         -0.077      NA
##      .Q3          0.352      NA
##      .Q4          0.525      NA
##      .Q5          0.591      NA

```



```

##      .Q6                0.616        NA
##      .Q7                0.381        NA
##      .Q8                0.267        NA
##      .Q9                0.775        NA
##      .Q10              0.554        NA
##      G                  1.000
##      xi1                1.000
##      xi2                1.000
##      xi3                1.000
##
## lavaan 0.6-11 ended normally after 50 iterations
##
##      Estimator                      ML
##      Optimization method          NLMINB
##      Number of model parameters    35
##
##      Number of observations        3368
##
## Model Test User Model:
##
##      Test statistic                206.078
##      Degrees of freedom            20
##      P-value (Chi-square)          0.000
##
## Model Test Baseline Model:
##
##      Test statistic                16315.763
##      Degrees of freedom            45
##      P-value                      0.000
##
## User Model versus Baseline Model:
##
##      Comparative Fit Index (CFI)    0.989
##      Tucker-Lewis Index (TLI)      0.974
##
## Loglikelihood and Information Criteria:
##
##      Loglikelihood user model (H0)  -42422.024
##      Loglikelihood unrestricted model (H1) -42318.985
##
##      Akaike (AIC)                  84914.049
##      Bayesian (BIC)                 85128.321
##      Sample-size adjusted Bayesian (BIC) 85017.110
##
## Root Mean Square Error of Approximation:
##
##      RMSEA                        0.053
##      90 Percent confidence interval - lower 0.046

```

```

## 90 Percent confidence interval - upper      0.059
## P-value RMSEA <= 0.05                      0.248
##
## Standardized Root Mean Square Residual:
##
## SRMR                                         0.019
##
## Parameter Estimates:
##
## Standard errors                          Standard
## Information                              Expected
## Information saturated (h1) model          Structured
##
## Latent Variables:
##      Estimate Std.Err z-value P(>|z|)
## G =~
## Q1          0.322   0.078   4.111   0.000
## Q2          0.196   0.090   2.179   0.029
## Q3          0.675   0.067  10.054   0.000
## Q4          0.595   0.051  11.684   0.000
## Q5          0.570   0.047  12.094   0.000
## Q6          0.416   0.034  12.317   0.000
## Q7          0.449   0.062   7.268   0.000
## Q8          0.454   0.069   6.600   0.000
## Q9          0.351   0.057   6.129   0.000
## Q10         0.419   0.043   9.672   0.000
## xi1 =~
## Q1          0.519   0.038  13.840   0.000
## Q2          1.018   0.034  29.844   0.000
## Q3          0.640   0.070   9.131   0.000
## Q4          0.476   0.061   7.862   0.000
## xi2 =~
## Q5          0.577   0.048  12.054   0.000
## Q6          0.262   0.031   8.332   0.000
## Q7          0.747   0.039  19.267   0.000
## Q8          0.836   0.038  21.881   0.000
## Q9          0.660   0.034  19.400   0.000
## xi3 =~
## Q10         0.840   0.061  13.857   0.000
## Q6          0.201   0.028   7.161   0.000
## Q1          0.313   0.037   8.487   0.000
##
## Covariances:
##      Estimate Std.Err z-value P(>|z|)
## G ~~
## xi1          0.000
## xi2          0.000
## xi3          0.000

```

```
## xi1 ~~
## xi2          0.696    0.036   19.354    0.000
## xi3          0.410    0.058    7.090    0.000
## xi2 ~~
## xi3          0.429    0.052    8.236    0.000
##
## Variances:
##           Estimate Std.Err z-value P(>|z|)
## .Q1           0.572   0.023   24.759   0.000
## .Q2           0.073   0.064    1.129   0.259
## .Q3           0.297   0.016   19.145   0.000
## .Q4           0.492   0.017   29.527   0.000
## .Q5           0.588   0.017   33.891   0.000
## .Q6           0.604   0.017   36.143   0.000
## .Q7           0.379   0.012   30.469   0.000
## .Q8           0.274   0.012   22.434   0.000
## .Q9           0.776   0.021   37.111   0.000
## .Q10          0.527   0.097    5.405   0.000
## G             1.000
## xi1           1.000
## xi2           1.000
## xi3           1.000
```

```
#reliability, unidimensionality
```

```
{
  twofrel = round(semTools::reliability(bifactor_cfa)[-5,],2)
  twofrel = cbind(twofrel,NA)
  twofrel = cbind(twofrel, "two-factor")
  colnames(twofrel) = c("G", "$\\xi_1$", "$\\xi_2$", "$\\xi_3$", "model")
  rownames(twofrel) = c("$\\alpha$", "$\\omega$", "$\\omega_2$", "$\\omega_3$")

  threefrel = round(semTools::reliability(bifactor_cfa_4)[-5,],2)
  threefrel = cbind(threefrel,"three-factor")
  colnames(threefrel) = colnames(twofrel)
  rownames(threefrel) = rownames(twofrel)

  write.csv(rbind(twofrel,threefrel),
            file="composite_reliability.csv")

}
```

```
#measurement invariance
```

```
{
  #fit MI models
  gender_configural = cfa(bifactor_model,
                          data=df_clean[,c(SCS_vars,"gender")],
```

```

                                group="gender")
gender_weak = cfa(bifactor_model,
                  data=df_clean[,c(SCS_vars,"gender")],
                  group="gender",
                  group.equal=c("loadings") )
gender_strong = cfa(bifactor_model,
                   data=df_clean[,c(SCS_vars,"gender")],
                   group="gender",
                   group.equal=c("loadings","intercepts") )
gender_strict = cfa(bifactor_model,
                   data=df_clean[,c(SCS_vars,"gender")],
                   group="gender",
                   group.equal=c("loadings","intercepts","residuals") )

mi_gender_modelcomp = anova(gender_configural,
                           gender_weak,
                           gender_strong,
                           gender_strict)

mi_gender_compout=data.frame(cbind(DF=mi_gender_modelcomp$Df,
                                   AIC=mi_gender_modelcomp$AIC,
                                   BIC=mi_gender_modelcomp$BIC,
                                   Chisq=mi_gender_modelcomp$Chisq,
                                   Chisq_diff=mi_gender_modelcomp$`Chisq diff`,
                                   DF_diff=mi_gender_modelcomp$`Df diff`,
                                   p=mi_gender_modelcomp$`Pr(>Chisq)`))
rownames(mi_gender_compout) = c("configural MI",
                                "weak MI",
                                "strong MI",
                                "strict MI")
write.csv(mi_gender_compout,"mi_gender_compout.csv")

df_agegroups = df_clean[,c(SCS_vars)]
df_agegroups$age = df_clean$age > median(df_clean$age)
age_configural = cfa(bifactor_model_4,
                    data=df_agegroups[,c(SCS_vars,"age")],
                    group="age")
age_weak = cfa(bifactor_model_4,
               data=df_agegroups[,c(SCS_vars,"age")],
               group="age",
               group.equal=c("loadings") )
age_strong = cfa(bifactor_model_4,
                 data=df_agegroups[,c(SCS_vars,"age")],

```

```

        group="age",
        group.equal=c("loadings","intercepts") )
age_strict = cfa(bifactor_model_4,
        data=df_agegroups[,c(SCS_vars,"age")],
        group="age",
        group.equal=c("loadings","intercepts","residuals") )

mi_age_modelcomp = anova(age_configural,
        age_weak,
        age_strong,
        age_strict)

mi_age_compout=data.frame(cbind(DF=mi_age_modelcomp$Df,
        AIC=mi_age_modelcomp$AIC,
        BIC=mi_age_modelcomp$BIC,
        Chisq=mi_age_modelcomp$Chisq,
        Chisq_diff=mi_age_modelcomp$`Chisq diff`,
        DF_diff=mi_age_modelcomp$`Df diff`,
        p=mi_age_modelcomp$`Pr(>Chisq)`))
rownames(mi_age_compout) = c("configural MI",
        "weak MI",
        "strong MI",
        "strict MI")
write.csv(mi_age_compout,"mi_age_compout.csv")

#factor structure (just a reminder for
#modification index interpretation):
# G =~ Q1+Q2+Q3+Q4+Q5+Q6+Q7+Q8+Q9+Q10
# xi1 =~ Q1+Q2+Q3+Q4
# xi2 =~ Q5+Q6+Q7+Q8+Q9
# xi3 =~ Q10+Q6+Q1

#look at mod. indices of strong invariance model
modindices(gender_weak)
modindices(age_weak)

}

```

```

## Warning in lav_object_post_check(object): lavaan WARNING: some estimated ov
## variances are negative

```

##	lhs	op	rhs	block	group	level	mi	epc	sepc.lv	sepc.all	sepc.nox
## 1	G =~		Q1	1	1	1	2.533	0.187	0.058	0.054	0.054
## 11	xi1 =~		Q1	1	1	1	4.204	0.128	0.069	0.065	0.065
## 15	xi2 =~		Q5	1	1	1	0.308	0.031	0.018	0.016	0.016
## 20	xi3 =~		Q10	1	1	1	3.005	-0.329	-0.272	-0.231	-0.231

## 23	G ~~ xi1	1	1	1	0.212	0.004	0.022	0.022	0.022
## 24	G ~~ xi2	1	1	1	1.025	0.008	0.044	0.044	0.044
## 25	G ~~ xi3	1	1	1	0.960	-0.015	-0.059	-0.059	-0.059
## 57	G == Q1	2	2	1	2.533	-0.187	-0.057	-0.052	-0.052
## 67	xi1 == Q1	2	2	1	4.203	-0.128	-0.068	-0.062	-0.062
## 71	xi2 == Q5	2	2	1	0.308	-0.031	-0.018	-0.016	-0.016
## 76	xi3 == Q10	2	2	1	3.005	0.329	0.289	0.243	0.243
## 79	G ~~ xi1	2	2	1	0.212	-0.003	-0.021	-0.021	-0.021
## 80	G ~~ xi2	2	2	1	1.025	-0.008	-0.043	-0.043	-0.043
## 81	G ~~ xi3	2	2	1	0.960	0.014	0.054	0.054	0.054
## 131	xi1 == Q5	1	1	1	1.045	0.054	0.029	0.026	0.026
## 132	xi1 == Q6	1	1	1	0.238	-0.024	-0.013	-0.013	-0.013
## 133	xi1 == Q7	1	1	1	1.586	-0.054	-0.029	-0.027	-0.027
## 134	xi1 == Q8	1	1	1	1.263	-0.050	-0.027	-0.025	-0.025
## 135	xi1 == Q9	1	1	1	7.657	0.147	0.079	0.070	0.070
## 136	xi1 == Q10	1	1	1	1.571	-0.172	-0.093	-0.079	-0.079
## 137	xi2 == Q1	1	1	1	0.061	0.012	0.007	0.007	0.007
## 138	xi2 == Q2	1	1	1	0.971	-0.072	-0.042	-0.039	-0.039
## 139	xi2 == Q3	1	1	1	0.097	0.015	0.009	0.008	0.008
## 140	xi2 == Q4	1	1	1	0.475	0.035	0.021	0.020	0.020
## 141	xi2 == Q10	1	1	1	0.392	-0.076	-0.045	-0.038	-0.038
## 142	xi3 == Q2	1	1	1	8.805	0.178	0.147	0.137	0.137
## 143	xi3 == Q3	1	1	1	0.001	0.001	0.001	0.001	0.001
## 144	xi3 == Q4	1	1	1	9.231	-0.111	-0.092	-0.088	-0.088
## 145	xi3 == Q5	1	1	1	17.922	0.147	0.122	0.110	0.110
## 146	xi3 == Q7	1	1	1	5.465	-0.070	-0.058	-0.054	-0.054
## 147	xi3 == Q8	1	1	1	9.288	-0.092	-0.076	-0.071	-0.071
## 148	xi3 == Q9	1	1	1	9.341	0.114	0.095	0.084	0.084
## 149	Q1 ~~ Q2	1	1	1	1.406	0.029	0.029	0.130	0.130
## 150	Q1 ~~ Q3	1	1	1	10.168	0.048	0.048	0.123	0.123
## 151	Q1 ~~ Q4	1	1	1	5.387	-0.037	-0.037	-0.070	-0.070
## 152	Q1 ~~ Q5	1	1	1	0.362	0.010	0.010	0.018	0.018
## 153	Q1 ~~ Q6	1	1	1	4.030	0.039	0.039	0.068	0.068
## 154	Q1 ~~ Q7	1	1	1	0.303	0.007	0.007	0.016	0.016
## 155	Q1 ~~ Q8	1	1	1	1.043	-0.014	-0.014	-0.037	-0.037
## 156	Q1 ~~ Q9	1	1	1	3.514	-0.032	-0.032	-0.052	-0.052
## 157	Q1 ~~ Q10	1	1	1	12.948	-0.176	-0.176	-0.326	-0.326
## 158	Q2 ~~ Q3	1	1	1	8.296	-0.058	-0.058	-0.371	-0.371
## 159	Q2 ~~ Q4	1	1	1	0.221	-0.008	-0.008	-0.040	-0.040
## 160	Q2 ~~ Q5	1	1	1	15.002	0.069	0.069	0.305	0.305
## 161	Q2 ~~ Q6	1	1	1	2.346	-0.026	-0.026	-0.110	-0.110
## 162	Q2 ~~ Q7	1	1	1	1.935	-0.020	-0.020	-0.108	-0.108
## 163	Q2 ~~ Q8	1	1	1	3.146	-0.026	-0.026	-0.177	-0.177
## 164	Q2 ~~ Q9	1	1	1	3.594	0.033	0.033	0.132	0.132
## 165	Q2 ~~ Q10	1	1	1	12.915	0.115	0.115	0.527	0.527
## 166	Q3 ~~ Q4	1	1	1	10.138	0.073	0.073	0.200	0.200
## 167	Q3 ~~ Q5	1	1	1	23.476	-0.077	-0.077	-0.197	-0.197
## 168	Q3 ~~ Q6	1	1	1	0.202	0.007	0.007	0.017	0.017

## 169	Q3	~~	Q7	1	1	1	0.064	-0.003	-0.003	-0.010	-0.010
## 170	Q3	~~	Q8	1	1	1	0.692	0.010	0.010	0.037	0.037
## 171	Q3	~~	Q9	1	1	1	6.855	0.041	0.041	0.095	0.095
## 172	Q3	~~	Q10	1	1	1	3.141	-0.035	-0.035	-0.092	-0.092
## 173	Q4	~~	Q5	1	1	1	0.344	-0.010	-0.010	-0.019	-0.019
## 174	Q4	~~	Q6	1	1	1	5.426	-0.038	-0.038	-0.071	-0.071
## 175	Q4	~~	Q7	1	1	1	3.333	0.024	0.024	0.058	0.058
## 176	Q4	~~	Q8	1	1	1	3.433	0.023	0.023	0.067	0.067
## 177	Q4	~~	Q9	1	1	1	5.016	-0.039	-0.039	-0.066	-0.066
## 178	Q4	~~	Q10	1	1	1	1.993	-0.029	-0.029	-0.057	-0.057
## 179	Q5	~~	Q6	1	1	1	5.724	0.040	0.040	0.069	0.069
## 180	Q5	~~	Q7	1	1	1	0.235	0.007	0.007	0.016	0.016
## 181	Q5	~~	Q8	1	1	1	0.104	-0.005	-0.005	-0.013	-0.013
## 182	Q5	~~	Q9	1	1	1	7.773	-0.050	-0.050	-0.080	-0.080
## 183	Q5	~~	Q10	1	1	1	10.007	0.066	0.066	0.121	0.121
## 184	Q6	~~	Q7	1	1	1	1.943	-0.019	-0.019	-0.042	-0.042
## 185	Q6	~~	Q8	1	1	1	3.008	-0.023	-0.023	-0.061	-0.061
## 186	Q6	~~	Q9	1	1	1	16.160	0.070	0.070	0.110	0.110
## 187	Q6	~~	Q10	1	1	1	5.934	0.078	0.078	0.141	0.141
## 188	Q7	~~	Q8	1	1	1	1.739	0.022	0.022	0.074	0.074
## 189	Q7	~~	Q9	1	1	1	2.023	-0.023	-0.023	-0.045	-0.045
## 190	Q7	~~	Q10	1	1	1	2.820	-0.030	-0.030	-0.068	-0.068
## 191	Q8	~~	Q9	1	1	1	1.236	0.018	0.018	0.044	0.044
## 192	Q8	~~	Q10	1	1	1	3.091	-0.031	-0.031	-0.087	-0.087
## 193	Q9	~~	Q10	1	1	1	3.263	0.040	0.040	0.067	0.067
## 194	xi1	==	Q5	2	2	1	0.918	0.051	0.027	0.024	0.024
## 195	xi1	==	Q6	2	2	1	3.480	-0.096	-0.051	-0.052	-0.052
## 196	xi1	==	Q7	2	2	1	1.019	0.044	0.023	0.022	0.022
## 197	xi1	==	Q8	2	2	1	0.427	-0.030	-0.016	-0.015	-0.015
## 198	xi1	==	Q9	2	2	1	0.256	-0.028	-0.015	-0.013	-0.013
## 199	xi1	==	Q10	2	2	1	8.131	0.411	0.217	0.183	0.183
## 200	xi2	==	Q1	2	2	1	1.218	-0.061	-0.036	-0.033	-0.033
## 201	xi2	==	Q2	2	2	1	2.488	0.118	0.070	0.065	0.065
## 202	xi2	==	Q3	2	2	1	0.571	-0.037	-0.022	-0.020	-0.020
## 203	xi2	==	Q4	2	2	1	0.002	-0.002	-0.001	-0.001	-0.001
## 204	xi2	==	Q10	2	2	1	2.068	0.200	0.118	0.099	0.099
## 205	xi3	==	Q2	2	2	1	0.122	0.019	0.017	0.016	0.016
## 206	xi3	==	Q3	2	2	1	0.024	-0.005	-0.004	-0.004	-0.004
## 207	xi3	==	Q4	2	2	1	0.690	-0.027	-0.024	-0.023	-0.023
## 208	xi3	==	Q5	2	2	1	14.860	0.123	0.108	0.096	0.096
## 209	xi3	==	Q7	2	2	1	2.194	-0.041	-0.036	-0.034	-0.034
## 210	xi3	==	Q8	2	2	1	4.792	-0.063	-0.055	-0.051	-0.051
## 211	xi3	==	Q9	2	2	1	7.454	0.099	0.087	0.073	0.073
## 212	Q1	~~	Q2	2	2	1	0.385	0.015	0.015	0.060	0.060
## 213	Q1	~~	Q3	2	2	1	18.073	-0.065	-0.065	-0.150	-0.150
## 214	Q1	~~	Q4	2	2	1	4.546	0.033	0.033	0.062	0.062
## 215	Q1	~~	Q5	2	2	1	0.366	0.010	0.010	0.017	0.017
## 216	Q1	~~	Q6	2	2	1	1.699	0.025	0.025	0.042	0.042

##	217	Q1	~~	Q7	2	2	1	1.502	0.017	0.017	0.035	0.035
##	218	Q1	~~	Q8	2	2	1	0.393	0.009	0.009	0.021	0.021
##	219	Q1	~~	Q9	2	2	1	15.322	-0.072	-0.072	-0.103	-0.103
##	220	Q1	~~	Q10	2	2	1	6.670	0.129	0.129	0.246	0.246
##	221	Q2	~~	Q3	2	2	1	5.119	0.044	0.044	0.236	0.236
##	222	Q2	~~	Q4	2	2	1	1.559	-0.022	-0.022	-0.094	-0.094
##	223	Q2	~~	Q5	2	2	1	0.003	-0.001	-0.001	-0.004	-0.004
##	224	Q2	~~	Q6	2	2	1	0.516	-0.012	-0.012	-0.044	-0.044
##	225	Q2	~~	Q7	2	2	1	0.045	0.003	0.003	0.015	0.015
##	226	Q2	~~	Q8	2	2	1	1.951	-0.021	-0.021	-0.115	-0.115
##	227	Q2	~~	Q9	2	2	1	5.733	0.043	0.043	0.143	0.143
##	228	Q2	~~	Q10	2	2	1	0.387	-0.018	-0.018	-0.080	-0.080
##	229	Q3	~~	Q4	2	2	1	0.234	0.010	0.010	0.026	0.026
##	230	Q3	~~	Q5	2	2	1	0.042	0.003	0.003	0.008	0.008
##	231	Q3	~~	Q6	2	2	1	0.741	0.013	0.013	0.030	0.030
##	232	Q3	~~	Q7	2	2	1	1.239	-0.014	-0.014	-0.040	-0.040
##	233	Q3	~~	Q8	2	2	1	0.229	-0.006	-0.006	-0.019	-0.019
##	234	Q3	~~	Q9	2	2	1	2.274	0.026	0.026	0.049	0.049
##	235	Q3	~~	Q10	2	2	1	3.517	0.036	0.036	0.092	0.092
##	236	Q4	~~	Q5	2	2	1	0.889	0.016	0.016	0.029	0.029
##	237	Q4	~~	Q6	2	2	1	14.131	-0.060	-0.060	-0.110	-0.110
##	238	Q4	~~	Q7	2	2	1	0.996	0.014	0.014	0.031	0.031
##	239	Q4	~~	Q8	2	2	1	0.968	0.013	0.013	0.033	0.033
##	240	Q4	~~	Q9	2	2	1	5.102	-0.041	-0.041	-0.064	-0.064
##	241	Q4	~~	Q10	2	2	1	0.556	-0.015	-0.015	-0.030	-0.030
##	242	Q5	~~	Q6	2	2	1	13.149	0.061	0.061	0.100	0.100
##	243	Q5	~~	Q7	2	2	1	0.581	0.012	0.012	0.024	0.024
##	244	Q5	~~	Q8	2	2	1	1.793	-0.021	-0.021	-0.049	-0.049
##	245	Q5	~~	Q9	2	2	1	21.614	-0.091	-0.091	-0.127	-0.127
##	246	Q5	~~	Q10	2	2	1	8.134	0.057	0.057	0.108	0.108
##	247	Q6	~~	Q7	2	2	1	0.012	-0.002	-0.002	-0.003	-0.003
##	248	Q6	~~	Q8	2	2	1	10.651	-0.046	-0.046	-0.105	-0.105
##	249	Q6	~~	Q9	2	2	1	16.568	0.076	0.076	0.106	0.106
##	250	Q6	~~	Q10	2	2	1	10.481	-0.103	-0.103	-0.190	-0.190
##	251	Q7	~~	Q8	2	2	1	13.549	0.065	0.065	0.186	0.186
##	252	Q7	~~	Q9	2	2	1	11.459	-0.059	-0.059	-0.102	-0.102
##	253	Q7	~~	Q10	2	2	1	7.700	-0.049	-0.049	-0.113	-0.113
##	254	Q8	~~	Q9	2	2	1	3.783	0.035	0.035	0.069	0.069
##	255	Q8	~~	Q10	2	2	1	2.726	-0.030	-0.030	-0.078	-0.078
##	256	Q9	~~	Q10	2	2	1	18.440	0.099	0.099	0.157	0.157

12 References

- Crocker, Linda, and James Algina. 2008. *Introduction to Classical and Modern Test Theory*. Cengage Learning.
- Guan, Ng Chong, and Muhamad Saiful Bahri Yusoff. 2011. “Missing Values in Data Analysis: Ignore or Impute?” *Education in Medicine Journal* 3 (1).
- Kalichman, Seth C, and David Rompa. 1995. “Sexual Sensation Seeking and Sexual Compulsivity Scales: Validity, and Predicting HIV Risk Behavior.” *Journal of Personality Assessment* 65 (3): 586–601.
- . 2001. “The Sexual Compulsivity Scale: Further Development and Use with HIV-Positive Persons.” *Journal of Personality Assessment* 76 (3): 379–95.
- Magis, David, Sébastien Béland, Francis Tuerlinckx, and Paul De Boeck. 2010. “A General Framework and an r Package for the Detection of Dichotomous Differential Item Functioning.” *Behavior Research Methods* 42 (3): 847–62.
- Mair, Patrick, and Reinhold Hatzinger. 2007. “Extended Rasch Modeling: The eRm Package for the Application of IRT Models in r.”
- Novick, Melvin R. 1965. “The Axioms and Principal Results of Classical Test Theory.” *ETS Research Bulletin Series* 1965 (1): i–31.
- Rasch, Georg. 1960. “Studies in Mathematical Psychology: I. Probabilistic Models for Some Intelligence and Attainment Tests.”
- Reynolds, Cecil R, and RA Livingston. 2021. *Mastering Modern Psychological Testing*. Springer.
- Rizopoulos, Dimitris. 2006. “Ltm: An r Package for Latent Variable Modelling and Item Response Theory Analyses.” *Journal of Statistical Software* 17 (5): 1–25. <https://doi.org/10.18637/jss.v017.i05>.
- Rosseel, Yves. 2012. “Lavaan: An r Package for Structural Equation Modeling.” *Journal of Statistical Software* 48: 1–36.
- Smyth, Rachael. 2022. “Item Response Theory for Polytomous Items.” <https://www.uwo.ca/fhs/tc/labs/12.PolytomousIRT/>
- Templin, Jonathan. 2022. “IRT Estimation with r Packages Mirt and Lavaan.” <https://jonathantemplin.com/irt-estimation-packages-mirt-lavaan/>.
- Van de Schoot, Rens, Peter Lugtig, and Joop Hox. 2012. “A Checklist for Testing Measurement Invariance.” *European Journal of Developmental Psychology* 9 (4): 486–92.
- Van der Linden, Wim J, and RK Hambleton. 1997. “Handbook of Item Response Theory.” *Taylor & Francis Group*. 1 (7): 8.
- Xu, Kate. 2012. “Multiple Group Measurement: Invariance Analysis in Lavaan.” *Cambridge Mass* 37.