

1

2

## Contents

3	<b>1 MC samples and selection</b>	<b>1</b>
4	1.1 Data and MC samples . . . . .	1
5	1.1.1 MC samples and cross-sections . . . . .	1
6	1.2 Z vertex reweighting . . . . .	3
7	1.2.1 Multijet background . . . . .	3
8	1.3 W analysis event selection and control plots . . . . .	8
9	1.3.1 Event selection . . . . .	8
10	1.3.2 $\sqrt{s} = 13$ TeV dataset control plots . . . . .	9
11	1.3.3 $\sqrt{s} = 5$ TeV dataset control plots . . . . .	19
12	Bibliography . . . . .	26

13

## List of Figures

14	11 Distributions for the 5 TeV low- $\mu$ dataset in a $Z/\gamma^* \rightarrow \mu\mu$ (top row) and a $Z/\gamma^* \rightarrow ee$	
15	(bottom row) selection. The data (points) is compared to $Z/\gamma^* \rightarrow \mu\mu$ or $Z/\gamma^* \rightarrow ee$ signal	
16	MC, respectively. The left and middle plots show the actual $\mu$ in a coarsely-binned and a	
17	finely-binned version. The right plot shows the number of reconstructed primary vertices	
18	$N_{PV}$ . . . . .	2

19	12	Distributions for the 13 TeV low- $\mu$ datasets taken in 2017 and 2018 in a $Z/\gamma^* \rightarrow \mu\mu$ (top row) and a $Z/\gamma^* \rightarrow ee$ (bottom row) selection. The data (points) is compared to $Z/\gamma^* \rightarrow \mu\mu$ or $Z/\gamma^* \rightarrow ee$ signal MC, respectively. All distributions are (roughly) normalised to the same number of selected events in the 2017 dataset. The left and middle plots show the actual $\mu$ in a coarsely-binned and a finely-binned version. The right plot shows the number of reconstructed primary vertices $N_{PV}$ . . . . .	2
25	13	Distributions for the 5 TeV (left) and 13 TeV (right) low- $\mu$ dataset(s) in a $Z/\gamma^* \rightarrow \mu\mu$ (top row) and a $Z/\gamma^* \rightarrow ee$ (bottom row) selection. The data (points) is compared to $Z/\gamma^* \rightarrow \mu\mu$ or $Z/\gamma^* \rightarrow ee$ signal MC, respectively. The distributions of the z-position of the primary vertex selected as the hard interaction are compared for the dataset(s) and the MC simulation before ("no $z_{\text{vtx}}$ rwgt", blue, only 13 TeV) and after reweighting (black). For the 13 TeV data the 2017 and 2018 data are shown separately and all distributions are (roughly) normalised to the same number of selected events in the 2017 dataset. . . . .	8
32	14	$\Sigma \vec{E}_T$ distribution in the muon and electron channel for the $\sqrt{s} = 13$ TeV dataset. . . . .	12
33	15	$\Sigma E_T$ distribution in the muon and electron channel for the $\sqrt{s} = 13$ TeV dataset. . . . .	13
34	16	$\vec{E}_T^{\text{miss}}$ distribution in the muon and electron channel for the $\sqrt{s} = 13$ TeV dataset. . . . .	14
35	17	Transverse mass distribution of the W boson in the muon and electron channel for the $\sqrt{s} = 13$ TeV dataset. . . . .	15
37	18	Lepton pseudorapidity distribution in the muon and electron channel for the $\sqrt{s} = 13$ TeV dataset. . . . .	16
39	19	Lepton transverse momentum distribution in the muon and electron channel for the $\sqrt{s} = 13$ TeV dataset. . . . .	17
41	110	W transverse momentum distribution in the muon and electron channel for the $\sqrt{s} = 13$ TeV dataset. . . . .	18
43	111	$\Sigma \vec{E}_T$ distribution in the muon and electron channel for the $\sqrt{s} = 5$ TeV dataset. . . . .	19
44	112	$\Sigma E_T$ distribution in the muon and electron channel for the $\sqrt{s} = 5$ TeV dataset. . . . .	20
45	113	$\vec{E}_T^{\text{miss}}$ distribution in the muon and electron channel for the $\sqrt{s} = 5$ TeV dataset. . . . .	21
46	114	Transverse mass distribution of the W boson in the muon and electron channel for the $\sqrt{s} = 5$ TeV dataset. . . . .	22
48	115	Lepton pseudorapidity distribution in the muon and electron channel for the $\sqrt{s} = 5$ TeV dataset. . . . .	23
50	116	Lepton transverse momentum distribution in the muon and electron channel for the $\sqrt{s} = 5$ TeV dataset. . . . .	24

52	117	W transverse momentum distribution in the muon and electron channel for the $\sqrt{s} = 5$ TeV dataset. . . . .	25
53			

54

## List of Tables

55	11	Monte Carlo samples at $\sqrt{s} = 13$ TeV. Given is a short description of the process, the ATLAS MC data set number (DSID), the names and version numbers of the MC generator(s), the used value of the higher order cross section times any branching and filter efficiencies ( $\sigma \cdot \text{BR} \cdot \epsilon_{\text{filter}}$ ) with the theoretical uncertainty in percent (“th. unc.”), and finally the number of events analysed after skimming at derivation production ( $N_{\text{evt}}^{\text{skim}}$ ) as well as the number of events originally processed and simulated ( $N_{\text{evt}}^{\text{unskim}}$ ). In the case of $Z \rightarrow \ell\ell$ samples, the given $\epsilon_{\text{filter}} > 1$ is related to the fact, that the cross sections were calculated for $66 < m_{\ell\ell} < 116$ GeV, but the generated mass range is larger. The last section of $t\bar{t}$ samples refers to variation samples for systematics studies. The MC equivalent luminosity $N_{\text{evt}}^{\text{unskim}}/(\sigma \cdot \text{BR} \cdot \epsilon_{\text{filter}})$ is generally above $3\text{fb}^{-1}$ for signal and significant backgrounds, the exception are Powheg $W \rightarrow \tau\nu$ and $Z \rightarrow \tau\tau$ samples, that have about $0.45\text{fb}^{-1}$ only. . . . .	4
56			
57			
58			
59			
60			
61			
62			
63			
64			
65			
66	12	Alternative signal $Z \rightarrow \ell\ell$ Monte Carlo samples at $\sqrt{s} = 13$ TeV produced with SHERPA. General description of the table see Table 11. The samples are split into a long list of orthogonal slices based on “max(pTV,HT)” and filtered further into “b/c/light-jet” subcomponents. For the purpose of this analysis, the number of events in each slice is such that the samples are about $2\text{fb}^{-1}$ each (after application of a penalty factor for negative weight events) and an “inclusive sample” is restored after merging the slices. . . . .	5
67			
68			
69			
70			
71			
72	13	Alternative signal $W \rightarrow \ell\nu$ Monte Carlo samples at $\sqrt{s} = 13$ TeV produced with SHERPA. See Table 12 for a description of the table. The samples are split into a long list of orthogonal slices based on “max(pTV,HT)” and filtered further into “b/c/light-jet” subcomponents. For the purpose of this analysis, the number of events in each slice is such that the samples are about $1\text{fb}^{-1}$ each (after application of a penalty factor for negative weight events) and an “inclusive sample” is restored after merging the slices. . . . .	6
73			
74			
75			
76			
77			
78	14	Monte Carlo samples at $\sqrt{s} = 5$ TeV. The table follows the same format as Table 11. The MC equivalent luminosity $N_{\text{evt}}^{\text{unskim}}/(\sigma \cdot \text{BR} \cdot \epsilon_{\text{filter}})$ is generally above $2.5\text{fb}^{-1}$ for signal and significant backgrounds, the exception are Powheg $W \rightarrow \tau\nu$ and $Z \rightarrow \tau\tau$ samples, that have about $0.20\text{fb}^{-1}$ and $0.45\text{fb}^{-1}$ only. . . . .	7
79			
80			
81			
82	15	Analysis cut flow for $W^+ \rightarrow e^+\nu$ 5 TeV signal selection. Lepton $p_T$ is required to be over 18 GeV before the final cut. . . . .	9
83			

84	16	Analysis cut flow for $W^+ \rightarrow e^+ \nu$ 13 TeV signal selection. Lepton $p_T$ is required to be over	
85		18 GeVbefore the final cut. . . . .	9
86	17	Analysis cut flow for $W^+ \rightarrow \mu^+ \nu$ 5 TeV signal selection. Lepton $p_T$ is required to be over	
87		18 GeVbefore the final cut. . . . .	10
88	18	Analysis cut flow for $W^+ \rightarrow \mu^+ \nu$ 13 TeV signal selection. Lepton $p_T$ is required to be over	
89		18 GeVbefore the final cut. . . . .	10
90	19	Analysis cut flow for $W^- \rightarrow e^- \nu$ 5 TeV signal selection. Lepton $p_T$ is required to be over	
91		18 GeVbefore the final cut. . . . .	10
92	110	Analysis cut flow for $W^- \rightarrow e^- \nu$ 13 TeV signal selection. Lepton $p_T$ is required to be over	
93		18 GeVbefore the final cut. . . . .	11
94	111	Analysis cut flow for $W^- \rightarrow \mu^- \nu$ 5 TeV signal selection. Lepton $p_T$ is required to be over	
95		18 GeVbefore the final cut. . . . .	11
96	112	Analysis cut flow for $W^- \rightarrow \mu^- \nu$ 13 TeV signal selection. Lepton $p_T$ is required to be over	
97		18 GeVbefore the final cut. . . . .	11
98	113	Observed and Expected yield comparison for all signal selections. . . . .	11

# 1

## MC samples and selection

*“Potentielle citation sans aucun rapport avec le sujet”*

— Personne inconnue, contexte à déterminer

### 1.1 Data and MC samples

The data and MC samples for this study were collected under special beam conditions that ensure low pile-up. The data samples were collected in three runs:

- $\sqrt{s} = 5.02\text{TeV}$  data taken in November 2017, ATLAS data period M, preliminary calibrated luminosity  $256.827\text{pb}^{-1}$  with an uncertainty of  $\pm 1.6\%$
- $\sqrt{s} = 13\text{TeV}$  data taken in November 2017, ATLAS data period N, preliminary online luminosity  $146.6\text{pb}^{-1}$
- $\sqrt{s} = 13\text{TeV}$  data taken in June 2018, ATLAS data period G4+J, preliminary online luminosity  $193.2\text{pb}^{-1}$


The runs of November 2017 and the run of June 2018 had the same bunch spacing of 25 ns, but a different filling scheme. The two main differences from the high- $\mu$  data collection are the following:

- In order to optimize topo-cluster response for the Hadronic Recoil (HR) lower topo-cluster thresholds were used.
- Single  $e$  and  $\mu$  triggers with significantly lower thresholds and looser identification criteria are run without prescale, most notably HLT\_e15\_1h1oose\_nod0\_L1EM12 and HLT\_mu14.


At the beginning of 5 TeV fills the pile-up reached  $\mu \backslash 5$ , slowly descending to  $\mu \backslash 1$  by the end of the run. In the case of 13 TeV the luminosity was levelled at  $\mu = 2$  in the course of the run. The corresponding distributions for  $\mu$  and  $N_{PV}$  for the 5 TeV and 13 TeV runs are shown in Fig. 11 and Fig. 12.

#### 1.1.1 MC samples and cross-sections

Signal and background processes are modelled using fully simulated and reconstructed using Monte-Carlo (MC) samples, specifically tuned for the special run conditions, namely the pileup conditions,



**Figure 11:** Distributions for the 5 TeV low- $\mu$  dataset in a  $Z/\gamma^* \rightarrow \mu\mu$  (top row) and a  $Z/\gamma^* \rightarrow ee$  (bottom row) selection. The data (points) is compared to  $Z/\gamma^* \rightarrow \mu\mu$  or  $Z/\gamma^* \rightarrow ee$  signal MC, respectively. The left and middle plots show the actual  $\mu$  in a coarsely-binned and a finely-binned version. The right plot shows the number of reconstructed primary vertices  $N_{PV}$ .



**Figure 12:** Distributions for the 13 TeV low- $\mu$  datasets taken in 2017 and 2018 in a  $Z/\gamma^* \rightarrow \mu\mu$  (top row) and a  $Z/\gamma^* \rightarrow ee$  (bottom row) selection. The data (points) is compared to  $Z/\gamma^* \rightarrow \mu\mu$  or  $Z/\gamma^* \rightarrow ee$  signal MC, respectively. All distributions are (roughly) normalised to the same number of selected events in the 2017 dataset. The left and middle plots show the actual  $\mu$  in a coarsely-binned and a finely-binned version. The right plot shows the number of reconstructed primary vertices  $N_{PV}$ .

lower topo-cluster noise thresholds and adapter trigger menu. No pileup reweighting is performed. The information on the simulated samples and their properties is given in Tables 11, 12, 13, 14 [1]. The predicted event counts are normalized to the cross-sections quoted in the table. The primary signal event samples for  $W$  and  $Z$  production are obtained using POWHEG [2, 3, 4, 5] event generator with CT10 PDF, linked with PYTHIA8 [6] with AZNLO tune [7]. POWHEG+PYTHIA88 samples are interfaced to PHOTOS++ [8] for final state Quantum Electrodynamics (QED) effects simulation. A set of alternative samples at  $\sqrt{s} = 13\text{TeV}$  was prepared with SHERPA2.2.2 [9] using the NNPDF3.0 PDFs and merging  $V + 0, 1, 2$  at NLO accuracy with  $V + 3, 4$  at LO accuracy with the MEPS@NLO scheme. A similar set for  $\sqrt{s} = 5\text{TeV}$  was prepared with SHERPA2.2.5 with a setup similar to 13 TeV samples. Pileup is modelled by overlaying simulated soft events over the original hard-scattering event. These soft events were modelled using PYTHIA8 with NNPDF2.3LO set of PDFs [10] and the A3 tune [11]. The  $W$  and  $Z$  processes samples are normalized to NNLO calculations performed using the DYTURBO, an optimised version of DYNNLO [12, 13] using the MMHT2014nnlo PDF set [14]. Corresponding numerical values were taken from the corresponding ATLAS publications of the 2015 data at 13 TeV [15] and 5.02 TeV [16] are presented in Table 11 for 13 TeV and Table 14 for 5 TeV. The uncertainties on those cross-sections arise from the choice of PDF set, from factorization and renormalisation scale dependence, and the strong coupling constant  $\alpha_s$  uncertainty resulting in the total uncertainty estimate of 5%.

Backgrounds from top-quark pair-production  $t\bar{t}$  and single-top production ( $Wt$ ,  $t$ -channel,  $s$ -channel) were generated with POWHEG+PYTHIA8. The 5 TeV  $t\bar{t}$  cross section is taken as the top++ prediction observed by CMS [17]. Di-boson combinations  $VV$ ,  $V = W, Z$  are generated with SHERPA in all decay channels with a requirement of having at least one real lepton in the final state.

## 1.2 Z vertex reweighting

The 5 TeV MC samples have been generated to be perfectly matched to the data Although this is not the case for 13 TeV samples, which can be seen at Fig. 13. It is also seen from these plots that the 2017 and 2018 data were collected at two different runs under different beam conditions. To avoid possible impact on the acceptance the MC samples were reweighted to the data using  $Z \rightarrow ee$  and  $Z \rightarrow \mu\mu$  selections.

### 1.2.1 Multijet background

The estimate of the multijet background, which contain contributions from fake leptons produced in semi-leptonic decays of heavy quarks, in-flight pion decays, photon conversions, etc, is done using a data-driven techniques described in Reference [Xu:2657146]. TO BE ADDED.

Process	Data set	Generator	$\sigma \cdot \text{BR} \cdot \epsilon_{\text{filter}}$ [nb] (th. unc.)	$N_{\text{evt}}^{\text{skim}} [10^6]$	$N_{\text{evt}}^{\text{unskim}} [10^6]$
$W^+ \rightarrow e^+ \nu$	361100	POWHEG+PYTHIA8	11.61 (5%)	40	40
$W^+ \rightarrow \mu^+ \nu$	361101	POWHEG+PYTHIA8	11.61 (5%)	40	40
$W^+ \rightarrow \tau^+ \nu$	361102	POWHEG+PYTHIA8	11.61 (5%)	0.28	5.0
$W^- \rightarrow e^- \bar{\nu}$	361103	POWHEG+PYTHIA8	8.630 (5%)	30	30
$W^- \rightarrow \mu^- \bar{\nu}$	361104	POWHEG+PYTHIA8	8.630 (5%)	29	29
$W^- \rightarrow \tau^- \bar{\nu}$	361105	POWHEG+PYTHIA8	8.630 (5%)	0.24	4.0
$Z \rightarrow ee$	361106	POWHEG+PYTHIA8	$1.910 \times 1.03$ (5%)	10	10
$Z \rightarrow \mu\mu$	361107	POWHEG+PYTHIA8	$1.910 \times 1.025$ (5%)	10	10
$Z \rightarrow \tau\tau$	361108	POWHEG+PYTHIA8	$1.910 \times 1.025$ (5%)	0.12	1.0
$ZZ(q\bar{q}\ell\ell)$	363356	SHERPA 2.2.1	$0.01556 \times 0.141$ (10%)	0.0064	0.010
$WZ(q\bar{q}\ell\ell)$	363358	SHERPA 2.2.1	0.003433 (10%)	0.0063	0.010
$WW(q\bar{q}\ell\nu)$	363359	SHERPA 2.2.1	0.02472 (10%)	0.0093	0.020
$WW(\ell\nu q\bar{q})$	363360	SHERPA 2.2.1	0.02472 (10%)	0.0093	0.020
$WZ(\ell\nu q\bar{q})$	363489	SHERPA 2.2.1	0.01142 (10%)	0.0047	0.010
$ZZ(4\ell)$	364250	SHERPA 2.2.2	0.001252 (10%)	0.0057	0.010
$WZ(3\ell\nu)$	364253	SHERPA 2.2.2	0.004583 (10%)	0.0062	0.010
$WW(2\ell 2\nu)$	364254	SHERPA 2.2.2	0.01250 (10%)	0.0073	0.010
$WZ(\ell 3\nu)$	364255	SHERPA 2.2.2	0.003235 (10%)	0.0050	0.010
$Wt$	410013	POWHEG+PYTHIA8	0.03582 (10%)	0.0037	0.010
$W\bar{t}$	410014	POWHEG+PYTHIA8	0.03399 (10%)	0.0037	0.010
$t\bar{t}$ (nominal)	410470	POWHEG+PYTHIA8	$0.8318 \times 0.544$ (7%)	1.2	2.0
$t(t - \text{chan.} t)$	410642	POWHEG+PYTHIA8	0.03699 (10%)	0.016	0.030
$t(t - \text{chan.} \bar{t})$	410643	POWHEG+PYTHIA8	0.02217 (10%)	0.011	0.020
$t(s - \text{chan.} t)$	410644	POWHEG+PYTHIA8	0.002027 (10%)	0.0050	0.010
$t(s - \text{chan.} \bar{t})$	410645	POWHEG+PYTHIA8	0.001268 (10%)	0.0052	0.010
$t\bar{t}$ (syst.)	410480	POWHEG+PYTHIA8	$0.8318 \times 0.438$ (7%)	0.85	1.5
$t\bar{t}$ (syst.)	410482	POWHEG+PYTHIA8	$0.8318 \times 0.105$ (7%)	0.40	0.50
$t\bar{t}$ (syst.)	410557	POWHEG+PYTHIA8	$0.8318 \times 0.438$ (7%)	0.85	1.5
$t\bar{t}$ (syst.)	410558	POWHEG+PYTHIA8	$0.8318 \times 0.105$ (7%)	0.40	0.50

**Table 11:** Monte Carlo samples at  $\sqrt{s} = 13\text{TeV}$ . Given is a short description of the process, the ATLAS MC data set number (DSID), the names and version numbers of the MC generator(s), the used value of the higher order cross section times any branching and filter efficiencies ( $\sigma \cdot \text{BR} \cdot \epsilon_{\text{filter}}$ ) with the theoretical uncertainty in percent (“th. unc.”), and finally the number of events analysed after skimming at derivation production ( $N_{\text{evt}}^{\text{skim}}$ ) as well as the number of events originally processed and simulated ( $N_{\text{evt}}^{\text{unskim}}$ ). In the case of  $Z \rightarrow \ell\ell$  samples, the given  $\epsilon_{\text{filter}} > 1$  is related to the fact, that the cross sections were calculated for  $66 < m_{\ell\ell} < 116\text{GeV}$ , but the generated mass range is larger. The last section of  $t\bar{t}$  samples refers to variation samples for systematics studies. The MC equivalent luminosity  $N_{\text{evt}}^{\text{unskim}}/(\sigma \cdot \text{BR} \cdot \epsilon_{\text{filter}})$  is generally above  $3\text{fb}^{-1}$  for signal and significant backgrounds, the exception are Powheg  $W \rightarrow \tau\nu$  and  $Z \rightarrow \tau\tau$  samples, that have about  $0.45\text{fb}^{-1}$  only.



Process	Data set	Generator	$\sigma \cdot \text{BR} \cdot \epsilon_{\text{filter}}$ [nb] (th. unc.)	$N_{\text{evt}}^{\text{skim}} [10^6]$	$N_{\text{evt}}^{\text{unskim}} [10^6]$
$Z \rightarrow \mu\mu$	364100	SHERPA 2.2.1	$1.932 \times 0.822$ (5%)	8.0	8.0
$Z \rightarrow \mu\mu$	364101	SHERPA 2.2.1	$1.933 \times 0.114$ (5%)	1.5	1.5
$Z \rightarrow \mu\mu$	364102	SHERPA 2.2.1	$1.932 \times 0.0660$ (5%)	1.1	1.1
$Z \rightarrow \mu\mu$	364103	SHERPA 2.2.1	$0.1063 \times 0.690$ (5%)	1.5	1.5
$Z \rightarrow \mu\mu$	364104	SHERPA 2.2.1	$0.1062 \times 0.200$ (5%)	0.40	0.40
$Z \rightarrow \mu\mu$	364105	SHERPA 2.2.1	$0.1063 \times 0.114$ (5%)	0.25	0.25
$Z \rightarrow \mu\mu$	364106	SHERPA 2.2.1	$0.03889 \times 0.593$ (5%)	0.20	0.20
$Z \rightarrow \mu\mu$	364107	SHERPA 2.2.1	$0.03885 \times 0.235$ (5%)	0.060	0.060
$Z \rightarrow \mu\mu$	364108	SHERPA 2.2.1	$0.03889 \times 0.156$ (5%)	0.035	0.035
$Z \rightarrow \mu\mu$	364109	SHERPA 2.2.1	$0.008310 \times 0.561$ (5%)	0.020	0.020
$Z \rightarrow \mu\mu$	364110	SHERPA 2.2.1	$0.008310 \times 0.266$ (5%)	0.010	0.010
$Z \rightarrow \mu\mu$	364111	SHERPA 2.2.1	$0.008320 \times 0.177$ (5%)	0.0050	0.0050
$Z \rightarrow \mu\mu$	364112	SHERPA 2.2.1	$0.001740$ (5%)	0.0050	0.0050
$Z \rightarrow \mu\mu$	364113	SHERPA 2.2.1	$0.0001400$ (5%)	0.0050	0.0050
$Z \rightarrow ee$	364114	SHERPA 2.2.1	$1.933 \times 0.821$ (5%)	8.0	8.0
$Z \rightarrow ee$	364115	SHERPA 2.2.1	$1.932 \times 0.114$ (5%)	1.5	1.5
$Z \rightarrow ee$	364116	SHERPA 2.2.1	$1.932 \times 0.0658$ (5%)	1.1	1.1
$Z \rightarrow ee$	364117	SHERPA 2.2.1	$0.1080 \times 0.694$ (5%)	1.5	1.5
$Z \rightarrow ee$	364118	SHERPA 2.2.1	$0.1077 \times 0.191$ (5%)	0.40	0.40
$Z \rightarrow ee$	364119	SHERPA 2.2.1	$0.1078 \times 0.119$ (5%)	0.25	0.25
$Z \rightarrow ee$	364120	SHERPA 2.2.1	$0.03964 \times 0.616$ (5%)	0.20	0.20
$Z \rightarrow ee$	364121	SHERPA 2.2.1	$0.03967 \times 0.233$ (5%)	0.060	0.060
$Z \rightarrow ee$	364122	SHERPA 2.2.1	$0.04068 \times 0.150$ (5%)	0.035	0.035
$Z \rightarrow ee$	364123	SHERPA 2.2.1	$0.008460 \times 0.569$ (5%)	0.020	0.020
$Z \rightarrow ee$	364124	SHERPA 2.2.1	$0.008450 \times 0.266$ (5%)	0.010	0.010
$Z \rightarrow ee$	364125	SHERPA 2.2.1	$0.008470 \times 0.177$ (5%)	0.0050	0.0050
$Z \rightarrow ee$	364126	SHERPA 2.2.1	$0.001760$ (5%)	0.0050	0.0050
$Z \rightarrow ee$	364127	SHERPA 2.2.1	$0.0001451$ (5%)	0.0050	0.0050


**Table 12:** Alternative signal  $Z \rightarrow \ell\ell$  Monte Carlo samples at  $\sqrt{s} = 13\text{TeV}$  produced with SHERPA. General description of the table see Table 11. The samples are split into a long list of orthogonal slices based on “max(pTV,HT)” and filtered further into “b/c/light-jet” subcomponents. For the purpose of this analysis, the number of events in each slice is such that the samples are about  $2\text{fb}^{-1}$  each (after application of a penalty factor for negative weight events) and an “inclusive sample” is restored after merging the slices.

Process	Data set	Generator	$\sigma \cdot \text{BR} \cdot \epsilon_{\text{filter}}$ [nb] (th. unc.)	$N_{\text{evt}}^{\text{skim}} [10^6]$	$N_{\text{evt}}^{\text{unskim}} [10^6]$
$W \rightarrow \mu\nu$	364156	SHERPA 2.2.1	$18.58 \times 0.825$ (5%)	31	31
$W \rightarrow \mu\nu$	364157	SHERPA 2.2.1	$18.57 \times 0.131$ (5%)	8.1	8.1
$W \rightarrow \mu\nu$	364158	SHERPA 2.2.1	$18.57 \times 0.0433$ (5%)	2.6	2.6
$W \rightarrow \mu\nu$	364159	SHERPA 2.2.1	$0.9173 \times 0.674$ (5%)	6.3	6.3
$W \rightarrow \mu\nu$	364160	SHERPA 2.2.1	$0.9172 \times 0.244$ (5%)	2.1	2.1
$W \rightarrow \mu\nu$	364161	SHERPA 2.2.1	$0.9163 \times 0.0847$ (5%)	0.23	0.23
$W \rightarrow \mu\nu$	364162	SHERPA 2.2.1	$0.3296 \times 0.600$ (5%)	0.80	0.80
$W \rightarrow \mu\nu$	364163	SHERPA 2.2.1	$0.3297 \times 0.293$ (5%)	0.27	0.27
$W \rightarrow \mu\nu$	364164	SHERPA 2.2.1	$0.3295 \times 0.111$ (5%)	0.099	0.099
$W \rightarrow \mu\nu$	364165	SHERPA 2.2.1	$0.06993 \times 0.548$ (5%)	0.068	0.068
$W \rightarrow \mu\nu$	364166	SHERPA 2.2.1	$0.06995 \times 0.320$ (5%)	0.034	0.034
$W \rightarrow \mu\nu$	364167	SHERPA 2.2.1	$0.06991 \times 0.125$ (5%)	0.014	0.014
$W \rightarrow \mu\nu$	364168	SHERPA 2.2.1	$0.01456$ (5%)	0.020	0.020
$W \rightarrow \mu\nu$	364169	SHERPA 2.2.1	$0.001200$ (5%)	0.004	0.004
$W \rightarrow e\nu$	364170	SHERPA 2.2.1	$18.58 \times 0.825$ (5%)	31	31
$W \rightarrow e\nu$	364171	SHERPA 2.2.1	$18.57 \times 0.131$ (5%)	8.3	8.3
$W \rightarrow e\nu$	364172	SHERPA 2.2.1	$18.57 \times 0.0448$ (5%)	2.5	2.5
$W \rightarrow e\nu$	364173	SHERPA 2.2.1	$0.9168 \times 0.675$ (5%)	6.4	6.4
$W \rightarrow e\nu$	364174	SHERPA 2.2.1	$0.9176 \times 0.244$ (5%)	2.1	2.1
$W \rightarrow e\nu$	364175	SHERPA 2.2.1	$0.9173 \times 0.0851$ (5%)	0.79	0.79
$W \rightarrow e\nu$	364176	SHERPA 2.2.1	$0.3295 \times 0.599$ (5%)	0.76	0.76
$W \rightarrow e\nu$	364177	SHERPA 2.2.1	$0.3297 \times 0.288$ (5%)	0.28	0.28
$W \rightarrow e\nu$	364178	SHERPA 2.2.1	$0.3295 \times 0.111$ (5%)	0.10	0.10
$W \rightarrow e\nu$	364179	SHERPA 2.2.1	$0.06993 \times 0.548$ (5%)	0.070	0.070
$W \rightarrow e\nu$	364180	SHERPA 2.2.1	$0.06996 \times 0.320$ (5%)	0.034	0.034
$W \rightarrow e\nu$	364181	SHERPA 2.2.1	$0.06994 \times 0.137$ (5%)	0.014	0.014
$W \rightarrow e\nu$	364182	SHERPA 2.2.1	$0.01460$ (5%)	0.020	0.020
$W \rightarrow e\nu$	364183	SHERPA 2.2.1	$0.001200$ (5%)	0.0050	0.0050

**Table 13:** Alternative signal  $W \rightarrow \ell\nu$  Monte Carlo samples at  $\sqrt{s} = 13\text{TeV}$  produced with SHERPA. See Table 12 for a description of the table. The samples are split into a long list of orthogonal slices based on “max(pTV,HT)” and filtered further into “b/c/light-jet” subcomponents. For the purpose of this analysis, the number of events in each slice is such that the samples are about  $1\text{fb}^{-1}$  each (after application of a penalty factor for negative weight events) and an “inclusive sample” is restored after merging the slices.

Process	Data set	Generator	$\sigma \cdot \text{BR} \cdot \epsilon_{\text{filter}}$ [nb] (th. unc.)	$N_{\text{evt}}^{\text{skim}} [10^6]$	$N_{\text{evt}}^{\text{unskim}} [10^6]$
$W^+ \rightarrow e^+ \nu$	361100	POWHEG+PYTHIA8	4.357 (5%)	11	11
$W^+ \rightarrow \mu^+ \nu$	361101	POWHEG+PYTHIA8	4.357 (5%)	11	11
$W^+ \rightarrow \tau^+ \nu$	361102	POWHEG+PYTHIA8	4.357 (5%)	0.065	0.94
$W^- \rightarrow e^- \bar{\nu}$	361103	POWHEG+PYTHIA8	2.902 (5%)	7.0	7.0
$W^- \rightarrow \mu^- \bar{\nu}$	361104	POWHEG+PYTHIA8	2.902 (5%)	7.0	7.0
$W^- \rightarrow \tau^- \bar{\nu}$	361105	POWHEG+PYTHIA8	2.902 (5%)	0.039	0.59
$Z \rightarrow ee$	361106	POWHEG+PYTHIA8	$0.6600 \times 1.025$ (5%)	6.3	6.3
$Z \rightarrow \mu\mu$	361107	POWHEG+PYTHIA8	$0.6600 \times 1.025$ (5%)	3.4	3.4
$Z \rightarrow \tau\tau$	361108	POWHEG+PYTHIA8	$0.6600 \times 1.025$ (5%)	0.039	0.29
$Z \rightarrow ee$	364381	SHERPA 2.2.5	$0.6600 \times 1.12$ (5%)	5.0	5.0
$Z \rightarrow \mu\mu$	364382	SHERPA 2.2.5	$0.6600 \times 1.12$ (5%)	5.0	5.0
$Z \rightarrow \tau\tau$	364383	SHERPA 2.2.5	$0.6600 \times 1.12$ (5%)	1.5	1.5
$W \rightarrow e\nu$	364384	SHERPA 2.2.5	7.259 (5%)	25	25
$W \rightarrow \mu\nu$	364385	SHERPA 2.2.5	7.259 (5%)	25	25
$W \rightarrow \tau\nu$	364386	SHERPA 2.2.5	7.259 (5%)	6.0	6.0
$ZZ(4\ell)$	361063	SHERPA 2.1	0.004624 (10%)	0.017	0.049
$WZ(\ell\ell\ell^- \nu \text{SF})$	361064	SHERPA 2.1	0.0005324 (10%)	0.0073	0.015
$WZ(\ell\ell\ell^- \nu \text{OF})$	361065	SHERPA 2.1	0.001041 (10%)	0.012	0.030
$WZ(\ell\ell\ell^+ \nu \text{SF})$	361066	SHERPA 2.1	0.0008433 (10%)	0.010	0.020
$WZ(\ell\ell\ell^+ \nu \text{OF})$	361067	SHERPA 2.1	0.001633 (10%)	0.016	0.039
$WW(2\ell 2\nu)$	361068	SHERPA 2.1	0.003356 (10%)	0.068	0.090
$WW(q\bar{q}\ell\nu)$	361091	SHERPA 2.1	0.006059 (10%)	0.078	0.15
$WW(\ell\nu q\bar{q})$	361092	SHERPA 2.1	0.006082 (10%)	0.14	0.26
$WZ(\ell\nu q\bar{q})$	361093	SHERPA 2.1	0.002503 (10%)	0.039	0.075
$WZ(q\bar{q}\ell\ell)$	361094	SHERPA 2.1	0.0007518 (10%)	0.017	0.025
$ZZ(q\bar{q}\ell\ell)$	361096	SHERPA 2.1	$0.003789 \times 0.148$ (10%)	0.0070	0.010
$t\bar{t}$	410470	POWHEG+PYTHIA8	$0.06890 \times 0.544$ (7%)	1.8	2.8
$t(\text{s-channel.}t)$	410644	POWHEG+PYTHIA8	0.0005400 (10%)	0.028	0.050
$t(\text{s-channel.}\bar{t})$	410645	POWHEG+PYTHIA8	0.0002751 (10%)	0.028	0.050
$Wt$	410646	POWHEG+PYTHIA8	0.002990 (10%)	0.018	0.050
$W\bar{t}$	410647	POWHEG+PYTHIA8	0.002983 (10%)	0.019	0.050
$t(\text{t-channel.}t)$	410658	POWHEG+PYTHIA8	0.005414 (10%)	0.028	0.050
$t(\text{t-channel.}\bar{t})$	410659	POWHEG+PYTHIA8	0.002682 (10%)	0.028	0.050

**Table 14:** Monte Carlo samples at  $\sqrt{s} = 5\text{TeV}$ . The table follows the same format as Table 11. The MC equivalent luminosity  $N_{\text{evt}}^{\text{unskim}}/(\sigma \cdot \text{BR} \cdot \epsilon_{\text{filter}})$  is generally above  $2.5\text{fb}^{-1}$  for signal and significant backgrounds, the exception are Powheg  $W \rightarrow \tau\nu$  and  $Z \rightarrow \tau\tau$  samples, that have about  $0.20\text{fb}^{-1}$  and  $0.45\text{fb}^{-1}$  only.



**Figure 13:** Distributions for the 5 TeV (left) and 13 TeV (right) low- $\mu$  dataset(s) in a  $Z/\gamma^* \rightarrow \mu\mu$  (top row) and a  $Z/\gamma^* \rightarrow ee$  (bottom row) selection. The data (points) is compared to  $Z/\gamma^* \rightarrow \mu\mu$  or  $Z/\gamma^* \rightarrow ee$  signal MC, respectively. The distributions of the  $z$ -position of the primary vertex selected as the hard interaction are compared for the dataset(s) and the MC simulation before (“no  $z_{vtx}$  rwgt”, blue, only 13 TeV) and after reweighting (black). For the 13 TeV data the 2017 and 2018 data are shown separately and all distributions are (roughly) normalised to the same number of selected events in the 2017 dataset.

### 1.3 W analysis event selection and control plots

#### 1.3.1 Event selection

Both in case of 5 and 13 TeV events with  $W \rightarrow \ell\nu$  candidate were selected base on a single-lepton trigger requirement. The trigger for  $W \rightarrow e\nu$  event candidate HLT\_e15\_1h1oose\_nod0\_L1EM12 require at least one reconstructed electron with  $E_T$  larger than 15 GeV passing *loose* identification requirements. Candidates for  $W \rightarrow \mu\nu$  were triggered by HLT\_mu14 trigger requiring one muon with  $E_T$  larger than 14 GeV.

Events are required to contain exactly one lepton (muon or electron) candidate having  $p_T > 25\text{GeV}$ .

Electrons are required to have  $|\eta| < 2.47$  excluding transition region  $1.37 < |\eta| < 1.52$ . Muons Events with additional leptons of the same flavour with transverse momentum greater than 20 GeV satisfying some ID criteria are discarded, to better reject the  $Z$  background. The ID point is *medium* for the muon channel, and *loose* for the electron channel. There is no requirement on the number of leptons with different flavour than the channel under study.

To suppress background, in particular from multijet processes, events are required to have  $E_T^{\text{miss}}$  greater than 25 GeV. The  $W$  boson transverse mass  $m_T$  is demanded to be larger than 50 GeV. This transverse mass is defined as follows:

$$m_T = \sqrt{2p_T^\nu p_T^l (1 - \cos \Delta\phi^\nu)} \quad (1.1)$$

The tables 15,17,19,111 contain signal selection event yields for the  $W^\pm \rightarrow \ell^\pm \nu$  at  $\sqrt{s} = 5$  TeV low- $\mu$  dataset. Similarly the tables 16,18,110,112 contain the corresponding numbers for the 13 TeV low- $\mu$  dataset. Table 113 provides a comparison between observed and expected yields. Events denoted as  $W \rightarrow \ell \nu$  in the tables and the plots contain the sum of background events coming from  $W \rightarrow \tau \nu$  and other  $W$  leptonic decays other than the signal.

Cut	Data	Signal	$W^\pm \rightarrow \ell^\pm \nu$ BG	$Z \rightarrow \ell\ell$	Top	Diboson	Multijet
One electron	1993720	643610 $\pm$ 260	32940 $\pm$ 190	44338 $\pm$ 71	1754.4 $\pm$ 3.9	772.2 $\pm$ 3.7	-
Electron trig matched	1907724	612940 $\pm$ 250	30790 $\pm$ 190	42100 $\pm$ 69	1698.5 $\pm$ 3.8	741.1 $\pm$ 3.6	-
Isolation	1438941	610320 $\pm$ 250	30590 $\pm$ 190	41923 $\pm$ 69	1663.6 $\pm$ 3.8	722.5 $\pm$ 3.6	-
$p_T^e > 25\text{GeV}$	720284	482240 $\pm$ 220	14790 $\pm$ 130	31955 $\pm$ 53	1464.5 $\pm$ 3.5	592.1 $\pm$ 3.2	-
$E_T^{\text{miss}} > 25\text{GeV}$	440605	421510 $\pm$ 210	9650 $\pm$ 100	1336 $\pm$ 20	1223 $\pm$ 3.2	420.8 $\pm$ 2.4	-
$m_T > 50\text{GeV}$	430620	417430 $\pm$ 210	8800 $\pm$ 96	1047 $\pm$ 16	944.3 $\pm$ 2.9	373.5 $\pm$ 2.2	3030 $\pm$ 550

**Table 15:** Analysis cut flow for  $W^+ \rightarrow e^+ \nu$  5 TeV signal selection. Lepton  $p_T$  is required to be over 18 GeV before the final cut.

Cut	Data	Signal	$W^\pm \rightarrow \ell^\pm \nu$ BG	$Z \rightarrow \ell\ell$	Top	Diboson	Multijet
One electron	7915023	1797340 $\pm$ 390	92520 $\pm$ 270	147490 $\pm$ 140	63207 $\pm$ 89	3069 $\pm$ 63	-
Electron trig matched	7840239	1709140 $\pm$ 380	86370 $\pm$ 260	139760 $\pm$ 140	61110 $\pm$ 88	2967 $\pm$ 62	-
Isolation	5413483	1698430 $\pm$ 380	85560 $\pm$ 260	138890 $\pm$ 140	59834 $\pm$ 87	2939 $\pm$ 61	-
$p_T^e > 25\text{GeV}$	2452868	1342200 $\pm$ 330	44450 $\pm$ 190	106270 $\pm$ 110	53811 $\pm$ 82	2565 $\pm$ 58	-
$E_T^{\text{miss}} > 25\text{GeV}$	1275513	1136520 $\pm$ 310	28580 $\pm$ 150	8313 $\pm$ 46	45707 $\pm$ 75	1990 $\pm$ 53	-
$m_T > 50\text{GeV}$	1207776	1117560 $\pm$ 310	24760 $\pm$ 130	6443 $\pm$ 36	34580 $\pm$ 65	1718 $\pm$ 50	28000 $\pm$ 1800

**Table 16:** Analysis cut flow for  $W^+ \rightarrow e^+ \nu$  13 TeV signal selection. Lepton  $p_T$  is required to be over 18 GeV before the final cut.

### 1.3.2 $\sqrt{s} = 13$ TeV dataset control plots

Control plots for the 13 TeV low- $\mu$  dataset are provided here after applying all corrections described in section ??, and after applying the selection described above in this section. In each figure, the right(left)-hand column shows distributions for the  $W^+$  ( $W^-$ ) process. The top (bottom) row shows the muon (electron) decay channel. In the ratio panels, the grey band is the total systematic uncertainty,

Cut	Data	Signal	$W^\pm \rightarrow \ell^\pm \nu$ BG	$Z \rightarrow \ell\ell$	Top	Diboson	Multijet
One muon	2434459	760980 $\pm$ 280	35090 $\pm$ 200	37015 $\pm$ 82	2025.3 $\pm$ 4.1	864.7 $\pm$ 3.7	-
Muon trig matched	2353403	664100 $\pm$ 260	30610 $\pm$ 190	32554 $\pm$ 76	1725.6 $\pm$ 3.8	746.6 $\pm$ 3.4	-
Isolation	1186616	659200 $\pm$ 260	30400 $\pm$ 190	32303 $\pm$ 76	1574.6 $\pm$ 3.7	710.1 $\pm$ 3.3	-
$p_T^\mu > 25\text{GeV}$	632016	508270 $\pm$ 230	13900 $\pm$ 130	22556 $\pm$ 57	1335.3 $\pm$ 3.4	568.2 $\pm$ 2.9	-
$E_T^{\text{miss}} > 25\text{GeV}$	470856	442600 $\pm$ 210	8700 $\pm$ 100	9959 $\pm$ 31	1111.8 $\pm$ 3	424.5 $\pm$ 2.5	-
$m_T > 50\text{GeV}$	457053	438280 $\pm$ 210	7879 $\pm$ 97	9649 $\pm$ 27	879.7 $\pm$ 2.8	381.7 $\pm$ 2.3	720 $\pm$ 190

**Table 17:** Analysis cut flow for  $W^+ \rightarrow \mu^+ \nu$  5 TeV signal selection. Lepton  $p_T$  is required to be over 18 GeV before the final cut.

Cut	Data	Signal	$W^\pm \rightarrow \ell^\pm \nu$ BG	$Z \rightarrow \ell\ell$	Top	Diboson	Multijet
One muon	9570104	2100770 $\pm$ 410	83110 $\pm$ 270	2019400 $\pm$ 2200	71602 $\pm$ 94	3442 $\pm$ 63	-
Muon trig matched	9382783	1840550 $\pm$ 390	72820 $\pm$ 250	1750400 $\pm$ 2000	61519 $\pm$ 87	2956 $\pm$ 59	-
Isolation	3905612	1821750 $\pm$ 380	71780 $\pm$ 250	595700 $\pm$ 1100	56849 $\pm$ 84	2916 $\pm$ 59	-
$p_T^\mu > 25\text{GeV}$	1930655	1393330 $\pm$ 340	34470 $\pm$ 170	170840 $\pm$ 490	49338 $\pm$ 78	2471 $\pm$ 54	-
$E_T^{\text{miss}} > 25\text{GeV}$	1321407	1173860 $\pm$ 310	21450 $\pm$ 140	51090 $\pm$ 180	41956 $\pm$ 72	1930 $\pm$ 49	-
$m_T > 50\text{GeV}$	1244892	1153800 $\pm$ 310	18270 $\pm$ 130	38304 $\pm$ 81	32375 $\pm$ 63	1705 $\pm$ 44	9040 $\pm$ 800

**Table 18:** Analysis cut flow for  $W^+ \rightarrow \mu^+ \nu$  13 TeV signal selection. Lepton  $p_T$  is required to be over 18 GeV before the final cut.

Cut	Data	Signal	$W^\pm \rightarrow \ell^\pm \nu$ BG	$Z \rightarrow \ell\ell$	Top	Diboson	Multijet
One electron	1724472	374900 $\pm$ 200	24150 $\pm$ 160	41995 $\pm$ 70	1590.5 $\pm$ 2.9	684.8 $\pm$ 4	-
Electron trig matched	1645694	359010 $\pm$ 200	22070 $\pm$ 160	39854 $\pm$ 68	1539.9 $\pm$ 2.9	655.7 $\pm$ 3.9	-
Isolation	1176976	357660 $\pm$ 200	21920 $\pm$ 160	39686 $\pm$ 68	1504.6 $\pm$ 2.8	640.7 $\pm$ 3.8	-
$p_T^e > 25\text{GeV}$	529183	302070 $\pm$ 180	11920 $\pm$ 110	30214 $\pm$ 52	1330.8 $\pm$ 2.6	532.9 $\pm$ 3.5	-
$E_T^{\text{miss}} > 25\text{GeV}$	281957	266750 $\pm$ 170	8084 $\pm$ 90	1293 $\pm$ 20	1112.5 $\pm$ 2.4	380 $\pm$ 3	-
$m_T > 50\text{GeV}$	274329	264540 $\pm$ 170	7317 $\pm$ 84	994 $\pm$ 16	855.2 $\pm$ 2.1	338.1 $\pm$ 2.9	2400 $\pm$ 500

**Table 19:** Analysis cut flow for  $W^- \rightarrow e^- \nu$  5 TeV signal selection. Lepton  $p_T$  is required to be over 18 GeV before the final cut.

whilst the brown band adds the MC statistical uncertainty in quadrature on top of it. In regions of the distributions insensitive to the modelling of  $p_T^W$  there is generally good agreement between data and predictions. The bulk of the  $m_T$  distribution is a typical example of distribution that is mostly insensitive to the modeling of  $p_T^W$ . The  $u_T$  distribution is an exception, and it can therefore be concluded that the baseline simulation is not modeling  $p_T^W$  satisfactorily.

Cut	Data	Signal	$W^\pm \rightarrow \ell^\pm \nu$ BG	$Z \rightarrow \ell\ell$	Top	Diboson	Multijet
One electron	7471742	1323710 $\pm$ 330	78230 $\pm$ 230	140980 $\pm$ 140	61951 $\pm$ 86	3059 $\pm$ 58	-
Electron trig matched	7402574	1267710 $\pm$ 330	72240 $\pm$ 230	133580 $\pm$ 140	59950 $\pm$ 85	2968 $\pm$ 57	-
Isolation	4949352	1260540 $\pm$ 330	71550 $\pm$ 230	132740 $\pm$ 140	58689 $\pm$ 84	2937 $\pm$ 57	-
$p_T^e > 25\text{GeV}$	2113364	1053510 $\pm$ 300	39660 $\pm$ 160	101350 $\pm$ 110	52923 $\pm$ 79	2544 $\pm$ 53	-
$E_T^{\text{miss}} > 25\text{GeV}$	1008915	900640 $\pm$ 280	25900 $\pm$ 130	7954 $\pm$ 45	45065 $\pm$ 73	1962 $\pm$ 48	-
$m_T > 50\text{GeV}$	949362	887810 $\pm$ 270	22400 $\pm$ 120	6052 $\pm$ 35	34177 $\pm$ 64	1695 $\pm$ 44	27400 $\pm$ 2000

**Table 110:** Analysis cut flow for  $W^- \rightarrow e^- \nu$  13 TeV signal selection. Lepton  $p_T$  is required to be over 18 GeV before the final cut.

Cut	Data	Signal	$W^\pm \rightarrow \ell^\pm \nu$ BG	$Z \rightarrow \ell\ell$	Top	Diboson	Multijet
One muon	2075709	440560 $\pm$ 220	22510 $\pm$ 170	34440 $\pm$ 80	1835.6 $\pm$ 3.1	751.5 $\pm$ 3.3	-
Muon trig matched	2002955	383720 $\pm$ 200	19640 $\pm$ 160	30277 $\pm$ 75	1561.6 $\pm$ 2.9	648 $\pm$ 3.1	-
Isolation	883078	381010 $\pm$ 200	19450 $\pm$ 160	30046 $\pm$ 74	1411 $\pm$ 2.7	616.9 $\pm$ 2.9	-
$p_T^\mu > 25\text{GeV}$	426119	314370 $\pm$ 180	9370 $\pm$ 110	20749 $\pm$ 56	1202.1 $\pm$ 2.5	505 $\pm$ 2.5	-
$E_T^{\text{miss}} > 25\text{GeV}$	298992	276060 $\pm$ 170	5893 $\pm$ 89	8716 $\pm$ 29	1004.2 $\pm$ 2.3	372.6 $\pm$ 2	-
$m_T > 50\text{GeV}$	287870	273710 $\pm$ 170	5158 $\pm$ 82	8408 $\pm$ 26	788.2 $\pm$ 2	335.6 $\pm$ 1.9	760 $\pm$ 160

**Table 111:** Analysis cut flow for  $W^- \rightarrow \mu^- \nu$  5 TeV signal selection. Lepton  $p_T$  is required to be over 18 GeV before the final cut.

Cut	Data	Signal	$W^\pm \rightarrow \ell^\pm \nu$ BG	$Z \rightarrow \ell\ell$	Top	Diboson	Multijet
One muon	8773414	1518070 $\pm$ 360	64930 $\pm$ 230	201900 $\pm$ 2200	70580 $\pm$ 90	3230 $\pm$ 60	-
Muon trig matched	8597493	1322980 $\pm$ 330	56520 $\pm$ 210	1750300 $\pm$ 2000	60579 $\pm$ 84	2806 $\pm$ 56	-
Isolation	3298569	1310310 $\pm$ 330	55680 $\pm$ 210	593700 $\pm$ 1100	55949 $\pm$ 80	2751 $\pm$ 55	-
$p_T^\mu > 25\text{GeV}$	1561721	1069770 $\pm$ 300	28230 $\pm$ 150	166810 $\pm$ 490	48544 $\pm$ 75	2362 $\pm$ 52	-
$E_T^{\text{miss}} > 25\text{GeV}$	1030406	910150 $\pm$ 280	17380 $\pm$ 120	47370 $\pm$ 180	41259 $\pm$ 69	1842 $\pm$ 46	-
$m_T > 50\text{GeV}$	963568	896850 $\pm$ 270	14710 $\pm$ 110	34572 $\pm$ 80	31772 $\pm$ 61	1598 $\pm$ 43	9050 $\pm$ 620

**Table 112:** Analysis cut flow for  $W^- \rightarrow \mu^- \nu$  13 TeV signal selection. Lepton  $p_T$  is required to be over 18 GeV before the final cut.

Selection	Observed	Expected
5TeV $W^+ \rightarrow e^+ \nu$	430620	431620 $\pm$ 600
5TeV $W^+ \rightarrow \mu^+ \nu$	457053	457790 $\pm$ 300
5TeV $W^- \rightarrow e^- \nu$	274329	276450 $\pm$ 530
5TeV $W^- \rightarrow \mu^- \nu$	287870	289160 $\pm$ 250
13TeV $W^+ \rightarrow e^+ \nu$	1207776	1213000 $\pm$ 1800
13TeV $W^+ \rightarrow \mu^+ \nu$	1244892	1253490 $\pm$ 870
13TeV $W^- \rightarrow e^- \nu$	949362	979500 $\pm$ 2000
13TeV $W^- \rightarrow \mu^- \nu$	963568	988560 $\pm$ 690

**Table 113:** Observed and Expected yield comparison for all signal selections.




Figure 14:  $\Sigma E_T$  distribution in the muon and electron channel for the  $\sqrt{s} = 13$  TeV dataset.






Figure 15:  $\Sigma E_T$  distribution in the muon and electron channel for the  $\sqrt{s} = 13$  TeV dataset.







Figure 16:  $\vec{E}_T^{\text{miss}}$  distribution in the muon and electron channel for the  $\sqrt{s} = 13 \text{ TeV}$  dataset.




**Figure 17:** Transverse mass distribution of the W boson in the muon and electron channel for the  $\sqrt{s} = 13$  TeV dataset.



**Figure 18:** Lepton pseudorapidity distribution in the muon and electron channel for the  $\sqrt{s} = 13$  TeV dataset.




**Figure 19:** Lepton transverse momentum distribution in the muon and electron channel for the  $\sqrt{s} = 13$  TeV dataset.



**Figure 110:**  $W$  transverse momentum distribution in the muon and electron channel for the  $\sqrt{s} = 13$  TeV dataset.

1.3.3  $\sqrt{s} = 5$  TeV dataset control plots

Control plots for the 5 TeV low- $\mu$  dataset are provided here after applying all corrections described in section ??, and after applying the selection described above in this section. In each figure, the right(left)-hand column shows distributions for the  $W^+$  ( $W^-$ ) process. The top (bottom) row shows the muon (electron) decay channel. In the ratio panels, the grey band is the total systematic uncertainty, whilst the brown band adds the MC statistical uncertainty in quadrature on top of it. In regions of the distributions insensitive to the modelling of  $p_T^W$  there is generally good agreement between data and predictions. The bulk of the  $m_T$  distribution is a typical example of distribution that is mostly insensitive to the modeling of  $p_T^W$ . Compared to the 13 TeV situation, the  $u_T$  distribution seems to indicate that the baseline simulation models  $p_T^W$  more satisfactorily.



**Figure 111:**  $\Sigma E_T$  distribution in the muon and electron channel for the  $\sqrt{s} = 5$  TeV dataset.




Figure 112:  $\Sigma E_T$  distribution in the muon and electron channel for the  $\sqrt{s} = 5$  TeV dataset.









Figure 113:  $\vec{E}_T^{\text{miss}}$  distribution in the muon and electron channel for the  $\sqrt{s} = 5 \text{ TeV}$  dataset.




**Figure 114:** Transverse mass distribution of the W boson in the muon and electron channel for the  $\sqrt{s} = 5$  TeV dataset.



**Figure 115:** Lepton pseudorapidity distribution in the muon and electron channel for the  $\sqrt{s} = 5$  TeV dataset.



**Figure 116:** Lepton transverse momentum distribution in the muon and electron channel for the  $\sqrt{s} = 5$  TeV dataset.



**Figure 117:** W transverse momentum distribution in the muon and electron channel for the  $\sqrt{s} = 5$  TeV dataset.

## Bibliography

- [1] Jan Kretzschmar. *Samples and Physics modelling for low pile-up runs taken in 2017 and 2018*. Tech. rep. ATL-COM-PHYS-2019-075. Geneva: CERN, Feb. 2019. URL: <https://cds.cern.ch/record/2657141>.
- [2] Paolo Nason. “A New method for combining NLO QCD with shower Monte Carlo algorithms”. In: *JHEP* 11 (2004), p. 040. DOI: 10.1088/1126-6708/2004/11/040. arXiv: hep-ph/0409146.
- [3] Stefano Frixione, Paolo Nason, and Carlo Oleari. “Matching NLO QCD computations with Parton Shower simulations: the POWHEG method”. In: *JHEP* 11 (2007), p. 070. DOI: 10.1088/1126-6708/2007/11/070. arXiv: 0709.2092 [hep-ph].
- [4] Simone Alioli et al. “NLO vector-boson production matched with shower in POWHEG”. In: *JHEP* 0807 (2008), p. 060. DOI: 10.1088/1126-6708/2008/07/060. arXiv: 0805.4802 [hep-ph].
- [5] Simone Alioli et al. “A general framework for implementing NLO calculations in shower Monte Carlo programs: the POWHEG BOX”. In: *JHEP* 06 (2010), p. 043. DOI: 10.1007/JHEP06(2010)043. arXiv: 1002.2581 [hep-ph].
- [6] T. Sjöstrand, S. Mrenna, and P. Skands. “Brief Introduction to PYTHIA 8.1”. In: *Comput. Phys. Comm.* 178 (2008), p. 85. DOI: 10.1016/j.cpc.2008.01.036. arXiv: 0710.3820v1 [hep-ph].
- [7] ATLAS Collaboration. “Measurement of the  $Z/\gamma^*$  boson transverse momentum distribution in  $pp$  collisions at  $\sqrt{s} = 7$  TeV with the ATLAS detector”. In: *JHEP* 09 (2014), p. 145. DOI: 10.1007/JHEP09(2014)145. arXiv: 1406.3660 [hep-ex].
- [8] Piotr Golonka and Zbigniew Was. “PHOTOS Monte Carlo: A Precision tool for QED corrections in  $Z$  and  $W$  decays”. In: *Eur. Phys. J. C* 45 (2006), pp. 97–107. DOI: 10.1140/epjc/s2005-02396-4. arXiv: hep-ph/0506026.
- [9] Stefan Höche et al. “NLO matrix elements and truncated showers”. In: *JHEP* 1108 (2011), p. 123. DOI: 10.1007/JHEP08(2011)123. arXiv: 1009.1127 [hep-ph].
- [10] Richard D. Ball et al. “Parton distributions with LHC data”. In: *Nucl. Phys. B* 867 (2013), p. 244. DOI: 10.1016/j.nuclphysb.2012.10.003. arXiv: 1207.1303 [hep-ph].
- [11] ATLAS Collaboration. *The Pythia 8 A3 tune description of ATLAS minimum bias and inelastic measurements incorporating the Donnachie–Landshoff diffractive model*. ATL-PHYS-PUB-2016-017. 2016. URL: <https://cds.cern.ch/record/2206965>.
- [12] S. Catani and M. Grazzini. “An NNLO subtraction formalism in hadron collisions and its application to Higgs boson production at the LHC”. In: *Phys. Rev. Lett.* 98 (2007), p. 222002. DOI: 10.1103/PhysRevLett.98.222002. arXiv: hep-ph/0703012 [hep-ph].
- [13] S. Catani et al. “Vector boson production at hadron colliders: A Fully exclusive QCD calculation at NNLO”. In: *Phys. Rev. Lett.* 103 (2009), p. 082001. DOI: 10.1103/PhysRevLett.103.082001. arXiv: 0903.2120 [hep-ph].

- 232 [14] L.A. Harland-Lang, A. D. Martin, P. Motylinski, R. S. Thorne. “Parton distributions in the LHC  
 233 era: MMHT 2014 PDFs”. In: *Eur. Phys. J. C* 75.5 (2015), p. 204. doi: 10.1140/epjc/s10052-015-  
 234 3397-6. arXiv: 1412.3989 [hep-ph].
- 235 [15] ATLAS Collaboration. “Measurement of  $W^\pm$  and  $Z$ -boson production cross sections in  $pp$  col-  
 236 lisions at  $\sqrt{s} = 13$  TeV with the ATLAS detector”. In: *Phys. Lett. B* 759 (2016), p. 601. doi:  
 237 10.1016/j.physletb.2016.06.023. arXiv: 1603.09222 [hep-ex].
- 238 [16] ATLAS Collaboration. “Measurements of  $W$  and  $Z$  boson production in  $pp$  collisions at  $\sqrt{s} =$   
 239  $5.02$  TeV with the ATLAS detector”. In: *Eur. Phys. J. C* 79 (2019), p. 128. doi: 10.1140/epjc/  
 240 s10052-019-6622-x. arXiv: 1810.08424 [hep-ex].
- 241 [17] CMS Collaboration. “Measurement of the inclusive  $t\bar{t}$  cross section in  $pp$  collisions at  $\sqrt{s} = 5.02$   
 242 TeV using final states with at least one charged lepton”. In: *JHEP* 03 (2018), p. 115. doi: 10.1007/  
 243 JHEP03(2018)115. arXiv: 1711.03143 [hep-ex].