

Department of Economics, Delhi School of Economics, University of Delhi

Forecasting Macroeconomic Variables Using Large Datasets: Dynamic Factor Model versus Large-Scale BVARs

Author(s): RANGAN GUPTA and ALAIN KABUNDI

Source: *Indian Economic Review*, Vol. 46, No. 1 (January - June 2011), pp. 23-40

Published by: Department of Economics, Delhi School of Economics, University of Delhi; Springer

Stable URL: <https://www.jstor.org/stable/23266415>

Accessed: 07-10-2019 10:28 UTC

JSTOR is a not-for-profit service that helps scholars, researchers, and students discover, use, and build upon a wide range of content in a trusted digital archive. We use information technology and tools to increase productivity and facilitate new forms of scholarship. For more information about JSTOR, please contact support@jstor.org.

Your use of the JSTOR archive indicates your acceptance of the Terms & Conditions of Use, available at <https://about.jstor.org/terms>



Springer, Department of Economics, Delhi School of Economics, University of Delhi are collaborating with JSTOR to digitize, preserve and extend access to *Indian Economic Review*

Forecasting Macroeconomic Variables Using Large Datasets: Dynamic Factor Model versus Large-Scale BVARs

RANGAN GUPTA¹

*Department of Economics,
University of Pretoria,
Pretoria*

AND

ALAIN KABUNDI

*Department of Economics,
University of Johannesburg,
Johannesburg*

ABSTRACT

This paper uses two-types of large-scale models, namely the Dynamic Factor Model (DFM) and Bayesian Vector Autoregressive (BVAR) Models based on alternative hyperparameters specifying the prior, which accommodates 267 macroeconomic time series, to forecast key macroeconomic variables of a small open economy. Using South Africa as a case study and per capita growth rate, inflation rate, and the short-term nominal interest rate as our variables of interest, we estimate the two-types of models over the period 1980Q1 to 2006Q4, and forecast one- to four-quarters-ahead over the 24-quarters out-of-sample horizon of 2001Q1 to 2006Q4. The forecast performances of the two large-scale models are compared with each other, and also with unrestricted three-variable Vector Autoregressive (VAR) and BVAR models, with identical hyperparameter values as the large-scale BVARs. The results, based on the average Root Mean Squared Errors (RMSEs), indicate that the large-scale models are better-suited for forecasting the three macroeconomic variables of our choice, and amongst the two types of large-scale models, DFM holds the edge.

JEL Classification: C11, C13, C33, C53.

Keywords: Dynamic Factor Model, BVAR, Forecast Accuracy.

1. INTRODUCTION

This paper exploits information contained in a large cross-section of time series, 267 to be specific, to forecast three key macroeconomic variables, namely, per capita growth rate, Consumer Price Index (CPI) inflation, and 91 days Treasury Bill rate for the South African economy, using a Dynamic Factor Model (DFM) and alternative Bayesian Vector Autoregressive (BVAR) models, based on different values of the hyperparameters specifying

¹ Corresponding Author: University of Pretoria, Department of Economics, Pretoria, 0002, South Africa.
Email: Rangan.Gupta@up.ac.za.

the prior. The two types of model are first estimated over the period of 1980:01 to 2000:04 using quarterly data, and then used to generate one- to four-quarters-ahead out-of-sample forecasts over a 24 quarter forecasting horizon of 2001:01 to 2006:04. The performance of these two large-scale models are also compared with an unrestricted Vector Autoregressive (VAR) model and BVAR models, with identical hyperparameter values as the large-scale BVARs, but including only three variables that we are concerned about. At this stage, it must be emphasized that choice of South Africa, as our country of interest, emerges purely from the motivation of this study discussed below, and, hence, there is no reason that the current work cannot be conducted for any other economy, especially given the general nature of the econometric models we use here.

The main motivation for this current piece of work emanates from two recent studies carried out for the South African economy by Gupta and Kabundi (2008) and Das et al. (2008). Gupta and Kabundi (2008) used a DFM to forecast growth per capita, inflation based on Gross Domestic Product deflator and 91 days Treasury Bill rate. When the forecast performance of the model was compared with an unrestricted VAR, alternative BVARs and a New Keynesian Dynamic Stochastic General Equilibrium (DSGE) model, estimated using the three variables of interest, the authors found that DFM outperformed all the models in terms of forecasting the interest rate, while it did no worse than the VAR and the BVARs in forecasting the other two variables. On the other hand, Das et al. (2008), forecasted regional house price inflation in five major metropolitan areas of South Africa based on the same DFM developed by Gupta and Kabundi (2008), which now also included house price inflation of different house size category² for the five metropolises under consideration. When the authors compared the forecasting performance of the DFM with spatial and non-spatial BVARs, besides, an unrestricted VAR, based on only the house price inflation, the DFM was found to outperform the other models in 12 of the 15 cases. Both Gupta and Kabundi (2008) and Das et al. (2008) attribute the better performance of the DFM to its ability to efficiently handle large amounts of information, and, hence, its capability to forecast more accurately.

In such a backdrop, this paper, using the same panel of 267 time series as used by Gupta and Kabundi (2008), tries to check for the validity of such a claim, by comparing the forecasting performance of a DFM, for three variables, with that of BVAR models, which based on their estimation method³ can also accommodate a panel as large as the one used in the DFM. Our analysis is quite similar in the spirit of two recent studies by Banbura et al. (2007) and Bloor and Matheson (2008). Banbura et al. (2007), using a data set of 131 monthly series for the US, shows that large-scale VAR models with the Bayesian shrinkage set in relation to the cross-sectional dimension, is much better equipped in forecasting than small-scale monetary VARs. Bloor and Matheson (2008), while repeating

2 The houses were categorized as small, medium and large based on the square meters of area covered. See Burger and van Rensburg (2008), Gupta and Das (2008) and Das et al. (2008) for further details regarding the housing market in South Africa.

3 See Section 4 laying out the basics of the BVAR models for further details.

the same analysis for New Zealand obtain similar results.

Our study is more in line with the work of Bloor and Matheson (2008), in the sense that, just like them, we also consider an emerging economy. We also allow for different degree of interaction amongst domestic and foreign variables⁴, in our data, to account for the small open economy structure of South Africa having a role in the forecasting results, and, hence, incorporate a bit of theory, in these otherwise atheoretical models.⁵ Note, Bloor and Matheson (2008), assumed the foreign variables to be exogenous in the system. To our knowledge, this is the first attempt to use large-scale BVAR models, as an alternative to a DFM, to forecast key macroeconomic variables in South Africa.

In general, the motivation to use a large data set to forecast an economy originates not only from the fact that such data is now available at lower cost, but also because, and, perhaps, more importantly, the increased power of computation has facilitated in using such huge amount of information to estimate and forecast with econometric models. Traditional forecasting models such VAR, known for their better predictability, are unable to handle information contained in a large dataset without facing the degrees of freedom problems. Dynamic factor models (DFM) on the other hand can efficiently handle large amounts of information and therefore help improve the forecasting performance of econometric models. VAR that uses information contained in few fundamental variables seems limited and insufficient to mimic complex economic relations and forecast the future. Today modern econometricians have the ability to extract important information from a large dataset and accurately forecast the future. In addition, central bankers, policymakers, and academics agree that economic agents monitor hundreds of economic variables in their decision-making process (Bernanke and Boivin, 2003). As Stock and Watson (2005) agreeably put it, the DFM transforms the *curse of dimensionality* into *blessing of dimensionality*.

The original dynamic factor models of Sargent and Sims (1977), Geweke (1977), Chamberlain (1983) and Chamberlain, and Rothschild (1983) have been improved recently through advances in estimation techniques proposed by Stock and Watson (2002), Forni et al. (2005), and Kapetanios and Marcellino (2004). The success of DFM is due to the fact that only few extracted common factors can explain to a large extent the variation of variables in a large cross-section of time series. Several empirical researches provide evidence of improvement in forecasting performance of macroeconomic variables using factor analysis (Giannone and Matheson, 2007; Van Nieuwenhuyze, 2007; Cristadoro et al., 2005; Forni et al., 2005; Schneider and Spitzer, 2004, Kabundi, 2004; Forni et al., 2001; and Stock and Waston, 2002a, 2002b, 1999, 1991, and 1989).

Unlike the unrestricted VAR, the Bayesian VAR (BVAR) can be seen as a valid alternative to the DFM as it can equally accommodate large number of predictors without

4 See Section 3 containing the discussion of data for further details.

5 See Section 4 for further details.

facing the risk of losing degrees of freedom associated with unrestricted VAR. By imposing restrictions related to the distribution of coefficients, BVAR models avoid the over parameterization and over fitting of the model that are inherent in a VAR framework. Hence, a BVAR is based on fast exploration of the model space and, as part of the Bayesian methodology it does not need to rely on asymptotic theory as in the case of an unrestricted VAR. Furthermore, given their estimation procedures, the DFM and the BVAR are capable of forecasting simultaneously a large number of time series other than the key variables of interest. In other words, the two models are ideal in using large amount of information to forecast the economy, and, hence, could be considered to be on an even ground.

The remainder of the paper is organized as follows: The following section briefly discusses the DFM. Section 3 outlines the basics of the VAR and Minnesota-type BVARs. Section 4 discusses the data used to estimate the DFM and BVARs, while Section 5 presents the results from the forecasting exercise. Finally, section 6 concludes.

2. THE BASICS OF DFM

This study uses the Dynamic Factor Model (DFM) developed by Forni et al. (2005) to extract common components between macroeconomics series, and then these common components are used to forecast the three key macroeconomic variables. In the VAR models, since all variables are used in forecasting, the number of parameters to estimate depend on the number of variables n . With such a large information set, n , the estimation of a large number of parameters leads to a curse of dimensionality. The DFM uses information set accounted by few factors $q \ll n$, which transforms the curse of dimensionality into a blessing of dimensionality.

The DFM expresses individual times series as the sum of two unobserved components: a common component driven by a small number of common factors and an idiosyncratic component, which are specific to each variable. The relevance of the method is that the DFM is able to extract the few factors that explain the comovement of all South African macroeconomic variables. Forni et al. (2005) demonstrated that when the number of factors is small relative to the number of variables and the panel is heterogeneous, the factors can be recovered from the present and past observations.

Consider a $n \times 1$ covariance stationary process $Y_t = (y_{1t}, \dots, y_{nt})'$. Suppose that X_t is the standardized version of Y_t , i.e. X_t has a mean zero and a variance equal to one. Under DFM proposed by Forni, Hallin, Lippi, and Reichlin (2005) (henceforth FHLR) X_t is described by a factor model, it can be written as the sum of two orthogonal components:

$$x_{it} = b_i(L)f_t + \xi_{it} \quad \dots(1)$$

or, in vector notation:

$$X_t = B(L)f_t + \xi_t \quad \dots(2)$$

where f_t is a $q \times 1$ vector of dynamic factors, $B(L) = B_0 + B_1L + \dots + B_sL^s$ is an $n \times q$ matrix of factor loadings of order s , ξ_t is an $n \times 1$ vector of idiosyncratic components. The number of static factors r is computed as $r = q(s + 1)$. However, in more general framework $r \geq q$, instead of the more restrictive $r = q(s + 1)$. In a DFM, f_t and ξ_{it} are mutually orthogonal stationary process, while χ_{it} is the common component.

Since dynamic common factors are latent, they need to be estimated. Forni et al. (2005) estimate dynamic factors through the use of dynamic principal component analysis. It involves estimating the eigenvalues and eigenvectors decomposition of spectral density matrix of X_t , which is a generalization of the orthogonalization process in case of static principal components.⁶

FHLR is a weighted version of the principal components estimator by Stock and Watson (2002b) based on dynamic PCA, which exploits information of leading and lagging variables where time series are converted to the frequency domain. However dynamic PC is two-sided filter. This causes a problem at the end of the sample, making it difficult to estimate and forecast the common component since no future observations are available. FHLR solves this problem by proposing a two-step approach. In the first step, it relies on the dynamic approach in the estimation of the covariance matrices of the common and idiosyncratic component (at all leads and lags) through an inverse Fourier transform of the spectral density matrices. It involves estimating the eigenvalues and eigenvectors decomposition of spectral density matrix of X_t , $\hat{\Sigma}(\theta)$ which has rank q , corresponding to q largest eigenvalues. For each frequency, θ , the spectral density matrix of X_t , which is estimated using the frequency $-\pi < \theta < \pi$, can be decomposed into spectral densities of the common and the idiosyncratic component, $\Sigma(\theta) = \Sigma_\chi + \Sigma_\xi$. Hence, the spectral density matrix of common component $\hat{\Sigma}_\chi$ is estimated. In the second step, this information is used to compute the factor space by r linear combinations of X_t that maximizes the estimated contemporaneous covariance matrices, explained by the common factors, estimated in the first step. These r linear combinations are the solutions from a dynamic principal component problem and have the efficient property of reducing the idiosyncratic component in the common factor space to a minimum, by selecting the variables with the highest common/idiosyncratic variance ratio. Importantly, this one-

6. See Gupta and Kabundi (2008) for a detailed description of the model.

sided approach is only used to estimate and forecast the common component.

Following Boivin and Ng (2005) we represent the idiosyncratic errors as AR(p) processes. Therefore, the forecasting equation to predict y_t is given by:

$$y_{t+h} = \alpha_0 + \alpha_1(L)' \hat{F}_t + \alpha_2(L)' y_t + \varepsilon_{t+h} \quad \dots(3)$$

where h is the forecasting horizon, $\alpha_i(L)$ are lag polynomials, which can be estimated restrictedly using Bayesian approach or unrestrictedly.

3. ALTERNATIVE FORECASTING MODELS: VAR AND BVAR⁷

The Vector Autoregressive (VAR) model, though 'atheoretical', is particularly useful for forecasting purposes. An unrestricted VAR model, as suggested by Sims (1980), can be written as follows:

$$y_t = A_0 + A(L)y_t + \varepsilon_t \quad \dots(4)$$

where y is a $(n \times 1)$ vector of variables being forecasted; $A(L)$ is a $(n \times n)$ polynomial matrix in the backshift operator L with lag length p , i.e., $A(L) = A_1L + A_2L^2 + \dots + A_pL^p$; A_0 is a $(n \times 1)$ vector of constant terms, and ε is a $(n \times 1)$ vector of error terms. In our case, we assume that $\varepsilon \sim N(0, \sigma^2 I_n)$, where I_n is a $n \times n$ identity matrix.

Note that VAR model generally uses equal lag length for all the variables of the model. One drawback of VAR models is that many parameters need to be estimated, some of which may be insignificant. This problem of over parameterization, resulting in multicollinearity and a loss of degrees of freedom, leads to inefficient estimates and possibly large out-of-sample forecasting errors. One solution, often adapted, is simply to exclude the insignificant lags based on statistical tests. Another approach is to use a near VAR, which specifies an unequal number of lags for the different equations.

However, an alternative approach to overcoming this over parameterization, as described in Litterman (1981), Doan et al. (1984), Todd (1984), Litterman (1986), and Spencer (1993), is to use a BVAR model. Instead of eliminating longer lags, the Bayesian method imposes restrictions on these coefficients by assuming that they are more likely to be near zero than the coefficients on shorter lags. However, if there are strong effects from less important variables, the data can override this assumption. The restrictions are imposed by specifying normal prior distributions with zero means and small standard deviations for all coefficients with the standard deviation decreasing as the lags increase. The exception to this is that the coefficient on the first own lag of a variable has a mean

⁷ This section relies heavily on the discussion available on VAR and BVAR in Dua and Ray (1995), LeSage (1999), Gupta and Sichei (2006), Gupta (2006, 2007) and Gupta and Das (2008a,b).

of unity. Litterman (1981) used a diffuse prior for the constant. This is popularly referred to as the 'Minnesota prior' due to its development at the University of Minnesota and the Federal Reserve Bank at Minneapolis.

Formally, as discussed above, the means and variances of the Minnesota prior take the following form:

$$\beta_i \sim N(1, \sigma_{\beta_i}^2) \text{ and } \beta_j \sim N(0, \sigma_{\beta_j}^2) \quad \dots(5)$$

where β_i denotes the coefficients associated with the lagged dependent variables in each equation of the VAR, while β_j represents any other coefficient. In the belief that lagged dependent variables are important explanatory variables, the prior means corresponding to them are set to unity, given that, as seen from (5), the distribution *a priori* is centered around a random walk. However, for all the other coefficients, β_j 's, in a particular equation of the VAR, a prior mean of zero is assigned to suggest that these variables are less important to the model.

The prior variances $\sigma_{\beta_i}^2$ and $\sigma_{\beta_j}^2$, specify uncertainty about the prior means $\bar{\beta}_i = 1$, and $\bar{\beta}_j = 0$, respectively. Because of the over parameterization of the VAR, Doan et al. (1984) suggested a formula to generate standard deviations as a function of small numbers of hyperparameters: w , d , and a weighting matrix $f(i, j)$. This approach allows the forecaster to specify individual prior variances for a large number of coefficients based on only a few hyperparameters. The specification of the standard deviation of the distribution of the prior imposed on variable j in equation i at lag m , for all i, j and m , defined as $\sigma_{i,j,m}$, can be specified as follows:

$$\sigma_{i,j,m} = [w \times g(m) \times f(i, j)] \frac{\hat{\sigma}_i}{\hat{\sigma}_j} \quad \dots(6)$$

with $f(i, j) = 1$, if $i = j$ and k_{ij} otherwise, with $(0 \leq k_{ij} \leq 1)$, $g(m) = m^{-d}$, $d > 0$. Note that $\hat{\sigma}_i$ is the estimated standard error of the univariate autoregression for variable i . The ratio $\hat{\sigma}_i / \hat{\sigma}_j$ scales the variables to account for differences in the units of measurement and, hence, causes specification of the prior without consideration of the magnitudes of the variables. The term w indicates the overall tightness and is also the standard deviation on the first own lag, with the prior getting tighter as we reduce the value. The parameter $g(m)$ measures the tightness on lag m with respect to lag 1, and is assumed to have a harmonic shape with a decay factor of d , which tightens the prior on increasing lags. The parameter $f(i, j)$ represents the tightness of variable j in equation i relative to variable i , and by increasing the interaction, i.e., the value of k_{ij} , we can loosen the prior.⁸ Note, in

8 For an illustration, see Dua and Ray (1995).

the standard Minnesota-type prior, the overall tightness (w) takes the values of 0.1, 0.2 and 0.3, while, the lag decay (d) is generally chosen to be equal to 0.5, 1.0 and 2.0. The interaction parameter (k_{ij}) is set to = 0.5 by following Le Sage (1999) and Doan (2007). However, as Doan (2008) points out, for a system with more than six equations, this value of k_{ij} is too loose, and suggests the use of an asymmetric interaction parameter by accounting for the important and unimportant variables. Thus, given that, we have domestic as well as foreign and world variables within our dataset, and realizing the South Africa is a small open economy, and, hence, domestic variables would have minimal, if any, effect on foreign and world variables, while the latter set of variables is sure to have an influence on the South African variables, we use the following weighting scheme for k_{ij} . Borrowing from the BVAR models used for regional forecasting, involving both regional and national variables, and following Kinal and Ratner (1986) and Shoesmith (1992), the weight of a foreign or world variable in a foreign or world equation, as well as a domestic equation, is set at 0.6. The weight of a domestic variable in other domestic equation is fixed at 0.1 and that in a foreign or world equation at 0.01. Finally, the weight of the domestic variable in its own equation is 1.0. These weights are in line with Litterman's circle-star structure. Star (foreign or world) variables affect both star and circle (domestic) variables, while circle variables primarily influence only other circle variables.⁹ In addition to using values of 0.1, 0.2 and 0.3 for w , we follow Banbura et al. (2007), Bloor and Matheson (2008) and De Mol et al. (2008) in setting the value of the overall tightness parameter to obtain a desired average fit for the three variables of interest in the in-sample period (1980Q1 to 2000Q4). The optimal value of $w(Fit)$ ($=0.012$)¹⁰ obtained in this fashion is then retained for the entire evaluation period. Specifically, for a desired Fit , w is chosen as follows:

$$w(Fit) = \arg \min_w \left| Fit - \frac{1}{3} \sum_{i=1}^3 \frac{MSE_i^w}{MSE_i^0} \right| \quad \dots(7)$$

where $MSE_i^w = \sqrt{\frac{1}{T_0 - p - 1} \sum_{t=p}^{T_0-2} (y_{i,t+1|t}^w - y_{i,t+1})^2}$, i.e., the one-step-ahead mean squared error

(MSE) evaluated using the training sample $t = 1, \dots, T_0 - 1$, with T_0 being the beginning of the sample period and p being the order of the VAR, MSE_i^0 is the MSE of variable i with the prior restriction imposed exactly ($w=0$), while, the baseline Fit is defined as the average relative MSE from an OLS-estimated VAR containing the three variables, i.e.,:

9 We also experimented by assigning higher and lower interaction values, in comparison to those specified above, to the star variables in both the star and circle equations, but, the rank ordering of the alternative forecasts remained the same.

10 Note in this case, following Banbura et al. (2007) and Bloor and Matheson (2008), the value of $d=1$ and k_{ij} follows Litterman's circle-star approach. A higher value of $d = 2$ was also used, but the forecasting results were found to deteriorate.

$$Fit = \frac{1}{3} \sum_{i=1}^3 \frac{MSE_i^{\infty}}{MSE_i^0} \quad \dots(8)$$

Finally, once the priors have been specified, the alternative BVARs, whether based on the 3 variables or all the 267 variables (symmetric or asymmetric), are estimated using Theil's (1971) mixed estimation technique. Specifically, suppose we denote a single equation of the VAR model as: $y_t = XA + \varepsilon_t$, with $Var(\varepsilon_t) = \sigma^2 I$ and where the matrix X denotes the lagged values of $y_{i,t}$ and the vector A denotes the coefficients $a_{i,j,m}(l)$, with l being the lag operator, on the lagged values of $y_{i,t}$, then the stochastic prior restrictions for this single equation can be written as:

$$\begin{bmatrix} M_{111} \\ M_{112} \\ \cdot \\ \cdot \\ \cdot \\ M_{nnp} \end{bmatrix} = \begin{bmatrix} \sigma/\sigma_{111} & 0 & \cdot & \cdot & \cdot & 0 \\ 0 & \sigma/\sigma_{112} & 0 & \cdot & \cdot & 0 \\ \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\ 0 & \cdot & \cdot & \cdot & \cdot & 0 \\ 0 & 0 & \cdot & \cdot & 0 & \sigma/\sigma_{nnp} \end{bmatrix} \begin{bmatrix} a_{111} \\ a_{112} \\ \cdot \\ \cdot \\ \cdot \\ a_{nnp} \end{bmatrix} + \begin{bmatrix} u_{111} \\ u_{112} \\ \cdot \\ \cdot \\ \cdot \\ u_{nnp} \end{bmatrix} \quad \dots(9)$$

Note, $Var(u) = \sigma^2 I$ and the prior means M_{ijm} and the prior standard deviation σ_{ijm} take the forms shown in (5) and (6). With (9) written as:

$$r = RA + u \quad \dots(10)$$

and the estimates for a typical equation are derived as follows:

$$\hat{A} = (X'X + R'R)^{-1} (X'y_t + R'r) \quad \dots(11)$$

Essentially then, the method involves supplementing the data with prior information on the distribution of the coefficients. The number of observations and degrees of freedom are increased by one in an artificial way, for each restriction imposed on the parameter estimates. The loss of degrees of freedom due to over-parameterization associated with a classical VAR model is, therefore, not a concern in the BVARs. Note, we compute point forecasts using the posterior mean of the parameters. Specifically, the point estimate of the one-step-ahead forecast is computed as:

$$\hat{y}_{t+1/t}^{(w,mo)} = \hat{A}_0^{(w,mo)} + \hat{A}_1^{(w,mo)} y_t + \dots + \hat{A}_p^{(w,mo)} y_{t-p+1}, \text{ with the } h\text{-steps ahead forecasts being}$$

computed recursively, where $\hat{A}_i^{(w,d,k_y,mo)}$, $i=0,\dots,p$ is the posterior mean of the constant and the autoregressive coefficients corresponding to a model mo , given the hyperparameters of the models.

4. DATA

It is imperative in factor analysis framework to extract common components from a data rich environment. After extracting common components of per capita growth rate, inflation, and nominal interest rates of South Africa, we make out-of-sample forecast for one, two, three, and four quarters ahead.

The data set contains 267 quarterly series of South Africa, ranging from real, nominal, and financial sectors. We also have intangible variables, such as confidence indices, and survey variables. In addition to national variables, the paper uses a set of global variables such as commodity industrial inputs price index and crude oil prices. The data also comprises series of major trading partners such as Germany, the United Kingdom (UK), and the United States (US) of America. The in-sample period contains data from 1980Q1 to 2000Q4. All series are seasonally adjusted and covariance stationary. The more powerful DFGLS test of Elliott, Rothenberg, and Stock (1996), instead of the most popular ADF test, is used to assess the degree of integration of all series. All non-stationary series are made stationary through differencing. The Schwarz information criterion is used in selecting the appropriate lag length in such a way that no serial correction is left in the stochastic error term. Where there were doubts about the presence of unit root, the KPSS test proposed by Kwiatkowski, Phillips, Schmidt, and Shin (1992), with the null hypothesis of stationarity, was applied. All series are standardized to have a mean of zero and a constant variance. It must, however be pointed out that, non-stationarity is not an issue with the BVAR, since Sims et al. (1990) indicate that with the Bayesian approach entirely based on the likelihood function, the associated inference does not need to take special account of non-stationarity, because the likelihood function has the same Gaussian shape regardless of the presence of non-stationarity. Hence, for the sake of comparison amongst the VARs, both classical and Bayesian, we make no attempt to make the variables stationary, unlike in the DFM.¹¹ However, following Banbura et al. (2007) and Bloor and Matheson (2008), for the variables in the panel that are characterized by mean-reversion, we set a white-noise prior, i.e., $\bar{\beta}_i = 0$, otherwise, we impose the random walk prior implying that: $\bar{\beta}_i = 1$. The in-sample period contains data from 1980Q1 to 2000Q4, while the out-of-sample set is 2001Q1-2006Q4.¹²

There are various statistical approaches in determining the number of factors in the DFM. For example, Bai and Ng (2002) developed some criteria guiding the selection of the number of factors in large dimensional panels. The principal component analysis (PCA) can also be used in establishing the number of factors in the DFM. The PCA suggests that the selection of a number of factors q be based on the first eigenvalues of the spectral density matrix of X_t . Then, the principal components are added until the

¹¹ See Dua and Ray (1995) for further details.

¹² Details about data and their statistical treatment of the variables used to estimate the DFM are available upon request.

increase in the explained variance is less than a specific $\alpha = 0.05$. The Bai and Ng (2002) approach proposes five static factors, while Bai and Ng (2007) suggests two primitive or dynamic factors. Similar to the latter method, the principal component technique, as proposed by Forni et al. (2000) suggests two dynamic factors. The first two dynamic principal components explain approximately 99 percent of variation, while the eigenvalue of the third component is $0.005 < 0.05$.

5. EVALUATION OF FORECAST ACCURACY

Given the specification of the priors above, we estimate a VAR and the small- and large-scale BVARs (both symmetric and asymmetric) over the period 1980:01 to 2000:04, based on quarterly data. Then we compute the out-of-sample one- through four-quarters-ahead forecasts for the period 2001:Q1 to 2006:Q4, and compare the forecast accuracy, relative to that of the forecasts generated by the benchmark DFM model. The different types of VARs are estimated with 5 lags¹³ of each variable. The VAR and the BVARs, for an initial prior, are estimated for the period of 1980:Q1 to 2000:Q4 and, then, we forecast from 2001:Q1 through to 2006:Q4. Since we use five lags, the initial five quarters of the sample, 1980:Q1 to 1981:Q4, are used to feed the lags. We generate dynamic forecasts, as would naturally be achieved in actual forecasting practice. The models are re-estimated each quarter over the out-of-sample forecast horizon in order to update the estimate of the coefficients, before producing the 4-quarters-ahead forecasts. This iterative estimation and 4-steps-ahead forecast procedure was carried out for 24 quarters, with the first forecast beginning in 2001:Q1. This experiment produced a total of 24 one-quarter-ahead forecasts, 24 two-quarters-ahead forecasts, and so on, up to 24 4-step-ahead forecasts. The RMSEs¹⁴ for the 24, quarter 1 ahead to quarter 4 forecasts are then calculated for per capita growth, CPI inflation and short-term nominal interest rate. The values of the RMSE statistic for one- to four-quarters-ahead forecasts for the period 2001:Q1 to 2006:Q4 are then examined. The model, DFM or any of the VARs, which produces the lowest average value for the RMSE is selected, as the 'optimal' model for a specific variable.

To evaluate the accuracy of forecasts generated by the DFM, we need alternative forecasts. To make the RMSEs comparable with the DFM, we report the same set of

¹³ The choice of 5 lags is based on the unanimity of the sequential modified LR test statistic, Akaike information criterion (AIC), the final prediction error (FPE) criterion, Schwarz information criterion and the Hannan-Quinn (HQ) information criterion applied to a stable VAR estimated with the three variables of concern. Note, stability, as usual, implies that no roots were found to lie outside the unit circle.

¹⁴ Note that if A_{t+n} denotes the actual value of a specific variable in period $t + n$ and F_{t+n} is the forecast made in period

t for $t + n$, the RMSE statistic can be defined as: $\sqrt{\frac{1}{N} \sum (A_{t+n} - F_{t+n})^2} \times 100$. For $n = 1$, the summation runs from 2001:Q1 to 2006:Q4, and for $n = 2$, the same covers the period of 2001:Q2 to 2006:Q4, and so on.

statistics for the out-of-sample forecasts generated from an unrestricted classical VAR, the three-variable BVARs and the asymmetric BVARs based on 267 variables. In Tables 1 to 3, we compare the RMSEs of one- to four-quarters-ahead out-of-sample-forecasts for the period 2001:Q1 to 2006:Q4, generated by the abovementioned models. At this stage, a few words need to be said regarding the choice of the evaluation criterion for the out-of-sample forecasts generated from Bayesian models. As Zellner (1986) points out the “optimal” Bayesian forecasts will differ depending upon the loss function employed and the form of predictive probability density function. In other words, Bayesian forecasts are sensitive to the choice of the measure used to evaluate the out-of-sample forecast errors. However, Zellner (1986) points out that the use of the mean of the predictive probability density function for a series, is optimal relative to a squared error loss function and the Mean Squared Error (MSE), and, hence, the RMSE is an appropriate measure to evaluate performance of forecasts, when the mean of the predictive probability density function is used. This is exactly what we do below in Tables 1 through 3, when we use the average RMSEs over the one- to four-quarter-ahead forecasting horizon. The conclusions, regarding each of the three variables, based on the average one- to four-quarters-ahead RMSEs, from these tables can be summarized as follows:

TABLE 1
RMSE FOR PER CAPITA GROWTH RATE (2001:01-2006:04)

		1	2	3	4	Average
	DFM	0.4556	0.4856	0.5219	0.7869	0.5625
	VAR	0.2733	0.2167	0.1148	0.0599	0.1662
w=0.3,d=0.5	SBVAR	0.4473	0.2111	0.1985	0.0029	0.2149
	LBVAR	0.5013	0.1428	0.3264	0.3314	0.3255
w=0.2,d=1	SBVAR	0.4573	0.1857	0.1554	0.0154	0.2034
	LBVAR	0.4365	0.0004	0.2979	0.2987	0.2584
w=0.1,d=1	SBVAR	0.4498	0.1945	0.1481	0.0263	0.2047
	LBVAR	0.3776	0.0109	0.1693	0.2050	0.1907
w=0.2,d=2	SBVAR	0.4846	0.1562	0.2732	0.1277	0.2604
	LBVAR	0.3520	0.2204	0.2390	0.3031	0.2786
w=0.1,d=2	SBVAR	0.4579	0.1941	0.2198	0.1345	0.2516
	LBVAR	0.5013	0.1428	0.3264	0.3314	0.3255
w(Fit)=0.012, d=1	LBVAR	0.1181	0.0096	0.0886	0.0367	0.0632

Note: SBVAR: 3-Variable BVAR; LBVAR: Large Asymmetric BVAR

TABLE 2
RMSE FOR CPI INFLATION (2001:01-2006:04)

		1	2	3	4	Average
	DFM	0.0111	0.0112	0.0112	0.0112	0.0112
	VAR	0.1695	0.0826	1.1857	0.7337	0.5429
w=0.3,d=0.5	SBVAR	0.0437	0.0206	1.2979	0.7093	0.5179
	LBVAR	0.0680	0.2534	0.6183	0.5343	0.3685
w=0.2,d=1	SBVAR	0.5768	0.1676	1.6281	1.0162	0.8472
	LBVAR	0.2777	0.1419	1.1231	0.5669	0.5274
w=0.1,d=1	SBVAR	0.5339	0.2034	1.5672	1.0458	0.8376
	LBVAR	0.2395	0.2457	1.0539	0.7546	0.5734
w=0.2,d=2	SBVAR	1.0603	0.0546	1.9936	0.9676	1.0190
	LBVAR	0.7323	0.4874	1.2126	0.8873	0.8299
w=0.1,d=2	SBVAR	0.9343	0.1142	1.7622	1.0941	0.9762
	LBVAR	0.6059	0.5223	1.2071	0.7793	0.7786
w(Fit)=0.012, d=1	LBVAR	0.3247	0.2285	0.4098	0.3351	0.3245

Note: SBVAR: 3-Variable BVAR; LBVAR: Large Asymmetric BVAR

TABLE 3
RMSE FOR 91 DAYS TREASURY BILL RATE (2001:01-2006:04)

		1	2	3	4	Average
	DFM	0.0169	0.0121	0.0116	0.0104	0.0127
	VAR	0.0395	0.1756	0.6709	0.8999	0.4465
w=0.3,d=0.5	SBVAR	0.1262	0.2213	0.1173	0.3531	0.2045
	LBVAR	0.1728	0.1113	0.1610	0.1775	0.1557
w=0.2,d=1	SBVAR	0.1456	0.1901	0.2735	0.4678	0.2693
	LBVAR	0.1455	0.2013	0.1506	0.2412	0.1846
w=0.1,d=1	SBVAR	0.1957	0.1505	0.4869	0.4935	0.3317
	LBVAR	0.1252	0.1858	0.0823	0.1111	0.1261
w=0.2,d=2	SBVAR	0.1425	0.1981	0.2585	0.4579	0.2642
	LBVAR	0.1252	0.1858	0.0823	0.1111	0.1261
w=0.1,d=2	SBVAR	0.1837	0.1789	0.4377	0.4835	0.3209
	LBVAR	0.1518	0.2750	0.1266	0.1495	0.1757
w(Fit)=0.012, d=1	LBVAR	0.0431	0.1203	0.0135	0.0001	0.0442

Note: SBVAR: 3-Variable BVAR; LBVAR: Large Asymmetric BVAR

- i. Per Capita Growth Rate: The DFM is outperformed by all the VAR models, small or large-scale, classical or Bayesian. The three variable classical VAR performs better than all the small-scale BVARs, as well as, the large-scale BVARs, except for the asymmetric BVAR based on $d = 1$ and $w = 0.012$. Or in other words, the asymmetric BVAR based on a decay factor of 1 and an overall weight of 0.012, determined via the procedure outlined by Banbura *et al.* (2007), Bloor and Matheson (2008) and De Mol *et al.* (2008) is best-suited in forecasting per capita growth rate.
- ii. CPI Inflation: For CPI inflation, the DFM outperforms all the other models. Amongst the VARs, the small-scale classical VAR outperforms all the three-variable BVAR except for the case where the priors are most loose ($w = 0.3$ and $d = 0.5$). Further, except for three alternative specification of the prior, specifically when $w = 0.3$ and $d = 0.5$, $w = 0.2$ and $d = 1.0$, and $d = 1$ and $w = 0.012$ the VAR, in general, is also better suited than the large-scale asymmetric BVARs. But, overall amongst the VARs, it is the large-scale asymmetric BVAR model with $w = 0.012$, $d = 1.0$ that is best-suited for forecasting CPI inflation;
- iii. 91-days Treasury bill rate: As with the CPI inflation, the DFM stands out in forecasting the Treasury bill rate, when compared to other alternative models. Amongst the VARs, the small-scale classical VAR is outperformed by all the small-scale BVARs and the large-scale asymmetric BVARs. Further, as with the CPI inflation, it is the large-scale asymmetric BVAR model with $w = 0.012$, $d = 1.0$ that is best-suited for forecasting the short-term interest rate amongst the VARs.

So, to summarize, we find that the DFM outperforms the VARs, small or large-scale, classical or Bayesian, by quite a margin in terms of the average RMSEs for one- to four-quarters-ahead forecasts for the CPI inflation and the Treasury Bill rate, but, in turn, is outperformed by the large-scale asymmetric BVAR with $w = 0.012$ and $d = 1.0$. Furthermore, setting the DFM aside for a moment, we observe that the next best performing model for forecasting the CPI inflation and the Treasury bill rate is the large-scale asymmetric BVAR with a relatively tight prior.

6. CONCLUSIONS

This paper exploits information contained in a large cross-section of time series to forecast three key macroeconomic variables, namely, per capita growth rate, the Consumer Price Index (CPI) inflation, and the 91 days Treasury Bill rate for the South African economy, using a DFM and BVARs, based on alternative values of the hyperparameters specifying the prior. The two-types of models are first estimated over the period of

1980:01 to 2000:04 using quarterly data, and are then used to generate one- to four-quarters-ahead out-of-sample forecasts over a 24 quarter forecasting horizon of 2001:01 to 2006:04. The performance of these two large-scale models are also compared with each other and with an unrestricted Vector Autoregressive (VAR) model and BVAR models, with identical hyperparameter values as the large-scale BVARs, but including only the three variables we are concerned of. As stressed in the introduction, given the modeling strategies, the current study can be easily generalized to any other country and should not be dubbed as specific to the South African economy.

In general, we find that the DFM outperforms all the alternative forms of the VARs by quite a distance in terms of the average RMSEs for one- to four-quarters-ahead forecasts for the CPI inflation and the Treasury bill rate, but, in turn, is outperformed by a large-scale asymmetric BVAR with tight priors. But, besides the DFM, we observe that the next best performing model for forecasting the CPI inflation and the Treasury bill rate is the large-scale asymmetric BVAR with relatively tight priors. Based on these results, what is of more importance, is the observation that, whether it is the DFM or the BVAR, it is always a large-scale model that is best-suited for forecasting the three variables of our concern. But, it seems that, perhaps, the DFM might have an edge over the large-scale BVARs, especially when one realizes that there are at least two major limitations to using a Bayesian approach for forecasting. The two shortcomings of the Bayesian models are as follows: Firstly, as it is clear from Tables 1 to 3, the forecast accuracy is sensitive to the choice of the priors. So if the prior is not well specified, an alternative model used for forecasting may perform better. Secondly, in case of the Bayesian models, one requires to specify an objective function, for example the average MSE or RMSEs, to search for the 'optimal' priors, which, in turn, needs to be optimized over the period for which we compute the in-sample or the out-of-sample forecasts. However, there is no guarantee that the chosen parameter values specifying the prior will continue to be 'optimal' beyond the period for which it was selected. Nevertheless, the importance of large-scale BVARs cannot be ignored¹⁵, especially when one realizes that they are the best possible alternative to the DFM, as far as accommodating large number of time series is concerned. Further, it is also important to check for the robustness of our conclusions, by redoing the exercise with BVARs based on modified form(s) of the Minnesota prior, by allowing for inverse-Wishart, sums of coefficients, and co-persistence priors, as suggested by Robertson and Tallman (1999).

REFERENCES

- Bai, Jushan, and Ng, Serena (2002), Determining the Number of Factors in Approximate Factor Models, *Econometrica*, Vol.70, 191–221.
- Bai, Jushan, and Ng, Serena (2007), Determining the Number of Primitive Shocks in Factor Models, *Journal of Business and Economic Statistics*, Vol.25, 52–60.

¹⁵ Please refer to Banbura et al. (2007) and Bloor and Matheson (2008) for further details.

- Banbura, Marta, Gianonne, Domenico and Reichlin, Lucrezia (2007), Bayesian VARs with Large Panels, *Centre for Economic Policy Research (CEPR), Discussion Paper*, 6326.
- Bernanke, Ben, and Boivin, Jean (2003), Monetary Policy in a Data-Rich Environment, *Journal of Monetary Economics*, Vol.50, 525–546.
- Bloor, Chris and Matheson, Troy (2008), Analysing Shock Transmission in a Data-Rich Environment : A Large BVAR for New Zealand, *Reserve Bank of New Zealand, Discussion Paper Series DP2008/09*.
- Boivin, Jean & Ng, Serena (2005), Understanding and Comparing Factor-Based Forecasts, *International Journal of Central Banking*, Vol. 1, 117–151
- Burger, Philippe and Van Rensburg, Lizelle J. (2008), Metropolitan House Prices in South Africa: Do they Converge?, *South African Journal of Economics*, Vol.76, 291–297.
- Chamberlain, Gary (1983), Funds, Factors, and Diversification in Arbitrage Pricing Models, *Econometrica*, 5 Vol.1, 1281–1304.
- Chamberlain, Gary, and M. Rothschild (1983), Arbitrage, Factor Structure and Mean-Variance Analysis in Large Markets, *Econometrica*, Vol.51, 1305–1324.
- Cristadoro, Ricardo, Forni, Mario, Reichlin, Lucrezia, and Giovanni, Veronese (2005), A Core Inflation Indicator for the Euro Area, *Journal of Money, Credit and Banking*, Vol.37, 539–560.
- Das, Sonali, Gupta, Rangan and Kabundi, Alain (2008), Is a DFM Well-Suited for Forecasting Regional House Price Inflation?, Forthcoming *Journal of Forecasting*.
- De Mol, Christine, Giannone, Domenico and Reichlin, Lucrezia (2008), Forecasting Using a Large Number of Predictors: Is Bayesian Regression a Valid Alternative to Principal Components?, Forthcoming *Journal of Econometrics*.
- Doan, Thomas, Robert, A., Litterman, B. and Sims, Christopher A. (1984), Forecasting and Conditional Projections Using Realistic Prior Distributions, *Econometric Reviews*, Vol. 3, 1–100.
- Doan, Thomas, A (2007), *RATS Version 7.0 User's Guide*, ESTIMA.
- Dua, Pami, and Ray, Subhash C. (1995), A BVAR Model for the Connecticut Economy, *Journal of Forecasting*, Vol.14, 167–180.
- Forni, Mario, Hallin, Marc, Lippi, Marco, and Reichlin, Lucrezia (2000), The Generalized Dynamic Factor Model: Identification and Estimation, *Review of Economics and Statistics*, Vol.82, 540–554.
- Forni, Mario, Hallin, Marc, Lippi, Marco, and Reichlin, Lucrezia (2001), Coincident and Leading Indicators for the Euro Area, *The Economic Journal*, Vol. 111, 62–85.
- Forni, Mario, Hallin, Marc, Lippi, Marco, and Reichlin, Lucrezia (2005), The Generalized Dynamic Factor Model, One Sided Estimation and Forecasting, *Journal of the American Statistical Association*, Vol. 100, 830–840.
- Geweke, John (1977), The Dynamic Factor Analysis of Economic Time Series, in D.J. Aigner and A. Goldberger, eds. *Latent Variables in Socio-Economic Models*, 365–383, Amsterdam: North Holland.
- Giannone, Domenico, and Matheson, Troy D. (2007), A New Core Inflation Indicator for New Zealand, *International Journal of Central Banking*, Vol. 3, 145–180.

- Gupta, Rangan, and Sichei, Moses (2006), A BVAR Model for the South African Economy, *South African Journal of Economics*, Vol. 74, 391-409.
- Gupta, Rangan (2006), Forecasting the South African Economy with VARs and VECMs, *South African Journal of Economics*, Vol. 74, 611-628.
- Gupta, Rangan (2007), Forecasting the South African Economy with Gibbs Sampled BVECMs, *South African Journal of Economics*, Vol. 75, 631-643.
- Gupta, Rangan, and Das, Sonali (2008a), Spatial Bayesian Methods for Forecasting House Prices in Six Metropolitan Areas of South Africa, *South African Journal of Economics*, Vol. 76, 298-313.
- Gupta, Rangan (2008b), Bayesian Methods of Forecasting Inventory Investment in South Africa, Forthcoming *South African Journal of Economics*.
- Gupta, Rangan and Kabundi, Alain (2008), A Dynamic Factor Model for Forecasting Macroeconomic Variables in South Africa, Forthcoming *International Journal of Forecasting*.
- Kabundi, Alain (2004), Estimation of Economic Growth Using Business Survey Data, *International Monetary Fund*, Working Paper 04/69.
- Kinal, Tennessee, and Ratner, Jonathan (1986), A VAR Forecasting Model of a Regional Economy: Its Construction and Comparison, *International Regional Science Review*, Vol. 10, 113-126.
- Kwiatowski, Denis, Phillips, Peter C.B., Schmidt, Peter and Shin, Yongcheol (1992), Testing the Null Hypothesis of Stationarity Against the Alternative of a Unit Root: How Sure Are We That Economic Time Series Have a Unit Root?, *Journal of Econometrics*, Vol. 54, 159-178.
- LeSage, James P. (1990), A Comparison of the Forecasting Ability of ECM and VAR Models, *The Review of Economics and Statistics*, Vol. 72, 664-671.
- LeSage, James. P. (1999), *Applied Econometrics Using MATLAB*, www.spatial-econometrics.com.
- Litterman, Robert B. (1981), A Bayesian Procedure for Forecasting with Vector Autoregressions, *Working Paper*, Federal Reserve Bank of Minneapolis.
- Litterman, Robert B. (1986), Forecasting with Bayesian Vector Autoregressions – Five Years of Experience, *Journal of Business and Economic Statistics*, Vol. 4, 25-38.
- Robertson, John C., and Tallman, Ellis W. (1999), Vector Autoregressions: Forecasting and Reality, *Federal Reserve Bank of Atlanta, Economic Review*, First Quarter, 4-18.
- Sargent, Thomas J., and Sims, Christopher A. (1977), Business Cycle Modelling Without Pretending to Have Too Much A Priori Economic Theory, in C. Sims, eds. *New Methods in Business Research*, Federal Reserve Bank of Minneapolis.
- Shoosmith, Gary L. (1992), Co-integration, Error Correction and Medium-Term Regional VAR Forecasting, *Journal of Forecasting*, Vol. 11, 91-109.
- Sims, Christopher A., Stock, James H. and Watson, Mark W. (1990), Inference in Linear Time Series Models with Some Unit Roots, *Econometrica*, Vol. 58, 113-144.
- Sims, Christopher A. (1980), Macroeconomics and Reality, *Econometrica*, Vol. 48, 1-48.
- Spencer, David E. (1993), Developing a Bayesian Vector Autoregression Model, *International Journal of Forecasting*, Vol. 9, 407-421.
- Stock, James H., and Watson, Mark W. (1989), New Indexes of Coincident and Leading Economic Indicators, *NBER Macroeconomics Annual*, 351-393.

- Stock, James H., and Watson, Mark W. (1991), A Probability Model of the Coincident Indicators, in K. Lahiri, and G. Moore, eds. *Leading Economic Indicators: New Approaches and Forecasting Record*, 63–95, Cambridge: Cambridge University Press.
- Stock, James H., and Watson, Mark W. (1999), Forecasting Inflation, *Journal of Monetary Economics*, Vol. 44, 293–335.
- Stock, James H., and Watson, Mark W. (2002a), Forecasting Using Principal Components from a Large Number of Predictors, *Journal of the American Statistical Association*, Vol. 97, 147–162.
- Stock, James H., and Watson, Mark W. (2002b), Macroeconomic Forecasting Using Diffusion Indexes, *Journal of Business and Economics Statistics*, Vol. 20, 147–162.
- Stock, James H., and Watson, Mark W. (2005), Implications of Dynamic Factor Models for VAR Analysis, *NBER Working Paper 11467*.
- Theil, Henri (1971), *Principles of Econometrics*, New York, John Wiley.
- Todd, Richard M. (1984), Improving Economic Forecasting with Bayesian Vector Autoregression, *Quarterly Review*, Federal Reserve Bank of Minneapolis, Fall, 18–29.
- Van Nieuwenhuyze, Christophe (2007), A Generalized Dynamic Factor Model for the Belgian Economy Identification of the Business Cycle and GDP Growth Forecasts, *Journal of Business Cycle Measurement and Analysis*, Vol. 2, 213–248.
- Zellner, Arnold (1986), A Tale of Forecasting 1001 Series: The Bayesian Knight Strikes Again, *International Journal of Forecasting*, Vol. 2, 494–494.