

Geographic Variation in Opioid Mortality by Race/Ethnicity, 1999–2016

Identifying epidemic hotspots

Mathew Kiang¹ Monica Alexander² Zhe Zhang³ Jarvis Chen¹

¹Department of Social and Behavioral Sciences
Harvard TH Chan School of Public Health

²Department of Demography
University of California, Berkeley

³Heinz College
Carnegie Mellon University

PAA 2018: Denver, CO

Introduction

- Opioid-related mortality is a massive public health issue

Introduction

- Opioid-related mortality is a massive public health issue
 - Over 42,000 opioid-related deaths in 2016

Introduction

- Opioid-related mortality is a massive public health issue
 - Over 42,000 opioid-related deaths in 2016
 - More than deaths due to motor vehicle accidents

Introduction

- Opioid-related mortality is a massive public health issue
 - Over 42,000 opioid-related deaths in 2016
 - More than deaths due to motor vehicle accidents
 - More than deaths due to firearms

Introduction

- Opioid-related mortality is a massive public health issue
 - Over 42,000 opioid-related deaths in 2016
 - More than deaths due to motor vehicle accidents
 - More than deaths due to firearms
- More alarming than the magnitude is the trajectory:

Introduction

- Opioid-related mortality is a massive public health issue
 - Over 42,000 opioid-related deaths in 2016
 - More than deaths due to motor vehicle accidents
 - More than deaths due to firearms
- More alarming than the magnitude is the trajectory:
 - Current rate of increase similar to peak of HIV epidemic

Introduction

- Opioid-related mortality is a massive public health issue
 - Over 42,000 opioid-related deaths in 2016
 - More than deaths due to motor vehicle accidents
 - More than deaths due to firearms
- More alarming than the magnitude is the trajectory:
 - Current rate of increase similar to peak of HIV epidemic
 - “Triple-wave” epidemic¹

¹For examples, see: Unick *et al.* 2013; Compton *et al.* 2016; Ciccarone 2017

Introduction

- Opioid-related mortality is a massive public health issue
 - Over 42,000 opioid-related deaths in 2016
 - More than deaths due to motor vehicle accidents
 - More than deaths due to firearms
- More alarming than the magnitude is the trajectory:
 - Current rate of increase similar to peak of HIV epidemic
 - “Triple-wave” epidemic¹
 - Not one, but multiple, overlapping epidemics


¹For examples, see: Unick *et al.* 2013; Compton *et al.* 2016; Ciccarone 2017

Introduction


- Opioid-related mortality is a massive public health issue
 - Over 42,000 opioid-related deaths in 2016
 - More than deaths due to motor vehicle accidents
 - More than deaths due to firearms
- More alarming than the magnitude is the trajectory:
 - Current rate of increase similar to peak of HIV epidemic
 - “Triple-wave” epidemic¹
 - Not one, but multiple, overlapping epidemics
 - Variation by type of opioid

¹For examples, see: Unick *et al.* 2013; Compton *et al.* 2016; Ciccarone 2017


Increased over 2,500% since 1980




Wave 1: Prescription Opioids, 1990–2010



Wave 2: Heroin, 2010–current



Wave 3: Synthetic Opioids, 2013–current



Substantial variation: By race/ethnicity

Substantial variation: By geography

Aims of the paper

- 1 Systematically describe the opioid epidemic from 1999 to 2016 by state, race/ethnicity, and opioid type.

Aims of the paper

- 1 Systematically describe the opioid epidemic from 1999 to 2016 by state, race/ethnicity, and opioid type.
- 2 Systematically describe the *current* epidemic.

Aims of the paper

- ➊ Systematically describe the opioid epidemic from 1999 to 2016 by state, race/ethnicity, and opioid type.
- ➋ Systematically describe the *current* epidemic.
- ➌ Identify “epidemic hotspots” — areas with both high mortality rates and rapid increases in mortality.

- Multiple cause of death data from NCHS

- Multiple cause of death data from NCHS
 - 1999–2015, restricted-access

- Multiple cause of death data from NCHS
 - 1999–2015, restricted-access
 - 2016, public-access (pending request to restricted file)

- Multiple cause of death data from NCHS
 - 1999–2015, restricted-access
 - 2016, public-access (pending request to restricted file)
- Age-standardized rates by state, race/ethnicity, and opioid type

- Multiple cause of death data from NCHS
 - 1999–2015, restricted-access
 - 2016, public-access (pending request to restricted file)
- Age-standardized rates by state, race/ethnicity, and opioid type
 - Underlying cause: X40–X44, X60–X64, X85, Y10–Y14

- Multiple cause of death data from NCHS
 - 1999–2015, restricted-access
 - 2016, public-access (pending request to restricted file)
- Age-standardized rates by state, race/ethnicity, and opioid type
 - Underlying cause: X40–X44, X60–X64, X85, Y10–Y14
 - Contributory cause: T40.0–T40.4 and T40.6

- Multiple cause of death data from NCHS
 - 1999–2015, restricted-access
 - 2016, public-access (pending request to restricted file)
- Age-standardized rates by state, race/ethnicity, and opioid type
 - Underlying cause: X40–X44, X60–X64, X85, Y10–Y14
 - Contributory cause: T40.0–T40.4 and T40.6
- Joinpoint regression (stratified by race/ethnicity, state, opioid type)

Methods

- Multiple cause of death data from NCHS
 - 1999–2015, restricted-access
 - 2016, public-access (pending request to restricted file)
- Age-standardized rates by state, race/ethnicity, and opioid type
 - Underlying cause: X40–X44, X60–X64, X85, Y10–Y14
 - Contributory cause: T40.0–T40.4 and T40.6
- Joinpoint regression (stratified by race/ethnicity, state, opioid type)
 - Segmented weighted least squares regression

- Multiple cause of death data from NCHS
 - 1999–2015, restricted-access
 - 2016, public-access (pending request to restricted file)
- Age-standardized rates by state, race/ethnicity, and opioid type
 - Underlying cause: X40–X44, X60–X64, X85, Y10–Y14
 - Contributory cause: T40.0–T40.4 and T40.6
- Joinpoint regression (stratified by race/ethnicity, state, opioid type)
 - Segmented weighted least squares regression
 - Grid-search to find optimal joinpoint locations

Methods

- Multiple cause of death data from NCHS
 - 1999–2015, restricted-access
 - 2016, public-access (pending request to restricted file)
- Age-standardized rates by state, race/ethnicity, and opioid type
 - Underlying cause: X40–X44, X60–X64, X85, Y10–Y14
 - Contributory cause: T40.0–T40.4 and T40.6
- Joinpoint regression (stratified by race/ethnicity, state, opioid type)
 - Segmented weighted least squares regression
 - Grid-search to find optimal joinpoint locations
 - Permutation test to find optimal joinpoint model (0-3 joins)

Methods

- Multiple cause of death data from NCHS
 - 1999–2015, restricted-access
 - 2016, public-access (pending request to restricted file)
- Age-standardized rates by state, race/ethnicity, and opioid type
 - Underlying cause: X40–X44, X60–X64, X85, Y10–Y14
 - Contributory cause: T40.0–T40.4 and T40.6
- Joinpoint regression (stratified by race/ethnicity, state, opioid type)
 - Segmented weighted least squares regression
 - Grid-search to find optimal joinpoint locations
 - Permutation test to find optimal joinpoint model (0-3 joins)
- Defined statistical significance as $P < 0.01$

Example Joinpoint Results: Maryland

Example Joinpoint Results: All opioids

Current Mortality

Current Trajectory

Epidemic Hotspots

Results

- There were no decreases in opioid mortality, by type, for either race/ethnicity, anywhere

Results

- There were no decreases in opioid mortality, by type, for either race/ethnicity, anywhere
- Where there were increases, it is just as bad (or worse) for the black population

Results

- There were no decreases in opioid mortality, by type, for either race/ethnicity, anywhere
- Where there were increases, it is just as bad (or worse) for the black population
- Increases are driven by heroin and synthetic opioids

Results

- There were no decreases in opioid mortality, by type, for either race/ethnicity, anywhere
- Where there were increases, it is just as bad (or worse) for the black population
- Increases are driven by heroin and synthetic opioids
 - Heroin APC: 30-34% for both populations

Results

- There were no decreases in opioid mortality, by type, for either race/ethnicity, anywhere
- Where there were increases, it is just as bad (or worse) for the black population
- Increases are driven by heroin and synthetic opioids
 - Heroin APC: 30-34% for both populations
 - Synthetic opioid APC: 70% for whites, 150% for blacks

Results

- There were no decreases in opioid mortality, by type, for either race/ethnicity, anywhere
- Where there were increases, it is just as bad (or worse) for the black population
- Increases are driven by heroin and synthetic opioids
 - Heroin APC: 30-34% for both populations
 - Synthetic opioid APC: 70% for whites, 150% for blacks
 - Synthetic opioids are doubling in 12 states for whites and 18 for blacks

Results

- There were no decreases in opioid mortality, by type, for either race/ethnicity, anywhere
- Where there were increases, it is just as bad (or worse) for the black population
- Increases are driven by heroin and synthetic opioids
 - Heroin APC: 30-34% for both populations
 - Synthetic opioid APC: 70% for whites, 150% for blacks
 - Synthetic opioids are doubling in 12 states for whites and 18 for blacks
- Strong geographical clustering in heroin and synthetic opioid mortality

Results

- There were no decreases in opioid mortality, by type, for either race/ethnicity, anywhere
- Where there were increases, it is just as bad (or worse) for the black population
- Increases are driven by heroin and synthetic opioids
 - Heroin APC: 30-34% for both populations
 - Synthetic opioid APC: 70% for whites, 150% for blacks
 - Synthetic opioids are doubling in 12 states for whites and 18 for blacks
- Strong geographical clustering in heroin and synthetic opioid mortality
 - Little evidence this is due to differences in opioid use

Results

- There were no decreases in opioid mortality, by type, for either race/ethnicity, anywhere
- Where there were increases, it is just as bad (or worse) for the black population
- Increases are driven by heroin and synthetic opioids
 - Heroin APC: 30-34% for both populations
 - Synthetic opioid APC: 70% for whites, 150% for blacks
 - Synthetic opioids are doubling in 12 states for whites and 18 for blacks
- Strong geographical clustering in heroin and synthetic opioid mortality
 - Little evidence this is due to differences in opioid use
 - Source of heroin/synthetic opioids is important

Conclusions

- Interventions must be local and tailored to region, race/ethnicity, and opioid type

Conclusions

- Interventions must be local and tailored to region, race/ethnicity, and opioid type
- Supply-side interventions need to be balanced with harm reduction interventions

Conclusions

- Interventions must be local and tailored to region, race/ethnicity, and opioid type
- Supply-side interventions need to be balanced with harm reduction interventions
 - Syringe exchange programs

Conclusions

- Interventions must be local and tailored to region, race/ethnicity, and opioid type
- Supply-side interventions need to be balanced with harm reduction interventions
 - Syringe exchange programs
 - Medication assisted treatment

Conclusions

- Interventions must be local and tailored to region, race/ethnicity, and opioid type
- Supply-side interventions need to be balanced with harm reduction interventions
 - Syringe exchange programs
 - Medication assisted treatment
 - Increased naloxone access

Conclusions

- Interventions must be local and tailored to region, race/ethnicity, and opioid type
- Supply-side interventions need to be balanced with harm reduction interventions
 - Syringe exchange programs
 - Medication assisted treatment
 - Increased naloxone access
 - Drug testing on-site and point-of-site

Conclusions

- Interventions must be local and tailored to region, race/ethnicity, and opioid type
- Supply-side interventions need to be balanced with harm reduction interventions
 - Syringe exchange programs
 - Medication assisted treatment
 - Increased naloxone access
 - Drug testing on-site and point-of-site
 - Supervised consumption sites

Conclusions

- Interventions must be local and tailored to region, race/ethnicity, and opioid type
- Supply-side interventions need to be balanced with harm reduction interventions
 - Syringe exchange programs
 - Medication assisted treatment
 - Increased naloxone access
 - Drug testing on-site and point-of-site
 - Supervised consumption sites
- Surveillance of illicit markets needs to be dramatically improved

Conclusions

- Interventions must be local and tailored to region, race/ethnicity, and opioid type
- Supply-side interventions need to be balanced with harm reduction interventions
 - Syringe exchange programs
 - Medication assisted treatment
 - Increased naloxone access
 - Drug testing on-site and point-of-site
 - Supervised consumption sites
- Surveillance of illicit markets needs to be dramatically improved
 - Better measurement of potency of drugs

Conclusions

- Interventions must be local and tailored to region, race/ethnicity, and opioid type
- Supply-side interventions need to be balanced with harm reduction interventions
 - Syringe exchange programs
 - Medication assisted treatment
 - Increased naloxone access
 - Drug testing on-site and point-of-site
 - Supervised consumption sites
- Surveillance of illicit markets needs to be dramatically improved
 - Better measurement of potency of drugs
 - Types of drugs

Conclusions

- Interventions must be local and tailored to region, race/ethnicity, and opioid type
- Supply-side interventions need to be balanced with harm reduction interventions
 - Syringe exchange programs
 - Medication assisted treatment
 - Increased naloxone access
 - Drug testing on-site and point-of-site
 - Supervised consumption sites
- Surveillance of illicit markets needs to be dramatically improved
 - Better measurement of potency of drugs
 - Types of drugs
 - Cost and availability

Thank you

Code and interactive results explorer:

`https://tiny.cc/paa2018`

mathewkiang.com | : mkiang | : @mathewkiang

Epidemic Hotspots