

## **§5.1: CONSTRUCTING ACCURATE GRAPHS OF ANTIDERIVATIVES**

---

Dr. Mike Janssen

April 14, 2021

## ANNOUNCEMENTS

- CoL 3: November 23 (Monday before Thanksgiving)
- CoL 3 interviews: done over Zoom the last week of class

## A SUBTHEME OF THE COURSE

**Question:** Given the derivative of an unknown function  $f$ , how much information can we obtain about  $f$  itself?

## A SUBTHEME OF THE COURSE

**Question:** Given the derivative of an unknown function  $f$ , how much information can we obtain about  $f$  itself?

Some answers:

## A SUBTHEME OF THE COURSE

**Question:** Given the derivative of an unknown function  $f$ , how much information can we obtain about  $f$  itself?

Some answers:

Given real numbers  $a \leq b$ ,

$$f(b) = f(a) + \int_a^b f'(x) dx.$$

## ACTIVITY 5.1.2

## TAKEAWAY

If  $F$  and  $G$  are antiderivatives of the same function, then  $F - G$  must be a constant.

## TAKEAWAY

If  $F$  and  $G$  are antiderivatives of the same function, then  $F - G$  must be a constant.

**Example:** Consider  $f(x) = 2x$ .



## ACTIVITY 5.1.3

## FUNCTIONS DEFINED BY INTEGRALS

We can carry this further for continuous functions  $f$ :

$$A(x) = \int_a^x f(t) dt$$

is known as an **integral function**.

## FUNCTIONS DEFINED BY INTEGRALS

We can carry this further for continuous functions  $f$ :

$$A(x) = \int_a^x f(t) dt$$

is known as an **integral function**. What information does it give us?

## ACTIVITY 5.1.4