

§4.4: THE FUNDAMENTAL THEOREM OF CALCULUS

Dr. Mike Janssen

April 9, 2021

ANNOUNCEMENTS

-

PREVIEW ACTIVITY DISCUSSION

THE CONNECTION

Given a 1-dimensional velocity described by $v(t)$ corresponding to a displacement function $s(t)$, we have seen that the distance traveled between $t = a$ and $t = b$ is given by

$$s(b) - s(a)$$

THE CONNECTION

Given a 1-dimensional velocity described by $v(t)$ corresponding to a displacement function $s(t)$, we have seen that the distance traveled between $t = a$ and $t = b$ is given by

$$s(b) - s(a) = \int_a^b v(t) dt$$

THE CONNECTION

Given a 1-dimensional velocity described by $v(t)$ corresponding to a displacement function $s(t)$, we have seen that the distance traveled between $t = a$ and $t = b$ is given by

$$s(b) - s(a) = \int_a^b v(t) dt$$

Theorem (Fundamental Theorem of Calculus)

If f is continuous on $[a, b]$ and F is any antiderivative of f , then

$$\int_a^b f(x) dx = F(b) - F(a)$$

THE CONNECTION

Given a 1-dimensional velocity described by $v(t)$ corresponding to a displacement function $s(t)$, we have seen that the distance traveled between $t = a$ and $t = b$ is given by

$$s(b) - s(a) = \int_a^b v(t) dt$$

Theorem (Fundamental Theorem of Calculus)

If f is continuous on $[a, b]$ and F is any antiderivative of f , then

$$\int_a^b f(x) dx = F(b) - F(a) = F(x) \Big|_a^b$$

THE CONNECTION

Given a 1-dimensional velocity described by $v(t)$ corresponding to a displacement function $s(t)$, we have seen that the distance traveled between $t = a$ and $t = b$ is given by

$$s(b) - s(a) = \int_a^b v(t) dt$$

Theorem (Fundamental Theorem of Calculus)

If f is continuous on $[a, b]$ and F is any antiderivative of f , then

$$\int_a^b f(x) dx = F(b) - F(a) = F(x) \Big|_a^b$$

The Antiderivative Problem: given a function f , find all antiderivatives F of f .

ACTIVITY 4.4.2-4.4.3

THE TOTAL CHANGE THEOREM

In the context of an object moving with velocity v , the integral gives the total displacement of the object. That is, it gives the **total change** in the antiderivative.

THE TOTAL CHANGE THEOREM

In the context of an object moving with velocity v , the integral gives the total displacement of the object. That is, it gives the **total change** in the antiderivative.

Theorem (Total Change Theorem)

If f is continuously differentiable on $[a, b]$ with derivative f' then

$$f(b) - f(a) = \int_a^b f'(x) dx.$$

THE TOTAL CHANGE THEOREM

In the context of an object moving with velocity v , the integral gives the total displacement of the object. That is, it gives the **total change** in the antiderivative.

Theorem (Total Change Theorem)

If f is continuously differentiable on $[a, b]$ with derivative f' then

$$f(b) - f(a) = \int_a^b f'(x) dx.$$

That is, the definite integral of the rate of change of a function on $[a, b]$ is the total change of the function itself on $[a, b]$.

ACTIVITY 4.4.4