

## **§2.6: DERIVATIVES OF INVERSE FUNCTIONS**

---

Dr. Mike Janssen  
February 26, 2021

# ANNOUNCEMENTS

-

## PREVIEW ACTIVITY DISCUSSION

# INVERSE FUNCTIONS

**Notation:** by  $f : A \rightarrow B$  we mean that  $f$  is a function with *domain*  $A$  and *codomain*  $B$ .

# INVERSE FUNCTIONS

**Notation:** by  $f : A \rightarrow B$  we mean that  $f$  is a function with *domain*  $A$  and *codomain*  $B$ .

The **inverse of**  $f$  (if it exists) is a function  $g : B \rightarrow A$  such that  $g(f(a)) = a$  for every possible choice of  $a$  in the domain  $A$  and  $f(g(b)) = b$  for every possible choice of  $b$  in  $B$ .

# INVERSE FUNCTIONS

**Notation:** by  $f : A \rightarrow B$  we mean that  $f$  is a function with *domain*  $A$  and *codomain*  $B$ .

The **inverse of**  $f$  (if it exists) is a function  $g : B \rightarrow A$  such that  $g(f(a)) = a$  for every possible choice of  $a$  in the domain  $A$  and  $f(g(b)) = b$  for every possible choice of  $b$  in  $B$ .

We usually denote the inverse of  $f$  as  $f^{-1}$ . Observe that  $y = f(x)$  if and only if  $x = f^{-1}(y)$ .

# INVERSE FUNCTIONS

**Notation:** by  $f : A \rightarrow B$  we mean that  $f$  is a function with *domain*  $A$  and *codomain*  $B$ .

The **inverse of  $f$**  (if it exists) is a function  $g : B \rightarrow A$  such that  $g(f(a)) = a$  for every possible choice of  $a$  in the domain  $A$  and  $f(g(b)) = b$  for every possible choice of  $b$  in  $B$ .

We usually denote the inverse of  $f$  as  $f^{-1}$ . Observe that  $y = f(x)$  if and only if  $x = f^{-1}(y)$ .

**Big Important Fact:**  $f : A \rightarrow B$  has an inverse if and only if  $f$  passes the vertical and horizontal line tests on  $A$

## EXAMPLES OF INVERSES

On appropriate domains:

- $y = a^x$  if and only if  $x = \log_a(y)$
- $y = x^n$  if and only if  $x = \sqrt[n]{y}$
- $y = \sin x$  if and only if  $x = \sin^{-1} y = \arcsin y$
- etc



# THE NATURAL LOG AND ITS DERIVATIVE

Let  $f(x) = \ln x$ . Let's find  $f'(x)$ . We know  $e^{f(x)} =$

## ACTIVITY 2.6.2

## THE DERIVATIVE OF ARCSIN X

**Question:** On what domain is  $f(x) = \sin x$  invertible?

## ACTIVITIES 2.6.3-2.6.4

## CODA: THE INVERSE FUNCTION THEOREM

### **Theorem (Inverse Functions)**

*Suppose that  $f$  is differentiable with inverse  $g$  and that  $(x_0, y_0)$  is a point that lies on the graph of  $f$  at which  $f'(x_0) \neq 0$ . Then*

$$g'(y_0) = \frac{1}{f'(x_0)}$$

*More generally, for any  $z$  in the domain of  $g'$ , we have  $g'(z) = 1/f'(g(z))$ .*