

§5.2: THE SECOND FUNDAMENTAL THEOREM

Dr. Mike Janssen

April 16, 2021

ANNOUNCEMENTS

-

PREVIOUSLY

FTC:

$$\int_a^b f(x) dx = F(b) - F(a),$$

so:

PREVIOUSLY

FTC:

$$\int_a^b f(x) dx = F(b) - F(a),$$

so:

1. (\rightarrow) If we have a graph of f and we can compute the area bounded by f on $[a, b]$, we can compute the change in an antiderivative F over the interval
2. (\leftarrow) If we can find an algebraic formula for F , we can compute the net signed area bounded by f

erf

PREVIEW ACTIVITY DISCUSSION

FROM INTEGRAL FUNCTIONS TO FTC II

Theorem (Fundamental Theorem of Calculus II)

If f is a continuous function and c is any constant, then f has a unique antiderivative A that satisfies $A(c) = 0$, and that antiderivative is given by the rule

$$A(x) = \int_c^x f(t) dt.$$

ACTIVITY 5.2.2

UNDERSTANDING INTEGRAL FUNCTIONS

Suppose

$$F(x) = \int_c^x f(t) dt,$$

where $F(c) = 0$.

UNDERSTANDING INTEGRAL FUNCTIONS

Suppose

$$F(x) = \int_c^x f(t) dt,$$

where $F(c) = 0$. Then:

$$\frac{d}{dx}[F(x)] = \frac{d}{dx} \left[\int_c^x f(t) dt \right] = f(x).$$

UNDERSTANDING INTEGRAL FUNCTIONS

Suppose

$$F(x) = \int_c^x f(t) dt,$$

where $F(c) = 0$. Then:

$$\frac{d}{dx}[F(x)] = \frac{d}{dx} \left[\int_c^x f(t) dt \right] = f(x).$$

Thus, we can understand the graph of F via f , even if an algebraic formula for F eludes us.

ACTIVITY 5.2.3

INVERSE PROCESSES (ALMOST)

Recall that if

$$F(x) = \int_a^x f(t) dt,$$

where $F(a) = 0$, then:

$$\frac{d}{dx} \left[\int_a^x f(t) dt \right] = f(x).$$

INVERSE PROCESSES (ALMOST)

Recall that if

$$F(x) = \int_a^x f(t) dt,$$

where $F(a) = 0$, then:

$$\frac{d}{dx} \left[\int_a^x f(t) dt \right] = f(x).$$

On the other hand, what is

$$\int_a^x \frac{d}{dt} [f(t)] dt?$$

INVERSE PROCESSES (ALMOST)

Recall that if

$$F(x) = \int_a^x f(t) dt,$$

where $F(a) = 0$, then:

$$\frac{d}{dx} \left[\int_a^x f(t) dt \right] = f(x).$$

On the other hand, what is

$$\int_a^x \frac{d}{dt}[f(t)] dt?$$

Since f is an antiderivative of $\frac{d}{dt}[f(t)]$, we can apply FTC I:

$$\int_a^x \frac{d}{dt}[f(t)] dt = f(t) \Big|_a^x = f(x) - f(a).$$

INVERSE PROCESSES (ALMOST)

Recall that if

$$F(x) = \int_a^x f(t) dt,$$

where $F(a) = 0$, then:

$$\frac{d}{dx} \left[\int_a^x f(t) dt \right] = f(x).$$

On the other hand, what is

$$\int_a^x \frac{d}{dt}[f(t)] dt?$$

Since f is an antiderivative of $\frac{d}{dt}[f(t)]$, we can apply FTC I:

$$\int_a^x \frac{d}{dt}[f(t)] dt = f(t) \Big|_a^x = f(x) - f(a).$$

Thus, integration and differentiation are (almost) inverse processes, up to the constant $f(a)$.

ACTIVITY 5.2.4

INTEGRATION: THE BIG PICTURE

- Integrals give us net signed area of the functions we integrate over an interval
- **FTC I**: We can calculate exact values of the integral using any antiderivative
- **FTC II**: the integral function gives a unique antiderivative
- **FTC consequence**: integration and differentiation are **almost** inverse processes