

## **§1.7: LIMITS, CONTINUITY, AND DIFFERENTIABILITY**

---

Dr. Mike Janssen

February 1, 2021

## PREVIEW ACTIVITY DISCUSSION

## LEFT/RIGHT HAND LIMITS

One way a limit can fail to exist at a point is if the function seems to approach two different numbers depending on the side we're coming from.

## LEFT/RIGHTHAND LIMITS

One way a limit can fail to exist at a point is if the function seems to approach two different numbers depending on the side we're coming from.

### Definition

We say that  $f$  has limit  $L_1$  as  $x$  **approaches  $a$  from the left** and write

$$\lim_{x \rightarrow a^-} f(x) = L_1$$

provided we can make the value of  $f(x)$  as close as we like to  $L_1$  by taking  $x$  sufficiently close to  $a$  while always having  $x < a$ . We call  $L_1$  the **left-hand limit** of  $f$  as  $x$  approaches  $a$ .

## LEFT/RIGHTHAND LIMITS

One way a limit can fail to exist at a point is if the function seems to approach two different numbers depending on the side we're coming from.

### Definition

We say that  $f$  has limit  $L_1$  as  $x$  **approaches  $a$  from the left** and write

$$\lim_{x \rightarrow a^-} f(x) = L_1$$

provided we can make the value of  $f(x)$  as close as we like to  $L_1$  by taking  $x$  sufficiently close to  $a$  while always having  $x < a$ . We call  $L_1$  the **left-hand limit** of  $f$  as  $x$  approaches  $a$ . Similarly, we say that  $L_2$  is the **right-hand limit** of  $f$  as  $x$  approaches  $a$  and write

$$\lim_{x \rightarrow a^+} f(x) = L_2.$$

## EXAMPLE/BIG IMPORTANT FACT

**Example:**

110 110 (-100,-100)(100,100) (-100,-100)

## EXAMPLE/BIG IMPORTANT FACT

**Example:**

110 110 (-100,-100)(100,100) (-100,-100)

**Fact:**  $\lim_{x \rightarrow a} f(x) = L$  if and only if  $\lim_{x \rightarrow a^+} f(x) = \lim_{x \rightarrow a^-} f(x) = L$

## ACTIVITY 1.7.2

125 125 (-100,-100)(100,100) (-100,-100)



## Definition

A function  $f$  is **continuous at**  $x = a$  provided that

1.  $f$  has a limit as  $x \rightarrow a$ ,
2.  $f$  is defined at  $x = a$ , and
3.  $\lim_{x \rightarrow a} f(x) = f(a)$

# CONTINUITY

## Definition


A function  $f$  is **continuous at**  $x = a$  provided that

1.  $f$  has a limit as  $x \rightarrow a$ ,
2.  $f$  is defined at  $x = a$ , and
3.  $\lim_{x \rightarrow a} f(x) = f(a)$

What does this mean?

*cont. implies  $\lim = fcn$*

ACTIVITY 1.7.3



## THE RELATIONSHIP BETWEEN DIFFERENTIABILITY AND CONTINUITY

**Fact:** We have seen that a function  $f(x)$  can be continuous at  $x = a$  but fail to have  $f'(a)$  exist. That is, a function can be continuous without being differentiable.

What was that function  $f(x)$ ?

# THE RELATIONSHIP BETWEEN DIFFERENTIABILITY AND CONTINUITY

**Fact:** We have seen that a function  $f(x)$  can be continuous at  $x = a$  but fail to have  $f'(a)$  exist. That is, a function can be continuous without being differentiable.

What was that function  $f(x)$ ?

**Question:** Can a function be differentiable without being continuous?

# THE RELATIONSHIP BETWEEN DIFFERENTIABILITY AND CONTINUITY


**Fact:** We have seen that a function  $f(x)$  can be continuous at  $x = a$  but fail to have  $f'(a)$  exist. That is, a function can be continuous without being differentiable.

What was that function  $f(x)$ ?

**Question:** Can a function be differentiable without being continuous?

**No.**

ACTIVITY 1.7.4



# REMINDERS

-