

§1.8: THE TANGENT LINE APPROXIMATION

Dr. Mike Janssen

February 3, 2021

ANNOUNCEMENTS

-

PREVIEW ACTIVITY DISCUSSION

When $g(x) = -x^2 + 3x + 2$, $g'(x) = -2x + 3$. What do we need to write the equation of a line?

THE RETURN OF THE INFINITY PRINCIPLE

Infinity Principle: Chop a hard problem up into pieces that can be easily solved

THE RETURN OF THE INFINITY PRINCIPLE

Infinity Principle: Chop a hard problem up into pieces that can be easily solved

Question: what is $\sqrt{4.01}$? This is a hard problem!

THE RETURN OF THE INFINITY PRINCIPLE

Infinity Principle: Chop a hard problem up into pieces that can be easily solved

Question: what is $\sqrt{4.01}$? This is a hard problem!

IP: let's find the tangent line of $f(x)$ at $a = 2$ and find an approximate answer

LOCAL LINEARIZATION

- Equations of lines need a point and slope.

LOCAL LINEARIZATION

- Equations of lines need a point and slope.
- The slope of $f(x)$ at $(a, f(a))$ is $f'(a)$.

LOCAL LINEARIZATION

- Equations of lines need a point and slope.
- The slope of $f(x)$ at $(a, f(a))$ is $f'(a)$.
- The equation of the line tangent to $f(x)$ at $(a, f(a))$ is thus

$$y = f'(a)(x - a) + f(a)$$

LOCAL LINEARIZATION

- Equations of lines need a point and slope.
- The slope of $f(x)$ at $(a, f(a))$ is $f'(a)$.
- The equation of the line tangent to $f(x)$ at $(a, f(a))$ is thus

$$y = f'(a)(x - a) + f(a)$$

- Often called the **local linearization** and denoted by $L(x)$

EXAMPLE

Let's approximate $\sqrt{4.01}$.

$$f'(2) = 1/4$$

ACTIVITY 1.8.2

HOW GOOD AN APPROXIMATION IS THE LINEARIZATION?

We know $f(a) = L(a)$ and $f'(a) = L'(a)$. What about values of x near a ?

HOW GOOD AN APPROXIMATION IS THE LINEARIZATION?

We know $f(a) = L(a)$ and $f'(a) = L'(a)$. What about values of x near a ?

Consider the second derivative:

- $f''(a) > 0$:

HOW GOOD AN APPROXIMATION IS THE LINEARIZATION?

We know $f(a) = L(a)$ and $f'(a) = L'(a)$. What about values of x near a ?

Consider the second derivative:

- $f''(a) > 0$: f is concave up, so the tangent line lies entirely below the curve ($L(x) < f(x)$)

HOW GOOD AN APPROXIMATION IS THE LINEARIZATION?

We know $f(a) = L(a)$ and $f'(a) = L'(a)$. What about values of x near a ?

Consider the second derivative:

- $f''(a) > 0$: f is concave up, so the tangent line lies entirely below the curve ($L(x) < f(x)$)
- $f''(a) < 0$:

HOW GOOD AN APPROXIMATION IS THE LINEARIZATION?

We know $f(a) = L(a)$ and $f'(a) = L'(a)$. What about values of x near a ?

Consider the second derivative:

- $f''(a) > 0$: f is concave up, so the tangent line lies entirely below the curve ($L(x) < f(x)$)
- $f''(a) < 0$: f is concave down, so the tangent line lies entirely above the curve ($L(x) > f(x)$)

HOW GOOD AN APPROXIMATION IS THE LINEARIZATION?

We know $f(a) = L(a)$ and $f'(a) = L'(a)$. What about values of x near a ?

Consider the second derivative:

- $f''(a) > 0$: f is concave up, so the tangent line lies entirely below the curve ($L(x) < f(x)$)
- $f''(a) < 0$: f is concave down, so the tangent line lies entirely above the curve ($L(x) > f(x)$)
- $f''(a) = 0$:




HOW GOOD AN APPROXIMATION IS THE LINEARIZATION?

We know $f(a) = L(a)$ and $f'(a) = L'(a)$. What about values of x near a ?

Consider the second derivative:

- $f''(a) > 0$: f is concave up, so the tangent line lies entirely below the curve ($L(x) < f(x)$)
- $f''(a) < 0$: f is concave down, so the tangent line lies entirely above the curve ($L(x) > f(x)$)
- $f''(a) = 0$: the concavity is changing, and it depends on how (up to down, or down to up)

ACTIVITY 1.8.3



CLOSING REFLECTIONS ON CHAPTER 1

- We have spent three weeks exploring what the derivative *means*

CLOSING REFLECTIONS ON CHAPTER 1

- We have spent three weeks exploring what the derivative *means*
- This is incredibly important for critical thinking about derivatives. A computer can't think critically in the same way we can.

CLOSING REFLECTIONS ON CHAPTER 1

- We have spent three weeks exploring what the derivative *means*
- This is incredibly important for critical thinking about derivatives. A computer can't think critically in the same way we can.
- We're going to spend time starting Friday on derivative rules. There will be a lot of algebra, but the types of functions whose derivatives we can precisely calculate will expand a lot!

CLOSING REFLECTIONS ON CHAPTER 1

- We have spent three weeks exploring what the derivative *means*
- This is incredibly important for critical thinking about derivatives. A computer can't think critically in the same way we can.
- We're going to spend time starting Friday on derivative rules. There will be a lot of algebra, but the types of functions whose derivatives we can precisely calculate will expand a lot!
- Don't lose sight of what it all means. We'll try to sprinkle in some contextual problems as we go.