


1. A number line is shown.



Give the letter of the point that is closest to each value.

a. $2\pi - 9$

Point	C
--------------	---

b. $\sqrt{3}$

Point	T
--------------	---

c. $\sqrt[3]{9} - 8$

Point	L
--------------	---

2. A list of numbers is shown.

$$\frac{27}{4}, -3.2, -\sqrt[3]{60}, \sqrt{48}, 7.05$$

List the numbers from least to greatest.

Least to Greatest	$-\sqrt[3]{60}, -3.2, \frac{27}{4}, \sqrt{48}, 7.05$
--------------------------	--

3. For each equation, select all of the solutions.

		Solutions				
		$x = -4$	$x = 4$	$x = -8$	$x = 8$	not a real solution
a.	$x^2 = 64$	☐	☐	☒	☒	☐
b.	$x^3 = -64$	☒	☐	☐	☐	☐
c.	$x^2 = -64$	☐	☐	☐	☐	☒

4. Compare each pair of expressions using the symbols $<$, $>$, or $=$.

a. $8 - \pi$
☐ $<$
☐ $>$
☐ $=$
 $2(2.6)$

b. $\frac{9}{4}$
☐ $<$
☒ $>$
☐ $=$
 $\frac{3}{4} \cdot \sqrt[3]{8}$

c. $(4)^2$
☐ $<$
☐ $>$
☒ $=$
 $\sqrt{256}$

d. $\sqrt{78}$
☐ $<$
☐ $>$
☐ $=$
 3π

5. Complete the statements.

When x is any real number other than 0 and it is multiplied by itself, the

result will

☐ always

 be a

☐ positive

 value.

☐ sometimes

☐ negative

Therefore, equations in the form $x^2 = p$, where p is a whole number will

have

☐ only one solution.

☐ two solutions.

☐ three solutions.

6. A list of numbers is shown.

$$\frac{\sqrt{12}}{2}, \quad -0.1\sqrt{144}, \quad -\sqrt[3]{9}, \quad \sqrt{43}, \quad -0.23\pi, \quad \frac{57}{10}$$

Write each number in the appropriate column of the table based on the definitions of rational and irrational numbers.

Rational Number	Irrational Number
$-0.1\sqrt{144}, \quad \frac{57}{10}$	$\frac{\sqrt{12}}{2}, -\sqrt[3]{9}, \sqrt{43}, -0.23\pi$