


Algebra 1
Unit 3
Lesson 8 Practice

Name _____
Date _____
Period _____

A scatter plot representing the gold medal winning times for the mens' 100-meter freestyle swimming race since 1972 is shown.



The raw data used to construct the scatter plot is shown.


Year	1972	1976	1980	1984	1988	1992	1996
Winning Time (in seconds)	51.22	49.99	50.4	49.8	48.63	49.02	49.74
Year	2000	2004	2008	2012	2016	2020	
Winning Time (in seconds)	48.3	48.17	47.21	47.52	47.9	47.92	

The equation of the line of best fit that models this data is $f(x) = -0.07x + 50.597$, where f is the winning time of the mens' 100-meter freestyle even, and x is the number of years since 1972.

- How many years after 1972 was the 2012 Olympics?
- Using the equation of the line of best fit, find the predicted finishing time of the gold medalist in 2012.
- Identify the residual value of the point representing the 2012 winning time.

4. Identify one point on the scatter plot where the predicted value is above the actual value.
5. Find the residual value of the point selected for question 4.
6. Identify one point on the scatter plot where the predicted value is below the actual value.
7. Find the residual value of the point selected for question 6.
8. Does the data appear to have a strong linear correlation? Explain how you know.

The residual plot for the scatter plot is shown.



9. Explain how the residual values support the answer from question 8.