


MACHINE LEARNING DATASET OF SEP EVENTS FROM SOLAR CYCLES 22, 23 AND 24

Sumanth Rotti & Petrus Martens


Department of Physics and Astronomy, Georgia State University, Atlanta, GA

Data Preparation Schema



Sort and Classify GOES-05 to 15 EPS/EPEAD data

Visual Inspection





Reliable GOES Missions List


Yes



Description

- Period of study: 1986 to 2017
- Integral proton flux data from GOES-SEM^[1].
- Proton Channels: P2 to P7 (>10 MeV to >100 MeV)
- Visually inspected the temporal profiles for flux enhancements to identify which GOES satellite (primary or secondary) data has better sensory response.
- If the primary shows higher flux measurements compared to the secondary, then, the primary satellite data is considered as reliable. Else, the secondary.
- The process is repeated for all years to identify reliable GOES data.
- Linear imputation of data gaps.
- Homogenized the dataset across all GOES missions.

Events Slicing



Summary

- The GSEP catalog (Rotti et al., paper *in preparation*) integrates the events from Papaioannou et al. $(2016)^{[2]}$ and CDAW-SEP^[3] and cross-checked with the NOAA-SEP list^[4].
- There are 342 SEP events in the GSEP list. In that, 246 events have peak proton fluxes >10 pfu in the >10 MeV channel.
- The metadata consists of physical parameters and observables concerning the possible source solar eruptions such as flares and coronal mass ejections for each SEP event.
- The GSEP headers also carry relevant indicators (data quality, observed GOES instrument, and parallel reports.)
- The time series are sliced with respect to the event start and end times as reported in the GSEP metadata
 - O Start time: Event onset minus 12 hours
 - o End time: Observed event termination.
- The sliced dataset and the metadata will be publicly available on Harvard Dataverse.
- The purpose of this work is to provide a base for experimenting with statistical and machine learning models on SEP events.

Acknowledgment:

We acknowledge the use of data from NOAA-GOES missions and thank the team for the availability of particle data. We also thank the teams behind the SEP catalogs for the opportunity to utilize their work. SR thanks Dr. Manolis Georgoulis and Dr. Berkay Aydin for the many discussions during the project; and Dr. Hazel Bain of NOAA for information on the GOES primary and secondary observations. This work was supported by the NASA FINESST Grant 80NSSC21K1388.

References:

[1] Grubb, R.N., 1975. The SMS/GOES space environment monitor subsystem. *NASA STI/Recon Technical Report N*, 76, p.28260. [2] Papaioannou, A., Sandberg, I., Anastasiadis, A., Kouloumvakos, A., Georgoulis, M.K., Tziotziou, K., Tsiropoula, G., Jiggens, P. and Hilgers, A., 2016. Solar flares, coronal mass ejections and solar energetic particle event characteristics. *Journal of Space Weather and Space Climate*, 6, p.A42.

[3] https://cdaw.gsfc.nasa.gov/CME_list/sepe/ [4] https://umbra.nascom.nasa.gov/SEP/