


Bayesian polynomial chaos

A bayesian take on polynomial chaos — offering new and rigorous ways for engineers to interpret and work with simulation and measurement data.

1 Background & motivation

Polynomial chaos^{*} [1] seeks to quantify the impact of an **aleatory** uncertainty ρ in a model's inputs \mathbf{x} , on its outputs f . This is done by creating a polynomial surrogate of the model, where the basis terms are from a family of orthogonal polynomials (orthogonal to $\rho(\mathbf{x})$).



The model is evaluated at quadrature points (tensor, sparse and effectively subsampled in higher dimensions [2,3]), facilitating rapid estimates of the mean, variance and skewness in f based on the uncertainty.

While extremely successful — with significant industrial uptake — two limitations of the paradigm are:

- the inability to account for **epistemic** uncertainty in the model;
- the inability to exploit prior information or even experimental data (beyond those used as model boundary conditions).

2 Bayesian polynomial chaos

Bayesian polynomial chaos is a Gaussian process [4] with a mean, and two-point covariance function, both formed with orthogonal polynomials

$$g \sim \mathcal{N}(\mu_g(\mathbf{x}), \Sigma_g(\mathbf{x}, \mathbf{x}'))$$

MEAN
FUNCTION COVARIANCE
FUNCTION

where these quantities are given by

$$\mu_g(\mathbf{x}) = \mathbf{V}^T(\mathbf{x}) \boldsymbol{\mu}_\alpha$$


VANDERMONDE
TYPE MATRIX

$$\Sigma_g(\mathbf{x}, \mathbf{x}') = \mathbf{V}^T(\mathbf{x}) \boldsymbol{\Sigma}_\alpha \mathbf{V}^T(\mathbf{x}')$$

MEAN & COVARIANCE
OF COEFFICIENTS

Whilst this paradigm naturally addresses the epistemic uncertainty issue, it offers three key ideas.



- i. A **multi-fidelity framework**, where the posterior coefficients of a lower-fidelity polynomial, become the prior coefficients for a higher-fidelity polynomial.



- ii. Ability to **fuse additional linear operatic information**. (e.g., integrals and gradients) into the polynomial posterior, by creating a joint distribution, conditioning, and then marginalising over the additional information.
- iii. Ability to construct **coregional polynomials** for related outputs from the physics-based model — reducing number of model evaluations.

3 Preliminary results

A demonstration of the multi-fidelity framework is given below. The objective is to predict experimental surface integrated heat transfer (IHT), for variations in Mach number, Reynolds number and turbulence intensity for a VKI turbine blade [5].



Funding

Research supported through the Alan Turing Institute Strategic Priorities Fund. This fund is delivered by UK Research and Innovation, with this award managed by EPSRC (EP/Too1569/1). The authors also acknowledge the Lloyd's Register Foundation Data-Centric Engineering programme and Rolls-Royce plc.



Pranay Seshadri
Imperial College London;
The Alan Turing Institute;
Effective Quadratures.
psesh.com



Andrew Duncan
Imperial College London;
The Alan Turing Institute;
Effective Quadratures.




Ashley Scillitoe
The Alan Turing Institute;
Effective Quadratures.

Imperial College London

imperial.ac.uk

The Alan Turing Institute

turing.ac.uk



Effective Quadratures

effective-quadratures.org

References

- [1] Xiu & Karniadakis (2002). *SIAM J. Scientific Computing*. [Paper](#)
- [2] Constantine et al. (2012). *Computer Methods in App. Mechanics*. [Paper](#)
- [3] Seshadri et al. (2017). *SIAM J. Uncertainty Quantification*. [Paper](#)
- [4] Rasmussen and Williams (2005). *MIT Press*. [Book](#)
- [5] Arts et al. (1990). *Von Karman Institute for Fluid Dynamics*. [Report](#)