

Intro stats with mosaic

ggformula version

Essential R syntax

Names in R are *case sensitive*

Function and arguments

```
rflip(10)
```

Optional arguments

```
rflip(10, prob = 0.8)
```

Assignment

```
x <- rflip(10, prob = 0.8)
```

Getting help on any function

```
help(mean)
```

Loading packages

```
library(mosaic)
```

Arithmetic operations

+	-	*	/	basic operations
^				exponentiation
()				grouping
sqrt(x)				square root
abs(x)				absolute value
log10(x)				logarithm, base 10
log(x)				natural logarithm, base <i>e</i>
exp(x)				exponential function e^x
factorial(k)				$k! = k(k-1) \dots 1$

Logical operators

== is equal to (note double equal sign)

!= is not equal to

< is less than

<= is less than or equal to

> is greater than

>= is greater than or equal to

& **A & B** is **TRUE** if both **A** and **B** are **TRUE**

| **A | B** is **TRUE** if one or both of **A** and **B** are **TRUE**

%in% includes; for example

```
"C" %in% c("A", "B")
```

is **FALSE**

Formula interface

Use for graphics, statistics, inference, and modeling operations.

```
goal(y ~ x, data = mydata)
```

Read as "Calculate **goal** for **y** using **mydata** "broken down by" **x**, or "modeled by" **x**.

```
mean(age ~ sex, data = HELPrct)
```

For graphics:

```
goal(y ~ x | z, color = ~ w,
      data = mydata)
```


y : y-axis variable (*optional*)

x : x-axis variable (*required*)


z : panel-by variable (*optional*)

w : color-by variable (*optional*)


```
gf_boxplot(wage ~ sex,
            data = CPS85)
```



```
gf_point(wage ~ educ | sex,
         data = CPS85, color = "blue")
```



```
gf_point(wage ~ educ,
         color = ~ sex, data = CPS85)
```



Examining data

Print short summary of all variables

```
inspect(HELPrct)
```

Number of rows and columns

```
dim(HELPrct)
nrow(HELPrct)
ncol(HELPrct)
```

Print first rows or last rows

```
head(KidsFeet)
tail(KidsFeet, 10)
```

Names of variables

```
names(HELPrct)
```


One categorical variable

Counts by category

```
tally(~ sex, data = HELPrct)
```

Percentages by category

```
tally(~ sex, format =
      "percent", data = HELPrct)
gf_percents(~ sex, data =
            HELPrct, fill = "cyan",
            color = "black")
```



Tests and confidence intervals

Exact test

```
result1 <-
  binom.test(~ (homeless ==
                "homeless"), data = HELPrct)
```

Approximate test (large samples)

```
result2 <-
  prop.test(~ (homeless ==
                "homeless"), data = HELPrct)
```

Extract confidence intervals and *p*-values

```
confint(result1)
pval(result2)
```

One quantitative variable

Make output more readable

```
options(digits = 3)
```

Compute summary statistics

```
mean(~ cesd, data = HELPrct)
```

Other summary statistics work similarly

```
median() iqr() max() min()
fivenum() sd() var() sum()
```

Table of summary statistics

```
favstats(~ cesd, data = HELPrct)
```

Summary statistics by group


```
favstats(cesd ~ sex,
         data = HELPrct)
```

Quantiles

```
quantile(~ cesd, data = HELPrct,
         prob = c(0.25, 0.5, 0.8))
```

Histogram

```
gf_dhistogram(~ cesd, data =
              HELPrct, bins = 12)
```



Normal probability plot

```
gf_qq(~ cesd, data = HELPrct)
```

Density plot

```
gf_dens(~ cesd, data = HELPrct,
        color = "blue", size = 1.25)
```

One-sample *t*-test

```
result <- t.test(~ cesd,
                 mu = 34, data = HELPrct)
```

Extract confidence intervals and *p*-values

```
confint(result)
pval(result)
```

Two categorical variables

Contingency table with margins
`tally(~ substance + sex,
 margins = TRUE,
 data = HELPrct)`

Percentages by column
`tally(~ sex | substance,
 format = "percent",
 data = HELPrct)`

Mosaic plot
`mosaicplot(~ substance + sex,
 color = TRUE, data = HELPrct)`



Chi-square test
`xchisq.test(~ substance + sex,
 data = HELPrct,
 correct = FALSE)`

Distributions

Normal distribution function
`pnorm(13, mean = 10, sd = 2)`

Normal distribution function with graph
`xpnorm(1.645, mean = 0, sd = 1)`

Normal distribution quantiles
`qnorm(0.95) # mean = 0, sd = 1`

Normal distribution quantiles with graph
`xqnorm(0.85, mean = 10, sd = 2)`

Binomial density function ("size" means n)
`dbinom(5, size = 8, prob = 0.65)`

Binomial distribution function
`pbinom(5, size = 8, prob = 0.65)`


Central portion of distribution
`cdist("norm", 0.95)
 cdist("t", c(0.90, 0.99), df = 5)`

Plotting distributions
`plotDist("binom", size = 8,
 prob = 0.65, xlim = c(-1, 9))
 plotDist("norm", mean = 10,
 sd = 2)`

Two quantitative variables

Correlation coefficient
`cor(cesd ~ mcs, data = HELPrct)`

Scatterplot with regression line and smooth
`gf_point(cesd ~ mcs,
 data = HELPrct) %>%
 gf_smooth(linetype = "dashed",
 color = "red") %>%
 gf_lm(size = 1.5)`



Simple linear regression
`cesdmodel <- lm(cesd ~ mcs,
 data = HELPrct)
 msummary(cesdmodel)`

Prediction
`lm_fun <- makeFun(cesdmodel)
 lm_fun(mcs = 35)`

Extract useful quantities
`anova(cesdmodel)
 coef(cesdmodel)
 confint(cesdmodel)
 rsquared(cesdmodel)`

Diagnostics; plot residuals
`gf_dhistogram(~resid(cesdmodel)
 gf_qq(~resid(cesdmodel))`

Diagnostics; plot residuals vs. fitted
`gf_point(resid(cesdmodel) ~
 fitted(cesdmodel)) %>%
 gf_lm(size = 2)`

Categorical response, quantitative predictor

Logistic regression
`logit_mod <-
 glm(homeless ~ age,
 family = binomial, data = HELPrct)
 msummary(logit_mod)`

Odds ratios and confidence intervals
`exp(coef(logit_mod))
 exp(confint(logit_mod))`

Data manipulation

From `dplyr` package
 For details, see [Tidyverse cheatsheet](#)

Drop, rename, or reorder variables
`select()`

Create new variables from existing ones
`mutate()`

Retain specific rows from data
`filter()`

Sort data rows
`arrange()`

Compute summary statistics by group
`group_by()
 summarize()`

Importing data

Import data from file or URL
`MustangPrice <-
 read.file("C:/MustangPrice.csv")
 # NOTE: R uses forward slashes!
 Dome <-
 read.file("http://www.mosaic-
 web.org/go/datasets/Dome.csv")`

Randomization and simulation

Fix random number sequence
`set.seed(42)`

Toss coins
`rflip(10) # default prob is 0.5`

Do something repeatedly
`do(5) * rflip(10, prob = 0.75)`

Draw a simple random sample
`sample(LETTERS, 10)
 deal(Cards, 5) # poker hand`

Resample with replacement
`Small <- sample(KidsFeet, 10)
 resample(Small)`

Random permutation (shuffling)
`shuffle(Cards)`

Random values from distributions
`rbinom(5, size = 10, prob = 0.7)
 rnorm(5, mean = 10, sd = 2)`

Quantitative response, categorical predictor


Two-level predictor: two-sample t test

Numeric summaries

`favstats(~length | sex,
 data = KidsFeet)`

Graphic summaries

`gf_qq(~ length | sex,
 data = KidsFeet) %>%
 gf_labs(x = "Normal quantile",
 y = "Length (cm)") %>%
 gf_qqline()`



`gf_boxplot(cesd ~ substance,
 data = HELPrct)`

Two-sample t -test and confidence interval

`result <- t_test(cesd ~ sex,
 data = HELPrct)
 result # view results
 confint(result)`

More than two levels (Analysis of variance)

Numeric summaries

`favstats(cesd ~ substance,
 data = HELPrct)`

Fit and summarize model

`modsubstance <- lm(cesd ~ substance,
 data = HELPrct)
 anova(modsubstance)`

Which differences are significant?

`pairwise <- TukeyHSD(modsubstance)
 mplot(pairwise)`

