

# On line-parallelisms of $\text{PG}(3, q)$

Francesco Pavese

(joint work with P. Santonastaso)

Polytechnic University of Bari – Italy

*Finite Geometries 2025*

*Seventh Irsee Conference*

$q = p^h$  prime power

$q = p^h$  prime power

4-dimensional vector space over  $\mathbb{F}_q$

$q = p^h$  prime power

4-dimensional vector space over  $\mathbb{F}_q$

$\text{PG}(3, q)$  projective space

$q = p^h$  prime power  
4-dimensional vector space over  $\mathbb{F}_q$   
 $\text{PG}(3, q)$  projective space

line-spread  $\mathcal{S}$

$q = p^h$  prime power  
4-dimensional vector space over  $\mathbb{F}_q$   
 $\text{PG}(3, q)$  projective space

line-spread  $S$

*mutually skew lines partitioning the points of  $\text{PG}(3, q)$*

$q = p^h$  prime power  
4-dimensional vector space over  $\mathbb{F}_q$   
 $\text{PG}(3, q)$  projective space

line-spread  $\mathcal{S}$

*mutually skew lines partitioning the points of  $\text{PG}(3, q)$*

$$|\mathcal{S}| = q^2 + 1$$

$q = p^h$  prime power  
4-dimensional vector space over  $\mathbb{F}_q$   
 $\text{PG}(3, q)$  projective space

line-spread  $\mathcal{S}$

*mutually skew lines partitioning the points of  $\text{PG}(3, q)$*

$$|\mathcal{S}| = q^2 + 1$$

parallelism  $\Pi$



$q = p^h$  prime power  
4-dimensional vector space over  $\mathbb{F}_q$   
 $\text{PG}(3, q)$  projective space

line-spread  $\mathcal{S}$

*mutually skew lines partitioning the points of  $\text{PG}(3, q)$*

$$|\mathcal{S}| = q^2 + 1$$

parallelism  $\Pi$

*mutually skew line-spreads partitioning the lines of  $\text{PG}(3, q)$*

$q = p^h$  prime power  
4-dimensional vector space over  $\mathbb{F}_q$   
 $\text{PG}(3, q)$  projective space

line-spread  $\mathcal{S}$

*mutually skew lines partitioning the points of  $\text{PG}(3, q)$*

$$|\mathcal{S}| = q^2 + 1$$

parallelism  $\Pi$

*mutually skew line-spreads partitioning the lines of  $\text{PG}(3, q)$*

$$|\Pi| = q^2 + q + 1$$

# Motivations

*Kirkman's schoolgirl problem (1850)*

# Motivations

*Kirkman's schoolgirl problem (1850)*

*Finite Geometry*

# Motivations

*Kirkman's schoolgirl problem (1850)*

*Finite Geometry*

- N.L. Johnson, *Combinatorics of Spreads and Parallelisms*, Chapman & Hall, 2010.

# Motivations

*Kirkman's schoolgirl problem (1850)*

*Finite Geometry*

- N.L. Johnson, *Combinatorics of Spreads and Parallelisms*, Chapman & Hall, 2010.

*Design Theory*

# Motivations

*Kirkman's schoolgirl problem (1850)*

*Finite Geometry*

- N.L. Johnson, *Combinatorics of Spreads and Parallelisms*, Chapman & Hall, 2010.

*Design Theory*

*resolution of the block design formed by points and lines of*  
 $\text{PG}(3, q)$

# Motivations

*Kirkman's schoolgirl problem (1850)*

*Finite Geometry*

- N.L. Johnson, *Combinatorics of Spreads and Parallelisms*, Chapman & Hall, 2010.

*Design Theory*

*resolution of the block design formed by points and lines of*  
 $\text{PG}(3, q)$

*Coding Theory*



# Motivations

## *Kirkman's schoolgirl problem (1850)*

### *Finite Geometry*

- N.L. Johnson, *Combinatorics of Spreads and Parallelisms*, Chapman & Hall, 2010.

### *Design Theory*

*resolution of the block design formed by points and lines of  $\text{PG}(3, q)$*

### *Coding Theory*

- T. Etzion, N. Silberstein, *Codes and Designs Related to Lifted MRD Codes*, IEEE, 2013.

# Motivations

## *Kirkman's schoolgirl problem (1850)*

### *Finite Geometry*

- N.L. Johnson, *Combinatorics of Spreads and Parallelisms*, Chapman & Hall, 2010.

### *Design Theory*

*resolution of the block design formed by points and lines of  $\text{PG}(3, q)$*

### *Coding Theory*

- T. Etzion, N. Silberstein, *Codes and Designs Related to Lifted MRD Codes*, IEEE, 2013.
- M. Giulietti, A. Sabatini, M. Timpanella, *PIR Codes from Combinatorial Structures*, LNCS, 2023



$\ell_1, \ell_2, \ell_3$  three pairwise skew lines

$\ell_1, \ell_2, \ell_3$  three pairwise skew lines

*regulus*  $\mathcal{R}$

$\ell_1, \ell_2, \ell_3$  three pairwise skew lines

*regulus*  $\mathcal{R}$

$q + 1$  pairwise skew lines transversal to  $\ell_1, \ell_2, \ell_3$

$\ell_1, \ell_2, \ell_3$  three pairwise skew lines

*regulus  $\mathcal{R}$*

$q + 1$  pairwise skew lines transversal to  $\ell_1, \ell_2, \ell_3$

*opposite regulus  $\mathcal{R}^\circ$*

$\ell_1, \ell_2, \ell_3$  three pairwise skew lines

*regulus  $\mathcal{R}$*

$q + 1$  pairwise skew lines transversal to  $\ell_1, \ell_2, \ell_3$

*opposite regulus  $\mathcal{R}^\circ$*

$$\bigcup_{\ell \in \mathcal{R}} \ell = \bigcup_{\ell \in \mathcal{R}^\circ} \ell$$



$\ell_1, \ell_2, \ell_3$  three pairwise skew lines

*regulus  $\mathcal{R}$*

$q + 1$  pairwise skew lines transversal to  $\ell_1, \ell_2, \ell_3$

*opposite regulus  $\mathcal{R}^\circ$*

$$\bigcup_{\ell \in \mathcal{R}} \ell = \bigcup_{\ell \in \mathcal{R}^\circ} \ell$$

*regular spread  $\mathcal{S}$*

$\ell_1, \ell_2, \ell_3$  three pairwise skew lines

*regulus  $\mathcal{R}$*

$q + 1$  pairwise skew lines transversal to  $\ell_1, \ell_2, \ell_3$

*opposite regulus  $\mathcal{R}^\circ$*

$$\bigcup_{\ell \in \mathcal{R}} \ell = \bigcup_{\ell \in \mathcal{R}^\circ} \ell$$

*regular spread  $\mathcal{S}$*

$\forall r \notin \mathcal{S}, \{\ell \in \mathcal{S} : |\ell \cap r| = 1\}$  is a regulus

$\ell_1, \ell_2, \ell_3$  three pairwise skew lines

*regulus  $\mathcal{R}$*

$q + 1$  pairwise skew lines transversal to  $\ell_1, \ell_2, \ell_3$

*opposite regulus  $\mathcal{R}^\circ$*

$$\bigcup_{\ell \in \mathcal{R}} \ell = \bigcup_{\ell \in \mathcal{R}^\circ} \ell$$

*regular spread  $\mathcal{S}$*

$\forall r \notin \mathcal{S}, \{\ell \in \mathcal{S} : |\ell \cap r| = 1\}$  is a regulus

$$\mathcal{R} \subset \mathcal{S}$$

$\ell_1, \ell_2, \ell_3$  three pairwise skew lines

*regulus  $\mathcal{R}$*

$q + 1$  pairwise skew lines transversal to  $\ell_1, \ell_2, \ell_3$

*opposite regulus  $\mathcal{R}^\circ$*

$$\bigcup_{\ell \in \mathcal{R}} \ell = \bigcup_{\ell \in \mathcal{R}^\circ} \ell$$

*regular spread  $\mathcal{S}$*

$\forall r \notin \mathcal{S}, \{\ell \in \mathcal{S} : |\ell \cap r| = 1\}$  is a regulus

$$\mathcal{R} \subset \mathcal{S}$$

*new spread by switching  $\mathcal{R}$  with  $\mathcal{R}^\circ$*

$\ell_1, \ell_2, \ell_3$  three pairwise skew lines

*regulus  $\mathcal{R}$*

$q + 1$  pairwise skew lines transversal to  $\ell_1, \ell_2, \ell_3$

*opposite regulus  $\mathcal{R}^\circ$*

$$\bigcup_{\ell \in \mathcal{R}} \ell = \bigcup_{\ell \in \mathcal{R}^\circ} \ell$$

*regular spread  $\mathcal{S}$*

$\forall r \notin \mathcal{S}, \{\ell \in \mathcal{S} : |\ell \cap r| = 1\}$  is a regulus

$$\mathcal{R} \subset \mathcal{S}$$

*new spread by switching  $\mathcal{R}$  with  $\mathcal{R}^\circ$*

$$(\mathcal{S} \setminus \mathcal{R}) \cup \mathcal{R}^\circ$$



Embed  $\Sigma \simeq \text{PG}(3, q)$  in  $\text{PG}(3, q^2)$

Embed  $\Sigma \simeq \text{PG}(3, q)$  in  $\text{PG}(3, q^2)$   
 $\tau$  involution of  $\text{PG}(3, q^2)$  with  $\text{Fix}(\tau) = \Sigma$



Embed  $\Sigma \simeq \text{PG}(3, q)$  in  $\text{PG}(3, q^2)$   
 $\tau$  involution of  $\text{PG}(3, q^2)$  with  $\text{Fix}(\tau) = \Sigma$

$t$  line of  $\text{PG}(3, q^2)$ ,  $t \cap \Sigma = \emptyset$

Embed  $\Sigma \simeq \text{PG}(3, q)$  in  $\text{PG}(3, q^2)$   
 $\tau$  involution of  $\text{PG}(3, q^2)$  with  $\text{Fix}(\tau) = \Sigma$

$t$  line of  $\text{PG}(3, q^2)$ ,  $t \cap \Sigma = \emptyset$

$$t^\tau \cap \Sigma = \emptyset$$

Embed  $\Sigma \simeq \text{PG}(3, q)$  in  $\text{PG}(3, q^2)$   
 $\tau$  involution of  $\text{PG}(3, q^2)$  with  $\text{Fix}(\tau) = \Sigma$

$t$  line of  $\text{PG}(3, q^2)$ ,  $t \cap \Sigma = \emptyset$

$$t^\tau \cap \Sigma = \emptyset$$

$r_P = \langle P, P^\tau \rangle_{q^2}$ ,  $r_P \cap \Sigma$  is a line of  $\Sigma$

Embed  $\Sigma \simeq \text{PG}(3, q)$  in  $\text{PG}(3, q^2)$   
 $\tau$  involution of  $\text{PG}(3, q^2)$  with  $\text{Fix}(\tau) = \Sigma$

$t$  line of  $\text{PG}(3, q^2)$ ,  $t \cap \Sigma = \emptyset$

$$t^\tau \cap \Sigma = \emptyset$$

$r_P = \langle P, P^\tau \rangle_{q^2}$ ,  $r_P \cap \Sigma$  is a line of  $\Sigma$

$\mathcal{D} = \{r_P \cap \Sigma : P \in t\}$ ,  $q^2 + 1$  pairwise skew lines of  $\Sigma$

Embed  $\Sigma \simeq \text{PG}(3, q)$  in  $\text{PG}(3, q^2)$   
 $\tau$  involution of  $\text{PG}(3, q^2)$  with  $\text{Fix}(\tau) = \Sigma$

$t$  line of  $\text{PG}(3, q^2)$ ,  $t \cap \Sigma = \emptyset$

$$t^\tau \cap \Sigma = \emptyset$$

$r_P = \langle P, P^\tau \rangle_{q^2}$ ,  $r_P \cap \Sigma$  is a line of  $\Sigma$

$\mathcal{D} = \{r_P \cap \Sigma : P \in t\}$ ,  $q^2 + 1$  pairwise skew lines of  $\Sigma$

Desarguesian line-spread

Embed  $\Sigma \simeq \text{PG}(3, q)$  in  $\text{PG}(3, q^2)$   
 $\tau$  involution of  $\text{PG}(3, q^2)$  with  $\text{Fix}(\tau) = \Sigma$

$t$  line of  $\text{PG}(3, q^2)$ ,  $t \cap \Sigma = \emptyset$

$$t^\tau \cap \Sigma = \emptyset$$

$r_P = \langle P, P^\tau \rangle_{q^2}$ ,  $r_P \cap \Sigma$  is a line of  $\Sigma$

$\mathcal{D} = \{r_P \cap \Sigma : P \in t\}$ ,  $q^2 + 1$  pairwise skew lines of  $\Sigma$

Desarguesian line-spread

$t, t^\tau$  transversal lines of  $\mathcal{D}$

Embed  $\Sigma \simeq \text{PG}(3, q)$  in  $\text{PG}(3, q^2)$   
 $\tau$  involution of  $\text{PG}(3, q^2)$  with  $\text{Fix}(\tau) = \Sigma$

$t$  line of  $\text{PG}(3, q^2)$ ,  $t \cap \Sigma = \emptyset$

$$t^\tau \cap \Sigma = \emptyset$$

$r_P = \langle P, P^\tau \rangle_{q^2}$ ,  $r_P \cap \Sigma$  is a line of  $\Sigma$

$\mathcal{D} = \{r_P \cap \Sigma : P \in t\}$ ,  $q^2 + 1$  pairwise skew lines of  $\Sigma$

Desarguesian line-spread

$t, t^\tau$  transversal lines of  $\mathcal{D}$

*Desarguesian*  $\iff$  *regular*

Embed  $\Sigma \simeq \text{PG}(3, q)$  in  $\text{PG}(3, q^2)$   
 $\tau$  involution of  $\text{PG}(3, q^2)$  with  $\text{Fix}(\tau) = \Sigma$

$t$  line of  $\text{PG}(3, q^2)$ ,  $t \cap \Sigma = \emptyset$

$$t^\tau \cap \Sigma = \emptyset$$

$r_P = \langle P, P^\tau \rangle_{q^2}$ ,  $r_P \cap \Sigma$  is a line of  $\Sigma$

$\mathcal{D} = \{r_P \cap \Sigma : P \in t\}$ ,  $q^2 + 1$  pairwise skew lines of  $\Sigma$

Desarguesian line-spread

$t, t^\tau$  transversal lines of  $\mathcal{D}$

Desarguesian  $\iff$  regular

Hall spread



Embed  $\Sigma \simeq \text{PG}(3, q)$  in  $\text{PG}(3, q^2)$   
 $\tau$  involution of  $\text{PG}(3, q^2)$  with  $\text{Fix}(\tau) = \Sigma$

$t$  line of  $\text{PG}(3, q^2)$ ,  $t \cap \Sigma = \emptyset$

$$t^\tau \cap \Sigma = \emptyset$$

$r_P = \langle P, P^\tau \rangle_{q^2}$ ,  $r_P \cap \Sigma$  is a line of  $\Sigma$

$\mathcal{D} = \{r_P \cap \Sigma : P \in t\}$ ,  $q^2 + 1$  pairwise skew lines of  $\Sigma$

Desarguesian line-spread

$t, t^\tau$  transversal lines of  $\mathcal{D}$

Desarguesian  $\iff$  regular

Hall spread

$$\mathcal{R} \subset \mathcal{D}$$

Embed  $\Sigma \simeq \text{PG}(3, q)$  in  $\text{PG}(3, q^2)$   
 $\tau$  involution of  $\text{PG}(3, q^2)$  with  $\text{Fix}(\tau) = \Sigma$

$t$  line of  $\text{PG}(3, q^2)$ ,  $t \cap \Sigma = \emptyset$

$$t^\tau \cap \Sigma = \emptyset$$

$r_P = \langle P, P^\tau \rangle_{q^2}$ ,  $r_P \cap \Sigma$  is a line of  $\Sigma$

$\mathcal{D} = \{r_P \cap \Sigma : P \in t\}$ ,  $q^2 + 1$  pairwise skew lines of  $\Sigma$

Desarguesian line-spread

$t, t^\tau$  transversal lines of  $\mathcal{D}$

Desarguesian  $\iff$  regular

Hall spread

$$\mathcal{R} \subset \mathcal{D}$$

$$(\mathcal{D} \setminus \mathcal{R}) \cup \mathcal{R}^\circ$$

# *Regular parallelisms of $\text{PG}(3, q)$*

*Regular parallelisms of  $\text{PG}(3, q)$*

$$q \equiv 2 \pmod{3}$$

*Regular parallelisms of  $\text{PG}(3, q)$*

$$q \equiv 2 \pmod{3}$$

- Penttinen, Williams, *Regular packings of  $\text{PG}(3, q)$* , *EJC*, 1998.

## *Regular parallelisms of $\text{PG}(3, q)$*

$$q \equiv 2 \pmod{3}$$

- Penttinen, Williams, *Regular packings of  $\text{PG}(3, q)$* , *EJC*, 1998.

## *Derived parallelisms*

### *Regular parallelisms of $\text{PG}(3, q)$*

$$q \equiv 2 \pmod{3}$$

- Penttila, Williams, *Regular packings of  $\text{PG}(3, q)$* , *EJC*, 1998.

### *Derived parallelisms*

$\Pi$  parallelism

### *Regular parallelisms of $\text{PG}(3, q)$*

$$q \equiv 2 \pmod{3}$$

- Penttila, Williams, *Regular packings of  $\text{PG}(3, q)$* , *EJC*, 1998.

### *Derived parallelisms*

$\Pi$  parallelism

$$\mathcal{S}, \mathcal{S}' \in \Pi, \mathcal{R} \subset \mathcal{S}, \mathcal{R}^o \subset \mathcal{S}'$$



### *Regular parallelisms of $\text{PG}(3, q)$*

$$q \equiv 2 \pmod{3}$$

- Penttila, Williams, *Regular packings of  $\text{PG}(3, q)$* , *EJC*, 1998.

### *Derived parallelisms*

$\Pi$  parallelism

$$\mathcal{S}, \mathcal{S}' \in \Pi, \mathcal{R} \subset \mathcal{S}, \mathcal{R}^o \subset \mathcal{S}'$$

$$((\mathcal{S} \setminus \mathcal{R}) \cup \mathcal{R}^o) \cup ((\mathcal{S}' \setminus \mathcal{R}^o) \cup \mathcal{R}) \cup (\Pi \setminus \{\mathcal{S}, \mathcal{S}'\})$$

### Regular parallelisms of $\text{PG}(3, q)$

$$q \equiv 2 \pmod{3}$$

- Penttinen, Williams, *Regular packings of  $\text{PG}(3, q)$* , *EJC*, 1998.

### Derived parallelisms

$\Pi$  parallelism

$$\mathcal{S}, \mathcal{S}' \in \Pi, \mathcal{R} \subset \mathcal{S}, \mathcal{R}^\circ \subset \mathcal{S}'$$

$$((\mathcal{S} \setminus \mathcal{R}) \cup \mathcal{R}^\circ) \cup ((\mathcal{S}' \setminus \mathcal{R}^\circ) \cup \mathcal{R}) \cup (\Pi \setminus \{\mathcal{S}, \mathcal{S}'\})$$

- Johnson, *Subplane Covered Nets*, Marcel Dekker, 2000.

### Regular parallelisms of $\text{PG}(3, q)$

$$q \equiv 2 \pmod{3}$$

- Penttila, Williams, *Regular packings of  $\text{PG}(3, q)$* , *EJC*, 1998.

### Derived parallelisms

$\Pi$  parallelism

$$\mathcal{S}, \mathcal{S}' \in \Pi, \mathcal{R} \subset \mathcal{S}, \mathcal{R}^\circ \subset \mathcal{S}'$$

$$((\mathcal{S} \setminus \mathcal{R}) \cup \mathcal{R}^\circ) \cup ((\mathcal{S}' \setminus \mathcal{R}^\circ) \cup \mathcal{R}) \cup (\Pi \setminus \{\mathcal{S}, \mathcal{S}'\})$$

- Johnson, *Subplane Covered Nets*, Marcel Dekker, 2000.
- Denniston, *Some packings of projective spaces*, AANL, 1972.

### Regular parallelisms of $\text{PG}(3, q)$

$$q \equiv 2 \pmod{3}$$

- Penttinen, Williams, *Regular packings of  $\text{PG}(3, q)$* , *EJC*, 1998.

### Derived parallelisms

$\Pi$  parallelism

$$\mathcal{S}, \mathcal{S}' \in \Pi, \mathcal{R} \subset \mathcal{S}, \mathcal{R}^o \subset \mathcal{S}'$$

$$((\mathcal{S} \setminus \mathcal{R}) \cup \mathcal{R}^o) \cup ((\mathcal{S}' \setminus \mathcal{R}^o) \cup \mathcal{R}) \cup (\Pi \setminus \{\mathcal{S}, \mathcal{S}'\})$$

- Johnson, *Subplane Covered Nets*, Marcel Dekker, 2000.
- Denniston, *Some packings of projective spaces*, *AANL*, 1972.
- Beutelspacher, *On parallelisms in finite projective spaces*, *GD*, 1974.

### Regular parallelisms of $\text{PG}(3, q)$

$$q \equiv 2 \pmod{3}$$

- Penttila, Williams, *Regular packings of  $\text{PG}(3, q)$* , *EJC*, 1998.

### Derived parallelisms

$\Pi$  parallelism

$$\mathcal{S}, \mathcal{S}' \in \Pi, \mathcal{R} \subset \mathcal{S}, \mathcal{R}^o \subset \mathcal{S}'$$

$$((\mathcal{S} \setminus \mathcal{R}) \cup \mathcal{R}^o) \cup ((\mathcal{S}' \setminus \mathcal{R}^o) \cup \mathcal{R}) \cup (\Pi \setminus \{\mathcal{S}, \mathcal{S}'\})$$

- Johnson, *Subplane Covered Nets*, Marcel Dekker, 2000.
- Denniston, *Some packings of projective spaces*, *AANL*, 1972.
- Beutelspacher, *On parallelisms in finite projective spaces*, *GD*, 1974.
- Diaz, Johnson, Montinaro, *Coset switching in parallelisms*, *FFA*, 2008.

$$r \in \mathcal{D},$$

$$r \in \mathcal{D},$$

$$E \leq \text{Stab}_r(\text{Aut}(\mathcal{D})), E \text{ elementary abelian, } |E| = q^2$$

$$r \in \mathcal{D},$$

$$E \leq \text{Stab}_r(\text{Aut}(\mathcal{D})), E \text{ elementary abelian, } |E| = q^2$$

*most of the known parallelisms  $\Pi$  of  $\text{PG}(3, q)$*



$$r \in \mathcal{D},$$

$$E \leq \text{Stab}_r(\text{Aut}(\mathcal{D})), E \text{ elementary abelian, } |E| = q^2$$

*most of the known parallelisms  $\Pi$  of  $\text{PG}(3, q)$*

$$i) \mathcal{D} \in \Pi,$$

$$r \in \mathcal{D},$$

$$E \leq \text{Stab}_r(\text{Aut}(\mathcal{D})), E \text{ elementary abelian, } |E| = q^2$$

*most of the known parallelisms  $\Pi$  of  $\text{PG}(3, q)$*

$$i) \mathcal{D} \in \Pi,$$

$$ii) E \leq \text{Aut}(\Pi).$$

$$r \in \mathcal{D},$$

$$E \leq \text{Stab}_r(\text{Aut}(\mathcal{D})), E \text{ elementary abelian, } |E| = q^2$$

*most of the known parallelisms  $\Pi$  of  $\text{PG}(3, q)$*

$$i) \mathcal{D} \in \Pi,$$

$$ii) E \leq \text{Aut}(\Pi).$$

$q$  even

$$r \in \mathcal{D},$$

$$E \leq \text{Stab}_r(\text{Aut}(\mathcal{D})), E \text{ elementary abelian, } |E| = q^2$$

*most of the known parallelisms  $\Pi$  of  $\text{PG}(3, q)$*

$$i) \mathcal{D} \in \Pi,$$

$$ii) E \leq \text{Aut}(\Pi).$$

$q$  even

*iii)  $\Pi$  contains  $\mathcal{D}$  and  $q^2 + q$  Hall spreads*

$$r \in \mathcal{D},$$

$$E \leq \text{Stab}_r(\text{Aut}(\mathcal{D})), E \text{ elementary abelian, } |E| = q^2$$

*most of the known parallelisms  $\Pi$  of  $\text{PG}(3, q)$*

$$i) \mathcal{D} \in \Pi,$$

$$ii) E \leq \text{Aut}(\Pi).$$

$q$  even

*iii)  $\Pi$  contains  $\mathcal{D}$  and  $q^2 + q$  Hall spreads*

*a Hall spread is obtained from a Desarguesian spread  $\mathcal{S}$*

$$r \in \mathcal{D},$$

$$E \leq \text{Stab}_r(\text{Aut}(\mathcal{D})), E \text{ elementary abelian, } |E| = q^2$$

*most of the known parallelisms  $\Pi$  of  $\text{PG}(3, q)$*

$$i) \mathcal{D} \in \Pi,$$

$$ii) E \leq \text{Aut}(\Pi).$$

$q$  even

*iii)  $\Pi$  contains  $\mathcal{D}$  and  $q^2 + q$  Hall spreads*

*a Hall spread is obtained from a Desarguesian spread  $\mathcal{S}$*

*$\mathcal{S} \cap \mathcal{D}$  is one of the  $q^2 + q$  reguli of  $\mathcal{D}$  through  $r$*

$$\bigcup_{P \in t} \langle P, P^T \rangle_{q^2} = \bigcup_i (\Sigma_i \cup \Sigma_i^T) \cup \Sigma \cup t \cup t^T$$

$\text{PG}(3, q^2)$

$$\bigcup_{P \in t} \langle P, P^T \rangle_{q^2} = \bigcup_i (\Sigma_i \cup \Sigma_i^T) \cup \Sigma \cup t \cup t^T$$

$\text{PG}(3, q^2)$



$\Sigma \simeq \text{PG}(3, q)$



$$\bigcup_{P \in t} \langle P, P^T \rangle_{q^2} = \bigcup_i (\Sigma_i \cup \Sigma_i^T) \cup \Sigma \cup t \cup t^T$$

$t$

$\text{PG}(3, q^2)$



$\Sigma \simeq \text{PG}(3, q)$

$$\bigcup_{P \in t} \langle P, P^T \rangle_{q^2} = \bigcup_i (\Sigma_i \cup \Sigma_i^T) \cup \Sigma \cup t \cup t^T$$

$t$


$\text{PG}(3, q^2)$




$\Sigma \simeq \text{PG}(3, q)$

$t^T$


$$\bigcup_{P \in t} \langle P, P^T \rangle_{q^2} = \bigcup_i (\Sigma_i \cup \Sigma_i^T) \cup \Sigma \cup t \cup t^T$$




$$\bigcup_{P \in t} \langle P, P^T \rangle_{q^2} = \bigcup_i (\Sigma_i \cup \Sigma_i^T) \cup \Sigma \cup t \cup t^T$$




$$\bigcup_{P \in t} \langle P, P^T \rangle_{q^2} = \bigcup_i (\Sigma_i \cup \Sigma_i^T) \cup \Sigma \cup t \cup t^T$$




$$\bigcup_{P \in t} \langle P, P^T \rangle_{q^2} = \bigcup_i (\Sigma_i \cup \Sigma_i^T) \cup \Sigma \cup t \cup t^T$$



$$\bigcup_{P \in t} \langle P, P^T \rangle_{q^2} = \bigcup_i (\Sigma_i \cup \Sigma_i^T) \cup \Sigma \cup t \cup t^T$$




$$\bigcup_{P \in t} \langle P, P^T \rangle_{q^2} = \bigcup_i (\Sigma_i \cup \Sigma_i^T) \cup \Sigma \cup t \cup t^T$$







$$\bigcup_{P \in t} \langle P, P^T \rangle_{q^2} = \bigcup_i (\Sigma_i \cup \Sigma_i^T) \cup \Sigma \cup t \cup t^T$$




$$\bigcup_{P \in t} \langle P, P^T \rangle_{q^2} = \bigcup_i (\Sigma_i \cup \Sigma_i^T) \cup \Sigma \cup t \cup t^T$$




$$\bigcup_{P \in t} \langle P, P^T \rangle_{q^2} = \bigcup_i (\Sigma_i \cup \Sigma_i^T) \cup \Sigma \cup t \cup t^T$$




$$\bigcup_{P \in t} \langle P, P^T \rangle_{q^2} = \bigcup_i (\Sigma_i \cup \Sigma_i^T) \cup \Sigma \cup t \cup t^T$$




$$\bigcup_{P \in t} \langle P, P^T \rangle_{q^2} = \bigcup_i (\Sigma_i \cup \Sigma_i^T) \cup \Sigma \cup t \cup t^T$$




$$\bigcup_{P \in t} \langle P, P^\tau \rangle_{q^2} = \bigcup_i (\Sigma_i \cup \Sigma_i^\tau) \cup \Sigma \cup t \cup t^\tau$$





$$\bigcup_{P \in t} \langle P, P^T \rangle_{q^2} = \bigcup_i (\Sigma_i \cup \Sigma_i^T) \cup \Sigma \cup t \cup t^T$$




























$\mathcal{S}_\ell$  Desarguesian spread of  $\Sigma$ ,  $\ell, \ell^\tau$  transversals of  $\mathcal{S}_\ell$






$\mathcal{S}_\ell$  Desarguesian spread of  $\Sigma$ ,  $\ell, \ell^\tau$  transversals of  $\mathcal{S}_\ell$




$\mathcal{S}_\ell$  Desarguesian spread of  $\Sigma$ ,  $\ell, \ell^\tau$  transversals of  $\mathcal{S}_\ell$

$\mathcal{D} \cap \mathcal{S}_\ell$  is a regulus  $\mathcal{R}_\ell \iff \ell \cap \Sigma_i$  is a subline.




$\mathcal{S}_\ell$  Desarguesian spread of  $\Sigma$ ,  $\ell, \ell^\tau$  transversals of  $\mathcal{S}_\ell$

$\mathcal{D} \cap \mathcal{S}_\ell$  is a regulus  $\mathcal{R}_\ell \iff \ell \cap \Sigma_i$  is a subline.




$$(\mathcal{S}_\ell \setminus \mathcal{R}_\ell) \cup \mathcal{R}_\ell^o \text{ Hall spread of } \Sigma$$




$(\mathcal{S}_\ell \setminus \mathcal{R}_\ell) \cup \mathcal{R}_\ell^\circ$  Hall spread of  $\Sigma$

$\mathcal{D}, (\mathcal{S}_\ell \setminus \mathcal{R}_\ell) \cup \mathcal{R}_\ell^\circ$  are pairwise disjoint spreads




$r \in \mathcal{D}, E \leq \text{Stab}_r(\text{Aut}(\mathcal{D})), E \text{ elementary abelian}, |E| = q^2$



$r \in \mathcal{D}, E \leq \text{Stab}_r(\text{Aut}(\mathcal{D})), E \text{ elementary abelian}, |E| = q^2$


$\ell$  line of  $\text{PG}(3, q^2)$ ,  $\ell \cap \Sigma = \emptyset$ ,  $\ell \cap \Sigma_i$  is a subline



$r \in \mathcal{D}, E \leq \text{Stab}_r(\text{Aut}(\mathcal{D})), E \text{ elementary abelian}, |E| = q^2$

$\ell$  line of  $\text{PG}(3, q^2)$ ,  $\ell \cap \Sigma = \emptyset$ ,  $\ell \cap \Sigma_i$  is a subline

$\ell^E \cup \{r\}$  is a Baer pencil  $p(P_{\alpha u}, \pi_{\alpha v})$






$r \in \mathcal{D}, E \leq \text{Stab}_r(\text{Aut}(\mathcal{D})), E \text{ elementary abelian}, |E| = q^2$

$\ell$  line of  $\text{PG}(3, q^2)$ ,  $\ell \cap \Sigma = \emptyset$ ,  $\ell \cap \Sigma_i$  is a subline


$\ell^E \cup \{r\}$  is a Baer pencil  $p(P_{\alpha u}, \pi_{\alpha v})$



$r \in \mathcal{D}, E \leq \text{Stab}_r(\text{Aut}(\mathcal{D})), E \text{ elementary abelian}, |E| = q^2$

$\ell$  line of  $\text{PG}(3, q^2)$ ,  $\ell \cap \Sigma = \emptyset$ ,  $\ell \cap \Sigma_i$  is a subline


$\ell^E \cup \{r\}$  is a Baer pencil  $p(P_{\alpha u}, \pi_{\alpha v})$



$r \in \mathcal{D}, E \leq \text{Stab}_r(\text{Aut}(\mathcal{D})), E \text{ elementary abelian}, |E| = q^2$

$\ell$  line of  $\text{PG}(3, q^2)$ ,  $\ell \cap \Sigma = \emptyset$ ,  $\ell \cap \Sigma_i$  is a subline


$\ell^E \cup \{r\}$  is a Baer pencil  $p(P_{\alpha u}, \pi_{\alpha v})$




$r \in \mathcal{D}, E \leq \text{Stab}_r(\text{Aut}(\mathcal{D})), E \text{ elementary abelian}, |E| = q^2$

$\ell$  line of  $\text{PG}(3, q^2)$ ,  $\ell \cap \Sigma = \emptyset$ ,  $\ell \cap \Sigma_i$  is a subline


$\ell^E \cup \{r\}$  is a Baer pencil  $p(P_{\alpha u}, \pi_{\alpha v})$




$\{(\mathcal{S}_\ell \setminus \mathcal{R}_\ell) \cup \mathcal{R}_\ell^o : \ell \in p(P_{\alpha u, \alpha v}) \setminus \{r\}\}$  set of  $q$  Hall spreads of  $\Sigma$



$\{(\mathcal{S}_\ell \setminus \mathcal{R}_\ell) \cup \mathcal{R}_\ell^\circ : \ell \in p(P_{\alpha u, \alpha v}) \setminus \{r\}\}$  set of  $q$  Hall spreads of  $\Sigma$   
 $\{\mathcal{D}\} \cup \{(\mathcal{S}_\ell \setminus \mathcal{R}_\ell) \cup \mathcal{R}_\ell^\circ : \ell \in p(P_{\alpha u, \alpha v}) \setminus \{r\}\}$  are pairwise disjoint




$$\Lambda \subset \mathbb{F}_{q^2}, |\Lambda| = q - 1, \{\alpha^{q+1} : \alpha \in \Lambda\} = \mathbb{F}_q \setminus \{0\}$$



$$\Lambda \subset \mathbb{F}_{q^2}, |\Lambda| = q - 1, \{\alpha^{q+1} : \alpha \in \Lambda\} = \mathbb{F}_q \setminus \{0\}$$

$$\mathcal{I} \subset \Lambda, |\mathcal{I}| = \lfloor \frac{q-2}{2} \rfloor, \mathcal{U} = \{u \in \mathbb{F}_{q^2} \mid u^{q+1} = 1\}$$






$$\Lambda \subset \mathbb{F}_{q^2}, |\Lambda| = q - 1, \{\alpha^{q+1} : \alpha \in \Lambda\} = \mathbb{F}_q \setminus \{0\}$$

$$\mathcal{I} \subset \Lambda, |\mathcal{I}| = \lfloor \frac{q-2}{2} \rfloor, \mathcal{U} = \{u \in \mathbb{F}_{q^2} \mid u^{q+1} = 1\}$$

$$P_{\alpha u} = (1, 0, \alpha u, 0), \pi_{\alpha v} : X_4 = \alpha v X_2, \alpha \in \mathcal{I}, u, v \in \mathcal{U}$$



$$P_{\alpha u} = (1, 0, \alpha u, 0), \alpha \in \mathcal{I}, u \in \mathcal{U}$$

$$P_{\alpha u} = (1, 0, \alpha u, 0), \alpha \in \mathcal{I}, u \in \mathcal{U}$$

$$\pi_{\alpha v} : X_4 = \alpha v X_2, \alpha \in \mathcal{I}, v \in \mathcal{U}$$

$$P_{\alpha u} = (1, 0, \alpha u, 0), \alpha \in \mathcal{I}, u \in \mathcal{U}$$

$$\pi_{\alpha v} : X_4 = \alpha v X_2, \alpha \in \mathcal{I}, v \in \mathcal{U}$$

$$p(P_{\alpha u}, \pi_{\alpha v}) \setminus \{r\}$$

$$P_{\alpha u} = (1, 0, \alpha u, 0), \alpha \in \mathcal{I}, u \in \mathcal{U}$$

$$\pi_{\alpha v} : X_4 = \alpha v X_2, \alpha \in \mathcal{I}, v \in \mathcal{U}$$

$$p(P_{\alpha u}, \pi_{\alpha v}) \setminus \{r\}$$

$$\mathcal{P} = \{(P_{\alpha_1 u_1}, \pi_{\alpha_1 v_1}), \dots, (P_{\alpha_{q+1} u_{q+1}}, \pi_{\alpha_{q+1} v_{q+1}})\}$$

$$P_{\alpha u} = (1, 0, \alpha u, 0), \alpha \in \mathcal{I}, u \in \mathcal{U}$$

$$\pi_{\alpha v} : X_4 = \alpha v X_2, \alpha \in \mathcal{I}, v \in \mathcal{U}$$

$$p(P_{\alpha u}, \pi_{\alpha v}) \setminus \{r\}$$

$$\mathcal{P} = \{(P_{\alpha_1 u_1}, \pi_{\alpha_1 v_1}), \dots, (P_{\alpha_{q+1} u_{q+1}}, \pi_{\alpha_{q+1} v_{q+1}})\}$$

$$\mathcal{L}_{\mathcal{P}} = \{p(P_{\alpha_i u_i}, \pi_{\alpha_i v_i}) \setminus \{r\} : (P_{\alpha_i u_i}, \pi_{\alpha_i v_i}) \in \mathcal{P}\}$$

$$P_{\alpha u} = (1, 0, \alpha u, 0), \alpha \in \mathcal{I}, u \in \mathcal{U}$$

$$\pi_{\alpha v} : X_4 = \alpha v X_2, \alpha \in \mathcal{I}, v \in \mathcal{U}$$

$$p(P_{\alpha u}, \pi_{\alpha v}) \setminus \{r\}$$

$$\mathcal{P} = \{(P_{\alpha_1 u_1}, \pi_{\alpha_1 v_1}), \dots, (P_{\alpha_{q+1} u_{q+1}}, \pi_{\alpha_{q+1} v_{q+1}})\}$$

$$\mathcal{L}_{\mathcal{P}} = \{p(P_{\alpha_i u_i}, \pi_{\alpha_i v_i}) \setminus \{r\} : (P_{\alpha_i u_i}, \pi_{\alpha_i v_i}) \in \mathcal{P}\}$$

$$\Pi_{\mathcal{P}} = \{(\mathcal{S}_{\ell} \setminus \mathcal{R}_{\ell}) \cup \mathcal{R}_{\ell}^{\circ} \mid \ell \in \mathcal{L}_{\mathcal{P}}\} \cup \{\mathcal{D}\}$$

$$P_{\alpha u} = (1, 0, \alpha u, 0), \alpha \in \mathcal{I}, u \in \mathcal{U}$$

$$\pi_{\alpha v} : X_4 = \alpha v X_2, \alpha \in \mathcal{I}, v \in \mathcal{U}$$

$$p(P_{\alpha u}, \pi_{\alpha v}) \setminus \{r\}$$

$$\mathcal{P} = \{(P_{\alpha_1 u_1}, \pi_{\alpha_1 v_1}), \dots, (P_{\alpha_{q+1} u_{q+1}}, \pi_{\alpha_{q+1} v_{q+1}})\}$$

$$\mathcal{L}_{\mathcal{P}} = \{p(P_{\alpha_i u_i}, \pi_{\alpha_i v_i}) \setminus \{r\} : (P_{\alpha_i u_i}, \pi_{\alpha_i v_i}) \in \mathcal{P}\}$$

$$\Pi_{\mathcal{P}} = \{(\mathcal{S}_{\ell} \setminus \mathcal{R}_{\ell}) \cup \mathcal{R}_{\ell}^{\circ} \mid \ell \in \mathcal{L}_{\mathcal{P}}\} \cup \{\mathcal{D}\}$$

*How to choose  $\mathcal{P}$  such that  $\Pi_{\mathcal{P}}$  is a parallelism?*



$$\ell_i \in p(P_{\alpha u_i}, \pi_{\alpha v_i}), \ell_j \in p(P_{\beta u_j}, \pi_{\beta v_j})$$

$$\ell_i \in p(P_{\alpha u_i}, \pi_{\alpha v_i}), \ell_j \in p(P_{\beta u_j}, \pi_{\beta v_j})$$

$$(\alpha, u_i, v_i) \neq (\beta, u_j, v_j)$$

$$\ell_i \in p(P_{\alpha u_i}, \pi_{\alpha v_i}), \ell_j \in p(P_{\beta u_j}, \pi_{\beta v_j})$$

$$(\alpha, u_i, v_i) \neq (\beta, u_j, v_j)$$

*Property ①*

$$\mathcal{R}_{\ell_i} \neq \mathcal{R}_{\ell_j} \iff u_i v_j - u_j v_i \neq 0.$$

$$\ell_i \in p(P_{\alpha u_i}, \pi_{\alpha v_i}), \ell_j \in p(P_{\beta u_j}, \pi_{\beta v_j}) \\ (\alpha, u_i, v_i) \neq (\beta, u_j, v_j)$$

Property ①

$$\mathcal{R}_{\ell_i} \neq \mathcal{R}_{\ell_j} \iff u_i v_j - u_j v_i \neq 0.$$

Property ②

$$(\mathcal{S}_{\ell_i} \setminus \mathcal{R}_{\ell_i}) \cap (\mathcal{S}_{\ell_j} \setminus \mathcal{R}_{\ell_j}) = \emptyset \iff \alpha u_i (\beta v_j)^q - (\alpha v_i)^q \beta u_j \neq 0.$$

$$\ell_i \in p(P_{\alpha u_i}, \pi_{\alpha v_i}), \ell_j \in p(P_{\beta u_j}, \pi_{\beta v_j}) \\ (\alpha, u_i, v_i) \neq (\beta, u_j, v_j)$$

Property ①

$$\mathcal{R}_{\ell_i} \neq \mathcal{R}_{\ell_j} \iff u_i v_j - u_j v_i \neq 0.$$

Property ②

$$(\mathcal{S}_{\ell_i} \setminus \mathcal{R}_{\ell_i}) \cap (\mathcal{S}_{\ell_j} \setminus \mathcal{R}_{\ell_j}) = \emptyset \iff \alpha u_i (\beta v_j)^q - (\alpha v_i)^q \beta u_j \neq 0.$$

Good set

$$\mathcal{P} = \{(P_{\alpha_1 u_1}, \pi_{\alpha_1 v_1}), \dots, (P_{\alpha_{q+1} u_{q+1}}, \pi_{\alpha_{q+1} v_{q+1}})\}$$

$$\ell_i \in p(P_{\alpha u_i}, \pi_{\alpha v_i}), \ell_j \in p(P_{\beta u_j}, \pi_{\beta v_j}) \\ (\alpha, u_i, v_i) \neq (\beta, u_j, v_j)$$

Property ①

$$\mathcal{R}_{\ell_i} \neq \mathcal{R}_{\ell_j} \iff u_i v_j - u_j v_i \neq 0.$$

Property ②

$$(\mathcal{S}_{\ell_i} \setminus \mathcal{R}_{\ell_i}) \cap (\mathcal{S}_{\ell_j} \setminus \mathcal{R}_{\ell_j}) = \emptyset \iff \alpha u_i (\beta v_j)^q - (\alpha v_i)^q \beta u_j \neq 0.$$

Good set

$$\mathcal{P} = \{(P_{\alpha_1 u_1}, \pi_{\alpha_1 v_1}), \dots, (P_{\alpha_{q+1} u_{q+1}}, \pi_{\alpha_{q+1} v_{q+1}})\}$$

$$\forall (P_{\alpha_i u_i}, \pi_{\alpha_i v_i}), (P_{\alpha_j u_j}, \pi_{\alpha_j v_j}) \in \mathcal{P}, (\alpha_i, u_i, v_i) \neq (\alpha_j, u_j, v_j),$$

$$\ell_i \in p(P_{\alpha u_i}, \pi_{\alpha v_i}), \ell_j \in p(P_{\beta u_j}, \pi_{\beta v_j}) \\ (\alpha, u_i, v_i) \neq (\beta, u_j, v_j)$$

Property ①

$$\mathcal{R}_{\ell_i} \neq \mathcal{R}_{\ell_j} \iff u_i v_j - u_j v_i \neq 0.$$

Property ②

$$(\mathcal{S}_{\ell_i} \setminus \mathcal{R}_{\ell_i}) \cap (\mathcal{S}_{\ell_j} \setminus \mathcal{R}_{\ell_j}) = \emptyset \iff \alpha u_i (\beta v_j)^q - (\alpha v_i)^q \beta u_j \neq 0.$$

Good set

$$\mathcal{P} = \{(P_{\alpha_1 u_1}, \pi_{\alpha_1 v_1}), \dots, (P_{\alpha_{q+1} u_{q+1}}, \pi_{\alpha_{q+1} v_{q+1}})\}$$

$$\forall (P_{\alpha_i u_i}, \pi_{\alpha_i v_i}), (P_{\alpha_j u_j}, \pi_{\alpha_j v_j}) \in \mathcal{P}, (\alpha_i, u_i, v_i) \neq (\alpha_j, u_j, v_j),$$

$$\textcircled{1} \quad u_i v_j - v_i u_j \neq 0,$$

$$\ell_i \in p(P_{\alpha u_i}, \pi_{\alpha v_i}), \ell_j \in p(P_{\beta u_j}, \pi_{\beta v_j}) \\ (\alpha, u_i, v_i) \neq (\beta, u_j, v_j)$$

Property ①

$$\mathcal{R}_{\ell_i} \neq \mathcal{R}_{\ell_j} \iff u_i v_j - u_j v_i \neq 0.$$

Property ②

$$(\mathcal{S}_{\ell_i} \setminus \mathcal{R}_{\ell_i}) \cap (\mathcal{S}_{\ell_j} \setminus \mathcal{R}_{\ell_j}) = \emptyset \iff \alpha u_i (\beta v_j)^q - (\alpha v_i)^q \beta u_j \neq 0.$$

Good set

$$\mathcal{P} = \{(P_{\alpha_1 u_1}, \pi_{\alpha_1 v_1}), \dots, (P_{\alpha_{q+1} u_{q+1}}, \pi_{\alpha_{q+1} v_{q+1}})\}$$

$$\forall (P_{\alpha_i u_i}, \pi_{\alpha_i v_i}), (P_{\alpha_j u_j}, \pi_{\alpha_j v_j}) \in \mathcal{P}, (\alpha_i, u_i, v_i) \neq (\alpha_j, u_j, v_j),$$

$$\textcircled{1} \quad u_i v_j - v_i u_j \neq 0,$$

$$\textcircled{2} \quad \alpha_i u_i (\alpha_j v_j)^q - (\alpha_i v_i)^q \alpha_j u_j \neq 0.$$



## Theorem

$$\mathcal{P} = \{(P_{\alpha_1 u_1}, \pi_{\alpha_1 v_1}), \dots, (P_{\alpha_{q+1} u_{q+1}}, \pi_{\alpha_{q+1} v_{q+1}})\},$$

## Theorem

$$\mathcal{P} = \{(P_{\alpha_1 u_1}, \pi_{\alpha_1 v_1}), \dots, (P_{\alpha_{q+1} u_{q+1}}, \pi_{\alpha_{q+1} v_{q+1}})\},$$

$$\mathcal{L}_{\mathcal{P}} = \{p(P_{\alpha_i u_i}, \pi_{\alpha_i v_i}) \setminus \{r\} \mid (P_{\alpha_i u_i}, \pi_{\alpha_i v_i}) \in \mathcal{P}\},$$

## Theorem

$$\mathcal{P} = \{(P_{\alpha_1 u_1}, \pi_{\alpha_1 v_1}), \dots, (P_{\alpha_{q+1} u_{q+1}}, \pi_{\alpha_{q+1} v_{q+1}})\},$$

$$\mathcal{L}_{\mathcal{P}} = \{p(P_{\alpha_i u_i}, \pi_{\alpha_i v_i}) \setminus \{r\} \mid (P_{\alpha_i u_i}, \pi_{\alpha_i v_i}) \in \mathcal{P}\},$$

$$\Pi_{\mathcal{P}} = \{(S_{\ell} \setminus \mathcal{R}_{\ell}) \cup \mathcal{R}_{\ell}^{\circ} \mid \ell \in \mathcal{L}_{\mathcal{P}}\} \cup \{\mathcal{D}\},$$

## Theorem

$$\mathcal{P} = \{(P_{\alpha_1 u_1}, \pi_{\alpha_1 v_1}), \dots, (P_{\alpha_{q+1} u_{q+1}}, \pi_{\alpha_{q+1} v_{q+1}})\},$$

$$\mathcal{L}_{\mathcal{P}} = \{p(P_{\alpha_i u_i}, \pi_{\alpha_i v_i}) \setminus \{r\} \mid (P_{\alpha_i u_i}, \pi_{\alpha_i v_i}) \in \mathcal{P}\},$$

$$\Pi_{\mathcal{P}} = \{(S_{\ell} \setminus \mathcal{R}_{\ell}) \cup \mathcal{R}_{\ell}^{\circ} \mid \ell \in \mathcal{L}_{\mathcal{P}}\} \cup \{\mathcal{D}\},$$

$\Pi_{\mathcal{P}}$  is a line-parallelism of  $\Sigma \iff \mathcal{P}$  is a good set.

### Theorem

$$\mathcal{P} = \{(P_{\alpha_1 u_1}, \pi_{\alpha_1 v_1}), \dots, (P_{\alpha_{q+1} u_{q+1}}, \pi_{\alpha_{q+1} v_{q+1}})\},$$

$$\mathcal{L}_{\mathcal{P}} = \{p(P_{\alpha_i u_i}, \pi_{\alpha_i v_i}) \setminus \{r\} \mid (P_{\alpha_i u_i}, \pi_{\alpha_i v_i}) \in \mathcal{P}\},$$

$$\Pi_{\mathcal{P}} = \{(\mathcal{S}_{\ell} \setminus \mathcal{R}_{\ell}) \cup \mathcal{R}_{\ell}^{\circ} \mid \ell \in \mathcal{L}_{\mathcal{P}}\} \cup \{\mathcal{D}\},$$

$\Pi_{\mathcal{P}}$  is a line-parallelism of  $\Sigma \iff \mathcal{P}$  is a good set.

### Theorem

$\Pi$  parallelism of  $\Sigma$  such that

### Theorem

$$\mathcal{P} = \{(P_{\alpha_1 u_1}, \pi_{\alpha_1 v_1}), \dots, (P_{\alpha_{q+1} u_{q+1}}, \pi_{\alpha_{q+1} v_{q+1}})\},$$

$$\mathcal{L}_{\mathcal{P}} = \{p(P_{\alpha_i u_i}, \pi_{\alpha_i v_i}) \setminus \{r\} \mid (P_{\alpha_i u_i}, \pi_{\alpha_i v_i}) \in \mathcal{P}\},$$

$$\Pi_{\mathcal{P}} = \{(\mathcal{S}_{\ell} \setminus \mathcal{R}_{\ell}) \cup \mathcal{R}_{\ell}^{\circ} \mid \ell \in \mathcal{L}_{\mathcal{P}}\} \cup \{\mathcal{D}\},$$

$\Pi_{\mathcal{P}}$  is a line-parallelism of  $\Sigma \iff \mathcal{P}$  is a good set.

### Theorem

$\Pi$  parallelism of  $\Sigma$  such that

i)  $\mathcal{D} \in \Pi$ ,

### Theorem

$$\mathcal{P} = \{(P_{\alpha_1 u_1}, \pi_{\alpha_1 v_1}), \dots, (P_{\alpha_{q+1} u_{q+1}}, \pi_{\alpha_{q+1} v_{q+1}})\},$$

$$\mathcal{L}_{\mathcal{P}} = \{p(P_{\alpha_i u_i}, \pi_{\alpha_i v_i}) \setminus \{r\} \mid (P_{\alpha_i u_i}, \pi_{\alpha_i v_i}) \in \mathcal{P}\},$$

$$\Pi_{\mathcal{P}} = \{(\mathcal{S}_{\ell} \setminus \mathcal{R}_{\ell}) \cup \mathcal{R}_{\ell}^{\circ} \mid \ell \in \mathcal{L}_{\mathcal{P}}\} \cup \{\mathcal{D}\},$$

$\Pi_{\mathcal{P}}$  is a line-parallelism of  $\Sigma \iff \mathcal{P}$  is a good set.

### Theorem

$\Pi$  parallelism of  $\Sigma$  such that

i)  $\mathcal{D} \in \Pi$ ,

ii)  $E \leq \text{Aut}(\Pi)$ .

### Theorem

$$\mathcal{P} = \{(P_{\alpha_1 u_1}, \pi_{\alpha_1 v_1}), \dots, (P_{\alpha_{q+1} u_{q+1}}, \pi_{\alpha_{q+1} v_{q+1}})\},$$

$$\mathcal{L}_{\mathcal{P}} = \{p(P_{\alpha_i u_i}, \pi_{\alpha_i v_i}) \setminus \{r\} \mid (P_{\alpha_i u_i}, \pi_{\alpha_i v_i}) \in \mathcal{P}\},$$

$$\Pi_{\mathcal{P}} = \{(\mathcal{S}_{\ell} \setminus \mathcal{R}_{\ell}) \cup \mathcal{R}_{\ell}^{\circ} \mid \ell \in \mathcal{L}_{\mathcal{P}}\} \cup \{\mathcal{D}\},$$

$\Pi_{\mathcal{P}}$  is a line-parallelism of  $\Sigma \iff \mathcal{P}$  is a good set.

### Theorem

$\Pi$  parallelism of  $\Sigma$  such that

i)  $\mathcal{D} \in \Pi$ ,

ii)  $E \leq \text{Aut}(\Pi)$ .

iii)  $\Pi$  contains  $\mathcal{D}$  and  $q^2 + q$  Hall spread



### Theorem

$$\mathcal{P} = \{(P_{\alpha_1 u_1}, \pi_{\alpha_1 v_1}), \dots, (P_{\alpha_{q+1} u_{q+1}}, \pi_{\alpha_{q+1} v_{q+1}})\},$$

$$\mathcal{L}_{\mathcal{P}} = \{p(P_{\alpha_i u_i}, \pi_{\alpha_i v_i}) \setminus \{r\} \mid (P_{\alpha_i u_i}, \pi_{\alpha_i v_i}) \in \mathcal{P}\},$$

$$\Pi_{\mathcal{P}} = \{(S_{\ell} \setminus \mathcal{R}_{\ell}) \cup \mathcal{R}_{\ell}^{\circ} \mid \ell \in \mathcal{L}_{\mathcal{P}}\} \cup \{\mathcal{D}\},$$

$\Pi_{\mathcal{P}}$  is a line-parallelism of  $\Sigma \iff \mathcal{P}$  is a good set.

### Theorem

$\Pi$  parallelism of  $\Sigma$  such that

i)  $\mathcal{D} \in \Pi$ ,

ii)  $E \leq \text{Aut}(\Pi)$ .

iii)  $\Pi$  contains  $\mathcal{D}$  and  $q^2 + q$  Hall spread

a Hall spread is obtained from a Desarguesian spread  $S_{\ell}$

### Theorem

$$\mathcal{P} = \{(P_{\alpha_1 u_1}, \pi_{\alpha_1 v_1}), \dots, (P_{\alpha_{q+1} u_{q+1}}, \pi_{\alpha_{q+1} v_{q+1}})\},$$

$$\mathcal{L}_{\mathcal{P}} = \{p(P_{\alpha_i u_i}, \pi_{\alpha_i v_i}) \setminus \{r\} \mid (P_{\alpha_i u_i}, \pi_{\alpha_i v_i}) \in \mathcal{P}\},$$

$$\Pi_{\mathcal{P}} = \{(S_{\ell} \setminus \mathcal{R}_{\ell}) \cup \mathcal{R}_{\ell}^{\circ} \mid \ell \in \mathcal{L}_{\mathcal{P}}\} \cup \{\mathcal{D}\},$$

$\Pi_{\mathcal{P}}$  is a line-parallelism of  $\Sigma \iff \mathcal{P}$  is a good set.

### Theorem

$\Pi$  parallelism of  $\Sigma$  such that

i)  $\mathcal{D} \in \Pi$ ,

ii)  $E \leq \text{Aut}(\Pi)$ .

iii)  $\Pi$  contains  $\mathcal{D}$  and  $q^2 + q$  Hall spread

a Hall spread is obtained from a Desarguesian spread  $S_{\ell}$

$S_{\ell} \cap \mathcal{D}$  is one of the  $q^2 + q$  reguli of  $\mathcal{D}$  through  $r$

### Theorem

$$\mathcal{P} = \{(P_{\alpha_1 u_1}, \pi_{\alpha_1 v_1}), \dots, (P_{\alpha_{q+1} u_{q+1}}, \pi_{\alpha_{q+1} v_{q+1}})\},$$

$$\mathcal{L}_{\mathcal{P}} = \{p(P_{\alpha_i u_i}, \pi_{\alpha_i v_i}) \setminus \{r\} \mid (P_{\alpha_i u_i}, \pi_{\alpha_i v_i}) \in \mathcal{P}\},$$

$$\Pi_{\mathcal{P}} = \{(S_{\ell} \setminus \mathcal{R}_{\ell}) \cup \mathcal{R}_{\ell}^{\circ} \mid \ell \in \mathcal{L}_{\mathcal{P}}\} \cup \{\mathcal{D}\},$$

$\Pi_{\mathcal{P}}$  is a line-parallelism of  $\Sigma \iff \mathcal{P}$  is a good set.

### Theorem

$\Pi$  parallelism of  $\Sigma$  such that

i)  $\mathcal{D} \in \Pi$ ,

ii)  $E \leq \text{Aut}(\Pi)$ .

iii)  $\Pi$  contains  $\mathcal{D}$  and  $q^2 + q$  Hall spread

a Hall spread is obtained from a Desarguesian spread  $S_{\ell}$

$S_{\ell} \cap \mathcal{D}$  is one of the  $q^2 + q$  reguli of  $\mathcal{D}$  through  $r$

$$\iff \Pi = \Pi_{\mathcal{P}}, \text{ for some good set } \mathcal{P}.$$

$$\mathcal{P} = \{(P_{\alpha_1 u_1}, \pi_{\alpha_1 v_1}), \dots, (P_{\alpha_{q+1} u_{q+1}}, \pi_{\alpha_{q+1} v_{q+1}})\}$$

$$\mathcal{P} = \{(P_{\alpha_1 u_1}, \pi_{\alpha_1 v_1}), \dots, (P_{\alpha_{q+1} u_{q+1}}, \pi_{\alpha_{q+1} v_{q+1}})\}$$

$$\{(1, \alpha_i u_i, \alpha_i v_i) \mid (P_{\alpha_i u_i}, \pi_{\alpha_i v_i}) \in \mathcal{P}\}, \alpha_i \in \mathcal{I}, u_i, v_i \in \mathcal{U}$$

$$\mathcal{P} = \{(P_{\alpha_1 u_1}, \pi_{\alpha_1 v_1}), \dots, (P_{\alpha_{q+1} u_{q+1}}, \pi_{\alpha_{q+1} v_{q+1}})\}$$

$$\{(1, \alpha_i u_i, \alpha_i v_i) \mid (P_{\alpha_i u_i}, \pi_{\alpha_i v_i}) \in \mathcal{P}\}, \alpha_i \in \mathcal{I}, u_i, v_i \in \mathcal{U}$$

# good sets:

$$\left(\frac{q-2}{2}\right)^{q+1} (q+1)! \quad \text{if } q \text{ is even,}$$

$$\left(\frac{(q+1)(q-3)}{16}\right)^{\frac{q+1}{2}} \prod_{i=0}^{\frac{q-1}{2}} (q+1-2i)^2 \quad \text{if } q \equiv 1 \pmod{4},$$

$$\left(\frac{q-1}{4}\right)^{q+1} \prod_{i=0}^{\frac{q-1}{2}} (q+1-2i)^2 \quad \text{if } q \equiv 3 \pmod{4}.$$



$$\mathcal{P} \neq \mathcal{P}' \implies \Pi_{\mathcal{P}} \neq \Pi_{\mathcal{P}'}$$



$$\mathcal{P} \neq \mathcal{P}' \implies \Pi_{\mathcal{P}} \neq \Pi_{\mathcal{P}'}$$

*line-parallelisms  $\Pi$  and  $\Pi'$  of  $\text{PG}(3, q)$  are equivalent*

$$\mathcal{P} \neq \mathcal{P}' \implies \Pi_{\mathcal{P}} \neq \Pi_{\mathcal{P}'}$$

*line-parallelisms  $\Pi$  and  $\Pi'$  of  $\text{PG}(3, q)$  are equivalent*

$$\Pi^g = \Pi', g \in \text{PTL}(4, q)$$

$$\mathcal{P} \neq \mathcal{P}' \implies \Pi_{\mathcal{P}} \neq \Pi_{\mathcal{P}'}$$

*line-parallelisms  $\Pi$  and  $\Pi'$  of  $\text{PG}(3, q)$  are equivalent*

$$\Pi^g = \Pi', g \in \text{PTL}(4, q)$$

*$\Pi_{\mathcal{P}}, \Pi_{\mathcal{P}'}$  may be equivalent*

$$\mathcal{P} \neq \mathcal{P}' \implies \Pi_{\mathcal{P}} \neq \Pi_{\mathcal{P}'}$$

*line-parallelisms  $\Pi$  and  $\Pi'$  of  $\text{PG}(3, q)$  are equivalent*

$$\Pi^g = \Pi', g \in \text{PTL}(4, q)$$

*$\Pi_{\mathcal{P}}, \Pi_{\mathcal{P}'}$  may be equivalent*

*even if  $\mathcal{P} \neq \mathcal{P}'$*

$$\mathcal{P} \neq \mathcal{P}' \implies \Pi_{\mathcal{P}} \neq \Pi_{\mathcal{P}'}$$

*line-parallelisms  $\Pi$  and  $\Pi'$  of  $\text{PG}(3, q)$  are equivalent*

$$\Pi^g = \Pi', \quad g \in \text{P}\Gamma\text{L}(4, q)$$

*$\Pi_{\mathcal{P}}, \Pi_{\mathcal{P}'}$  may be equivalent*

*even if  $\mathcal{P} \neq \mathcal{P}'$*

$\Gamma$  the subgroup of  $\text{P}\Gamma\text{L}(4, q)$  that stabilizes  $\mathcal{D}$

$$\mathcal{P} \neq \mathcal{P}' \implies \Pi_{\mathcal{P}} \neq \Pi_{\mathcal{P}'}$$

*line-parallelisms  $\Pi$  and  $\Pi'$  of  $\text{PG}(3, q)$  are equivalent*

$$\Pi^g = \Pi', \quad g \in \text{P}\Gamma\text{L}(4, q)$$

*$\Pi_{\mathcal{P}}, \Pi_{\mathcal{P}'}$  may be equivalent*

*even if  $\mathcal{P} \neq \mathcal{P}'$*

$\Gamma$  the subgroup of  $\text{P}\Gamma\text{L}(4, q)$  that stabilizes  $\mathcal{D}$

$\Gamma_r$  the subgroup of  $\Gamma$  that stabilizes the line  $r$

$$\mathcal{P} \neq \mathcal{P}' \implies \Pi_{\mathcal{P}} \neq \Pi_{\mathcal{P}'}$$

*line-parallelisms  $\Pi$  and  $\Pi'$  of  $\text{PG}(3, q)$  are equivalent*

$$\Pi^g = \Pi', \quad g \in \text{P}\Gamma\text{L}(4, q)$$

*$\Pi_{\mathcal{P}}, \Pi_{\mathcal{P}'}$  may be equivalent*

*even if  $\mathcal{P} \neq \mathcal{P}'$*

$\Gamma$  the subgroup of  $\text{P}\Gamma\text{L}(4, q)$  that stabilizes  $\mathcal{D}$

$\Gamma_r$  the subgroup of  $\Gamma$  that stabilizes the line  $r$

$$|\Gamma_r| = 2hq^2(q^2 - 1)(q + 1)$$

$$\mathcal{P} \neq \mathcal{P}' \implies \Pi_{\mathcal{P}} \neq \Pi_{\mathcal{P}'}$$

*line-parallelisms  $\Pi$  and  $\Pi'$  of  $\text{PG}(3, q)$  are equivalent*

$$\Pi^g = \Pi', \quad g \in \text{P}\Gamma\text{L}(4, q)$$

*$\Pi_{\mathcal{P}}, \Pi_{\mathcal{P}'}$  may be equivalent*

*even if  $\mathcal{P} \neq \mathcal{P}'$*

$\Gamma$  the subgroup of  $\text{P}\Gamma\text{L}(4, q)$  that stabilizes  $\mathcal{D}$

$\Gamma_r$  the subgroup of  $\Gamma$  that stabilizes the line  $r$

$$|\Gamma_r| = 2hq^2(q^2 - 1)(q + 1)$$

$\Pi_{\mathcal{P}}, \Pi_{\mathcal{P}'}$  line-parallelisms of  $\text{PG}(3, q)$ , where  $\mathcal{P}, \mathcal{P}'$  are good sets.



$$\mathcal{P} \neq \mathcal{P}' \implies \Pi_{\mathcal{P}} \neq \Pi_{\mathcal{P}'}$$

*line-parallelisms  $\Pi$  and  $\Pi'$  of  $\text{PG}(3, q)$  are equivalent*

$$\Pi^g = \Pi', g \in \text{P}\Gamma\text{L}(4, q)$$

*$\Pi_{\mathcal{P}}, \Pi_{\mathcal{P}'}$  may be equivalent*

*even if  $\mathcal{P} \neq \mathcal{P}'$*

$\Gamma$  the subgroup of  $\text{P}\Gamma\text{L}(4, q)$  that stabilizes  $\mathcal{D}$

$\Gamma_r$  the subgroup of  $\Gamma$  that stabilizes the line  $r$

$$|\Gamma_r| = 2hq^2(q^2 - 1)(q + 1)$$

$\Pi_{\mathcal{P}}, \Pi_{\mathcal{P}'}$  line-parallelisms of  $\text{PG}(3, q)$ , where  $\mathcal{P}, \mathcal{P}'$  are good sets.

$$\Pi_{\mathcal{P}}^g = \Pi_{\mathcal{P}'}, g \in \text{P}\Gamma\text{L}(4, q) \implies g \in \Gamma_r.$$

$$\mathcal{T} = \{\Pi_{\mathcal{P}} \mid \mathcal{P} \text{ is a good set}\}$$

$$\mathcal{T} = \{\Pi_{\mathcal{P}} \mid \mathcal{P} \text{ is a good set}\}$$

Theorem

*# mutually not equivalent line-parallelisms of  $\text{PG}(3, q)$  in  $\mathcal{T}$  is at least*

$$\mathcal{T} = \{\Pi_{\mathcal{P}} \mid \mathcal{P} \text{ is a good set}\}$$

### Theorem

# mutually not equivalent line-parallelisms of  $\text{PG}(3, q)$  in  $\mathcal{T}$  is at least

$$\left(\frac{q-1}{2}\right)^{q+1} \frac{(q-2)!}{2hq(q+1)} \quad \text{if } q \text{ is even,}$$

$$\left(\frac{q-3}{4}\right)^{q+1} \frac{(q^2-1)^{\frac{q-1}{2}}}{2hq^2(q+1)} \quad \text{if } q \equiv 1 \pmod{4},$$

$$\left(\frac{(q-1)(q-5)}{16}\right)^{\frac{q+1}{2}} \frac{(q^2-1)^{\frac{q-1}{2}}}{2hq^2(q+1)} \quad \text{if } q \equiv 3 \pmod{4}.$$

*THANK YOU*