

A skew polynomial framework for semifields and MRD codes

Paolo Santonastaso

Polytechnic University of Bari

Finite Geometries 2025
Seventh Irsee Conference

31 August - 6 September, 2025

“Coding theory is the theory of subsets of a metric space (X, d) ”

Rank-metric space

“Coding theory is the theory of subsets of a metric space (X, d) ”

Rank-metric space:

- $X = M_{m \times n}(\mathbb{F}_q)$
- $d_R(A, B) = rk(A - B)$, where $A, B \in M_{m \times n}(\mathbb{F}_q)$.

Rank-metric space

“Coding theory is the theory of subsets of a metric space (X, d) ”

Rank-metric space:

- $X = M_{m \times n}(\mathbb{F}_q)$
- $d_R(A, B) = rk(A - B)$, where $A, B \in M_{m \times n}(\mathbb{F}_q)$.

A **rank metric code** \mathcal{C} is a subset of $(M_{m \times n}(\mathbb{F}_q), d_R)$.

Rank-metric space

“Coding theory is the theory of subsets of a metric space (X, d) ”

Rank-metric space:

- $X = M_{m \times n}(\mathbb{F}_q)$
- $d_R(A, B) = rk(A - B)$, where $A, B \in M_{m \times n}(\mathbb{F}_q)$.

A **rank metric code** \mathcal{C} is a subset of $(M_{m \times n}(\mathbb{F}_q), d_R)$.

$$d(\mathcal{C}) = \min\{rk(X - Y) : X, Y \in \mathcal{C}, X \neq Y\}$$

Rank metric

Rank-metric space:

- $X = M_{m \times n}(\mathbb{F}_q)$
- $d_R(A, B) = rk(A - B)$, where $A, B \in M_{m \times n}(\mathbb{F}_q)$.

Singleton-like bound

$$|C| \leq q^{\max\{m,n\}(\min\{m,n\}-d+1)}$$

Rank metric

Rank-metric space:

- $X = M_{m \times n}(\mathbb{F}_q)$
- $d_R(A, B) = rk(A - B)$, where $A, B \in M_{m \times n}(\mathbb{F}_q)$.

Singleton-like bound

$$|\mathcal{C}| \leq q^{\max\{m,n\}(\min\{m,n\}-d+1)}$$

If $|\mathcal{C}| = q^{\max\{m,n\}(\min\{m,n\}-d+1)} \Rightarrow \mathcal{C}$ **Maximum Rank Distance (MRD) code**

Rank metric

Rank-metric space:

- $X = M_{m \times n}(\mathbb{F}_q)$
- $d_R(A, B) = rk(A - B)$, where $A, B \in M_{m \times n}(\mathbb{F}_q)$.

Singleton-like bound

$$|C| \leq q^{\max\{m,n\}(\min\{m,n\}-d+1)}$$

If $|C| = q^{\max\{m,n\}(\min\{m,n\}-d+1)} \Rightarrow C$ **Maximum Rank Distance (MRD) code**

Equivalence

$$C, C' \subseteq M_{m \times n}(\mathbb{F}_q)$$

$$C \sim C' \iff C' = A \cdot C^\tau \cdot B = \{AC^\tau B : C \in C\},$$

$$A \in GL(m, q), B \in GL(n, q), \tau \in Aut(\mathbb{F}_q)$$

Skew polynomial rings

$$n = m$$

Skew polynomial rings

$$n = m$$

- $\mathbb{F}_{q^n}/\mathbb{F}_q$ finite field extension

Skew polynomial rings

$$n = m$$

- $\mathbb{F}_{q^n}/\mathbb{F}_q$ finite field extension
- $\text{Gal}(\mathbb{F}_{q^n}/\mathbb{F}_q) = \langle \sigma \rangle$

Skew polynomial rings

$$n = m$$

- $\mathbb{F}_{q^n}/\mathbb{F}_q$ finite field extension
- $\text{Gal}(\mathbb{F}_{q^n}/\mathbb{F}_q) = \langle \sigma \rangle$

$$R = \left\{ \sum_{i=0}^t \alpha_i x^i : \alpha_i \in \mathbb{F}_{q^m} \right\}$$

$$\sum_i \alpha_i x^i + \sum_i \beta_i x^i = \sum_i (\alpha_i + \beta_i) x^i$$

$$x \cdot \alpha = \sigma(\alpha) \cdot x$$

Skew polynomial rings

$$n = m$$

- $\mathbb{F}_{q^n}/\mathbb{F}_q$ finite field extension
- $\text{Gal}(\mathbb{F}_{q^n}/\mathbb{F}_q) = \langle \sigma \rangle$

$$R = \left\{ \sum_{i=0}^t \alpha_i x^i : \alpha_i \in \mathbb{F}_{q^m} \right\}$$

$$\sum_i \alpha_i x^i + \sum_i \beta_i x^i = \sum_i (\alpha_i + \beta_i) x^i$$

$$x \cdot \alpha = \sigma(\alpha) \cdot x$$

$$(R, +, \cdot)$$

Skew polynomial ring



O. Ore: Theory of non-commutative polynomials, *Annals of Mathematics*, (1933)

Skew polynomial rings

$$R = \left\{ \sum_{i=0}^t \alpha_i x^i : \alpha_i \in \mathbb{F}_{q^n} \right\}$$

$$f = \sum_i \alpha_i x^i$$

$$\phi_f : \beta \in \mathbb{F}_{q^n} \mapsto \sum_i \alpha_i \sigma^i(\beta) \in \mathbb{F}_{q^n}$$

\mathbb{F}_q -linear map

Skew polynomial rings

$$R = \left\{ \sum_{i=0}^t \alpha_i x^i : \alpha_i \in \mathbb{F}_{q^n} \right\}$$

$$f = \sum_i \alpha_i x^i$$

$$\phi_f : \beta \in \mathbb{F}_{q^n} \mapsto \sum_i \alpha_i \sigma^i(\beta) \in \mathbb{F}_{q^n}$$

\mathbb{F}_q -linear map

$$\frac{R}{R(x^n - 1)} \cong \text{End}_{\mathbb{F}_q}(\mathbb{F}_{q^n}) \cong M_n(\mathbb{F}_q)$$

$$\text{rk}(f) = \dim_{\mathbb{F}_q}(\text{Im}(\phi_f))$$

Skew polynomial rings

Rank-metric space:

- $X = \frac{R}{R(x^n-1)} = \left\{ f = \sum_{i=0}^{n-1} \alpha_i x^i : \alpha_i \in \mathbb{F}_{q^n} \right\}$
- $rk(f) = \dim_{\mathbb{F}_q}(Im(\phi_f))$
- $d_R(f, g) = rk(f - g)$

$$(M_n(\mathbb{F}_q), d_R) \cong \left(\frac{R}{R(x^n-1)}, d_R \right)$$

$$(M_n(\mathbb{F}_q), d_R) \cong \left(\frac{R}{R(x^n - 1)}, d_R \right) \text{ rank-metric space}$$

(Generalized) Gabidulin codes	$\{\alpha_0 + \alpha_1 x + \dots + \alpha_{k-1} x^{k-1} : \alpha_i \in \mathbb{F}_{q^n}\}$
(Generalized) Twisted Gabidulin codes	$\left\{ \alpha_0 + \sum_{i=1}^{k-1} \alpha_i x^i + \rho(\alpha_0) \eta x^k : \alpha_i \in \mathbb{F}_{q^n} \right\}$
Codes from scattered polynomials	$\{\alpha_0 + \alpha_1 f(x) : \alpha_0, \alpha_1 \in \mathbb{F}_{q^n}\}$
Trombetti-Zhou codes	$\left\{ \alpha'_0 + \sum_{i=1}^{k-1} \alpha_i x^i + \gamma \alpha'_k x^k : \alpha_i \in \mathbb{F}_{q^n}, \alpha'_0, \alpha'_k \in \mathbb{F}_{q^{n/2}} \right\}$

MRD codes

(Generalized) Gabidulin codes	$\{\alpha_0 + \alpha_1 x + \dots + \alpha_{k-1} x^{k-1} : \alpha_i \in \mathbb{F}_{q^n}\}$
(Generalized) Twisted Gabidulin codes	$\left\{ \alpha_0 + \sum_{i=1}^{k-1} \alpha_i x^i + \rho(\alpha_0) \eta x^k : \alpha_i \in \mathbb{F}_{q^n} \right\}$
Codes from scattered polynomials	$\{\alpha_0 + \alpha_1 f(x) : \alpha_0, \alpha_1 \in \mathbb{F}_{q^n}\}$
Trombetti-Zhou codes	$\left\{ \alpha'_0 + \sum_{i=1}^{k-1} \alpha'_i x^i + \gamma \alpha'_k x^k : \alpha_i \in \mathbb{F}_{q^n}, \alpha'_0, \alpha'_k \in \mathbb{F}_{q^{n/2}} \right\}$

MRD conditions

$$\bullet \quad 1 \leq k < n$$



P. Delsarte: Bilinear forms over a finite field, with applications to coding theory, *Journal of Combinatorial Theory, Series A* (1978)



E. Gabidulin: Theory of codes with maximum rank distance, *Problems of information transmission*, (1985)



A. Kshevetskiy and E. Gabidulin: The new construction of rank codes, *International Symposium on Information Theory*, (2005)

MRD codes

(Generalized) Gabidulin codes	$\{\alpha_0 + \alpha_1 x + \dots + \alpha_{k-1} x^{k-1} : \alpha_i \in \mathbb{F}_{q^n}\}$
(Generalized) Twisted Gabidulin codes	$\left\{ \alpha_0 + \sum_{i=1}^{k-1} \alpha_i x^i + \rho(\alpha_0) \eta x^k : \alpha_i \in \mathbb{F}_{q^n} \right\}$
Codes from scattered polynomials	$\{\alpha_0 + \alpha_1 f(x) : \alpha_0, \alpha_1 \in \mathbb{F}_{q^n}\}$
Trombetti-Zhou codes	$\left\{ \alpha'_0 + \sum_{i=1}^{k-1} \alpha_i x^i + \gamma \alpha'_k x^k : \alpha_i \in \mathbb{F}_{q^n}, \alpha'_0, \alpha'_k \in \mathbb{F}_{q^{n/2}} \right\}$

MRD conditions

- $1 \leq k < n$
- $N_{q^n/q}(\eta) \neq (-1)^{nk}$



J. Sheekey: A new family of linear maximum rank distance codes, *Advances in Mathematics of Communications*, (2016)



G. Lunardon, R. Trombetti, and Y. Zhou: Generalized twisted Gabidulin codes, *Journal of Combinatorial Theory, Series A*, (2018)



K. Otal and F. Ozbudak: Additive rank metric codes, *IEEE Transactions on Information Theory*, (2016)

(Generalized) Gabidulin codes	$\{\alpha_0 + \alpha_1 x + \dots + \alpha_{k-1} x^{k-1} : \alpha_i \in \mathbb{F}_{q^n}\}$
(Generalized) Twisted Gabidulin codes	$\{\alpha_0 + \sum_{i=1}^{k-1} \alpha_i x^i + \rho(\alpha_0) \eta x^k : \alpha_i \in \mathbb{F}_{q^n}\}$
Codes from scattered polynomials	$\{\alpha_0 + \alpha_1 f(x) : \alpha_0, \alpha_1 \in \mathbb{F}_{q^n}\}$
Trombetti-Zhou codes	$\{\alpha'_0 + \sum_{i=1}^{k-1} \alpha_i x^i + \gamma \alpha'_k x^k : \alpha_i \in \mathbb{F}_{q^n}, \alpha'_0, \alpha'_k \in \mathbb{F}_{q^{n/2}}\}$



D. Bartoli, B. Csajbók, A. Giannoni, G. G. Grimaldi, G. Longobardi, G. Lunardon, G. Marino, M. Montanucci, A. Neri, O. Polverino, PS, V. Smaldore, M. Timpanella, R. Trombetti, C. Zanella, Y. Zhou, F. Zullo...

MRD codes

(Generalized) Gabidulin codes	$\{\alpha_0 + \alpha_1 x + \dots + \alpha_{k-1} x^{k-1} : \alpha_i \in \mathbb{F}_{q^n}\}$
(Generalized) Twisted Gabidulin codes	$\left\{ \alpha_0 + \sum_{i=1}^{k-1} \alpha_i x^i + \rho(\alpha_0) \eta x^k : \alpha_i \in \mathbb{F}_{q^n} \right\}$
Codes from scattered polynomials	$\{\alpha_0 + \alpha_1 f(x) : \alpha_0, \alpha_1 \in \mathbb{F}_{q^n}\}$
Trombetti-Zhou codes	$\left\{ \alpha'_0 + \sum_{i=1}^{k-1} \alpha_i x^i + \gamma \alpha'_k x^k : \alpha_i \in \mathbb{F}_{q^n}, \alpha'_0, \alpha'_k \in \mathbb{F}_{q^{n/2}} \right\}$

MRD conditions

- $1 \leq k < n$
- $N_{q^n/q}(\gamma) \notin \mathbb{F}_q^{(2)}$ (q odd)



R. Trombetti and Y. Zhou: A new family of MRD codes in $\mathbb{F}_q^{2n \times 2n}$ with right and middle nuclei \mathbb{F}_{q^n} , *IEEE Transactions on Information Theory*, (2018)

Division Algebras

Definition

- \mathbb{F} field
- \mathbb{A} vector space over \mathbb{F}
-

$$\star : \mathbb{A} \times \mathbb{A} \rightarrow \mathbb{A}, \quad (a, b) \mapsto a \star b$$

(\mathbb{F} -bilinear map)

- $a \in \mathbb{A},$

$$L_a : b \in \mathbb{A} \mapsto a \star b \in \mathbb{A} \quad \text{invertible}$$




($\mathbb{A}, +, \star$) division algebra

- $1 \in \mathbb{A} \Rightarrow \mathbb{A}$ unital division algebra
- \star associative $\Rightarrow \mathbb{A}$ associative division algebra
- \star commutative $\Rightarrow \mathbb{A}$ commutative division algebra

Division algebras


- Every field is a division algebra
- Hamilton's quaternion algebra (1843)



$\downarrow \times \rightarrow$	1	i	j	k
1	1	i	j	k
i	i	-1	k	-j
j	j	-k	-1	i
k	k	j	-i	-1

(non-commutative, associative division algebra)

- Graves's Octonion algebra (1843)



\times	i	j	k	E	I	J	K
i	-1	k	-j	I	-E	-K	J
j	-k	-1	i	J	K	-E	-I
k	j	-i	-1	K	-J	I	-E
E	-I	-J	-K	-1	i	j	k
I	E	-K	j	-i	-1	-k	j
J	K	E	-i	-j	k	-1	-i
K	-j	I	E	-k	-j	i	-1

(non-commutative, non-associative division algebra)

$$|\mathbb{A}| \text{ finite} \Rightarrow \mathbb{A} \text{ (pre-)semifield}$$

Wedderburn's little theorem

Every associative semifield is a field



L. E. Dickson: On commutative linear algebras in which division is always uniquely possible, *Transactions of the American Mathematical Society* (1906)

- Dickson
- Hughes-Kleinfeld
- Knuth
- Cohen-Ganley
- Coulter-Matthews
- Jha-Johnson
- Dempwolff
- Kantor
- Budaghyan-Helleseth
- various subsets of
[Ebert-Johnson-Marino-Polverino-
Trombetti-Lunardon-Lavrauw]
- Zha-Kyureghyan-Wang
- Bierbrauer
- Pott-Zhou
- Bartoli-Bierbrauer-Kyureghyan-Giulietti-
Marcugini-Pambianco
- Sheekey
- Gologlu-Kölsch, Kölsch
- Lobillo-PS-Sheekey
- ...

Why Semifields?

- projective planes;
- spreads;
- PN functions;
- relative difference sets;
- additive Hamming-metric codes;
- MRD codes with $d = n$.

Division Algebras

- $(\mathbb{A}, +, \star)$ semifield over \mathbb{F}_q

Division Algebras

- $(\mathbb{A}, +, \star)$ semifield over \mathbb{F}_q
- $a \in \mathbb{A},$

$$L_a : b \in \mathbb{A} \longmapsto a \star b \in \mathbb{A},$$

$$L_a \in \text{End}_{\mathbb{F}_q}(\mathbb{A})$$

-

$$\mathcal{C}(\mathbb{A}) := \{L_a : a \in \mathbb{A}\} \subset \text{End}_{\mathbb{F}_q}(\mathbb{A}) \cong M_n(\mathbb{F}_q), \quad \dim_{\mathbb{F}_q}(\mathbb{A}) = n$$

(spread set of \mathbb{A})

Division Algebras

- $(\mathbb{A}, +, \star)$ semifield over \mathbb{F}_q
- $a \in \mathbb{A},$

$$L_a : b \in \mathbb{A} \mapsto a \star b \in \mathbb{A},$$

$$L_a \in \text{End}_{\mathbb{F}_q}(\mathbb{A})$$

-

$$\mathcal{C}(\mathbb{A}) := \{L_a : a \in \mathbb{A}\} \subset \text{End}_{\mathbb{F}_q}(\mathbb{A}) \cong M_n(\mathbb{F}_q), \quad \dim_{\mathbb{F}_q}(\mathbb{A}) = n$$

(spread set of \mathbb{A})

Theorem (J. De la Cruz, M. Kiermaier, A. Wassermann, and W. Willems (2016) - A. Gruica, A. Ravagnani, J. Sheekey, and F. Zullo (2023))

- $\mathcal{C}(\mathbb{A}) \subseteq M_n(\mathbb{F}_q)$ is an MRD code with $n = d$
- There is a one-to-one correspondence between *isotopy classes of semifields* and *equivalence classes of MRD codes* in $M_n(\mathbb{F}_q)$ with minimum distance $d = n$



J. Sheekey: New semifields and new MRD codes from skew polynomial rings, *Journal of the London Mathematical Society*, (2020)



J. Sheekey: New semifields and new MRD codes from skew polynomial rings, *Journal of the London Mathematical Society*, (2020)

$$(M_n(\mathbb{F}_q), d_R) \cong \left(\frac{R}{R(x^n - 1)}, d_R \right)$$



J. Sheekey: New semifields and new MRD codes from skew polynomial rings, *Journal of the London Mathematical Society*, (2020)

$$(M_n(\mathbb{F}_q), d_R) \cong \left(\frac{R}{R(x^n - 1)}, d_R \right)$$

$x^n - 1$ central element of R

Codes from skew polynomial rings

Centre of skew polynomial rings

$$R = \left\{ \sum_{i=0}^t \alpha_i x^i : \alpha_i \in \mathbb{F}_{q^m} \right\}$$

$$Z(R) = \{ F(x^n) : F(y) \in \mathbb{F}_q[y] \} \cong \mathbb{F}_q[y]$$

Codes from skew polynomial rings

Centre of skew polynomial rings

$$R = \left\{ \sum_{i=0}^t \alpha_i x^i : \alpha_i \in \mathbb{F}_{q^m} \right\}$$

$$Z(R) = \{ F(x^n) : F(y) \in \mathbb{F}_q[y] \} \cong \mathbb{F}_q[y]$$

$y \neq F(y)$ irreducible $\implies RF(x^n)$ two-sided maximal ideal

Codes from skew polynomial rings

Centre of skew polynomial rings

$$R = \left\{ \sum_{i=0}^t \alpha_i x^i : \alpha_i \in \mathbb{F}_{q^m} \right\}$$

$$Z(R) = \{F(x^n) : F(y) \in \mathbb{F}_q[y]\} \cong \mathbb{F}_q[y]$$

$y \neq F(y)$ irreducible $\implies RF(x^n)$ two-sided maximal ideal

$F(y) \in \mathbb{F}_q[y]$ irreducible polynomial, $\deg(F) = s$

Codes from skew polynomial rings

Centre of skew polynomial rings

$$R = \left\{ \sum_{i=0}^t \alpha_i x^i : \alpha_i \in \mathbb{F}_{q^m} \right\}$$

$$Z(R) = \{F(x^n) : F(y) \in \mathbb{F}_q[y]\} \cong \mathbb{F}_q[y]$$

$y \neq F(y)$ irreducible $\implies RF(x^n)$ two-sided maximal ideal

$F(y) \in \mathbb{F}_q[y]$ irreducible polynomial, $\deg(F) = s$

\Downarrow

$R_F := \frac{R}{RF(x^n)}$ simple and left Artinian ring

Codes from skew polynomial rings

Centre of skew polynomial rings

$$R = \left\{ \sum_{i=0}^t \alpha_i x^i : \alpha_i \in \mathbb{F}_{q^m} \right\}$$

$$Z(R) = \{F(x^n) : F(y) \in \mathbb{F}_q[y]\} \cong \mathbb{F}_q[y]$$

$y \neq F(y)$ irreducible $\implies RF(x^n)$ two-sided maximal ideal

$F(y) \in \mathbb{F}_q[y]$ irreducible polynomial, $\deg(F) = s$

\Downarrow

$R_F := \frac{R}{RF(x^n)}$ simple and left Artinian ring

\Downarrow

$\varphi_F : R_F \cong M_n(\mathbb{F}_{q^s})$ (by Artin-Wedderburn Theorem)

Codes from skew polynomial rings

Centre of skew polynomial rings

$$R = \left\{ \sum_{i=0}^t \alpha_i x^i : \alpha_i \in \mathbb{F}_{q^m} \right\}$$

$$\varphi_F : R_F = \frac{R}{RF(x^n)} \cong M_n(\mathbb{F}_{q^s})$$

$$\Downarrow$$

$$a \in R_F, \quad rk(a) := rk(\varphi_F(a))$$

$$\Downarrow$$

$$(R_F, d_R) \cong (M_n(\mathbb{F}_{q^s}), d_R)$$

$$RF(x^n)$$

$$\Downarrow$$

$$\varphi_F : R_F \cong M_n(\mathbb{F}_{q^s}) \text{ (by Artin-Wedderburn Theorem)}$$

Codes from skew polynomial rings

Centre of skew polynomial rings

$$R = \left\{ \sum_{i=0}^t \alpha_i x^i : \alpha_i \in \mathbb{F}_{q^m} \right\}$$

$$\varphi_F : R_F = \frac{R}{RF(x^n)} \cong M_n(\mathbb{F}_{q^s})$$

$$\Downarrow$$

$$a \in R_F, \quad rk(a) := rk(\varphi_F(a))$$

$$\Downarrow$$

$$(R_F, d_R) \cong (M_n(\mathbb{F}_{q^s}), d_R)$$

$$RF(x^n)$$

$$F(y) = y - 1$$

$$R_F \cong \frac{R}{R(x^n - 1)} \cong M_n(\mathbb{F}_q)$$

MRD codes from skew polynomial rings

$$(R_F, d_R) \cong (M_n(\mathbb{F}_{q^s}), d_R)$$

Theorem (J. Sheekey, 2020)

- $\rho \in \text{Aut}(\mathbb{F}_{q^n})$
- $1 \leq k < n$
- $\eta \in \mathbb{F}_{q^n}$: $N_{q^n/q'}(\eta)N_{q/q'}((-1)^{sk(n-1)}F(0)^k) \neq 1$

$$S(F) := \left\{ \alpha_0 + \sum_{i=1}^{sk-1} \alpha_i x^i + \eta \rho(\alpha_0) x^{sk} : \alpha_i \in \mathbb{F}_{q^n} \right\} \subseteq R_F \cong M_n(\mathbb{F}_{q^s}).$$

\Downarrow

$S(F)$ is an MRD code in $M_n(\mathbb{F}_{q^s})$ with $|S(F)| = q^{nsk}$ and $d(S(F)) = n - k + 1$

MRD codes from skew polynomial rings

$$(R_F, d_R) \cong (M_n(\mathbb{F}_{q^s}), d_R)$$

Theorem (J. Sheekey, 2020)

$$S(F) := \left\{ \alpha_0 + \sum_{i=1}^{sk-1} \alpha_i x^i + \eta \rho(\alpha_0) x^{sk} : \alpha_i \in \mathbb{F}_{q^n} \right\} \subseteq R_F \cong M_n(\mathbb{F}_{q^s}).$$

MRD codes from skew polynomial rings

$$(R_F, d_R) \cong (M_n(\mathbb{F}_{q^s}), d_R)$$

Theorem (J. Sheekey, 2020)

$$S(F) := \left\{ \alpha_0 + \sum_{i=1}^{sk-1} \alpha_i x^i + \eta \rho(\alpha_0) x^{sk} : \alpha_i \in \mathbb{F}_{q^n} \right\} \subseteq R_F \cong M_n(\mathbb{F}_{q^s}).$$

$s = 1$, $F(y) = y - 1$ and $\eta = 0$

$$S(F) = \left\{ \alpha_0 + \sum_{i=1}^{k-1} \alpha_i x^i : \alpha_i \in \mathbb{F}_{q^n} \right\} \subseteq R_F = \frac{R}{R(x^n - 1)} \cong M_n(\mathbb{F}_q)$$

(Generalized) Gabidulin codes

MRD codes from skew polynomial rings

$$(R_F, d_R) \cong (M_n(\mathbb{F}_{q^s}), d_R)$$

Theorem (J. Sheekey, 2020)

$$S(F) := \left\{ \alpha_0 + \sum_{i=1}^{sk-1} \alpha_i x^i + \eta \rho(\alpha_0) x^{sk} : \alpha_i \in \mathbb{F}_{q^n} \right\} \subseteq R_F \cong M_n(\mathbb{F}_{q^s}).$$

$s = 1$, $F(y) = y - 1$ and $\eta \neq 0$

$$S(F) = \left\{ \alpha_0 + \sum_{i=1}^{k-1} \alpha_i x^i + \eta \rho(\alpha_0) : \alpha_i \in \mathbb{F}_{q^n} \right\} \subseteq R_F = \frac{R}{R(x^n - 1)} \cong M_n(\mathbb{F}_q)$$

(Generalized) Twisted Gabidulin codes

MRD codes from skew polynomial rings

$$(R_F, d_R) \cong (M_n(\mathbb{F}_{q^s}), d_R)$$

Theorem (J. Sheekey, 2020)

$$S(F) := \left\{ \alpha_0 + \sum_{i=1}^{sk-1} \alpha_i x^i + \eta \rho(\alpha_0) x^{sk} : \alpha_i \in \mathbb{F}_{q^n} \right\} \subseteq R_F \cong M_n(\mathbb{F}_{q^s}).$$

$k = 1$ and $\eta = 0$

$$S(F) = \left\{ \alpha_0 + \sum_{i=1}^{s-1} \alpha_i x^i : \alpha_i \in \mathbb{F}_{q^n} \right\} \subseteq R_F = \frac{R}{R(F(x^n))} \cong M_n(\mathbb{F}_{q^s})$$

Sandler's semifields/cyclic semifields

MRD codes from skew polynomial rings

$$(R_F, d_R) \cong (M_n(\mathbb{F}_{q^s}), d_R)$$

Theorem (J. Sheekey, 2020)

- $\rho \in \text{Aut}(\mathbb{F}_{q^n})$
- $1 \leq k < n$
- $N_{q^n/q'}(\eta)N_{q/q'}((-1)^{sk(n-1)}F(0)^k) \neq 1$

$$S(F) := \left\{ \alpha_0 + \sum_{i=1}^{sk-1} \alpha_i x^i + \eta \rho(\alpha_0) x^{sk} : \alpha_i \in \mathbb{F}_{q^n} \right\} \subseteq R_F \cong M_n(\mathbb{F}_{q^s}).$$

\Downarrow

$S(F)$ is an MRD code in $M_n(\mathbb{F}_{q^s})$

Theorem (J. Sheekey, 2020)

The family $S(F)$ contains new semifields and new MRD codes for infinite choices of s and n (and k).

Codes from skew polynomial rings

$$(R_F, d_R) \cong (M_n(\mathbb{F}_{q^s}), d_R)$$



F.J. Lobillo, PS, and J. Sheekey: Quotients of skew polynomial rings: new constructions of division algebras and MRD codes, *arXiv preprint arXiv:2502.13531*, (2025)

Codes from skew polynomial rings

$$(R_F, d_R) \cong (M_n(\mathbb{F}_{q^s}), d_R)$$

- n even
- q odd

Codes from skew polynomial rings

$$(R_F, d_R) \cong (M_n(\mathbb{F}_{q^s}), d_R)$$

- n even
- q odd

$$\left\{ \alpha'_0 + \sum_{i=1}^{k-1} \alpha_i x^i + \gamma \alpha'_k x^k : \alpha_i \in \mathbb{F}_{q^n}, \alpha'_0, \alpha'_k \in \mathbb{F}_{q^{n/2}} \right\} \subseteq \frac{R}{R(x^n - 1)}$$

Codes from skew polynomial rings

$$(R_F, d_R) \cong (M_n(\mathbb{F}_{q^s}), d_R)$$

- n even
- q odd

$$\left\{ \alpha'_0 + \sum_{i=1}^{k-1} \alpha_i x^i + \gamma \alpha'_k x^k : \alpha_i \in \mathbb{F}_{q^n}, \alpha'_0, \alpha'_k \in \mathbb{F}_{q^{n/2}} \right\} \subseteq \frac{R}{R(x^n - 1)}$$

Theorem (F.J. Lobillo, PS and J. Sheekey, 2025)

$$D(F) := \left\{ \alpha'_0 + \sum_{i=1}^{sk-1} \alpha_i x^i + \gamma \alpha'_{sk} x^{sk} : \alpha_i \in \mathbb{F}_{q^n}, \alpha'_0, \alpha'_{sk} \in \mathbb{F}_{q^{n/2}} \right\} \subseteq R_F \cong M_n(\mathbb{F}_{q^s}).$$

Codes from skew polynomial rings

$$(R_F, d_R) \cong (M_n(\mathbb{F}_{q^s}), d_R)$$

- n even
- q odd

$$\left\{ \alpha'_0 + \sum_{i=1}^{k-1} \alpha_i x^i + \gamma \alpha'_k x^k : \alpha_i \in \mathbb{F}_{q^n}, \alpha'_0, \alpha'_k \in \mathbb{F}_{q^{n/2}} \right\} \subseteq \frac{R}{R(x^n - 1)}$$

Theorem (F.J. Lobillo, PS and J. Sheekey, 2025)

$$D(F) := \left\{ \alpha'_0 + \sum_{i=1}^{sk-1} \alpha_i x^i + \gamma \alpha'_{sk} x^{sk} : \alpha_i \in \mathbb{F}_{q^n}, \alpha'_0, \alpha'_{sk} \in \mathbb{F}_{q^{n/2}} \right\} \subseteq R_F \cong M_n(\mathbb{F}_{q^s}).$$

$$(-1)^{ks} F(0)^k N_{\mathbb{F}_{q^n}/\mathbb{F}_q}(\gamma) \notin \mathbb{F}_q^{(2)}$$

Codes from skew polynomial rings

$$(R_F, d_R) \cong (M_n(\mathbb{F}_{q^s}), d_R)$$

- n even
- q odd

$$\left\{ \alpha'_0 + \sum_{i=1}^{k-1} \alpha_i x^i + \gamma \alpha'_k x^k : \alpha_i \in \mathbb{F}_{q^n}, \alpha'_0, \alpha'_k \in \mathbb{F}_{q^{n/2}} \right\} \subseteq \frac{R}{R(x^n - 1)}$$

Theorem (F.J. Lobillo, PS and J. Sheekey, 2025)

$$D(F) := \left\{ \alpha'_0 + \sum_{i=1}^{sk-1} \alpha_i x^i + \gamma \alpha'_{sk} x^{sk} : \alpha_i \in \mathbb{F}_{q^n}, \alpha'_0, \alpha'_{sk} \in \mathbb{F}_{q^{n/2}} \right\} \subseteq R_F \cong M_n(\mathbb{F}_{q^s}).$$

$$(-1)^{ks} F(0)^k N_{\mathbb{F}_{q^n}/\mathbb{F}_q}(\gamma) \notin \mathbb{F}_q^{(2)}$$

\Downarrow

$D(F)$ is an **MRD code** in $M_n(\mathbb{F}_{q^s})$ with $|D(F)| = q^{nsk}$ and $d(D(F)) = n - k + 1$

Codes from skew polynomial rings

Theorem (F.J. Lobillo, PS, J. Sheekey 2025)

$$D(F) := \left\{ \alpha'_0 + \sum_{i=1}^{sk-1} \alpha_i x^i + \gamma \alpha'_{sk} x^{sk} : \alpha_i \in \mathbb{F}_{q^n}, \alpha'_0, \alpha'_{sk} \in \mathbb{F}_{q^{n/2}} \right\} \subseteq R_F \cong M_n(\mathbb{F}_{q^s}).$$

$$(-1)^{ks} F(0)^k N_{\mathbb{F}_{q^n}/\mathbb{F}_q}(\gamma) \notin \mathbb{F}_q^{(2)}$$

\Downarrow

$D(F)$ is an **MRD code** in $M_n(\mathbb{F}_{q^s})$ with $|D(F)| = q^{nsk}$ and $d(D(F)) = n - k + 1$

Corollary (F.J. Lobillo, PS and J. Sheekey, 2025)

For $k = 1$, for every s , $D(F)$ defines a **semifield** over \mathbb{F}_q , with $|D(F)| = q^{sn}$.

Codes from skew polynomial rings

$$(R_F, d_R) \cong (M_n(\mathbb{F}_{q^s}), d_R)$$

Theorem (F.J. Lobillo, PS, and J. Sheekey, 2025)

$$D(F) := \left\{ \alpha'_0 + \sum_{i=1}^{sk-1} \alpha_i x^i + \gamma \alpha'_k x^{sk} : \alpha_i \in \mathbb{F}_{q^n}, \alpha'_0, \alpha'_k \in \mathbb{F}_{q^{n/2}} \right\} \subseteq R_F \cong M_n(\mathbb{F}_{q^s}).$$

$s = 1$ and $F(y) = y - 1$

$$D(F) = \left\{ \alpha'_0 + \sum_{i=1}^{k-1} \alpha_i x^i + \gamma \alpha'_k : \alpha_i \in \mathbb{F}_{q^n}, \alpha'_0, \alpha'_k \in \mathbb{F}_{q^{n/2}} \right\} \subseteq R_F = \frac{R}{R(x^n - 1)} \cong M_n(\mathbb{F}_q)$$

Trombetti-Zhou codes

Codes from skew polynomial rings

$$(R_F, d_R) \cong (M_n(\mathbb{F}_{q^s}), d_R)$$

Theorem (F.J. Lobillo, PS, J. Sheekey 2025)

$$D(F) := \left\{ \alpha'_0 + \sum_{i=1}^{sk-1} \alpha_i x^i + \gamma \alpha'_{sk} x^{sk} : \alpha_i \in \mathbb{F}_{q^n}, \alpha'_0, \alpha'_{sk} \in \mathbb{F}_{q^{n/2}} \right\} \subseteq R_F \cong M_n(\mathbb{F}_{q^s}).$$

$k = 1$, $F(y) = y - 1$ and $s = 1$

$$D(F) = \{ \alpha'_0 + \gamma \alpha'_1 : \alpha'_0, \alpha'_1 \in \mathbb{F}_{q^{n/2}} \} \subseteq R_F = \frac{R}{RF(x^n)} \cong M_n(\mathbb{F}_q)$$

Hughes-Kleinfeld semifields

Equivalence Issue

$(\mathbb{A}, +, \star)$ semifield

$$\mathbb{N}_l(\mathbb{A}) = \{a \in \mathbb{A} : a \star (b \star c) = (a \star b) \star c, \text{ for all } b, c \in \mathbb{A}\}$$

(Left nucleus)

$$\mathbb{N}_m(\mathbb{A}) = \{b \in \mathbb{A} : a \star (b \star c) = (a \star b) \star c, \text{ for all } a, c \in \mathbb{A}\},$$

(Middle nucleus)

$$\mathbb{N}_r(\mathbb{A}) = \{c \in \mathbb{A} : a \star (b \star c) = (a \star b) \star c, \text{ for all } a, b \in \mathbb{A}\},$$

(Right nucleus)

$$Z(\mathbb{A}) = \mathbb{N}_l(\mathbb{A}) \cap \mathbb{N}_m(\mathbb{A}) \cap \mathbb{N}_r(\mathbb{A}) \cap \{a \in \mathbb{A} : a \star b = b \star a \text{ for all } b \in \mathbb{A}\}.$$

(center)

Equivalence Issue

$$\mathcal{C} \subseteq M_n(\mathbb{F})$$

$$L(\mathcal{C}) = \{A \in M_n(\mathbb{F}) : A\mathcal{C} \subseteq \mathcal{C}\}$$

(Left idealiser)

$$R(\mathcal{C}) = \{B \in M_n(\mathbb{F}) : \mathcal{C}B \subseteq \mathcal{C}\},$$

(Right idealiser)

$$\text{Cen}(\mathcal{C}) = \{A \in M_n(\mathbb{F}) : AX = XA \text{ for every } X \in \mathcal{C}\},$$

(centraliser)

$$Z(\mathcal{C}) = L(\mathcal{C}) \cap R(\mathcal{C}).$$

(center)

Equivalence Issue

Theorem (G. Lunardon, R. Trombetti, and Y. Zhou, 2017- J. Sheekey, 2020)

- $(\mathbb{A}, +, \star)$ semifield, $\mathcal{C} = \mathcal{C}(\mathbb{A})$ spread set. Then

$$|L(\mathcal{C})| = |\mathbb{N}_l(\mathbb{A})|$$

$$|R(\mathcal{C})| = |\mathbb{N}_m(\mathbb{A})|$$

$$|Cen(\mathcal{C})| = |\mathbb{N}_r(\mathbb{A})|$$

$$|Z(\mathcal{C})| = |Z(\mathbb{A})|$$

- $I_n \in \mathcal{C}, \mathcal{C}' \subseteq M_n(\mathbb{F}_q)$. If $\mathcal{C} \sim \mathcal{C}'$, then

$$|L(\mathcal{C})| = |L(\mathcal{C}')|$$

$$|R(\mathcal{C})| = |R(\mathcal{C}')|$$

$$|Cen(\mathcal{C})| = |Cen(\mathcal{C}')|$$

$$|Z(\mathcal{C})| = |Z(\mathcal{C}')|$$

Equivalence Issue

Theorem (G. Lunardon, R. Trombetti, and Y. Zhou, 2017- J. Sheekey, 2020)

- $(\mathbb{A}, +, \star)$ semifield, $\mathcal{C} = \mathcal{C}(\mathbb{A})$ spread set. Then

$$|L(\mathcal{C})| = |\mathbb{N}_l(\mathbb{A})|$$

$$|R(\mathcal{C})| = |\mathbb{N}_m(\mathbb{A})|$$

$$|Cen(\mathcal{C})| = |\mathbb{N}_r(\mathbb{A})|$$

$$|Z(\mathcal{C})| = |Z(\mathbb{A})|$$

- $I_n \in \mathcal{C}, \mathcal{C}' \subseteq M_n(\mathbb{F}_q)$. If $\mathcal{C} \sim \mathcal{C}'$, then

$$(|\mathcal{C}|, |L(\mathcal{C})|, |R(\mathcal{C})|, |Cen(\mathcal{C})|, |Z(\mathcal{C})|)$$

Nuclear parameters of \mathcal{C}

$$|Cen(\mathcal{C})| = |Cen(\mathcal{C}')|$$

$$|Z(\mathcal{C})| = |Z(\mathcal{C}')|$$

Theorem (F.J. Lobillo, PS, and J. Sheekey, 2025)

$\mathcal{C} = D(F)$, $k \leq n/2$. Then

$$L(\mathcal{C}) \cong \mathbb{F}_{q^{n/2}} \quad R(\mathcal{C}) \cong \mathbb{F}_{q^{n/2}} \quad \text{Cen}(\mathcal{C}) \cong \mathbb{F}_{q^s} \quad Z(\mathcal{C}) \cong \mathbb{F}_q$$

Equivalence Issue

Theorem (F.J. Lobillo, PS, and J. Sheekey, 2025)

$\mathcal{C} = D(F)$, $k \leq n/2$. Then

$$L(\mathcal{C}) \cong \mathbb{F}_{q^{n/2}} \quad R(\mathcal{C}) \cong \mathbb{F}_{q^{n/2}} \quad \text{Cen}(\mathcal{C}) \cong \mathbb{F}_{q^s} \quad Z(\mathcal{C}) \cong \mathbb{F}_q$$

Theorem (F.J. Lobillo, PS, and J. Sheekey, 2025)

The family $D(F)$ contains **new semifields** and **new MRD codes** for infinite choices of s and n .

What's next?

Why $F(y)$ irreducible?!

What's next?

Why $F(y)$ irreducible?!

$$F(y) = F_1(y) \cdots F_t(y), \quad \gcd(F_i, F_j) = 1$$

F_i irreducible polynomial

What's next?

Why $F(y)$ irreducible?!

$$F(y) = F_1(y) \cdots F_t(y), \quad \gcd(F_i, F_j) = 1$$

F_i irreducible polynomial

\Downarrow

$$\frac{R}{RF(x^n)} \cong \bigoplus_{i=1}^t \frac{R}{RF_i(x^n)} \cong \bigoplus_{i=1}^t M_n(\mathbb{F}_{q^{s_i}}), \quad s_i = \deg(F_i).$$

What's next?

Why $F(y)$ irreducible?!

$$F(y) = F_1(y) \cdots F_t(y), \quad \gcd(F_i, F_j) = 1$$

F_i irreducible polynomial

\Downarrow

$$\frac{R}{RF(x^n)} \cong \bigoplus_{i=1}^t \frac{R}{RF_i(x^n)} \cong \bigoplus_{i=1}^t M_n(\mathbb{F}_{q^{s_i}}), \quad s_i = \deg(F_i).$$

\Downarrow

$$\left(\frac{R}{RF(x^n)}, \text{srk} \right) \cong \left(\bigoplus_{i=1}^t M_n(\mathbb{F}_{q^{s_i}}), \text{srk} \right)$$

What's next?

Why $F(y)$ irreducible?!

$$F(y) = F_1(y) \cdots F_t(y), \quad \gcd(F_i, F_j) = 1$$

F_i irreducible polynomial



A. Neri and PS: Sum-rank metric codes and additive MDS codes from quotients of skew polynomial rings, *in preparation*.

New constructions of MSRD codes and additive MDS codes

$$\left(\frac{R}{RF(x^n)}, \text{srk} \right) \cong \left(\bigoplus_{i=1}^t M_n(\mathbb{F}_{q^{s_i}}), \text{srk} \right)$$

What's next?

$$F(y) = F_1(y) \cdots F_t(y), \quad \gcd(F_i, F_j) = 1$$

F_i irreducible polynomial, $\deg(F_i) = s$

$$\left(\frac{R}{RF(x^n)}, \text{srk} \right) \cong \left(\bigoplus_{i=1}^t M_n(\mathbb{F}_{q^s}), \text{srk} \right)$$

Theorem (A. Neri, and PS)

Let $a \in R/RF(x^n)$. Then

$$\text{srk}(a) = tn - \frac{1}{s} \deg(\gcd(a, F(x^n)))$$

What's next?

$$\left(\frac{R}{RF(x^n)}, \text{srk} \right) \cong \left(\bigoplus_{i=1}^t M_n(\mathbb{F}_{q^s}), \text{srk} \right)$$

Theorem (A. Neri, and PS)

$$S(F) := \left\{ \alpha_0 + \sum_{i=1}^{sk-1} \alpha_i x^i + \eta \rho(\alpha_0) x^{sk} : \alpha_i \in \mathbb{F}_{q^n} \right\} \subseteq R_F \cong \bigoplus_{i=1}^t M_n(\mathbb{F}_{q^s})$$

What's next?

$$\left(\frac{R}{RF(x^n)}, \text{srk} \right) \cong \left(\bigoplus_{i=1}^t M_n(\mathbb{F}_{q^s}), \text{srk} \right)$$

Theorem (A. Neri, and PS)

$$S(F) := \left\{ \alpha_0 + \sum_{i=1}^{sk-1} \alpha_i x^i + \eta \rho(\alpha_0) x^{sk} : \alpha_i \in \mathbb{F}_{q^n} \right\} \subseteq R_F \cong \bigoplus_{i=1}^t M_n(\mathbb{F}_{q^s})$$

$$\mathbb{N}_{\mathbb{F}_{q^n}/\mathbb{K}'}(\eta) \mathbb{N}_{\mathbb{F}_q/\mathbb{K}'} \left((-1)^{sk(n-1)} \prod_{i=1}^t F_{i,0}^{j_i} \right) \neq 1, \quad \text{with } j_1 + \dots + j_t = k$$

What's next?

$$\left(\frac{R}{RF(x^n)}, \text{srk} \right) \cong \left(\bigoplus_{i=1}^t M_n(\mathbb{F}_{q^s}), \text{srk} \right)$$

Theorem (A. Neri, and PS)

$$S(F) := \left\{ \alpha_0 + \sum_{i=1}^{sk-1} \alpha_i x^i + \eta \rho(\alpha_0) x^{sk} : \alpha_i \in \mathbb{F}_{q^n} \right\} \subseteq R_F \cong \bigoplus_{i=1}^t M_n(\mathbb{F}_{q^s})$$

$$\mathbb{N}_{\mathbb{F}_{q^n}/\mathbb{K}'}(\eta) \mathbb{N}_{\mathbb{F}_q/\mathbb{K}'} \left((-1)^{sk(n-1)} \prod_{i=1}^t F_{i,0}^{j_i} \right) \neq 1, \quad \text{with } j_1 + \dots + j_t = k$$

\Downarrow

$$S(F) \text{ is an MSRD code in } \bigoplus_{i=1}^t M_n(\mathbb{F}_{q^s}) \text{ with } d(S(F)) = tn - k + 1$$



U. Martínez-Peñas: Skew and linearized Reed-Solomon codes and maximum sum rank distance codes over any division ring. *Journal of Algebra*, (2018) ([LRS codes](#))



A. Neri: Twisted linearized Reed-Solomon codes: A skew polynomial framework. *Journal of Algebra*, (2022) ([TLRS codes](#))

What's next?

$$\left(\frac{R}{RF(x^n)}, \text{srk} \right) \cong \left(\bigoplus_{i=1}^t M_n(\mathbb{F}_{q^s}), \text{srk} \right)$$

Theorem (A. Neri, and PS, 2025)

$$D(F) := \left\{ \alpha'_0 + \sum_{i=1}^{sk-1} \alpha_i x^i + \gamma \alpha'_{sk} x^{sk} : \alpha_i \in \mathbb{F}_{q^n}, \alpha'_0, \alpha'_{sk} \in \mathbb{F}_{q^{n/2}} \right\} \subseteq R_F \cong M_n(\mathbb{F}_{q^s}).$$

What's next?

$$\left(\frac{R}{RF(x^n)}, \text{srk} \right) \cong \left(\bigoplus_{i=1}^t M_n(\mathbb{F}_{q^s}), \text{srk} \right)$$

Theorem (A. Neri, and PS, 2025)

$$D(F) := \left\{ \alpha'_0 + \sum_{i=1}^{sk-1} \alpha_i x^i + \gamma \alpha'_{sk} x^{sk} : \alpha_i \in \mathbb{F}_{q^n}, \alpha'_0, \alpha'_{sk} \in \mathbb{F}_{q^{n/2}} \right\} \subseteq R_F \cong M_n(\mathbb{F}_{q^s}).$$

$$(-1)^{sk(n-1)} \prod_{i=1}^t F_{i,0}^{j_i} N_{\mathbb{F}_{q^n}/\mathbb{F}_q}(\gamma) \notin \mathbb{F}_q^{(2)}, \quad \text{with } j_1 + \dots + j_t = k.$$

What's next?

$$\left(\frac{R}{RF(x^n)}, \text{srk} \right) \cong \left(\bigoplus_{i=1}^t M_n(\mathbb{F}_{q^s}), \text{srk} \right)$$

Theorem (A. Neri, and PS, 2025)

$$D(F) := \left\{ \alpha'_0 + \sum_{i=1}^{sk-1} \alpha_i x^i + \gamma \alpha'_{sk} x^{sk} : \alpha_i \in \mathbb{F}_{q^n}, \alpha'_0, \alpha'_{sk} \in \mathbb{F}_{q^{n/2}} \right\} \subseteq R_F \cong M_n(\mathbb{F}_{q^s}).$$

$$(-1)^{sk(n-1)} \prod_{i=1}^t F_{i,0}^{j_i} N_{\mathbb{F}_{q^n}/\mathbb{F}_q}(\gamma) \notin \mathbb{F}_q^{(2)}, \quad \text{with } j_1 + \dots + j_t = k.$$

\Downarrow

$D(F)$ is an **MSRD code** in $\bigoplus_{i=1}^t M_n(\mathbb{F}_{q^s})$ with $d(D(F)) = tn - k + 1$



A. Neri: Twisted linearized Reed-Solomon codes: A skew polynomial framework. *Journal of Algebra*, (2022) (**TLRS codes of TZ-type**)

Thank you for your attention!