

Design switching on graphs

Robin Simoens

Ghent University & Universitat Politècnica de Catalunya



4 September 2025

Based on joint work with Ferdinand Ihringer (SUSTech)

Design switching on graphs

Definition

An (r, λ) -**design** is an incidence structure where

- every point is in r blocks,
- every two points are in λ blocks.




Figure: An $(r = 3, \lambda = 1)$ -design

Design switching on graphs

Definition

An (r, λ) -**design** is an incidence structure where

- every point is in r blocks,
- every two points are in λ blocks.




Figure: An $(r = 3, \lambda = 1)$ -design

Definition

Switching is a local graph operation, resulting in a *cospectral* graph.

Cospectral graphs

Definition

Cospectral graphs have the same adjacency spectrum.



Figure: Cospectral graphs. Both have spectrum $\{-2, 0, 0, 0, 2\}$.

Cospectral graphs

Definition

Cospectral graphs have the same adjacency spectrum.



Figure: Cospectral graphs. Both have spectrum $\{-2, 0, 0, 0, 2\}$.

Conjecture (van Dam and Haemers, 2003)

Almost all graphs are determined by their spectrum.

Cospectral graphs

Conjecture (van Dam and Haemers, 2003)

Almost all graphs are determined by their spectrum.

Cospectral graphs

Conjecture (van Dam and Haemers, 2003)

Almost all graphs are determined by their spectrum.

► Interesting for complexity theory




Figure: Is graph isomorphism an easy or hard problem?

Cospectral graphs

Conjecture (van Dam and Haemers, 2003)

Almost all graphs are determined by their spectrum.

- Interesting for complexity theory
- Interesting for chemistry




Figure: The molecular graph of acetaldehyde (ethanal).

Cospectral graphs

Conjecture (van Dam and Haemers, 2003)

Almost all graphs are determined by their spectrum.

- Interesting for complexity theory
- Interesting for chemistry

😊 Computational evidence [[Brouwer and Spence, 2009](#)]

n	3	4	5	6	7	8	9	10	11
ratio	1	1	0.941	0.936	0.895	0.861	0.814	0.787	0.789

Cospectral graphs

Conjecture (van Dam and Haemers, 2003)

Almost all graphs are determined by their spectrum.

- Interesting for complexity theory
- Interesting for chemistry

😊 Computational evidence [Brouwer and Spence, 2009]

😞 Almost all

- trees [Schwenk, 1973]
- strongly regular graphs [Fon-Der-Flaass, 2002]
- cographs [Wang and Huang, 2025]

are **not** determined by their spectrum.

Cospectral graphs

Conjecture (van Dam and Haemers, 2003)

Almost all graphs are determined by their spectrum.

- Interesting for complexity theory
- Interesting for chemistry

😊 Computational evidence [Brouwer and Spence, 2009]

😞 Almost all

- trees [Schwenk, 1973]
- strongly regular graphs [Fon-Der-Flaass, 2002]
- cographs [Wang and Huang, 2025]


are **not** determined by their spectrum.

😊 Exponentially many graphs are determined by their spectrum [Koval and Kwan, 2023]

How to find cospectral graphs

Theorem (GM_4 switching, Godsil and McKay, 1982)


Consider a graph with a regular subgraph C of size 4 such that every vertex $x \notin C$ has 0, 2 or 4 neighbours in C . If $x \notin C$ has 2 neighbours in C , reverse its adjacencies with C . The obtained graph is cospectral.



How to find cospectral graphs

Theorem (GM₄ switching, Godsil and McKay, 1982)


Consider a graph with a regular subgraph C of size 4 such that every vertex $x \notin C$ has 0, 2 or 4 neighbours in C . If $x \notin C$ has 2 neighbours in C , reverse its adjacencies with C . The obtained graph is cospectral.



How to find cospectral graphs

Theorem (GM₄ switching, Godsil and McKay, 1982)



Consider a graph with a regular subgraph C of size 4 such that every vertex $x \notin C$ has 0, 2 or 4 neighbours in C . If $x \notin C$ has 2 neighbours in C , reverse its adjacencies with C . The obtained graph is cospectral.



How to find cospectral graphs

Theorem (GM₄ switching, Godsil and McKay, 1982)

Consider a graph with a regular subgraph C of size 4 such that every vertex $x \notin C$ has 0, 2 or 4 neighbours in C . If $x \notin C$ has 2 neighbours in C , reverse its adjacencies with C . The obtained graph is cospectral.



How to find cospectral graphs

Theorem (GM₄ switching, Godsil and McKay, 1982)


Consider a graph with a regular subgraph C of size 4 such that every vertex $x \notin C$ has 0, 2 or 4 neighbours in C . If $x \notin C$ has 2 neighbours in C , reverse its adjacencies with C . The obtained graph is cospectral.

Proof.


$$\begin{pmatrix} A_{11} & A'_{12} \\ A'_{21} & A_{22} \end{pmatrix} = \begin{pmatrix} \frac{1}{2}J - I & O \\ O & I \end{pmatrix}^T \begin{pmatrix} A_{11} & A_{12} \\ A_{21} & A_{22} \end{pmatrix} \begin{pmatrix} \frac{1}{2}J - I & O \\ O & I \end{pmatrix}.$$




How to find cospectral graphs




How to find cospectral graphs




How to find cospectral graphs




How to find cospectral graphs




How to find cospectral graphs




How to find cospectral graphs




How to find cospectral graphs



How to find cospectral graphs




How to find cospectral graphs




$\text{AG}(2, 2)$

How to find cospectral graphs



How to find cospectral graphs




C

$\text{PG}(2, 2)$

Design switching

Theorem (Ihringer and Simoens, 2025+)


Consider a graph with a *certain* subgraph C whose vertices are identified as points of an (r, λ) -design such that every vertex $x \notin C$ is adjacent to the points of a block. Let π be a *certain* permutation of the blocks. If $x \notin C$ is adjacent to the points of B , make it adjacent to the points of $\pi(B)$. The obtained graph is cospectral.



Design switching

Theorem (Ihringer and Simoens, 2025+)


Consider a graph with a *certain* subgraph C whose vertices are identified as points of an (r, λ) -design such that every vertex $x \notin C$ is adjacent to the points of a block. Let π be a *certain* permutation of the blocks. If $x \notin C$ is adjacent to the points of B , make it adjacent to the points of $\pi(B)$. The obtained graph is cospectral.



Design switching

Theorem (Ihringer and Simoens, 2025+)

Consider a graph with a *certain* subgraph C whose vertices are identified as points of an (r, λ) -design such that every vertex $x \notin C$ is adjacent to the points of a block. Let π be a *certain* permutation of the blocks. If $x \notin C$ is adjacent to the points of B , make it adjacent to the points of $\pi(B)$. The obtained graph is cospectral.



Known switching methods

Definition


Switching is a local graph operation, resulting in a *cospectral graph*.

- GM-switching [Godsil and McKay, 1982]
- WQH-switching [Wang, Qiu and Hu, 2019]
- AH-switching [Abiad and Haemers, 2012]
 - Sun graph switching [Mao, Wang, Liu and Qiu, 2023]
 - Fano switching [Abiad, van de Berg and Simoens, 2025+]
 - Cube switching [Abiad, van de Berg and Simoens, 2025+]

Abiad and Haemers (2012):

Conjugation of the adjacency matrix A with $Q = \begin{pmatrix} R & O \\ O & I \end{pmatrix}$, where

$$R = \frac{1}{2} \begin{pmatrix} -1 & 1 & 1 & 0 & 1 & 0 & 0 \\ 0 & -1 & 1 & 1 & 0 & 1 & 0 \\ 0 & 0 & -1 & 1 & 1 & 0 & 1 \\ 1 & 0 & 0 & -1 & 1 & 1 & 0 \\ 0 & 1 & 0 & 0 & -1 & 1 & 1 \\ 1 & 0 & 1 & 0 & 0 & -1 & 1 \\ 1 & 1 & 0 & 1 & 0 & 0 & -1 \end{pmatrix}$$



$\text{PG}(2, 2)$


Fano switching

Theorem (Abiad, van de Berg and Simoens, 2025+)

Consider a graph with a subgraph C whose vertices are identified as points of the Fano plane such that:

- *C is edgeless or complete.*
- *Every vertex $x \notin C$ has 0, 3, 4 or 7 neighbours in C .*
 - *If x has 3 neighbours in C , they form a line.*
 - *If x has 4 neighbours in C , they form the complement of a line.*

Let π be a permutation of the lines. If $x \notin C$ is (non)adjacent to the vertices of ℓ , make it (non)adjacent to the vertices of $\pi(\ell)$. The obtained graph is cospectral.




Fano switching

Theorem (Abiad, van de Berg and Simoens, 2025+)

Consider a graph with a subgraph C whose vertices are identified as points of the Fano plane such that:

- ▶ C is edgeless or complete.
- ▶ Every vertex $x \notin C$ has 0, 3, 4 or 7 neighbours in C .
 - ▶ If x has 3 neighbours in C , they form a line.
 - ▶ If x has 4 neighbours in C , they form the complement of a line.

Let π be a permutation of the lines. If $x \notin C$ is (non)adjacent to the vertices of ℓ , make it (non)adjacent to the vertices of $\pi(\ell)$. The obtained graph is cospectral.




Fano switching

Theorem (Abiad, van de Berg and Simoens, 2025+)

Consider a graph with a subgraph C whose vertices are identified as points of the Fano plane such that:

- ▶ C is edgeless or complete.
- ▶ Every vertex $x \notin C$ has 0, 3, 4 or 7 neighbours in C .
 - ▶ If x has 3 neighbours in C , they form a line.
 - ▶ If x has 4 neighbours in C , they form the complement of a line.

Let π be a permutation of the lines. If $x \notin C$ is (non)adjacent to the vertices of ℓ , make it (non)adjacent to the vertices of $\pi(\ell)$. The obtained graph is cospectral.





Fano switching

Theorem (Abiad, van de Berg and Simoens, 2025+)



Consider a graph with a subgraph C whose vertices are identified as points of the Fano plane such that:

- ▶ C is edgeless or complete.
- ▶ Every vertex $x \notin C$ has 0, 3, 4 or 7 neighbours in C .
 - ▶ If x has 3 neighbours in C , they form a line.
 - ▶ If x has 4 neighbours in C , they form the complement of a line.



Let π be a permutation of the lines. If $x \notin C$ is (non)adjacent to the vertices of ℓ , make it (non)adjacent to the vertices of $\pi(\ell)$. The obtained graph is cospectral.



Fano switching



Fano switching



Both graphs have spectrum $\{(-\sqrt{5})^1, (-\sqrt{2})^2, (0)^3, (\sqrt{2})^2, (\sqrt{5})^1\}$.

Theorem (Ihringer and Simoens, 2025+)

Consider a graph with a *certain* subgraph C whose vertices are identified as points of an (r, λ) -design such that every vertex $x \notin C$ is adjacent to the points of a block. Let π be a *certain* permutation of the blocks. If $x \notin C$ is adjacent to the points of B , make it adjacent to the points of $\pi(B)$. The obtained graph is cospectral.

Theorem (Ihringer and Simoens, 2025+)

Consider a graph with an *edgeless or complete subgraph* C whose vertices are identified as points of an (r, λ) -design such that every vertex $x \notin C$ is adjacent to the points of a block. Let π be a *certain* permutation of the blocks. If $x \notin C$ is adjacent to the points of B , make it adjacent to the points of $\pi(B)$. The obtained graph is cospectral.


Theorem (Ihringer and Simoens, 2025+)

Consider a graph with an *edgeless or complete subgraph* C whose vertices are identified as points of an (r, λ) -*design* such that every vertex $x \notin C$ is adjacent to the points of a block. Let π be a permutation of the blocks such that for all blocks B_i, B_j ,











$$|B_i \cap B_j| = |\pi(B_i) \cap \pi(B_j)|.$$

If $x \notin C$ is adjacent to the points of B , make it adjacent to the points of $\pi(B)$. The obtained graph is cospectral.


Design switching













is an $(r = 3, \lambda = 1)$ -design with incidence matrix

						
	B_1	B_2	B_3	B_4	B_5	B_6
 p_1	1	1	1	0	0	0
 p_2	1	0	0	1	1	0
 p_3	0	1	0	1	0	1
 p_4	0	0	1	0	1	1

Design switching




is an $(r = 3, \lambda = 1)$ -design with incidence matrix











						
	B_1	B_2	B_3	B_4	B_5	B_6
 p_1	1	1	1	0	0	0
 p_2	1	0	0	1	1	0
 p_3	0	1	0	1	0	1
 p_4	0	0	1	0	1	1

$\pi : B_i \mapsto B_{7-i}$ preserves pairwise intersection

Design switching



is an $(r = 3, \lambda = 1)$ -design with incidence matrix


						
	B_1	B_2	B_3	B_4	B_5	B_6
 p_1	1	1	1	0	0	0
 p_2	1	0	0	1	1	0
 p_3	0	1	0	1	0	1
 p_4	0	0	1	0	1	1

$\pi : B_i \mapsto B_{7-i}$ preserves pairwise intersection

Theorem (GM₄ switching, Godsil and McKay, 1982)

Consider a graph with a regular subgraph C of size 4 such that every vertex $x \notin C$ has 0, 2 or 4 neighbours in C . If $x \notin C$ has 2 neighbours in C , reverse its adjacencies with C . The obtained graph is cospectral.

Design switching



is an $(r = 3, \lambda = 1)$ -design with incidence matrix


	B_1	B_2	B_3	B_4	B_5	B_6
p_1	1	1	1	0	0	0
p_2	1	0	0	1	1	0
p_3	0	1	0	1	0	1
p_4	0	0	1	0	1	1

$\pi : B_i \mapsto B_{7-i}$ preserves pairwise intersection

Theorem (GM₄ switching, Godsil and McKay, 1982)

Consider a graph with a **regular subgraph C of size 4** such that every vertex $x \notin C$ has 0, 2 or 4 neighbours in C . If $x \notin C$ has 2 neighbours in C , reverse its adjacencies with C . The obtained graph is cospectral.

Design switching



is an $(r = 3, \lambda = 1)$ -design with incidence matrix


$$\begin{array}{c}
\textcolor{red}{/} \quad \textcolor{red}{/} \quad \textcolor{red}{/} \quad \textcolor{red}{/} \quad \textcolor{red}{/} \quad \textcolor{red}{/} \\
B_1 \ B_2 \ B_3 \ B_4 \ B_5 \ B_6 \\
\bullet p_1 \left(\begin{array}{cccccc} 1 & 1 & 1 & 0 & 0 & 0 \\ 1 & 0 & 0 & 1 & 1 & 0 \\ 0 & 1 & 0 & 1 & 0 & 1 \\ 0 & 0 & 1 & 0 & 1 & 1 \end{array} \right) \\
\bullet p_2 \\
\bullet p_3 \\
\bullet p_4
\end{array}$$

$$\pi : B_i \mapsto B_{7-i} \text{ preserves pairwise intersection}$$

Theorem (GM₄ switching, Godsil and McKay, 1982)

Consider a graph with a regular subgraph C of size 4 such that every vertex $x \notin C$ has 0, 2 or 4 neighbours in C . If $x \notin C$ has 2 neighbours in C , reverse its adjacencies with C . The obtained graph is cospectral.

Design switching



is an $(r = 4, \lambda = 2)$ -design with incidence matrix


$$\begin{array}{c}
\textcolor{red}{\diagup} \quad \textcolor{red}{\diagdown} \quad \textcolor{red}{\diagup} \quad \textcolor{red}{\diagdown} \quad \textcolor{red}{\diagdown} \quad \textcolor{red}{\diagup} \\
B_1 \ B_2 \ B_3 \ B_4 \ B_5 \ B_6 \\
\bullet p_1 \left(\begin{array}{ccccccc|c} 0 & 1 & 1 & 1 & 0 & 0 & 0 & 1 \\ 0 & 1 & 0 & 0 & 1 & 1 & 0 & 1 \\ 0 & 0 & 1 & 0 & 1 & 0 & 1 & 1 \\ 0 & 0 & 0 & 1 & 0 & 1 & 1 & 1 \end{array} \right)
\end{array}$$

$$\pi : B_i \mapsto B_{7-i} \text{ preserves pairwise intersection}$$

Theorem (GM₄ switching, Godsil and McKay, 1982)

Consider a graph with a regular subgraph C of size 4 such that every vertex $x \notin C$ has 0, 2 or 4 neighbours in C . If $x \notin C$ has 2 neighbours in C , reverse its adjacencies with C . The obtained graph is cospectral.

Design switching




is an $(r = 3, \lambda = 1)$ -design with incidence matrix















$$\begin{array}{c} \text{ } \\ \text{ } \\ \text{ } \\ \text{ } \\ \text{ } \\ \text{ } \\ \text{ } \end{array} \begin{array}{ccccccc} \text{ } & \text{ } & \text{ } & \text{ } & \text{ } & \text{ } & \text{ } \\ B_1 & B_2 & B_3 & B_4 & B_5 & B_6 & B_7 \end{array} \begin{pmatrix} 1 & 1 & 1 & 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 1 & 1 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 & 1 & 1 \\ 0 & 1 & 0 & 1 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 & 1 & 0 & 1 \\ 0 & 0 & 1 & 0 & 1 & 1 & 0 \\ 0 & 0 & 1 & 1 & 0 & 0 & 1 \end{pmatrix}$$

Any permutation of the lines π preserves pairwise intersection

Design switching




is an $(r = 3, \lambda = 1)$ -design with incidence matrix

							
	B_1	B_2	B_3	B_4	B_5	B_6	B_7
 p_1	1	1	1	0	0	0	0
 p_2	1	0	0	1	1	0	0
 p_3	1	0	0	0	0	1	1
 p_4	0	1	0	1	0	1	0
 p_5	0	1	0	0	1	0	1
 p_6	0	0	1	0	1	1	0
 p_7	0	0	1	1	0	0	1

Any permutation of the lines π preserves pairwise intersection

➤ Fano switching

Design switching



is an $(r = 8, \lambda = 4)$ -design with incidence matrix

		B_1	B_2	B_3	B_4	B_5	B_6	B_7	$\overline{B_1}$	$\overline{B_2}$	$\overline{B_3}$	$\overline{B_4}$	$\overline{B_5}$	$\overline{B_6}$	$\overline{B_7}$
p_1		0	1	1	1	0	0	0	0	0	0	1	1	1	1
p_2		0	1	0	0	1	1	0	0	1	1	0	0	1	1
p_3		0	1	0	0	0	0	1	1	0	1	1	1	0	0
p_4		0	0	1	0	1	0	1	0	1	0	1	0	1	1
p_5		0	0	1	0	0	1	0	1	0	1	1	0	1	0
p_6		0	0	0	1	0	1	1	0	1	1	0	0	1	1
p_7		0	0	0	1	1	0	0	1	1	0	0	1	1	0

Any permutation of the lines π preserves pairwise intersection

➤ Fano switching

Design switching

Theorem (Ihringer and Simoens, 2025+)

Consider a graph with an *edgeless or complete subgraph* C whose vertices are identified as points of an (r, λ) -*design* such that every vertex $x \notin C$ is adjacent to the points of a block. Let π be a permutation of the blocks such that for all blocks B_i, B_j ,

$$|B_i \cap B_j| = |\pi(B_i) \cap \pi(B_j)|.$$

If $x \notin C$ is adjacent to the points of B , make it adjacent to the points of $\pi(B)$. The obtained graph is cospectral.

Design switching

Theorem (Ihringer and Simoens, 2025+)

Consider a graph with an **edgeless or complete subgraph C** whose vertices are identified as points of an **(r, λ) -design** such that every vertex **$x \notin C$ is adjacent to the points of a block**. Let π be a permutation of the blocks such that for all blocks B_i, B_j ,

$$|B_i \cap B_j| = |\pi(B_i) \cap \pi(B_j)|.$$

If $x \notin C$ is adjacent to the points of B , make it adjacent to the points of $\pi(B)$. The obtained graph is cospectral.

Proof. Define $R = \frac{1}{r-\lambda} (N(N^\pi)^T - \lambda J)$, where N^π is obtained from the incidence matrix N by permuting the columns with π .

$$\begin{pmatrix} A_{11} & A'_{12} \\ A'_{21} & A_{22} \end{pmatrix} = \begin{pmatrix} R & O \\ O & I \end{pmatrix}^T \begin{pmatrix} A_{11} & A_{12} \\ A_{21} & A_{22} \end{pmatrix} \begin{pmatrix} R & O \\ O & I \end{pmatrix}.$$



Design switching

Theorem (Ihringer and Simoens, 2025+)

Consider a graph with a subgraph C with adjacency matrix $A_{11} = R^T A_{11} R$ whose vertices are identified as points of an (r, λ) -design such that every vertex $x \notin C$ is adjacent to the points of a block. Let π be a permutation of the blocks such that for all blocks B_i, B_j ,

$$|B_i \cap B_j| = |\pi(B_i) \cap \pi(B_j)|.$$

If $x \notin C$ is adjacent to the points of B , make it adjacent to the points of $\pi(B)$. The obtained graph is cospectral.

Proof. Define $R = \frac{1}{r-\lambda} (N(N^\pi)^T - \lambda J)$, where N^π is obtained from the incidence matrix N by permuting the columns with π .

$$\begin{pmatrix} A_{11} & A'_{12} \\ A'_{21} & A_{22} \end{pmatrix} = \begin{pmatrix} R & O \\ O & I \end{pmatrix}^T \begin{pmatrix} A_{11} & A_{12} \\ A_{21} & A_{22} \end{pmatrix} \begin{pmatrix} R & O \\ O & I \end{pmatrix}.$$

□

Design switching

Theorem (Ihringer and Simoens, 2025+)

Consider a graph with a subgraph C with adjacency matrix $A_{11} = R^T A_{11} R$ whose vertices are identified as points of an (r, λ) -design such that every vertex $x \notin C$ is adjacent to the points of a block or its complement. Let π be a permutation of the blocks such that for all blocks B_i, B_j ,

$$|B_i \cap B_j| = |\pi(B_i) \cap \pi(B_j)|.$$

If $x \notin C$ is adjacent to the points of B , make it adjacent to the points of $\pi(B)$. The obtained graph is cospectral.

Proof. Define $R = \frac{1}{r-\lambda} (N(N^\pi)^T - \lambda J)$, where N^π is obtained from the incidence matrix N by permuting the columns with π .

$$\begin{pmatrix} A_{11} & A'_{12} \\ A'_{21} & A_{22} \end{pmatrix} = \begin{pmatrix} R & O \\ O & I \end{pmatrix}^T \begin{pmatrix} A_{11} & A_{12} \\ A_{21} & A_{22} \end{pmatrix} \begin{pmatrix} R & O \\ O & I \end{pmatrix}.$$



SMALL $2-(v, k, \lambda)$ -DESIGNS

Small $2-(v, k, \lambda)$ -designs

v	# methods	Method
4	1	GM_4 switching
5	0	
6	1	GM_6 switching
7	1	Fano switching
8	10	$AG(3, 2)$ -switching
9	≥ 2	$AG(2, 3)$ -switching
10	≥ 4	
11	≥ 77	Paley biplane switching
12	≥ 6	
13	≥ 187	$PG(3, 2)$ -switching

Table: Switching methods from small $2-(v, k, \lambda)$ -designs.

AN APPLICATION

An application

Definition

The **triangular graph** T_n has as vertices the 2-subsets of $\{1, \dots, n\}$, where two vertices are adjacent if they intersect.


$$T_n \cong L(K_n) \cong J(n, 2)$$

An application

Definition

The **triangular graph** T_n has as vertices the 2-subsets of $\{1, \dots, n\}$, where two vertices are adjacent if they intersect.

$$T_n \cong L(K_n) \cong J(n, 2)$$




The octahedral graph T_4

An application


Definition

The **triangular graph** T_n has as vertices the 2-subsets of $\{1, \dots, n\}$, where two vertices are adjacent if they intersect.

$$T_n \cong L(K_n) \cong J(n, 2)$$



The octahedral graph T_4



The Petersen graph $\overline{T_5}$

An application

Theorem (Chang and Hoffman, independently, 1959)

The triangular graph T_n is determined by its spectrum iff $n \neq 8$.

An application

Definition

The **q-triangular graph** $T_{q,n}$ has as vertices the **2-dimensional subspaces of \mathbb{F}_q^n** where two vertices are adjacent if they intersect.

An application

Definition

The **q-triangular graph** $T_{q,n}$ has as vertices the **lines of $\text{PG}(n-1, q)$** where two vertices are adjacent if they intersect.

An application

Definition

The **q -triangular graph** $T_{q,n}$ has as vertices the **lines of $\text{PG}(n-1, q)$** where two vertices are adjacent if they intersect.

Theorem (Ihringer and Munemasa, 2019)

The q -triangular graph $T_{q,n}$ is not determined by its spectrum if $n \geq 4$.

An application

Definition

The **q -triangular graph** $T_{q,n}$ has as vertices the **lines of $\text{PG}(n-1, q)$** where two vertices are adjacent if they intersect.

Theorem (Ihringer and Munemasa, 2019)

The q -triangular graph $T_{q,n}$ is not determined by its spectrum if $n \geq 4$.

Proof. Fix a subplane $\text{PG}(2, q) \subseteq \text{PG}(n-1, q)$ and let

$$\mathcal{P} = \{\text{lines of } \text{PG}(2, q)\}$$

$$\mathcal{B} = \{\text{point pencils of } \text{PG}(2, q)\}$$

Design switching on $(\mathcal{P}, \mathcal{B})$, using any permutation π of \mathcal{B} that is not an automorphism \Rightarrow maximal cliques of size $q^2 + q$. □

An application

Theorem (Ihringer and Simoens, 2025+)

There are at least $q!$ graphs with the same spectrum as $T_{q,n}$.

An application

Theorem (Ihringer and Simoens, 2025+)

There are at least $q!$ graphs with the same spectrum as $T_{q,n}$.

Proof. Let Γ_π be the graph obtained from design switching $T_{n,q}$ with π .

$$\Gamma_{\pi_1} \cong \Gamma_{\pi_2} \iff \pi_1, \pi_2 \in \text{same double coset of } \text{Aut}(D) \text{ in } \text{Sym}(\mathcal{B})$$

There are $\geq q!$ double cosets.



An application

Theorem (Ihringer and Simoens, 2025+)

There are at least $q!$ graphs with the same spectrum as $T_{q,n}$.

Proof. Let Γ_π be the graph obtained from design switching $T_{n,q}$ with π .

$$\Gamma_{\pi_1} \cong \Gamma_{\pi_2} \iff \pi_1, \pi_2 \in \text{same double coset of } \text{Aut}(D) \text{ in } \text{Sym}(\mathcal{B})$$

There are $\geq q!$ double cosets. □

➤ Many strongly regular graphs with the same parameters.

Corollary (Fon-Der-Flaass, 2002)

*Almost all strongly regular graphs are **not** determined by their spectrum.*

Concluding remarks

- Many new switching methods

Concluding remarks

- Many new switching methods
- Alternative proofs of cospectrality results:
 - q -triangular graphs [Ihringer, Munemasa, 2019]
 - Collinearity graphs of polar spaces [Brouwer, Ihringer, Kantor, 2022]
 - Collinearity graphs of generalised quadrangles [Guo, van Dam, 2022]

Concluding remarks

- Many new switching methods
- Alternative proofs of cospectrality results:
 - q -triangular graphs [Ihringer, Munemasa, 2019]
 - Collinearity graphs of polar spaces [Brouwer, Ihringer, Kantor, 2022]
 - Collinearity graphs of generalised quadrangles [Guo, van Dam, 2022]
- More general: two different designs

Thank you for listening!



F. Ihringer and R. Simoens,
Design switching on graphs, arXiv:2508.11523.