

Exploring Quasi-Hermitian Varieties: Properties and Applications

Angela Aguglia

Dipartimento di Meccanica, Matematica e Management
Politecnico di Bari, Italy

*based on joint works with
Cossidente, Giuzzi, Korchmáros, Longobardi, Montinaro, Siconolfi*

7th Irsee Conference

5th September 2025

Contents

- 1 Preliminaries
- 2 Constructions of Q -H varieties
- 3 Automorphism groups and equivalences
- 4 Applications
- 5 Open Problems

Quasi-Polar Spaces

Definition (Schillewaert, Van de Voorde (2022))

A **quasi-polar space** is a set of points \mathcal{S} in $\text{PG}(r, q)$, where $r \geq 2$ and q is a prime power, such that the intersection sizes with hyperplanes match those of a non-degenerate classical polar space \mathcal{P} embedded in $\text{PG}(r, q)$.

Quasi-Polar Spaces

Definition (Schillewaert, Van de Voorde (2022))

A **quasi-polar space** is a set of points \mathcal{S} in $\text{PG}(r, q)$, where $r \geq 2$ and q is a prime power, such that the intersection sizes with hyperplanes match those of a non-degenerate classical polar space \mathcal{P} embedded in $\text{PG}(r, q)$.

Quasi-quadrics were introduced in 2000 by
De Clerck, Hamilton, O'Keefe, and Penttila.

Quasi-Polar Spaces

Definition (Schillewaert, Van de Voorde (2022))

A **quasi-polar space** is a set of points \mathcal{S} in $\text{PG}(r, q)$, where $r \geq 2$ and q is a prime power, such that the intersection sizes with hyperplanes match those of a non-degenerate classical polar space \mathcal{P} embedded in $\text{PG}(r, q)$.

Quasi-quadrics were introduced in 2000 by
De Clerck, Hamilton, O'Keefe, and Penttila.

This idea traces back to **Segre**, who in 1954 defined an oval in a finite projective plane as a combinatorial abstraction of a conic in $\text{PG}(2, q)$.

Reference:

- B. Segre, Sulle ovali nei piani lineari finiti, *Rendiconti dell'Accademia Nazionale dei Lincei*, (1954).

Quasi-Polar Spaces

Definition (Schillewaert, Van de Voorde (2022))

A **quasi-polar space** is a set of points \mathcal{S} in $\text{PG}(r, q)$, where $r \geq 2$ and q is a prime power, such that the intersection sizes with hyperplanes match those of a non-degenerate classical polar space \mathcal{P} embedded in $\text{PG}(r, q)$.

Quasi-quadrics were introduced in 2000 by
De Clerck, Hamilton, O'Keefe, and Penttila.

This idea traces back to **Segre**, who in 1954 defined an oval in a finite projective plane as a combinatorial abstraction of a conic in $\text{PG}(2, q)$.

The analogous concept for Hermitian varieties, that is **quasi-Hermitian varieties**, was formally introduced in 2010 by **De Winter and Schillewaert.**

The Hermitian Case

A **non-singular Hermitian variety** $\mathcal{H}(r, q^2)$ in the projective space $\text{PG}(r, q^2)$ is defined as the set of absolute points of a non-degenerate unitary polarity ρ .

The Hermitian Case

A **non-singular Hermitian variety** $\mathcal{H}(r, q^2)$ in the projective space $\text{PG}(r, q^2)$ is defined as the set of absolute points of a non-degenerate unitary polarity ρ .

That is,

$$\mathcal{H}(r, q^2) = \{P \in \text{PG}(r, q^2) \mid P \in P^\rho\}$$

where P^ρ is the polar hyperplane of P under the unitary polarity ρ .

The Hermitian Case

A **non-singular Hermitian variety** $\mathcal{H}(r, q^2)$ in the projective space $\text{PG}(r, q^2)$ is defined as the set of absolute points of a non-degenerate unitary polarity ρ .

That is,

$$\mathcal{H}(r, q^2) = \{P \in \text{PG}(r, q^2) \mid P \in P^\rho\}$$

where P^ρ is the polar hyperplane of P under the unitary polarity ρ .

Property

A non-singular Hermitian variety $\mathcal{H}(r, q^2)$ is a hypersurface with equation:

$$(X_0^q, \dots, X_r^q)H(X_0, \dots, X_r)^T = 0,$$

where H is a non-singular Hermitian $(r+1) \times (r+1)$ matrix.

Projective Equivalence

Any non-singular Hermitian variety in $\text{PG}(r, q^2)$ can be mapped to any other non-singular Hermitian variety in $\text{PG}(r, q^2)$ via a projectivity.

Projective Equivalence

Any non-singular Hermitian variety in $\text{PG}(r, q^2)$ can be mapped to any other non-singular Hermitian variety in $\text{PG}(r, q^2)$ via a projectivity.

Special Case: The Plane

In the plane, the non-singular Hermitian curve $\mathcal{H}(2, q^2)$ is also known as the **classical or Hermitian unital**.

Projective Equivalence

Any non-singular Hermitian variety in $\text{PG}(r, q^2)$ can be mapped to any other non-singular Hermitian variety in $\text{PG}(r, q^2)$ via a projectivity.

Special Case: The Plane

In the plane, the non-singular Hermitian curve $\mathcal{H}(2, q^2)$ is also known as the **classical or Hermitian unital**.

A **unital embedded** in $\text{PG}(2, q^2)$ is a set of $q^3 + 1$ ($= |\mathcal{H}(2, q^2)|$) points such that every line of the plane intersects it in either 1 or $q + 1$ points.

Definition (De Winter, Schillewaert (2010))

A point set \mathcal{S} of $\text{PG}(r, q^2)$ is a **quasi-Hermitian variety** if it meets each hyperplane in either

$$|\mathcal{H}(r-1, q^2)| = \frac{(q^r + (-1)^{r-1})(q^{r-1} - (-1)^{r-1})}{q^2 - 1}, \text{ or}$$

$$|P_0\mathcal{H}(r-2, q^2)| = \frac{(q^r + (-1)^{r-1})(q^{r-1} - (-1)^{r-1})}{q^2 - 1} + (-1)^{r-1}q^{r-1}$$

points.

Definition (De Winter, Schillewaert (2010))

A point set \mathcal{S} of $\text{PG}(r, q^2)$ is a **quasi-Hermitian variety** if it meets each hyperplane in either

$$|\mathcal{H}(r-1, q^2)| = \frac{(q^r + (-1)^{r-1})(q^{r-1} - (-1)^{r-1})}{q^2 - 1}, \text{ or}$$

$$|P_0\mathcal{H}(r-2, q^2)| = \frac{(q^r + (-1)^{r-1})(q^{r-1} - (-1)^{r-1})}{q^2 - 1} + (-1)^{r-1}q^{r-1}$$

points.

$\mathcal{H}(r, q^2)$ is a quasi-Hermitian variety, called the *classical quasi-Hermitian variety*.

Quasi-Hermitian Varieties as Two-Character Sets

Property

A *quasi-Hermitian variety* in the projective space $\text{PG}(r, q^2)$ is a **two-character set**, meaning a point set with exactly two possible intersection sizes with hyperplanes.

Quasi-Hermitian Varieties as Two-Character Sets

Property

A *quasi-Hermitian variety* in the projective space $\text{PG}(r, q^2)$ is a **two-character set**, meaning a point set with exactly two possible intersection sizes with hyperplanes.

Why It Matters

Two-character sets have wide-ranging applications:

- They give rise to **strongly regular graphs**.
- They generate **two-weight linear codes**.

Quasi-Hermitian Varieties as Two-Character Sets

Property

A *quasi-Hermitian variety* in the projective space $\text{PG}(r, q^2)$ is a **two-character set**, meaning a point set with exactly two possible intersection sizes with hyperplanes.

Why It Matters

Two-character sets have wide-ranging applications:

- They give rise to **strongly regular graphs**.
- They generate **two-weight linear codes**.

Key references:

- R. Delsarte, Weights of linear codes and strongly regular normed spaces, *Discrete Math.*, **3** (1972)
- R. Calderbank, W. Kantor, The geometry of two-weight codes, *The Bulletin of the London Mathematical Society*, **18** (1986)

Cardinality of a quasi-Hermitian variety

Theorem 1 (Schillewaert, Van de Voorde (2022))

Let \mathcal{S} be a quasi-Hermitian variety in $\text{PG}(r, q^2)$ with $r \geq 3$, then:

$$|\mathcal{S}| = |\mathcal{H}(r, q^2)|.$$

If $\mathcal{S} \subseteq \text{PG}(2, q^2)$ is a point set such that every line intersects \mathcal{S} in either 1 or $q + 1$ points, then:

$$|\mathcal{S}| \in \{q^2 + q + 1, q^3 + 1\}.$$

Cardinality of a quasi-Hermitian variety

Theorem 1 (Schillewaert, Van de Voorde (2022))

Let \mathcal{S} be a quasi-Hermitian variety in $\text{PG}(r, q^2)$ with $r \geq 3$, then:

$$|\mathcal{S}| = |\mathcal{H}(r, q^2)|.$$

If $\mathcal{S} \subseteq \text{PG}(2, q^2)$ is a point set such that every line intersects \mathcal{S} in either 1 or $q + 1$ points, then:

$$|\mathcal{S}| \in \{q^2 + q + 1, q^3 + 1\}.$$

Perspective

We interpret a quasi-Hermitian variety \mathcal{S} as higher-dimensional generalization of a unital. Thus, $|\mathcal{S}| = |\mathcal{H}(r, q^2)|$ for any $r \geq 2$.

Contents

- 1 Preliminaries
- 2 Constructions of Q -H varieties
- 3 Automorphism groups and equivalences
- 4 Applications
- 5 Open Problems

Known Constructions of Q-H varieties

All Dimensions $r > 2$

- De Winter, Schillewaert (2010),

Known Constructions of Q-H varieties

All Dimensions $r > 2$

- De Winter, Schillewaert (2010),

All Dimensions

- Aguglia, Cossidente, Korchmáros (2012),
- Aguglia (2013),

Known Constructions of Q-H varieties

All Dimensions $r > 2$

- De Winter, Schillewaert (2010),

All Dimensions

- Aguglia, Cossidente, Korchmáros (2012),
- Aguglia (2013),

Odd Dimensions

- Pavese (2015),

Known Constructions of Q-H varieties

All Dimensions $r > 2$

- De Winter, Schillewaert (2010),

All Dimensions

- Aguglia, Cossidente, Korchmáros (2012),
- Aguglia (2013),

Odd Dimensions

- Pavese (2015),

Three-Dimensional Constructions

- Lia, Lavrauw, Pavese (2024),
- Lia, Sheekey (2024).

BM quasi-Hermitian Variety Construction

Fix a projective frame in $\text{PG}(r, q^2)$ with homogeneous coordinates (X_0, X_1, \dots, X_r) .

BM quasi-Hermitian Variety Construction

Fix a projective frame in $\text{PG}(r, q^2)$ with homogeneous coordinates (X_0, X_1, \dots, X_r) .

Consider the affine space $\text{AG}(r, q^2)$ whose infinite hyperplane Σ_∞ has equation $X_0 = 0$.

BM quasi-Hermitian Variety Construction

Fix a projective frame in $\text{PG}(r, q^2)$ with homogeneous coordinates (X_0, X_1, \dots, X_r) .

Consider the affine space $\text{AG}(r, q^2)$ whose infinite hyperplane Σ_∞ has equation $X_0 = 0$.

Then $\text{AG}(r, q^2)$ has affine coordinates (x_1, x_2, \dots, x_r) where $x_i = X_i/X_0$ for $i \in \{1, \dots, r\}$.

Take $a \in \text{GF}(q^2)$ and $b \in \text{GF}(q^2) \setminus \text{GF}(q)$ and consider the projective variety $\mathcal{B}_{a,b}$ of equation

$$\begin{aligned} X_r^q X_0^q - X_r X_0^{2q-1} + a^q (X_1^{2q} + \dots + X_{r-1}^{2q}) - a (X_1^2 + \dots + X_n^2) X_0^{2q-2} \\ = (b^q - b) (X_1^{q+1} + \dots + X_{r-1}^{q+1}) X_0^{q-1}. \end{aligned} \quad (1)$$

Take $a \in \text{GF}(q^2)$ and $b \in \text{GF}(q^2) \setminus \text{GF}(q)$ and consider the projective variety $\mathcal{B}_{a,b}$ of equation

$$\begin{aligned} X_r^q X_0^q - X_r X_0^{2q-1} + a^q (X_1^{2q} + \dots + X_{r-1}^{2q}) - a (X_1^2 + \dots + X_n^2) X_0^{2q-2} \\ = (b^q - b) (X_1^{q+1} + \dots + X_{r-1}^{q+1}) X_0^{q-1}. \end{aligned} \quad (1)$$

Let $\mathcal{F} \subset \Sigma_\infty$ be the Hermitian cone

$$\mathcal{F} := \{(0, x_1, \dots, x_r) \mid x_1^{q+1} + \dots + x_{r-1}^{q+1} = 0\}.$$

Take $a \in \text{GF}(q^2)$ and $b \in \text{GF}(q^2) \setminus \text{GF}(q)$ and consider the projective variety $\mathcal{B}_{a,b}$ of equation

$$\begin{aligned} X_r^q X_0^q - X_r X_0^{2q-1} + a^q (X_1^{2q} + \dots + X_{r-1}^{2q}) - a (X_1^2 + \dots + X_n^2) X_0^{2q-2} \\ = (b^q - b) (X_1^{q+1} + \dots + X_{r-1}^{q+1}) X_0^{q-1}. \end{aligned} \quad (1)$$

Let $\mathcal{F} \subset \Sigma_\infty$ be the Hermitian cone

$$\mathcal{F} := \{(0, x_1, \dots, x_r) \mid x_1^{q+1} + \dots + x_{r-1}^{q+1} = 0\}.$$

If $a = 0$ then $\mathcal{B}_{a,b}$ is a non-singular Hermitian variety.

Assumptions

Assume $a \neq 0$ and one of the following conditions:

Assumptions

Assume $a \neq 0$ and one of the following conditions:

- 1 r and q are both odd and $4a^{q+1} + (b^q - b)^2 \neq 0$;

Assumptions

Assume $a \neq 0$ and one of the following conditions:

- i r and q are both odd and $4a^{q+1} + (b^q - b)^2 \neq 0$;
- ii r is even, q is odd and $4a^{q+1} + (b^q - b)^2$ is a non-square in $\text{GF}(q)$;

Assumptions

Assume $a \neq 0$ and one of the following conditions:

- i r and q are both odd and $4a^{q+1} + (b^q - b)^2 \neq 0$;
- ii r is even, q is odd and $4a^{q+1} + (b^q - b)^2$ is a non-square in $\text{GF}(q)$;
- iii r and q are both even and $\text{tr}_{q/2}(\frac{a^{q+1}}{(b^q + b)^2}) = 0$;

Assumptions

Assume $a \neq 0$ and one of the following conditions:

- i r and q are both odd and $4a^{q+1} + (b^q - b)^2 \neq 0$;
- ii r is even, q is odd and $4a^{q+1} + (b^q - b)^2$ is a non-square in $\text{GF}(q)$;
- iii r and q are both even and $\text{tr}_{q/2}(\frac{a^{q+1}}{(b^q + b)^2}) = 0$;
- iv r is odd and q is even.

Assumptions

Assume $a \neq 0$ and one of the following conditions:


- i r and q are both odd and $4a^{q+1} + (b^q - b)^2 \neq 0$;
- ii r is even, q is odd and $4a^{q+1} + (b^q - b)^2$ is a non-square in $\text{GF}(q)$;
- iii r and q are both even and $\text{tr}_{q/2}(\frac{a^{q+1}}{(b^q + b)^2}) = 0$;
- iv r is odd and q is even.

If $r = 2$ then $\mathcal{B}_{a,b}$ is a non-classical Buekenhout-Metz unital.

Case $r = 3$ and q odd

$\mathcal{B}_{a,b} \cap \Sigma_\infty = \ell_1 \cup \ell_2$ where $\ell_1 : X_1 - \nu X_2 = 0 = X_0$, $\ell_2 : X_1 + \nu X_2 = 0 = X_0$ and $\nu \in \text{GF}(q^2) : \nu^2 = -1$.


$PG(3, q^2)$



Case $r = 3$ and q odd

$\mathcal{B}_{a,b} \cap \Sigma_\infty = \ell_1 \cup \ell_2$ where $\ell_1 : X_1 - \nu X_2 = 0 = X_0$, $\ell_2 : X_1 + \nu X_2 = 0 = X_0$ and $\nu \in \text{GF}(q^2) : \nu^2 = -1$.


$PG(3, q^2)$



Case $r = 3$ and q odd

$\mathcal{B}_{a,b} \cap \Sigma_\infty = \ell_1 \cup \ell_2$ where $\ell_1 : X_1 - \nu X_2 = 0 = X_0$, $\ell_2 : X_1 + \nu X_2 = 0 = X_0$ and $\nu \in \text{GF}(q^2) : \nu^2 = -1$.

$PG(3, q^2)$



For any $r \geq 2$, we define $\mathcal{M}_{a,b} := (\mathcal{B}_{a,b} \setminus \Sigma_\infty) \cup \mathcal{F}$

Assumptions

Assume $a \neq 0$ and one of the following conditions:

- i r and q are both odd and $4a^{q+1} + (b^q - b)^2 \neq 0$;
- ii r is even, q is odd and $4a^{q+1} + (b^q - b)^2$ is a non-square in $\text{GF}(q)$;
- iii r and q are both even and $\text{tr}_{q/2}(\frac{a^{q+1}}{(b^q + b)^2}) = 0$;
- iv r is odd and q is even.

Theorem 2 (A., Cossidente, Korchmáros (2012))

The set $\mathcal{M}_{a,b}$ consisting of the affine points of $\mathcal{B}_{a,b}$ plus the infinite points of \mathcal{F} is a non-classical quasi-Hermitian variety of $\text{PG}(r, q^2)$, $r \geq 2$.

Assumptions

Assume $a \neq 0$ and one of the following conditions:

- i r and q are both odd and $4a^{q+1} + (b^q - b)^2 \neq 0$;
- ii r is even, q is odd and $4a^{q+1} + (b^q - b)^2$ is a non-square in $\text{GF}(q)$;
- iii r and q are both even and $\text{tr}_{q/2}(\frac{a^{q+1}}{(b^q + b)^2}) = 0$;
- iv r is odd and q is even.

Theorem 2 (A., Cossidente, Korchmáros (2012))

The set $\mathcal{M}_{a,b}$ consisting of the affine points of $\mathcal{B}_{a,b}$ plus the infinite points of \mathcal{F} is a non-classical quasi-Hermitian variety of $\text{PG}(r, q^2)$, $r \geq 2$.

This quasi-Hermitian variety is called a *BM quasi-Hermitian variety*.

Crucial Tool: A Non-Standard Model of $\text{PG}(r, q^2)$

Fix $a \in \text{GF}(q^2)^*$.

Crucial Tool: A Non-Standard Model of $\text{PG}(r, q^2)$

Fix $a \in \text{GF}(q^2)^*$.

Define the quadric $\mathcal{Q}_a(m, d)$ for $m = (m_1, \dots, m_{r-1}) \in \text{GF}(q^2)^{r-1}$ and $d \in \text{GF}(q^2)$ as the set of points satisfying:

$$x_r = a(x_1^2 + \dots + x_{r-1}^2) + m_1x_1 + \dots + m_{r-1}x_{r-1} + d.$$

Crucial Tool: A Non-Standard Model of $\text{PG}(r, q^2)$

Fix $a \in \text{GF}(q^2)^*$.

Define the quadric $\mathcal{Q}_a(m, d)$ for $m = (m_1, \dots, m_{r-1}) \in \text{GF}(q^2)^{r-1}$ and $d \in \text{GF}(q^2)$ as the set of points satisfying:

$$x_r = a(x_1^2 + \dots + x_{r-1}^2) + m_1x_1 + \dots + m_{r-1}x_{r-1} + d.$$

Consider the pair (\mathcal{P}, Σ) , where:

Crucial Tool: A Non-Standard Model of $\text{PG}(r, q^2)$

Fix $a \in \text{GF}(q^2)^*$.

Define the quadric $\mathcal{Q}_a(m, d)$ for $m = (m_1, \dots, m_{r-1}) \in \text{GF}(q^2)^{r-1}$ and $d \in \text{GF}(q^2)$ as the set of points satisfying:

$$x_r = a(x_1^2 + \dots + x_{r-1}^2) + m_1x_1 + \dots + m_{r-1}x_{r-1} + d.$$

Consider the pair (\mathcal{P}, Σ) , where:

- **Points** $P \in \mathcal{P}$: all points of the affine geometry $\text{AG}(r, q^2)$;

Crucial Tool: A Non-Standard Model of $\text{PG}(r, q^2)$

Fix $a \in \text{GF}(q^2)^*$.

Define the quadric $\mathcal{Q}_a(m, d)$ for $m = (m_1, \dots, m_{r-1}) \in \text{GF}(q^2)^{r-1}$ and $d \in \text{GF}(q^2)$ as the set of points satisfying:

$$x_r = a(x_1^2 + \dots + x_{r-1}^2) + m_1x_1 + \dots + m_{r-1}x_{r-1} + d.$$

Consider the pair (\mathcal{P}, Σ) , where:


- **Points** $P \in \mathcal{P}$: all points of the affine geometry $\text{AG}(r, q^2)$;
- **Hyperplanes** $\pi \in \Sigma$:
 - all affine hyperplanes passing through the point at infinity $P_\infty = (0, 0, \dots, 0, 1)$;
 - all quadrics of the form $\mathcal{Q}_a(m, d)$.

Lemma 3 (A., Cossidente, Korchmáros, (2012))

For every non-zero $a \in \text{GF}(q^2)$, (\mathcal{P}, Σ) defines an incidence structure Π_a isomorphic to $\text{AG}(r, q^2)$.

Lemma 3 (A., Cossidente, Korchmáros, (2012))


For every non-zero $a \in \text{GF}(q^2)$, (\mathcal{P}, Σ) defines an incidence structure Π_a isomorphic to $\text{AG}(r, q^2)$.



Completing Π_a with its points at infinity in the usual way gives a projective space $\overline{\Pi}_a$ isomorphic to $\text{PG}(r, q^2)$.

Completing Π_a with its points at infinity in the usual way gives a projective space $\overline{\Pi}_a$ isomorphic to $\text{PG}(r, q^2)$.

Let $\mathcal{H}(r, q^2) : x_r^q - x_r = (b^q - b)(x_1^{q+1} + \dots + x_{r-1}^{q+1})$



BT Q-H Variety Construction

BT Q-H Variety Construction

Let e be an odd number, $r \geq 2$ and consider the projective space $\text{PG}(r, q^2)$ where $q = 2^e$.

BT Q-H Variety Construction

Let e be an odd number, $r \geq 2$ and consider the projective space $\text{PG}(r, q^2)$ where $q = 2^e$.

Choose now $\varepsilon \in \text{GF}(q^2) \setminus \text{GF}(q)$ such that $\text{tr}_{q^2/q}(\varepsilon) = 1$.

BT Q-H Variety Construction

Let e be an odd number, $r \geq 2$ and consider the projective space $\text{PG}(r, q^2)$ where $q = 2^e$.

Choose now $\varepsilon \in \text{GF}(q^2) \setminus \text{GF}(q)$ such that $\text{tr}_{q^2/q}(\varepsilon) = 1$.

The function $\sigma : x \rightarrow (x)^{2^{\frac{e+1}{2}}}$ is an automorphism of $\text{GF}(q)$.

BT Q-H Variety Construction

Let e be an odd number, $r \geq 2$ and consider the projective space $\text{PG}(r, q^2)$ where $q = 2^e$.

Choose now $\varepsilon \in \text{GF}(q^2) \setminus \text{GF}(q)$ such that $\text{tr}_{q^2/q}(\varepsilon) = 1$.

The function $\sigma : x \rightarrow (x)^{2^{\frac{e+1}{2}}}$ is an automorphism of $\text{GF}(q)$.

Now, let $\mathcal{V}_\varepsilon^r$ be the variety of $\text{PG}(r, q^2)$ represented by:

$$x_r^q + x_r = \Gamma_\varepsilon(x_1) + \dots + \Gamma_\varepsilon(x_{r-1}) \quad (2)$$

where

$$\Gamma_\varepsilon(x) = [x + (x^q + x)\varepsilon]^{\sigma+2} + (x^q + x)^\sigma + (x^{2q} + x^2)\varepsilon + x^{q+1} + x^2.$$

BT Q-H Variety Construction

Let e be an odd number, $r \geq 2$ and consider the projective space $\text{PG}(r, q^2)$ where $q = 2^e$.

Choose now $\varepsilon \in \text{GF}(q^2) \setminus \text{GF}(q)$ such that $\text{tr}_{q^2/q}(\varepsilon) = 1$.

The function $\sigma : x \rightarrow (x)^{2^{\frac{e+1}{2}}}$ is an automorphism of $\text{GF}(q)$.

Now, let $\mathcal{V}_\varepsilon^r$ be the variety of $\text{PG}(r, q^2)$ represented by:

$$x_r^q + x_r = \Gamma_\varepsilon(x_1) + \dots + \Gamma_\varepsilon(x_{r-1}) \quad (2)$$

where

$$\Gamma_\varepsilon(x) = [x + (x^q + x)\varepsilon]^{\sigma+2} + (x^q + x)^\sigma + (x^{2q} + x^2)\varepsilon + x^{q+1} + x^2.$$

We define

$$\mathcal{H}_\varepsilon^r := (\mathcal{V}_\varepsilon^r \setminus \Sigma_\infty) \cup \mathcal{F},$$

where

$$\mathcal{F} = \{(0, X_1, \dots, X_r) \mid X_1^{q+1} + \dots + X_{r-1}^{q+1} = 0\}.$$

Theorem [A. (2013)]

$\mathcal{H}_\varepsilon^r$ is a non-classical quasi-Hermitian variety in $\text{PG}(r, q^2)$ which, for $r = 2$, corresponds to a Buekenhout–Tits unital.

Theorem [A. (2013)]

\mathcal{H}_ϵ^r is a non-classical quasi-Hermitian variety in $\text{PG}(r, q^2)$ which, for $r = 2$, corresponds to a Buekenhout–Tits unital.

We refer to \mathcal{H}_ϵ^r as a *BT quasi-Hermitian variety* in $\text{PG}(r, q^2)$ for all $r \geq 3$.

Theorem [A. (2013)]

$\mathcal{H}_\varepsilon^r$ is a non-classical quasi-Hermitian variety in $\text{PG}(r, q^2)$ which, for $r = 2$, corresponds to a Buekenhout–Tits unital.

We refer to $\mathcal{H}_\varepsilon^r$ as a *BT quasi-Hermitian variety* in $\text{PG}(r, q^2)$ for all $r \geq 3$.

Automorphism 2-Group Result [A.(2013)]

The elementary abelian 2-group E of order q^r , generated by collineations with matrices of the form:

$$\begin{pmatrix} 1 & \gamma_1\varepsilon & \gamma_2\varepsilon & \cdots & \gamma_{r-1}\varepsilon & \gamma_r + (\sum_{i=1}^{r-1} \gamma_i)\sigma\varepsilon \\ 0 & 1 & 0 & \cdots & 0 & \gamma_1 + \gamma_1\varepsilon \\ 0 & 0 & 1 & \cdots & 0 & \gamma_2 + \gamma_2\varepsilon \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & 0 & \cdots & 1 & \gamma_{r-1} + \gamma_{r-1}\varepsilon \\ 0 & 0 & 0 & \cdots & 0 & 1 \end{pmatrix}$$

with $\gamma_i \in \mathbb{F}_q$ for $i = 1, \dots, r$, is a subgroup of $\text{Aut}(\mathcal{H}_\varepsilon^r)$.

A Classical Problem

Having only a few intersection numbers with hyperplanes is a strong combinatorial property however, this condition alone is not sufficient to characterize Hermitian varieties.

A Classical Problem

Having only a few intersection numbers with hyperplanes is a strong combinatorial property however, this condition alone is not sufficient to characterize Hermitian varieties.

Problem

Can we find a characterization of Hermitian varieties among quasi-Hermitian ones?

Some Known Characterizations

Theorem 4 (De Winter, Schillewaert (2010))

A quasi-Hermitian variety of $\text{PG}(r, q^2)$, $r \geq 3$ is classical if it has the same intersection numbers with respect to spaces of codimension 2 as a non-singular Hermitian variety.

Some Known Characterizations

Theorem 4 (De Winter, Schillewaert (2010))

A quasi-Hermitian variety of $\text{PG}(r, q^2)$, $r \geq 3$ is classical if it has the same intersection numbers with respect to spaces of codimension 2 as a non-singular Hermitian variety.

Theorem 5 (A., Bartoli, Storme, Weiner (2018))

A quasi-Hermitian variety of $\text{PG}(r, q^2)$, with $r = 3$ and $q = p^h > 4$, or $r \geq 4$, $q = p > 4$, or $r \geq 4$, $q = p^2$, $p > 3$ prime, is classical if and only if it is in the \mathbb{F}_p -code spanned by the hyperplanes of $\text{PG}(r, q^2)$.

Theorem 6 (Napolitano (2023))

Let \mathcal{H} be a set of points in $\text{PG}(3, q^2)$, with $q \neq 2$, such that:

- \mathcal{H} has the same size as the Hermitian surface;
- \mathcal{H} contains no plane;
- every line is either fully contained in \mathcal{H} or intersects \mathcal{H} in at most $q + 1$ points;
- every plane intersects \mathcal{H} in at least $q^3 + 1$ points.

Then \mathcal{H} is a **quasi-Hermitian variety**.

Moreover, if there is no external line, then \mathcal{H} is a **Hermitian surface**.

Research Question and Approach

Research Question and Approach

Problem

Can we characterize BM and BT quasi-Hermitian varieties based on their incidence properties or their automorphism groups?

Research Question and Approach

Problem

Can we characterize BM and BT quasi-Hermitian varieties based on their incidence properties or their automorphism groups?

Our Approach

- Classify these varieties up to projective equivalence.
- Determine their full automorphism groups.
- Derive group-theoretic characterizations.

Contents

- 1 Preliminaries
- 2 Constructions of Q -H varieties
- 3 Automorphism groups and equivalences
- 4 Applications
- 5 Open Problems

Projective equivalence classes of $\mathcal{M}_{a,b}$.

Projective equivalence classes of $\mathcal{M}_{a,b}$.

How many projectively inequivalent BM Q-H varieties can be obtained from varying the parameters (a, b) ?

Projective equivalence classes of $\mathcal{M}_{a,b}$.

How many projectively inequivalent BM Q-H varieties can be obtained from varying the parameters (a, b) ?

Theorem 7 (Baker and Ebert (1982), Ebert (1993))

Let $q = p^n \geq 4$ be a prime power. Then the number of projectively inequivalent BM unitals of $\text{PG}(2, q^2)$ is

$$\frac{1}{2n} \left[n_0 + \sum_{k|n} \Phi \left(\frac{2n}{k} \right) p^k \right],$$

where Φ is the Euler Φ -function and n_0 is the odd part of n if $p > 2$ or $n_0 = 0$ if $p = 2$.

Case $r = 3$

Theorem 8 (A., Giuzzi (2023))

Let $q = p^n$ with p an odd prime. Then the number of projectively inequivalent BM quasi-Hermitian varieties $\mathcal{M}_{a,b}$ of $\text{PG}(3, q^2)$ is

$$\frac{1}{n} \left(\sum_{k|n} \Phi \left(\frac{n}{k} \right) p^k \right) - 2,$$

where Φ is the Euler Φ -function.

Case $r = 3$

Theorem 8 (A., Giuzzi (2023))

Let $q = p^n$ with p an odd prime. Then the number of projectively inequivalent BM quasi-Hermitian varieties $\mathcal{M}_{a,b}$ of $\text{PG}(3, q^2)$ is

$$\frac{1}{n} \left(\sum_{k|n} \Phi \left(\frac{n}{k} \right) p^k \right) - 2,$$

where Φ is the Euler Φ -function.

Theorem 9 (A., Giuzzi, Montinaro, Siconolfi (2025))

All BM quasi-Hermitian varieties $\mathcal{M}_{a,b}$ of $\text{PG}(3, q^2)$, q even are equivalent.

Crucial tools

Assume q odd and set:

$$\ell_1 : X_1 - \nu X_2 = 0 = X_0, \quad \ell_2 : X_1 + \nu X_2 = 0 = X_0$$

where $\nu \in \text{GF}(q^2)$ such that $\nu^2 + 1 = 0$.

Theorem 10 (A., Giuzzi (2023))

Let $\mathcal{M}_{a,b}$ be a BM quasi-Hermitian variety of $\text{PG}(3, q^2)$, with $q \equiv 1 \pmod{4}$. Then,

- through each affine point of $\mathcal{M}_{a,b}$ pass exactly two lines of $\mathcal{M}_{a,b}$;
- through each point at infinity on the union $\ell_1 \cup \ell_2$ there pass $q + 1$ lines of a pencil contained in $\mathcal{M}_{a,b}$;
- through each point at infinity not on $\ell_1 \cup \ell_2$ there passes only one line of $\mathcal{M}_{a,b}$.

Theorem 11 (A., Giuzzi (2023))

Let $\mathcal{M}_{a,b}$ be a BM quasi-Hermitian variety of $\text{PG}(3, q^2)$, with $q \equiv 3 \pmod{4}$. Then,

- no line of $\mathcal{M}_{a,b}$ passes through any affine point of $\mathcal{M}_{a,b}$;
- through each point at infinity in $(\mathcal{M}_{a,b} \cap \Sigma_\infty) \setminus \{P_\infty\}$ there passes only one line of $\mathcal{M}_{a,b}$.
- through the point $P_\infty(0, 0, \dots, 1)$ there are $q + 1$ lines contained in $\mathcal{M}_{a,b}$

Suppose q even and set ℓ_∞ : $X_0 = X_1 + X_2 = 0$.

Suppose q even and set $\ell_\infty: X_0 = X_1 + X_2 = 0$.

Theorem 12 (A., Giuzzi, Montinaro, Siconolfi (2025))

Let $\mathcal{M}_{a,b}$ be the BM quasi-Hermitian variety of $\text{PG}(3, q^2)$, with q even. Then:

- *through each affine point of $\mathcal{M}_{a,b}$ there passes exactly one line of $\mathcal{M}_{a,b}$;*
- *through each point at infinity in $\mathcal{M}_{a,b} \cap \ell_\infty$ there pass $q + 1$ lines of a pencil contained in $\mathcal{M}_{a,b}$.*

Automorphism groups in $\text{PG}(3, q^2)$, q even prime power

Let ϕ_s , ψ_γ , μ_δ , and τ_e be the collineations associated with the following matrices, where:

- $s, e \in \text{GF}(q)$,
- $\delta \in \text{GF}(q)^*$,
- $\gamma = (\gamma_1, \gamma_2) \in \text{GF}(q^2)^2$.

$$\phi_s : \begin{pmatrix} 1 & 0 & 0 & s \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}, \psi_\gamma(a, b) : \begin{pmatrix} 1 & \gamma_1 & \gamma_2 & a(\gamma_1^2 + \gamma_2^2) + b(\gamma_1^{q+1} + \gamma_2^{q+1}) \\ 0 & 1 & 0 & (b + b^q)\gamma_1^q \\ 0 & 0 & 1 & (b + b^q)\gamma_2^q \\ 0 & 0 & 0 & 1 \end{pmatrix}$$

$$\mu_\delta : \text{diag}(1, \delta, \delta, \delta^2), \quad \tau_e : \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & e+1 & e & 0 \\ 0 & e & e+1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}$$

Theorem 13 (A., Giuzzi, Montinaro, Siconolfi (2025))

The stabilizer of $\mathcal{M}_{a,b}$ in $PGL_4(q^2)$, q even, is the group

$$G(a, b) = \langle \phi_s, \psi_\gamma(a, b), \tau_e, \mu_\delta : \gamma \in \text{GF}(q^2)^2, s, e, \delta \in \text{GF}(q), \delta \neq 0 \rangle$$

of order $q^6(q-1)$.

Theorem 13 (A., Giuzzi, Montinaro, Siconolfi (2025))

The stabilizer of $\mathcal{M}_{a,b}$ in $PGL_4(q^2)$, q even, is the group

$$G(a, b) = \langle \phi_s, \psi_\gamma(a, b), \tau_e, \mu_\delta : \gamma \in GF(q^2)^2, s, e, \delta \in GF(q), \delta \neq 0 \rangle$$

of order $q^6(q-1)$.

Theorem 14 (A., Giuzzi, Montinaro, Siconolfi (2025))

Let $\sigma \in PGL_4(q^2)$ be induced by a generator of $\text{Aut}(GF(q^2))$, q even, and let $\beta \in PGL_4(q^2)$ map $\mathcal{M}_{1,\epsilon}$ with $\text{tr}_{q^2/q}(\epsilon) = 1$ onto $\mathcal{M}_{a,b}$. Then the stabilizer in $PGL_4(q^2)$ of $\mathcal{M}_{a,b}$ is

$$\Gamma(a, b) = \langle \phi_s, \psi_\gamma(a, b), \tau_e, \mu_\delta, \sigma^\beta : \gamma \in GF(q^2)^2, s, e, \delta \in GF(q), \delta \neq 0 \rangle,$$

and its order is $q^6(q-1)\log_2 q$.

Projective Equivalence Classes of $\mathcal{H}_\varepsilon^r$

Case $r = 2$: The BT quasi-Hermitian varieties coincide with the BT unitals.

Projective Equivalence Classes of $\mathcal{H}_\varepsilon^r$

Case $r = 2$: The BT quasi-Hermitian varieties coincide with the BT unitals.

Result: All BT unitals are equivalent under the action of $P\Gamma L(3, q^2)$, as proven by **J. Faulkner** and **G. Van de Voorde** (2025), resolving an open problem posed by Barwick and Ebert in their book *Unitals in Projective Planes* (2008).

Projective Equivalence Classes of $\mathcal{H}_\varepsilon^r$

Case $r = 2$: The BT quasi-Hermitian varieties coincide with the BT unitals.

Result: All BT unitals are equivalent under the action of $P\Gamma L(3, q^2)$, as proven by **J. Faulkner** and **G. Van de Voorde** (2025), resolving an open problem posed by Barwick and Ebert in their book *Unitals in Projective Planes* (2008).

Problem

What occurs in higher-dimensional spaces?

Theorem 15 (A., Montinaro (submitted))

Let $\varepsilon_1, \varepsilon_2 \in \mathbb{F}_{q^2} : \varepsilon_i^2 + \varepsilon_i = \delta_i$ with $T(\delta_i) = 1$, $i = 1, 2$. Assume $\alpha = \varepsilon_2 - \varepsilon_1$, $B = \left(\frac{\alpha}{\delta_1}\right)^{\sigma/2}$, $\rho = \left(\frac{\delta_1}{\delta_2}\right)^{\sigma/2+1}$ and $r \geq 2$. The projectivity ξ represented by the $(r+1) \times (r+1)$ matrix A below, maps $\mathcal{H}_{\varepsilon_1}^r$ onto $\mathcal{H}_{\varepsilon_2}^r$

$$A = \begin{pmatrix} 1 & B\rho\frac{\varepsilon_2}{\varepsilon_1} & B\rho\frac{\varepsilon_2}{\varepsilon_1} & \cdots & B\rho\frac{\varepsilon_2}{\varepsilon_1} & (r-1)\rho^2 B^{\sigma+2} \frac{\varepsilon_2^{q+2}}{\varepsilon_1^{q+1}} \\ 0 & \frac{\rho\varepsilon_2}{\varepsilon_1} & 0 & \cdots & 0 & B\rho^2 \frac{\varepsilon_2^{q+1}}{\varepsilon_1^{q+1}} \\ 0 & 0 & \frac{\rho\varepsilon_2}{\varepsilon_1} & \cdots & 0 & B\rho^2 \frac{\varepsilon_2^{q+1}}{\varepsilon_1^{q+1}} \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & 0 & \cdots & \frac{\rho\varepsilon_2}{\varepsilon_1} & B\rho^2 \frac{\varepsilon_2^{q+1}}{\varepsilon_1^{q+1}} \\ 0 & 0 & 0 & \cdots & 0 & \frac{\rho^2 \varepsilon_2^{q+1}}{\varepsilon_1^{q+1}} \end{pmatrix}$$

Theorem 15 (A., Montinaro (submitted))

Let $\varepsilon_1, \varepsilon_2 \in \mathbb{F}_{q^2} : \varepsilon_i^2 + \varepsilon_i = \delta_i$ with $T(\delta_i) = 1$, $i = 1, 2$. Assume $\alpha = \varepsilon_2 - \varepsilon_1$, $B = (\frac{\alpha}{\delta_1})^{\sigma/2}$, $\rho = \left(\frac{\delta_1}{\delta_2}\right)^{\sigma/2+1}$ and $r \geq 2$. The projectivity ξ represented by the $(r+1) \times (r+1)$ matrix A below, maps $\mathcal{H}_{\varepsilon_1}^r$ onto $\mathcal{H}_{\varepsilon_2}^r$

$$A = \begin{pmatrix} 1 & B\rho\frac{\varepsilon_2}{\varepsilon_1} & B\rho\frac{\varepsilon_2}{\varepsilon_1} & \cdots & B\rho\frac{\varepsilon_2}{\varepsilon_1} & (r-1)\rho^2 B^{\sigma+2} \frac{\varepsilon_2^{q+2}}{\varepsilon_1^{q+1}} \\ 0 & \frac{\rho\varepsilon_2}{\varepsilon_1} & 0 & \cdots & 0 & B\rho^2 \frac{\varepsilon_2^{q+1}}{\varepsilon_1^{q+1}} \\ 0 & 0 & \frac{\rho\varepsilon_2}{\varepsilon_1} & \cdots & 0 & B\rho^2 \frac{\varepsilon_2^{q+1}}{\varepsilon_1^{q+1}} \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & 0 & \cdots & \frac{\rho\varepsilon_2}{\varepsilon_1} & B\rho^2 \frac{\varepsilon_2^{q+1}}{\varepsilon_1^{q+1}} \\ 0 & 0 & 0 & \cdots & 0 & \frac{\rho^2 \varepsilon_2^{q+1}}{\varepsilon_1^{q+1}} \end{pmatrix}$$

Lemma 16 (A., Montinaro (submitted))

A Sylow 2-subgroup of $\text{Aut}(\mathcal{H}_{\varepsilon}^r)$ fixes a unique incident point-hyperplane pair of $\text{PG}(r, q^2)$.

Collineation group of $\mathcal{H}_\varepsilon^3$

Theorem 17 (A., Montinaro (subm.))

The following hold:

- 1 $\text{Aut}(\mathcal{H}_\varepsilon^3) \cap \text{PGL}_4(q^2) = E \langle \vartheta \rangle$, with $\vartheta : (x_0, x_1, x_2, x_3) \rightarrow (x_0, x_2, x_1, x_3)$, is a group of order $2q^3$;
- 2 $\text{Aut}(\mathcal{H}_\varepsilon^3)$ preserves the triple $(P_\infty, \ell_\infty, \Sigma_\infty)$.

Collineation group of $\mathcal{H}_\varepsilon^3$

Theorem 17 (A., Montinaro (subm.))

The following hold:

- 1 $\text{Aut}(\mathcal{H}_\varepsilon^3) \cap \text{PGL}_4(q^2) = E \langle \vartheta \rangle$, with $\vartheta : (x_0, x_1, x_2, x_3) \rightarrow (x_0, x_2, x_1, x_3)$, is a group of order $2q^3$;
- 2 $\text{Aut}(\mathcal{H}_\varepsilon^3)$ preserves the triple $(P_\infty, \ell_\infty, \Sigma_\infty)$.

Theorem 18 (A., Montinaro (subm.))

$\text{Aut}(\mathcal{H}_\varepsilon^3) = E \langle \vartheta, \phi \rangle$, with $\vartheta : (x_0, x_1, x_2, x_3) \rightarrow (x_0, x_2, x_1, x_3)$ and

$$\phi : (x_0, x_1, x_2, x_3) \rightarrow (x_0^2, x_1^2, x_2^2, x_3^2) \begin{pmatrix} 1 & 1 & 1 & 0 \\ 0 & \delta^{\frac{\sigma}{2}} \varepsilon^q & 0 & \delta^{\frac{\sigma}{2}} \varepsilon^q \\ 0 & 0 & \delta^{\frac{\sigma}{2}} \varepsilon^q & \delta^{\frac{\sigma}{2}} \varepsilon^q \\ 0 & 0 & 0 & \delta^{\sigma+1} \end{pmatrix},$$

is a group of order $4eq^3$.

Tools: some geometric properties

Set $\ell_\infty : X_0 = X_1 + X_2 = 0$.

Tools: some geometric properties

Set $\ell_\infty : X_0 = X_1 + X_2 = 0$.

Theorem 19 (A., Montinaro (submitted))

Let $\mathcal{H}_\varepsilon^3$ be the BT quasi-Hermitian variety of $\text{PG}(3, q^2)$. Then the following statements hold:

- 1 each affine line contained in $\mathcal{H}_\varepsilon^3$ intersects ℓ_∞ at one among the points $L_\infty^{\varepsilon^q \alpha} = (0, 1, 1, \varepsilon^q \alpha)$, where $\alpha \in \text{GF}(q)$;
- 2 through each point $L_\infty^{\varepsilon^q \alpha}$, there exist exactly $q + 1$ coplanar lines contained in $\mathcal{H}_\varepsilon^3$, one of which is ℓ_∞ . Each such set of $q + 1$ lines forms a Hermitian cone $\Pi_0 \mathcal{H}(1, q^2)$.

Contents

- 1 Preliminaries
- 2 Constructions of Q -H varieties
- 3 Automorphism groups and equivalences
- 4 Applications**
- 5 Open Problems

Construction of MDS codes

Let $G_1(x), \dots, G_N(x)$ be N **multivariate polynomials** over $\text{GF}(q)$ and $\mathcal{W} \subset \text{GF}(q)^{n+1}$.

Construction of MDS codes

Let $G_1(x), \dots, G_N(x)$ be N **multivariate polynomials** over $\text{GF}(q)$ and $\mathcal{W} \subset \text{GF}(q)^{n+1}$.

Here, the evaluation code $\mathcal{C} := \mathcal{C}(G_1, \dots, G_N; \mathcal{W})$ defined by G_1, \dots, G_N over a set \mathcal{W} is the image of the map

$$ev_{G_1, \dots, G_N} : \begin{cases} \mathcal{W} \rightarrow \text{GF}(q)^N \\ x \rightarrow (G_1(x), \dots, G_N(x)). \end{cases}$$

Construction of MDS codes

Let $G_1(x), \dots, G_N(x)$ be N **multivariate polynomials** over $\text{GF}(q)$ and $\mathcal{W} \subset \text{GF}(q)^{n+1}$.

Here, the evaluation code $\mathcal{C} := \mathcal{C}(G_1, \dots, G_N; \mathcal{W})$ defined by G_1, \dots, G_N over a set \mathcal{W} is the image of the map

$$\text{ev}_{G_1, \dots, G_N} : \begin{cases} \mathcal{W} \rightarrow \text{GF}(q)^N \\ x \rightarrow (G_1(x), \dots, G_N(x)). \end{cases}$$

Assume that \mathcal{C} has q^t codewords and Hamming distance d .

The Singleton Bound

The Singleton bound

It is known that for a code C over $\text{GF}(q)$ with parameters $[N, q^t, d]$, the following holds:

$$q^t \leq q^{N-d+1}$$

The Singleton Bound

The Singleton bound

It is known that for a code C over $\text{GF}(q)$ with parameters $[N, q^t, d]$, the following holds:

$$q^t \leq q^{N-d+1}$$

When equality holds, the code is called a *Maximum Distance Separable* code (MDS).

The Singleton Bound

The Singleton bound

It is known that for a code C over $\text{GF}(q)$ with parameters $[N, q^t, d]$, the following holds:

$$q^t \leq q^{N-d+1}$$

When equality holds, the code is called a *Maximum Distance Separable* code (MDS).

If the evaluation code \mathcal{C} is MDS, then any $t = N - d + 1$ of the varieties $V(G_i) : G_i = 0$, for $i = 1, \dots, N$, must intersect in exactly one point in the ambient space \mathcal{W} .

The Singleton Bound

The Singleton bound

It is known that for a code C over $\text{GF}(q)$ with parameters $[N, q^t, d]$, the following holds:

$$q^t \leq q^{N-d+1}$$

When equality holds, the code is called a *Maximum Distance Separable* code (MDS).

If the evaluation code \mathcal{C} is MDS, then any $t = N - d + 1$ of the varieties $V(G_i) : G_i = 0$, for $i = 1, \dots, N$, must intersect in exactly one point in the ambient space \mathcal{W} .

Define $\mathcal{W} := \text{GF}(q^2) \times \text{GF}(q^2) \times T$, where T is a transversal of $\text{GF}(q)$ viewed as an additive subgroup of $\text{GF}(q^2)$.

Determinantal Condition

Consider the subset Ω of $\text{GF}(q^2)^2$ such that for each $(\omega_1^i, \omega_2^i) \in \Omega$ with $i \geq 5$, the following condition holds:

$$\det \begin{pmatrix} 1 & \omega_1^1 & \omega_2^1 & (\omega_1^1)^q & (\omega_2^1)^q \\ 1 & \omega_1^2 & \omega_2^2 & (\omega_1^2)^q & (\omega_2^2)^q \\ 1 & \omega_1^3 & \omega_2^3 & (\omega_1^3)^q & (\omega_2^3)^q \\ 1 & \omega_1^4 & \omega_2^4 & (\omega_1^4)^q & (\omega_2^4)^q \\ 1 & \omega_1^5 & \omega_2^5 & (\omega_1^5)^q & (\omega_2^5)^q \end{pmatrix} \neq 0 \quad (\text{DetCond})$$

Fix a basis $(1, \epsilon)$ of $\text{GF}(q^2)$ regarded as a vector space over $\text{GF}(q)$, with $\epsilon \in \text{GF}(q^2) \setminus \text{GF}(q) : \text{tr}_{q^2/q}(\epsilon) = 0$ for q odd or $\text{tr}_{q^2/q}(\epsilon) = 1$ for q even.

Fix a basis $(1, \epsilon)$ of $\text{GF}(q^2)$ regarded as a vector space over $\text{GF}(q)$, with $\epsilon \in \text{GF}(q^2) \setminus \text{GF}(q) : \text{tr}_{q^2/q}(\epsilon) = 0$ for q odd or $\text{tr}_{q^2/q}(\epsilon) = 1$ for q even.

Write $\omega_i^j = \omega_{i,0}^j + \omega_{i,1}^j \epsilon$ for all $i = 1, 2$ and $j = 1, \dots, 5$.

Fix a basis $(1, \epsilon)$ of $\text{GF}(q^2)$ regarded as a vector space over $\text{GF}(q)$, with $\epsilon \in \text{GF}(q^2) \setminus \text{GF}(q) : \text{tr}_{q^2/q}(\epsilon) = 0$ for q odd or $\text{tr}_{q^2/q}(\epsilon) = 1$ for q even.

Write $\omega_i^j = \omega_{i,0}^j + \omega_{i,1}^j \epsilon$ for all $i = 1, 2$ and $j = 1, \dots, 5$.

Then (DetCond) becomes

$$\det \begin{pmatrix} 1 & \omega_{1,0}^1 & \omega_{2,0}^1 & \omega_{1,1}^1 & \omega_{2,1}^1 \\ 1 & \omega_{1,0}^2 & \omega_{2,0}^2 & \omega_{1,1}^2 & \omega_{2,1}^2 \\ 1 & \omega_{1,0}^3 & \omega_{2,0}^3 & \omega_{1,1}^3 & \omega_{2,1}^3 \\ 1 & \omega_{1,0}^4 & \omega_{2,0}^4 & \omega_{1,1}^4 & \omega_{2,1}^4 \\ 1 & \omega_{1,0}^5 & \omega_{2,0}^5 & \omega_{1,1}^5 & \omega_{2,1}^5 \end{pmatrix} \neq 0 \quad (\text{DetCond}q)$$

for any choice of five elements in Ω .

Arcs in Projective Space

Condition (DetCond $_q$) states that the rows of the matrix represent the coordinates of points lying on an arc in $AG(4, q)$.

Arcs in Projective Space

Condition (DetCond q) states that the rows of the matrix represent the coordinates of points lying on an arc in $AG(4, q)$.

An arc in $PG(4, q)$, that is a set of points such that any five span the entire space, has maximum size $q + 1$.

Arcs in Projective Space

Condition (DetCond q) states that the rows of the matrix represent the coordinates of points lying on an arc in $AG(4, q)$.

An arc in $PG(4, q)$, that is a set of points such that any five span the entire space, has maximum size $q + 1$.

Therefore, we have the bound: $|\Omega| \leq q$.

Arcs in Projective Space

Condition (DetCond q) states that the rows of the matrix represent the coordinates of points lying on an arc in $AG(4, q)$.

An arc in $PG(4, q)$, that is a set of points such that any five span the entire space, has maximum size $q + 1$.

Therefore, we have the bound: $|\Omega| \leq q$.

Theorem 20 (Ball, Lavrauw (2019))

The only $(q + 1)$ -arcs in $PG(4, q)$ for $q > 13$ odd or $q \geq 8$ even, are projectively equivalent to the normal rational curve:

$$\Gamma_4 = \{(1, t, t^2, t^3, t^4) \mid t \in GF(q)\} \cup \{(0, 0, 0, 0, 1)\}.$$

Defining the Forms F_i

We can set

$$\Omega := \{(t + \varepsilon t^2, t^3 + \varepsilon t^4) : t \in \text{GF}(q)\}$$

so that it corresponds to an arc in $AG(4, q)$.

Defining the Forms F_i

We can set

$$\Omega := \{(t + \varepsilon t^2, t^3 + \varepsilon t^4) : t \in \text{GF}(q)\}$$

so that it corresponds to an arc in $AG(4, q)$.

Now, consider the following forms

$$\begin{aligned} F_i(X_0, X_1, X_2, X_3) = & (b - b^q)X_0^{q-1}(X_1^{q+1} + X_2^{q+1}) + \\ & X_0^q X_3^q - X_3 X_0^{2q-1} + a^q(X_1^{2q} + X_2^{2q}) - aX_0^{2q-2}(X_1^2 + X_2^2) + \\ & [2a^q(\omega_1^i)^q - (b^q - b)\omega_1^i]X_0^q X_1^q + [2a^q(\omega_2^i)^q - (b^q - b)\omega_2^i]X_0^q X_2^q + \\ & [2a\omega_1^i + (b - b^q)(\omega_1^i)^q]X_0^{2q-1} X_1 + [2a\omega_2^i + (b - b^q)(\omega_2^i)^q]X_0^{2q-1} X_2 \end{aligned}$$

where $(\omega_1^i, \omega_2^i) \in \Omega$ for $i = 1, \dots, q$.

These forms F_i define BM Q-H varieties such that any subset of five intersects at exactly one point in \mathcal{W} .

These forms F_i define BM Q-H varieties such that any subset of five intersects at exactly one point in \mathcal{W} .

Let $T_0 = \{x \in \text{GF}(q^2) : x^q + x = 0\}$.

These forms F_i define BM Q-H varieties such that any subset of five intersects at exactly one point in \mathcal{W} .

Let $T_0 = \{x \in \text{GF}(q^2) : x^q + x = 0\}$.

Set

$$\mathcal{C}(F_1, \dots, F_q; \mathcal{W}) = \\ \{(F_1(1, x, y, z), F_2(1, x, y, z), \dots, F_q(1, x, y, z)) | (x, y, z) \in \mathcal{W}\}.$$

These forms F_i define BM Q-H varieties such that any subset of five intersects at exactly one point in \mathcal{W} .

Let $T_0 = \{x \in \text{GF}(q^2) : x^q + x = 0\}$.

Set

$$\mathcal{C}(F_1, \dots, F_q; \mathcal{W}) = \{(F_1(1, x, y, z), F_2(1, x, y, z), \dots, F_q(1, x, y, z)) \mid (x, y, z) \in \mathcal{W}\}.$$

Theorem 21 (A., Giuzzi, Siconolfi (2025))

Let $q > 13$. The code $\mathcal{C}(F_1, \dots, F_q; \mathcal{W})$ is a $\text{GF}(q)$ -linear $[q, 5, q - 4]$ -MDS code over T_0 .

Some Equivalent Codes

Write $\varepsilon^q + \varepsilon = a_0$, so $\text{tr}_{q^2/q}(x_0 + \varepsilon x_1) = 2x_0 + a_0x_1$, for $x_0, x_1 \in \text{GF}(q)$.

Some Equivalent Codes

Write $\varepsilon^q + \varepsilon = a_0$, so $\text{tr}_{q^2/q}(x_0 + \varepsilon x_1) = 2x_0 + a_0x_1$, for $x_0, x_1 \in \text{GF}(q)$.

So $T_0 = \{x \in \text{GF}(q^2) | \text{tr}_{q^2/q}(x) = 0\} = \{(a_0 - 2\varepsilon)x_0 | x_0 \in \text{GF}(q)\}$.

Some Equivalent Codes

Write $\varepsilon^q + \varepsilon = a_0$, so $\text{tr}_{q^2/q}(x_0 + \varepsilon x_1) = 2x_0 + a_0x_1$, for $x_0, x_1 \in \text{GF}(q)$.

So $T_0 = \{x \in \text{GF}(q^2) | \text{tr}_{q^2/q}(x) = 0\} = \{(a_0 - 2\varepsilon)x_0 | x_0 \in \text{GF}(q)\}$.

Put $\theta = (a_0 - 2\varepsilon)$.

Some Equivalent Codes

Write $\varepsilon^q + \varepsilon = a_0$, so $\text{tr}_{q^2/q}(x_0 + \varepsilon x_1) = 2x_0 + a_0x_1$, for $x_0, x_1 \in \text{GF}(q)$.

So $T_0 = \{x \in \text{GF}(q^2) \mid \text{tr}_{q^2/q}(x) = 0\} = \{(a_0 - 2\varepsilon)x_0 \mid x_0 \in \text{GF}(q)\}$.

Put $\theta = (a_0 - 2\varepsilon)$.

Theorem 22 (A., Giuzzi, Siconolfi (2025))

The code $\mathcal{C}' := (\theta^{-1})\mathcal{C}(F_1, F_1, \dots, F_q; \mathcal{W})$ is equivalent to a q -ary Reed-Solomon code. In particular, it can be further extended to a $[q+1, 5, q-3]_q$ Reed-Solomon code.

Some Equivalent Codes

Write $\varepsilon^q + \varepsilon = a_0$, so $\text{tr}_{q^2/q}(x_0 + \varepsilon x_1) = 2x_0 + a_0x_1$, for $x_0, x_1 \in \text{GF}(q)$.

So $T_0 = \{x \in \text{GF}(q^2) | \text{tr}_{q^2/q}(x) = 0\} = \{(a_0 - 2\varepsilon)x_0 | x_0 \in \text{GF}(q)\}$.

Put $\theta = (a_0 - 2\varepsilon)$.

Theorem 22 (A., Giuzzi, Siconolfi (2025))

The code $\mathcal{C}' := (\theta^{-1})\mathcal{C}(F_1, F_1, \dots, F_q; \mathcal{W})$ is equivalent to a q -ary Reed-Solomon code. In particular, it can be further extended to a $[q+1, 5, q-3]_q$ Reed-Solomon code.

The extended Reed-Solomon code is obtained by \mathcal{C}' adding to each codeword the component $F_{q+1}(1, x, y, z)$ where

$$F_{q+1}(X_0, X_1, X_2, X_3) = (b^q - b)X_2^q X_0 + 2aX_2X_0^q$$

Non-degeneracy and Minimal Codes

Let \mathcal{C} be a q -ary linear $[n, r, d]$ code. The *support* of a codeword $\mathbf{c} = (c_1, \dots, c_n) \in \mathcal{C}$ is: $\text{supp}(\mathbf{c}) := \{i : c_i \neq 0\}$.

Non-degeneracy and Minimal Codes

Let \mathcal{C} be a q -ary linear $[n, r, d]$ code. The *support* of a codeword $\mathbf{c} = (c_1, \dots, c_n) \in \mathcal{C}$ is: $\text{supp}(\mathbf{c}) := \{i : c_i \neq 0\}$.

The *support* of the code \mathcal{C} is: $\text{supp}(\mathcal{C}) := \bigcup_{\mathbf{c} \in \mathcal{C}} \text{supp}(\mathbf{c})$.

Non-degeneracy and Minimal Codes

Let \mathcal{C} be a q -ary linear $[n, r, d]$ code. The *support* of a codeword $\mathbf{c} = (c_1, \dots, c_n) \in \mathcal{C}$ is: $\text{supp}(\mathbf{c}) := \{i : c_i \neq 0\}$.

The *support* of the code \mathcal{C} is: $\text{supp}(\mathcal{C}) := \bigcup_{\mathbf{c} \in \mathcal{C}} \text{supp}(\mathbf{c})$.

The code \mathcal{C} is *non-degenerate* if: $\text{supp}(\mathcal{C}) = \{1, \dots, n\}$.

Non-degeneracy and Minimal Codes

Let \mathcal{C} be a q -ary linear $[n, r, d]$ code. The *support* of a codeword $\mathbf{c} = (c_1, \dots, c_n) \in \mathcal{C}$ is: $\text{supp}(\mathbf{c}) := \{i : c_i \neq 0\}$.

The *support* of the code \mathcal{C} is: $\text{supp}(\mathcal{C}) := \bigcup_{\mathbf{c} \in \mathcal{C}} \text{supp}(\mathbf{c})$.

The code \mathcal{C} is *non-degenerate* if: $\text{supp}(\mathcal{C}) = \{1, \dots, n\}$.

A codeword $\mathbf{c} \in \mathcal{C}$ is *minimal* if for every non-zero codeword $\mathbf{c}' \in \mathcal{C}$ such that: $\text{supp}(\mathbf{c}') \subseteq \text{supp}(\mathbf{c})$, we have $\mathbf{c}' \in \langle \mathbf{c} \rangle$.

Non-degeneracy and Minimal Codes

Let \mathcal{C} be a q -ary linear $[n, r, d]$ code. The *support* of a codeword $\mathbf{c} = (c_1, \dots, c_n) \in \mathcal{C}$ is: $\text{supp}(\mathbf{c}) := \{i : c_i \neq 0\}$.

The *support* of the code \mathcal{C} is: $\text{supp}(\mathcal{C}) := \bigcup_{\mathbf{c} \in \mathcal{C}} \text{supp}(\mathbf{c})$.

The code \mathcal{C} is *non-degenerate* if: $\text{supp}(\mathcal{C}) = \{1, \dots, n\}$.

A codeword $\mathbf{c} \in \mathcal{C}$ is *minimal* if for every non-zero codeword $\mathbf{c}' \in \mathcal{C}$ such that: $\text{supp}(\mathbf{c}') \subseteq \text{supp}(\mathbf{c})$, we have $\mathbf{c}' \in \langle \mathbf{c} \rangle$.

The code \mathcal{C} is *minimal* if all its non-zero codewords are minimal.

Non-degeneracy and Minimal Codes

Let \mathcal{C} be a q -ary linear $[n, r, d]$ code. The *support* of a codeword $\mathbf{c} = (c_1, \dots, c_n) \in \mathcal{C}$ is: $\text{supp}(\mathbf{c}) := \{i : c_i \neq 0\}$.

The *support* of the code \mathcal{C} is: $\text{supp}(\mathcal{C}) := \bigcup_{\mathbf{c} \in \mathcal{C}} \text{supp}(\mathbf{c})$.

The code \mathcal{C} is *non-degenerate* if: $\text{supp}(\mathcal{C}) = \{1, \dots, n\}$.

A codeword $\mathbf{c} \in \mathcal{C}$ is *minimal* if for every non-zero codeword $\mathbf{c}' \in \mathcal{C}$ such that: $\text{supp}(\mathbf{c}') \subseteq \text{supp}(\mathbf{c})$, we have $\mathbf{c}' \in \langle \mathbf{c} \rangle$.

The code \mathcal{C} is *minimal* if all its non-zero codewords are minimal.

References:

- Alfarano, Borello, Neri (2022)
- Scotti (2024)
- Alon, Bishnoi, Das, Neri (2024)

Code \rightarrow Projective System

Let \mathcal{C} be a non-degenerate q -ary linear $[n, r, d]$ code. Its *projective system* is the (multi)set Ω of points in $\text{PG}(r - 1, q)$ corresponding to the columns of any generator matrix of \mathcal{C} .

Code \rightarrow Projective System

Let \mathcal{C} be a non-degenerate q -ary linear $[n, r, d]$ code. Its *projective system* is the (multi)set Ω of points in $\text{PG}(r - 1, q)$ corresponding to the columns of any generator matrix of \mathcal{C} .

Projective System \rightarrow Code

Any (multi)set Ω of points in $\text{PG}(r - 1, q)$ defines a code $\mathcal{C}(\Omega)$ whose generator matrix has as columns the coordinates of the points in Ω .

Code \rightarrow Projective System

Let \mathcal{C} be a non-degenerate q -ary linear $[n, r, d]$ code. Its *projective system* is the (multi)set Ω of points in $\text{PG}(r - 1, q)$ corresponding to the columns of any generator matrix of \mathcal{C} .

Projective System \rightarrow Code

Any (multi)set Ω of points in $\text{PG}(r - 1, q)$ defines a code $\mathcal{C}(\Omega)$ whose generator matrix has as columns the coordinates of the points in Ω .

Minimum Distance

For a code $\mathcal{C} = \mathcal{C}(\Omega)$, the minimum distance is given by:

$$d = |\Omega| - \max\{|\Omega \cap \Pi| : \Pi \text{ is a hyperplane of } \text{PG}(\langle \Omega \rangle)\}.$$

Characterization of Minimal Codes

Theorem 23 (Alfarano, Borello, Neri (2022))

Let Ω be a set of points in $\text{PG}(r, q)$ such that $\langle \Omega \rangle = \text{PG}(r, q)$. Then the code $\mathcal{C}(\Omega)$ is minimal if and only if for every hyperplane Π of $\text{PG}(r, q)$,

$$\langle \Pi \cap \Omega \rangle = \Pi,$$

i.e., Ω is a cutting set.

Characterization of Minimal Codes

Theorem 23 (Alfarano, Borello, Neri (2022))

Let Ω be a set of points in $\text{PG}(r, q)$ such that $\langle \Omega \rangle = \text{PG}(r, q)$. Then the code $\mathcal{C}(\Omega)$ is minimal if and only if for every hyperplane Π of $\text{PG}(r, q)$,

$$\langle \Pi \cap \Omega \rangle = \Pi,$$

i.e., Ω is a cutting set.

Sufficient Condition (Ashikhmin–Barg)(1994)

A q -ary linear code \mathcal{C} is minimal if $\frac{w_{\min}}{w_{\max}} > \frac{q-1}{q}$, where w_{\min} and w_{\max} are the minimum and maximum weights of non-zero codewords.

Minimality of the Code Associated with Q-H Varieties

If \mathcal{H} is a quasi-Hermitian variety in $\text{PG}(r, q^2)$, then the associated code $\mathcal{C}(\mathcal{H})$ has exactly **two distinct weights**.

Minimality of the Code Associated with Q-H Varieties

If \mathcal{H} is a quasi-Hermitian variety in $\text{PG}(r, q^2)$, then the associated code $\mathcal{C}(\mathcal{H})$ has exactly **two distinct weights**.

Although the sufficient condition for minimality does not hold in this case, the **Characterization Theorem** for minimal codes can still be applied.

Minimality of the Code Associated with Q-H Varieties

If \mathcal{H} is a quasi-Hermitian variety in $\text{PG}(r, q^2)$, then the associated code $\mathcal{C}(\mathcal{H})$ has exactly **two distinct weights**.

Although the sufficient condition for minimality does not hold in this case, the **Characterization Theorem** for minimal codes can still be applied.

Theorem 24 (A., Giuzzi, Ceria (2022))

Let \mathcal{H} be a quasi-Hermitian variety in $\text{PG}(r, q^2)$. Then, the code $\mathcal{C}(\mathcal{H})$ is minimal.

Codes from $\mathcal{V}_\varepsilon^r$

Let us consider the BT quasi-Hermitian variety:

$$\mathcal{H}_\varepsilon^r = (\mathcal{V}_\varepsilon^r \setminus \Sigma_\infty) \cup \mathcal{F}.$$

What can be said about the code $\mathcal{C}_\varepsilon^r := \mathcal{C}(\mathcal{V}_\varepsilon^r)$?

Codes from $\mathcal{V}_\varepsilon^r$

Let us consider the BT quasi-Hermitian variety:

$$\mathcal{H}_\varepsilon^r = (\mathcal{V}_\varepsilon^r \setminus \Sigma_\infty) \cup \mathcal{F}.$$

What can be said about the code $\mathcal{C}_\varepsilon^r := \mathcal{C}(\mathcal{V}_\varepsilon^r)$?

Key Properties

- It is a **few-weight code**.
- In some cases, the code is **minimal**

Codes from $\mathcal{V}_\varepsilon^r$

Let us consider the BT quasi-Hermitian variety:

$$\mathcal{H}_\varepsilon^r = (\mathcal{V}_\varepsilon^r \setminus \Sigma_\infty) \cup \mathcal{F}.$$

What can be said about the code $\mathcal{C}_\varepsilon^r := \mathcal{C}(\mathcal{V}_\varepsilon^r)$?

Key Properties

- It is a **few-weight code**.
- In some cases, the code is **minimal**

Theorem 25 (A., Giuzzi, Longobardi, Siconolfi (submitted))

The linear code $\mathcal{C}_\varepsilon^r$ generated by the projective points of $\mathcal{V}_\varepsilon^r$ in $\text{PG}(r, q^2)$ is a $(r+1)$ -dimensional minimal code for $r=3$ and $e \equiv 3 \pmod{4}$ or $r \geq 4$ and any odd integer e .

Key Lemma (A., Giuzzi, Longobardi, Siconolfi (submitted))

Let $r \geq 2$ and $q = 2^e$ with $e \geq 3$ odd. Then the Fermat hypersurface \mathcal{F}_n^r of degree $n = 2^{\frac{e-1}{2}} + 1$ in $\text{PG}(r, q^2)$, defined by:

$$\mathcal{F}_n^r : X_0^n + X_1^n + \cdots + X_r^n = 0$$

spans the entire projective space: $\langle \mathcal{F}_n^r \rangle = \text{PG}(r, q^2)$.

Key Lemma (A., Giuzzi, Longobardi, Siconolfi (submitted))

Let $r \geq 2$ and $q = 2^e$ with $e \geq 3$ odd. Then the Fermat hypersurface \mathcal{F}_n^r of degree $n = 2^{\frac{e-1}{2}} + 1$ in $\text{PG}(r, q^2)$, defined by:

$$\mathcal{F}_n^r : X_0^n + X_1^n + \cdots + X_r^n = 0$$

spans the entire projective space: $\langle \mathcal{F}_n^r \rangle = \text{PG}(r, q^2)$.

The intersection $\mathcal{V}_\varepsilon^r \cap \Sigma_\infty$ is represented by:

$$X_0 = X_1^{2^{\frac{e-1}{2}}+1} + X_2^{2^{\frac{e-1}{2}}+1} + \cdots + X_{r-1}^{2^{\frac{e-1}{2}}+1} = 0.$$

Key Lemma (A., Giuzzi, Longobardi, Siconolfi (submitted))

Let $r \geq 2$ and $q = 2^e$ with $e \geq 3$ odd. Then the Fermat hypersurface \mathcal{F}_n^r of degree $n = 2^{\frac{e-1}{2}} + 1$ in $\text{PG}(r, q^2)$, defined by:

$$\mathcal{F}_n^r : X_0^n + X_1^n + \cdots + X_r^n = 0$$

spans the entire projective space: $\langle \mathcal{F}_n^r \rangle = \text{PG}(r, q^2)$.

The intersection $\mathcal{V}_\varepsilon^r \cap \Sigma_\infty$ is represented by:

$$X_0 = X_1^{2^{\frac{e-1}{2}}+1} + X_2^{2^{\frac{e-1}{2}}+1} + \cdots + X_{r-1}^{2^{\frac{e-1}{2}}+1} = 0.$$

Remark

To determine the length and the weights of the projective linear code $\mathcal{C}_\varepsilon^r$, with $r \geq 3$, it is necessary to compute the number of $\text{GF}(q^2)$ -rational points of the Fermat hypersurface \mathcal{F}_n^{r-2} .

Proposition 26 (A., Giuzzi, Longobardi, Siconolfi (submitted))

The number N_{q^2} of $\text{GF}(q^2)$ -rational points of \mathcal{F}_n^r in $\text{PG}(r, q^2)$, $r \geq 2$, satisfies the following properties:

- ① *if $e \equiv 1 \pmod{4}$ then $N_{q^2} = \theta_{q^2}(r - 1)$;*
- ② *if $e \equiv 3 \pmod{4}$ then*

$$N_{q^2} \leq (n - 1)q^{2(r-1)} + nq^{2(r-2)} + \theta_{q^2}(r - 3).$$

- ③ *if $e = 3$ and $r = 2$ then $N_{q^2} = (q + 1)^2$.*

Proposition 26 (A., Giuzzi, Longobardi, Siconolfi (submitted))

The number N_{q^2} of $\text{GF}(q^2)$ -rational points of \mathcal{F}_n^r in $\text{PG}(r, q^2)$, $r \geq 2$, satisfies the following properties:

- ① *if $e \equiv 1 \pmod{4}$ then $N_{q^2} = \theta_{q^2}(r - 1)$;*
- ② *if $e \equiv 3 \pmod{4}$ then*

$$N_{q^2} \leq (n - 1)q^{2(r-1)} + nq^{2(r-2)} + \theta_{q^2}(r - 3).$$

- ③ *if $e = 3$ and $r = 2$ then $N_{q^2} = (q + 1)^2$.*

Proposition 27 (A., Giuzzi, Longobardi, Siconolfi (submitted))

If $e \equiv 3 \pmod{4}$, then the weights of the projective code $\mathcal{C}_\varepsilon^3 := \mathcal{C}(\mathcal{V}_\varepsilon^3)$ belong to the set

$$\{q^5, q^5 - q^3 + 3q^2, q^5 - q^3 + 2q^2, q^5 - q^3 + 3q^2 + q - 2, \\ q^5 - q^3 + 2q^2 + q - 2\}.$$

Theorem 28 (A., Giuzzi, Longobardi, Siconolfi (submitted))

The linear code $\mathcal{C}_\epsilon^4 = \mathcal{C}(\mathcal{V}_\epsilon^4)$ for $q = 2^e$ and $e > 1$ an odd integer is a 5-dimensional minimal code with the following parameters

- ① *If $e \equiv 1 \pmod{4}$, then \mathcal{C}_ϵ^4 has length $|\mathcal{V}_\epsilon^4| = q^7 + q^4 + q^2 + 1$ and weights*

$$\{q^7, q^7 - q^5 + q^4 + q^3 - q^2, q^7 - q^5 + q^4 + q^2, \\ q^7 - q^5 + q^4, q^7 - q^5 + q^4 - q^2, q^7 - q^5 + q^4 - 4q^2\};$$

- ② *If $e = 3$, then \mathcal{C}_ϵ^4 has length $|\mathcal{V}_\epsilon^4| = q^7 + q^4 + 2q^3 + q^2 + 1$ and weights*

$$\{q^7, q^7 - q^5 + q^4 + 3q^3 - q^2, q^7 - q^5 + q^4 + 2q^3 + q^2, \\ q^7 - q^5 + q^4 + 2q^3, q^7 - q^5 + q^4 + 2q^3 - q^2, q^7 - q^5 + q^4 + 2q^3 - 2q^2\}.$$

Cutting Gap: Measuring Cutting Sets

Cutting Gap: Measuring Cutting Sets

Definition 29 (Cutting Gap)

Let Ω be a non-empty set of points in $\text{PG}(r, q)$, and let $0 \leq k \leq r$. The *k-th cutting gap* of Ω is defined as:

$$\tau_k(\Omega) = k - \min \{ \dim(\langle \Pi \cap \Omega \rangle) : \dim(\Pi) = k \}.$$

Cutting Gap: Measuring Cutting Sets

Definition 29 (Cutting Gap)

Let Ω be a non-empty set of points in $\text{PG}(r, q)$, and let $0 \leq k \leq r$. The k -th cutting gap of Ω is defined as:

$$\tau_k(\Omega) = k - \min \{ \dim(\langle \Pi \cap \Omega \rangle) : \dim(\Pi) = k \}.$$

Ω is a k -fold strong blocking set if and only if $\tau_k(\Omega) = 0$, according to the definitions in

- A.A. Davydov, M. Giulietti, S. Marcugini, F. Pambianco, Linear nonbinary covering codes and saturating sets in projective spaces, *Adv. Math. Commun.* **5**, (2011)

Cutting Sets and Their Properties

We also refer to this property as being *k-cutting*. Thus, Ω is a *k-cutting set* if $\tau_k(\Omega) = 0$.

Cutting Sets and Their Properties

We also refer to this property as being *k-cutting*. Thus, Ω is a *k-cutting set* if $\tau_k(\Omega) = 0$.

In particular, Ω is a cutting set or, more in detail, a cutting set with respect to the hyperplanes, when $\tau_{r-1}(\Omega) = 0$.

Cutting Sets and Their Properties

We also refer to this property as being *k-cutting*. Thus, Ω is a *k-cutting* set if $\tau_k(\Omega) = 0$.

In particular, Ω is a cutting set or, more in detail, a cutting set with respect to the hyperplanes, when $\tau_{r-1}(\Omega) = 0$.

Proposition 30 (A. Giuzzi, Longobardi, Siconolfi, (submitted))

Let Ω be a k -cutting set in $\text{PG}(r, q)$. Then Ω is also ℓ -cutting for all $k \leq \ell \leq r$.

Cutting Gap of Hermitian Varieties

Proposition 31 (A., Giuzzi, Longobardi, Siconolfi, (submitted))

Let $\mathcal{H}^r = \mathcal{H}(r, q^2)$ be a non-degenerate Hermitian variety in $\text{PG}(r, q^2)$. Then the cutting gap satisfies:

$$\tau_{r-t}(\mathcal{H}^r) = \begin{cases} 0 & \text{if } t > \lfloor \frac{r}{2} \rfloor, \\ 1 & \text{if } t \leq \lfloor \frac{r}{2} \rfloor. \end{cases}$$

Contents

- 1 Preliminaries
- 2 Constructions of Q -H varieties
- 3 Automorphism groups and equivalences
- 4 Applications
- 5 Open Problems

Future Research Directions

Future Research Directions

- **Characterization via cutting gaps:** Investigate whether the cutting gaps listed in the last previous proposition characterize Hermitian varieties among quasi-Hermitian ones.

Future Research Directions

- **Characterization via cutting gaps:** Investigate whether the cutting gaps listed in the last previous proposition characterize Hermitian varieties among quasi-Hermitian ones.
- **Automorphism groups:** Complete the determination of the full automorphism group of BT and BM quasi-Hermitian varieties in $\text{PG}(r, q^2)$ for all dimensions $r \geq 4$ and any q .

Future Research Directions

- **Characterization via cutting gaps:** Investigate whether the cutting gaps listed in the last previous proposition characterize Hermitian varieties among quasi-Hermitian ones.
- **Automorphism groups:** Complete the determination of the full automorphism group of BT and BM quasi-Hermitian varieties in $\text{PG}(r, q^2)$ for all dimensions $r \geq 4$ and any q .
- **Sylow 2-subgroups:** Explore whether the property that any Sylow 2-subgroup of the collineation group of a BT Q-H variety fixes a unique incident point-hyperplane pair in $\text{PG}(r, q^2)$ can be used to characterize the BT Q-H varieties.

Thank you
for your attention!

Grazie!

These works were partially supported by the European Union under the Italian National Recovery and Resilience Plan (NRRP) of NextGenerationEU, partnership on "Telecommunications of the Future" (PE00000001 - program "RESTART", CUP: D93C22000910001).