

Boundary Effects in Stochastic Cyclic Competition Models on a Two-Dimensional Lattice

M. Lazarus Arnau, Shannon Serrao, Uwe C. Täuber


Department of Physics, Center for Soft Matter and Biological Physics
Virginia Tech

March 7, 2019



Motivation

Three species cyclic competition schemes motivated by examples in biology, population dynamics, and chemistry.



(a) Rock Paper Scissors

(b) May-Leonard




Models

Rock-Paper-Scissors/Cyclic Lotka-Volterra
(RPS) Model




$$X \in \{A, B, C\}, \quad Y \in \{A, B, C, \emptyset\}$$

May-Leonard (ML) Model




Combined System

- Periodic boundary conditions (i.e. toroidal topology)
- Maximum occupancy of one particle per lattice site.
- May-Leonard except for thin (width = 64) annular patch governed by RPS
- Random initial conditions.
- 512×512 lattice sites.
- $\alpha = 1.0$, $\epsilon_m = 5.0$. ϵ_r varied between simulations



System topology and layout


Combined System




Correlation Lengths and Permeation Distance

- Vertical correlation function $C(x, t, r)$ measured parallel to interface.
- $C(x, t, r)$ averaged over multiple runs and used to calculate the vertical correlation length $L(x)$.
- Plane-wave size and permeation distance are approximately equal for all for all ϵ_r tested.


$$C_{AA}(x, t, r) = \langle n_a(x, y, t)n_a(x, y + r, t) \rangle - \langle n_a(x, t) \rangle^2$$
$$C(x, t, r) = \frac{C_{AA}(x, t, r) + C_{BB}(x, t, r) + C_{CC}(x, t, r)}{3}$$



Well Mixing Effects



Average net population densities near and far from the RPS-ML interface for various RPS mobility rates.



Net population density vs. time of pure ML lattice starting from random initial condition

Conclusions

- Changing the microscopic rules from ML to RPS on a subsection of the lattice produces plane-waves emanating away from the interface.
- Size and permeation distance of plane-waves independent of RPS mobility
- Drop in net population density near the interface due to the system being more “well-mixed”.
- Suggests periodic, localized mixing as a potential method for disruption of steady-state pattern formation in cyclic competition models.

Acknowledgements:

Research was sponsored by the Army Research Office and was accomplished under Grant Number W911NF-17-1-0156.

Additional travel support from Society of Physics Students Travel Award